

Alena Vanžurová

On metrizable locally homogeneous affine 2-dimensional manifolds

*Archivum Mathematicum*, Vol. 49 (2013), No. 5, 347--357

Persistent URL: <http://dml.cz/dmlcz/143558>

## Terms of use:

© Masaryk University, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## ON METRIZABILITY OF LOCALLY HOMOGENEOUS AFFINE 2-DIMENSIONAL MANIFOLDS

ALENA VANŽUROVÁ

ABSTRACT. In [19] we proved a theorem which shows how to find, under particular assumptions guaranteeing metrizable (among others, recurrency of the curvature is necessary), all (at least local) pseudo-Riemannian metrics compatible with a given torsion-less linear connection without flat points on a two-dimensional affine manifold. The result has the form of an implication only; if there are flat points, or if curvature is not recurrent, we have no good answer in general, which can be also demonstrated by examples. Note that in higher dimension, the problem is not easy to solve.

Here we try to apply this apparatus to the two main types (A and B from [9], [1]) of torsion-less locally homogeneous connections defined in open domains of 2-manifolds. We prove that in dimension two a symmetric linear connection with constant Christoffels is metrizable if and only if it is locally flat. On the other hand, in the class of connections of type B there are even non-flat metrizable connections.

### 1. INTRODUCTION

To answer the question, under what conditions on initial data, a given linear connection on an  $n$ -dimensional manifold coincides with the Levi-Civita connection of some pseudo-Riemannian metric, is a difficult problem in general. Only the case  $n = 2$  is easy.

For the Riemannian manifold, probably the most effective method was offered by O. Kowalski in [5], [6], see also [17], [16], [18]. In [19] we developed a method which might be helpful in deciding whether a given linear connection on a 2-dimensional manifold arises as a Riemannian connection of some pseudo-Riemannian metric. In the simplest case, for nowhere flat affine two-manifolds, we formulated necessary and sufficient conditions for local metrizable in [19], and in favourable case all compatible metrics were described in terms of the Ricci tensor. Our aim is to show here application for the class of locally homogeneous connections in affine 2-manifolds, [14], [9], [1].

---

2010 *Mathematics Subject Classification*: primary 53B05; secondary 53B20.

*Key words and phrases*: manifold, connection, metric.

The author was supported by the grant from Grant Agency of Czech Republic GAČR no. P201/11/0356 and by the project of specific university research of the Brno University of Technology, No. FAST-S-13-2888.

DOI: 10.5817/AM2013-5-347

We apply simple methods suggested in [19] in the case of the two main classes of locally homogeneous linear connections, called Type A and Type B in [9], [1], [14]. Such connections play the key role in the Classification Theorem for locally homogeneous symmetric linear connections on two-dimensional manifolds. Let us consider the case  $n = 2$  in what follows.

**1.1. Affine two-manifolds.** First recall some notation. Let  $(M, \nabla)$  be an affine manifold with a linear connection  $\nabla$ , and let  $R$  denote the corresponding curvature tensor,  $R(X, Y)Z = [\nabla_X, \nabla_Y]Z - \nabla_{[X, Y]}Z$  for  $X, Y, Z$  from  $\mathcal{X}(M)$ .

As well known, in dimension two the curvature tensor  $R$  of type (1, 3) can be completely recovered from the Ricci tensor,  $R(X, Y)Z = \text{Ric}(Y, Z)X - \text{Ric}(X, Z)Y$  for  $X, Y, Z \in \mathcal{X}(M)$ . In local coordinates, as far as  $R^i_{hjj} = 0$  and  $R^i_{hij} = R_{jh}$  for  $j \neq i$ , we have explicit formulas

$$(1) \quad \begin{aligned} R_{11} &= -R^2_{112} = R^2_{121}, & R_{21} &= -R^1_{121} = R^1_{112}, \\ R_{12} &= -R^2_{212} = R^2_{221}, & R_{22} &= -R^1_{221} = R^1_{212}. \end{aligned}$$

Since  $\nabla R = \omega \otimes R$  is equivalent with  $\nabla \text{Ric} = \omega \otimes \text{Ric}$  provided  $n = 2$  we get:

**Lemma 1.** *The curvature tensor  $R$  of  $(M_2, \nabla)$  is recurrent if and only if the Ricci tensor is recurrent, and  $R = 0$  if and only if  $\text{Ric} = 0$ .*

The induced map  $Z \mapsto R(X, Y)Z$ ,  $R(X, Y): T_x M \rightarrow T_x M$  (for  $X, Y, Z$  from  $T_x M$ ) is linear and skew-symmetric. The Ricci tensor  $\text{Ric}$  of type (0, 2) is a trace of the endomorphism,  $\text{Ric}(Y, Z) = \text{Tr}\{X \mapsto R(X, Y)Z\}$ ,  $X, Y, Z \in \mathcal{X}(M)$  (hence it carries less information than  $R$  in general) and the formula  $\text{Tr} R(Y, Z) = \text{Ric}(Z, Y) - \text{Ric}(Y, Z)$  holds, [11, p. 14].

**1.2. The Ricci tensor of a pseudo-Riemannian manifold.** For a pseudo-Riemannian manifold  $(M, g)$  consider the curvature tensor of type (0, 4) introduced (up to sign) by  $\tilde{R}(X, Y, Z, W) = g(R(X, Y)Z, W)$ ; the relations  $\tilde{R}(X, Y, Z, W) = \tilde{R}(Z, W, X, Y) = -\tilde{R}(Y, X, Z, W) = -\tilde{R}(X, Y, W, Z)$  hold. In local coordinates components  $R_{hijk}$  of  $\tilde{R}$  and  $R^{\ell}_{ijk}$  of  $R$  satisfy  $R_{hijk} = R_{jkhi} = -R_{ihjk} = -R_{hikj}$ ,  $R_{hijk} = g_{hs}R^s_{ijk}$ , and  $g^{\ell h}R_{hijk} = R^{\ell}_{ijk}$ . We introduce also the Ricci tensor of type (1, 1) with components  $R^i_j = g^i_s R_{sj}$ . The scalar curvature  $\varrho$  is its trace,  $\varrho = \text{Tr Ric} = R^s_s = g^{ij}R_{ij}$ .

If  $n = 2$  the formula  $R_{hijk} = K(g_{hj}g_{ik} - g_{hi}g_{jk})$  holds ([3, p. 137]) and  $R^i_{hjk} = \delta^i_j R_{kh} - \delta^i_k R_{jh}$  is satisfied ([10], [15]). We then check easily  $R^i_{hjk} = K(\delta^i_k g_{hj} - \delta^i_j g_{hk})$ , and further  $\text{Ric} = \frac{1}{2}\varrho g = Kg$  (the Ricci tensor of a 2-dimensional pseudo-Riemannian manifold is proportional to the metric tensor ([10, p. 263], [15, p. 101]) which shows that an affine 2-manifold is an Einstein space, and its Ricci tensor is necessarily *symmetric*).

Obviously for a nowhere flat  $(M_2, g)$ , the Ricci tensor is *non-degenerate*. Moreover, any nowhere flat pseudo-Riemannian 2-manifold has a *recurrent curvature tensor* provided the sectional curvature does not vanish ([4, I, p. 280]):

**Lemma 2.** *The Ricci tensor of a nowhere flat 2-dimensional pseudo-Riemannian manifold is recurrent, and the corresponding 1-form is exact (gradient).*

**Proof.**  $R \neq 0$  is equivalent with  $K(x) \neq 0$  on  $M$  (from continuity,  $K$  is either positive, or negative);  $g = \alpha(x) \cdot \text{Ric}$  with  $\alpha(x) = \frac{1}{K(x)} \neq 0$ , and  $\nabla g = 0$ . Obviously  $\alpha(x) \cdot \text{Ric}$  is parallel, and  $\nabla \text{Ric} = d(-\ln |\alpha|) \otimes \text{Ric}$  holds, [19].  $\square$

Since in the case  $R = 0$ , the tensor  $\text{Ric}$  is in fact also recurrent ( $\omega = 0$ ), *recurrency is one of necessary conditions for metrizability of a two-manifold.*

**1.3. Metrizable of affine 2-manifolds.** We obtained two necessary conditions for local metrizable of a symmetric linear connection on a pseudo-Riemannian 2-manifold: the Ricci tensor must be symmetric, and must be also recurrent for some closed 1-form (proof of Lemma 2). Moreover,  $\text{Ric}$  may be degenerate only if  $R = 0$  holds, and if this is the case then  $\text{Ric} = 0$ . For the sake of global metrizable, the corresponding 1-form must be even exact. As well known, a flat connection is always globally metrizable; we can even prescribe a signature.

Now suppose that the Ricci tensor (or equivalently, the curvature) is non-vanishing in one fixed point  $x_0 \in M$ , and due to continuity, in some neighborhood of  $x_0$  (note that the subset of non-flat points is open).

**Theorem 1.** *Let  $(M_2, \nabla)$  be a 2-manifold with a symmetric linear connection such that the Ricci tensor is symmetric, recurrent, i.e.  $\nabla \text{Ric} = \omega \otimes \text{Ric}$  for some 1-form  $\omega$ , and regular ( $|R_{ij}| \neq 0$ ). Then at least locally, there is a metric compatible with the connection.*

**Proof.** Let  $x_0 \in M$ . Since  $|R_{ij}| \neq 0$  there is a pair  $(i, j)$  such that  $R_{ij} \neq 0$  about  $x_0$ . Due to the recurrency and regularity,  $d\omega = 0$ . Hence in some neighborhood of  $x_0$ , there is a function  $f$  such that  $\omega = df$ . It can be checked that  $e^{-f} \cdot \text{Ric}$  is parallel about  $x_0$ . Therefore  $g = e^{-f} \cdot \text{Ric}$  is a local metric compatible with  $\nabla$  in a neighborhood of  $x_0$ .  $\square$

**Theorem 2** ([19]). *Let  $(M_2, \nabla)$  be a two-dimensional manifold with a symmetric linear connection and the curvature  $R$  non-zero everywhere. If the Ricci tensor of  $\nabla$  is regular, symmetric and satisfies  $\nabla \text{Ric} = \omega \otimes \text{Ric}$  where  $\omega = df$  is an exact 1-form ( $f$  being some function) then  $g = e^{-f} \cdot \text{Ric}$  is a global metric tensor such that  $\nabla g = 0$ .*

Hence a now-where flat symmetric linear connection on  $M_2$  is metrizable if and only if its Ricci tensor is symmetric, regular and recurrent with the corresponding 1-form being exact. If this is the case and  $\nabla \text{Ric} = df \otimes \text{Ric}$  holds for some smooth function  $f$  then all global metrics corresponding to the connection  $\nabla$  form a 1-parameter family of homothetic metrics

$$(2) \quad g_b = \exp(-f + b) \cdot \text{Ric} \quad , \quad b \in \mathbb{R} \quad ,$$

(i.e. they differ from each other up to a scalar multiple, and each arises from the Ricci tensor as a multiple by a smooth function).

## 2. LOCALLY HOMOGENEOUS AFFINE 2-MANIFOLDS

As well known, a connection  $\nabla$  in a domain  $U$  of  $M_2$  is given uniquely by a family of components (Christoffel symbols) which are functions  $A(u, v), \dots, F(u, v)$  in two variables  $u, v$  where  $(u, v)$  are local coordinates in  $U$ ,

$$(3) \quad \begin{aligned} \nabla_{\partial_u} \partial_u &= A \partial_u + B \partial_v, & \nabla_{\partial_u} \partial_v &= C \partial_u + D \partial_v, \\ \nabla_{\partial_v} \partial_u &= \tilde{C} \partial_u + \tilde{D} \partial_v, & \nabla_{\partial_v} \partial_v &= E \partial_u + F \partial_v; \end{aligned}$$

to simplify the usual notation we denoted  $\partial_u = \frac{\partial}{\partial u}$ , similarly for  $v$ ,  $\Gamma_{11}^1 = A$ ,  $\Gamma_{11}^2 = B$ ,  $\Gamma_{12}^1 = C$ ,  $\Gamma_{12}^2 = D$ ,  $\Gamma_{21}^1 = \tilde{C}$ ,  $\Gamma_{21}^2 = \tilde{D}$ ,  $\Gamma_{22}^1 = E$ ,  $\Gamma_{22}^2 = F$ . An affine manifold  $(M, \nabla)$  is called (affine) *locally homogeneous* if for any points  $p, q$  in  $M$  there are neighborhoods  $U \ni p, V \ni q$  and an affine transformation  $f: (U, \nabla|_U) \rightarrow (V, \nabla|_V)$  sending  $p$  into  $q$ , [13].

A seemingly easy problem, to classify all locally homogeneous torsion-less linear connections in plain domains, was solved only recently, namely by a direct method in [14], by a group-theoretical method in [9], and in a more general setting, for connections with arbitrary torsion, in [1]; the proofs are based on the theory of Lie algebras of vector fields, [12]. Recall the result of B. Opozda from [14] which was a motivation for our contribution.

**Theorem 3.** *Let  $M_2$  be an affine locally homogeneous 2-manifold with a symmetric connection. Then for any point  $p \in M$  there is a neighborhood  $U_p$  such that either the restriction of the given connection is the Riemannian connection of a pseudo-Riemannian manifold of constant curvature, or there is a system  $(u, v)$  of local coordinates in  $U_p$  such that the restriction is  $\nabla$  with component expression (3) where  $A, \dots, F$  are constants (called connection of the Type A, with constant Christoffels), or the restriction is one of the symmetric connections  $\hat{\nabla}$  given by the formulas*

$$(4) \quad \begin{aligned} \text{TYPE B:} \quad \hat{\nabla}_{\partial_u} \partial_u &= u^{-1} (A \partial_u + B \partial_v), \\ \hat{\nabla}_{\partial_u} \partial_v &= u^{-1} (C \partial_u + D \partial_v), \\ \hat{\nabla}_{\partial_v} \partial_v &= u^{-1} (E \partial_u + F \partial_v) \end{aligned}$$

(called connections of the Type B) where  $A, \dots, F$  are real constants, not all of them are zero.

Formulation of the Theorem 3 might evoke an incorrect impression that it somehow separates metric connections from connections of Type A and Type B. A more general and more precisely formulated version can be found in [1, Theorem 2 (Classification Theorem), pp. 2,3] for connections of arbitrary torsion. Since our aim is to examine metric connections let us consider now only torsion-free (symmetric,  $C = \tilde{C}$ ,  $D = \tilde{D}$ ) connections in what follows. Components of the Ricci tensor of a

symmetric connection read:

$$\begin{aligned}
 R_{11} &= \text{Ric}(\partial_u, \partial_v) = B(F - C) + D(A - D) + B_v - D_u, \\
 R_{12} &= \text{Ric}(\partial_u, \partial_v) = CD - BE + D_v - F_u, \\
 R_{21} &= \text{Ric}(\partial_v, \partial_u) = CD - BE + C_u - A_v, \\
 R_{22} &= \text{Ric}(\partial_v, \partial_v) = E(A - D) + C(F - C) + E_u - C_v
 \end{aligned}
 \tag{5}$$

where the lower index means partial derivative according to the corresponding variable; Ric need not be symmetric, even for a symmetric connection.

**2.1. Type A connections, with constant Christoffels.** Let us examine metrizability of symmetric connections with constant Christoffels in open domains of 2-manifolds. We prove that a complete answer can be given as follows: *In dimension two, a symmetric linear connection with constant Christoffels is metrizable if and only if it is locally flat.* Though the metrizability problem means to solve a system of differential equations in general [2], in our case it is sufficient to solve a system of algebraic equations.

In the case of a connection with constant Christoffels defined in  $U \subset \mathbb{R}^2[u, v]$ , the curvature tensor  $R$  as well as the Ricci tensor are constant, the components being

$$\begin{aligned}
 R_{11} &= B(F - C) + D(A - D), \\
 R_{12} &= R_{21} = CD - BE \\
 R_{22} &= E(A - D) + C(F - C).
 \end{aligned}
 \tag{6}$$

Hence Ric is always symmetric, and the shape of  $R_{12}$  suggests that vanishing or non-vanishing of one of components  $B, C, D, E$  might have consequences. In what follows we distinguish from technical reasons the cases  $D = 0$  and  $D \neq 0$  (e.g. to be able to control our considerations by computations in Maple, though  $D$  is not a preferable coefficient by nature, we could start in fact with something else, e.g. with  $C$  or  $E$ ).

**Lemma 3.** *A symmetric linear connection  $\nabla$  in  $M_2 = \mathbb{R}^2[u, v]$  with constant Christoffels is locally flat just in the following cases:*

- (a)  $D = B = 0$  and  $A, C, E, F \in \mathbb{R}$  are related by  $AE - C^2 + CF = 0$ ;
- (b)  $D \neq 0, C = BE/D,$  and  $A, B, E, F \in \mathbb{R}$  are related by  $D^2A - EB^2 + DBF - D^3 = 0$ .

**Proof.** Suppose Ric = 0 holds. Let us distinguish the cases  $D = 0, D \neq 0$ . Let  $D = 0$ . Then the condition  $R_{12} = 0$  implies  $BE = 0$ . If  $E = 0$  as well then we get  $B(F - C) = 0 = C(F - C)$  from (6) and in the next step either  $B = C = 0$ , we have the subcase  $B = C = D = E = 0, A, F \in \mathbb{R}$ , or  $F = C$ , we obtain the subcase  $D = E = 0, F = C, A, B, C \in \mathbb{R}$ ; the condition from (a) obviously holds. On the other hand if  $B = 0$  then the equation  $C(F - C) + AE = 0$  is satisfied, and we have the case (a) again.

Assume  $D \neq 0$ . We get  $C = BE/D$  from  $R_{12} = 0$ . The other two components of the Ricci tensor vanish if and only if the following holds:

$$B(F - BE/D) + D(A - D) = 0, \quad (BE/D)(F - BE/D) + E(A - D) = 0.$$

This system is equivalent to the unique third-order algebraic equation  $D^2A - EB^2 + DBF - D^3 = 0$ , hence (b) is checked.  $\square$

The class of linear connections with constant Christoffels contains a three-parameter family of metrizable locally flat connections. To show that besides locally flat connections, among symmetric connections with constant Christoffels on 2-manifolds there are no other metrizable connections we can use Theorem 1. So let us examine recurrency of Ric as a necessary condition for metrizability. Constant components of the covariant derivative  $\nabla \text{Ric}$  are

$$\begin{aligned} -R_{11;1} &= 2(AR_{11} + BR_{12}), & -R_{11;2} &= 2(CR_{11} + DR_{12}), \\ -R_{12;1} &= CR_{11} + (A + D)R_{12} + BR_{22}, & -R_{12;2} &= ER_{11} + (C + F)R_{12} + DR_{22}, \\ -R_{22;1} &= 2(CR_{12} + DR_{22}), & -R_{22;2} &= 2(ER_{12} + FR_{22}). \end{aligned}$$

Hence a symmetric connection with constant Christoffels on  $M_2$  is recurrent if and only if there exist constants  $\varphi_1, \varphi_2$  such that the following system holds:

$$(7) \quad \begin{aligned} (\varphi_1 + 2A)R_{11} + 2BR_{12} &= 0, & CR_{11} + (\varphi_1 + A + D)R_{12} + BR_{22} &= 0, \\ 2CR_{12} + (\varphi_1 + 2D)R_{22} &= 0, & (\varphi_2 + 2C)R_{11} + 2DR_{12} &= 0, \\ ER_{11} + (\varphi_2 + C + F)R_{12} + DR_{22} &= 0, & 2ER_{12} + (\varphi_2 + 2F)R_{22} &= 0. \end{aligned}$$

**Lemma 4.** *If the Ricci tensor of a symmetric linear connection on  $M_2$  is recurrent, has constant Christoffels and satisfies  $R_{12} = 0$  then either Ric vanishes or is degenerate.*

**Proof.** Under our assumptions, the system (7) guaranteeing recurrency reduces to the system of algebraic equations

$$(8) \quad \begin{aligned} (\varphi_1 + 2A)R_{11} &= 0, & (\varphi_2 + 2C)R_{11} &= 0, \\ (\varphi_1 + 2D)R_{22} &= 0, & (\varphi_2 + 2F)R_{22} &= 0. \end{aligned}$$

Suppose that  $R_{11} \cdot R_{22} \neq 0$ . Then necessarily  $A = D$  and  $C = F$ , and by (6)  $R_{11} = R_{22} = 0$ , a contradiction.  $\square$

**Lemma 5.** *If the Ricci tensor of a symmetric linear connection with constant Christoffels on  $M_2$  is recurrent, non-vanishing and non-degenerate then the components must satisfy  $R_{11} \cdot R_{22} \neq 0$ .*

**Proof.** If  $R_{11} = 0$  then by (7),  $BR_{12} = DR_{12} = 0$ , hence either  $R_{12} = 0$  or  $B = D = 0$  which gives  $R_{12} = 0$  again, hence the matrix is singular and Ric degenerates. Similarly for  $R_{22} = 0$ .  $\square$

**Corollary 1.** *If a symmetric connection with constant Christoffels on  $M_2$  and non-vanishing Ricci tensor should have Ric recurrent and non-degenerate then necessarily  $R_{11} \cdot R_{12} \cdot R_{22} \neq 0$ .*

**Theorem 4.** *If  $(M, \nabla)$  is an affine 2-manifold with constant Christoffels such that the Ricci tensor is recurrent and all its components are non-vanishing on  $M$  then Ric is degenerate.*

**Proof.** Since  $R_{11} \neq 0$  we get from (7)

$$\begin{aligned} \varphi_1 &= -2A - 2B \frac{R_{12}}{R_{11}}, & \varphi_2 &= -2C - 2D \frac{R_{12}}{R_{11}}, \\ -2AR_{22} - 2B \frac{R_{12}R_{22}}{R_{11}} + 2CR_{12} + 2DR_{22} &= 0, \\ -2CR_{22} - 2D \frac{R_{12}R_{22}}{R_{11}} + 2ER_{12} + 2FR_{22} &= 0. \end{aligned}$$

We obtain

$$B \frac{R_{12}R_{22}}{R_{11}} = CR_{12} + (D - A)R_{22}, \quad D \frac{R_{12}R_{22}}{R_{11}} = ER_{12} + (F - C)R_{22} = 0.$$

First suppose that  $BD \neq 0$  and show that Ric is degenerate. Indeed, we get

$$(-D(D - A) + B(F - C))R_{22} - (CD - BE)R_{12} = 0.$$

Let us plug for  $R_{ij}$  from the expressions:  $0 = R_{11} \cdot R_{22} - R_{12}^2 = \det(R_{ij})$ .

Now suppose  $B = 0$ . Then

$$R_{11} = D(D - A), \quad R_{12} = CD, \quad R_{22} = C(F - C) + E(A - D)$$

and necessarily  $D \neq 0, C \neq 0$  (otherwise  $R_{12} = 0$ , and the Ricci tensor would vanish by Lemma 4). From the first of the above equations,  $(2A + \varphi_1)R_{11} = 0$  and due to  $R_{11} \neq 0, \varphi_1 = -2A$ . Hence plugging into the second equation we get  $CD(D - A + C) = 0$  which is equivalent to  $C = A - D$ , due to  $CD \neq 0$ . This yields  $R_{11} = R_{22}$ . Now plugging into the third equation we have  $(A - D)(R_{12} - R_{22}) = 0$ . Since  $A - D \neq 0$  (otherwise  $CD = 0$ , a contradiction) we obtain  $R_{12} = R_{22} = R_{11}$ . It follows  $\det(R_{ij}) = 0$ .

Finally, if  $D = 0$  then

$$R_{11} = B(F - C), \quad R_{12} = -BE, \quad R_{22} = C(F - C) + AE$$

and necessarily  $BE \neq 0, F \neq C$ . We get  $\varphi_2 = -2C$ ,

$$\begin{aligned} (F - C)R_{22} + ER_{12} &= 0, & R_{22} &= -\frac{E}{F - C}R_{12}, \\ (F - C)R_{12} + ER_{11} &= 0, & R_{11} &= -\frac{F - C}{E}R_{12}. \end{aligned}$$

It follows  $R_{11} \cdot R_{22} = R_{12}^2$ , hence  $\det(R_{ij}) = 0$  again. □

**Corollary 2.** *In the above notation, exactly the following choices of constants give rise to a metrizable connection  $\nabla$  on  $M_2$ :*

$$\begin{aligned} B = C = D = E = 0, \quad A, F \in \mathbb{R}; \\ D = E = 0, \quad F = C, \quad A, B, C \in \mathbb{R}; \end{aligned}$$



$$B = D = 0, \quad A, C, E, F \in \mathbb{R} \text{ satisfying } AE - C^2 + CF = 0;$$

$$D \neq 0, C = BE/D, \quad A, B, E, F \in \mathbb{R} \text{ satisfying } D^2A - EB^2 + DBF - D^3 = 0.$$

**Theorem 5.** *Locally flat connections are exactly the only metrizable connections among all symmetric connections with constant Christoffels defined on open domains of affine 2-manifolds.*

Note that the above result corresponds to the results obtained in [1].

**2.2. Connections of the Type B.** Now let us pay attention to metrizable of a symmetric linear connection  $\hat{\nabla}$  of the form (4) defined in  $\mathbb{R}^2[u, v] \setminus \{(0, v); v \in \mathbb{R}\}$ . It appears that the situation is quite different: there exist classes of metrizable non-flat connections which are of Type B.

Let us try the direct method based on Theorem 1. Let  $\hat{\Gamma}_{ij}^k$  be components of the connection  $\hat{\nabla}$  (i.e.  $\hat{\Gamma}_{11}^1 = A/u = \hat{A}$ ,  $\hat{\Gamma}_{11}^2 = B/u = \hat{B}$  etc.) and denote by  $R_{ij}$  components of the Ricci tensor  $\hat{\text{Ric}}$  of  $\hat{\nabla}$ . We obtain

$$(9) \quad \begin{aligned} R_{11} &= u^{-2}[B(F - C) + D(1 + A - D)], \\ R_{12} &= u^{-2}[CD - BE + F], \\ R_{21} &= u^{-2}[CD - BE - C], \\ R_{22} &= u^{-2}[E(A - D - 1) + C(F - C)]. \end{aligned}$$

Components of  $\hat{\nabla}\hat{\text{Ric}}$  are

$$\begin{aligned} -R_{11;1} &= u^{-1}((2A + 2)R_{11} + B(R_{12} + R_{21})), \\ -R_{12;1} &= u^{-1}(CR_{11} + (A + D + 2)R_{12} + BR_{22}), \\ -R_{21;1} &= u^{-1}(CR_{11} + (A + D + 2)R_{21} + BR_{22}), \\ -R_{22;1} &= u^{-1}(C(R_{12} + R_{21}) + (2D + 2)R_{22}), \\ -R_{11;2} &= u^{-1}(2CR_{11} + 2DR_{12}), \\ -R_{12;2} &= u^{-1}(ER_{11} + (C + F)R_{12} + DR_{22}), \\ -R_{21;2} &= u^{-1}(ER_{11} + (C + F)R_{21} + DR_{22}), \\ -R_{22;2} &= u^{-1}(2ER_{12} + 2FR_{22}). \end{aligned}$$

In what follows suppose that the connection  $\hat{\nabla}$  has symmetric Ricci tensor, equivalently  $F = -C$ .

Provided symmetry of Ricci is guaranteed, necessary and sufficient conditions for recurrency of  $\hat{\nabla}$  read: the system of algebraic equations

$$\begin{aligned}
 & \left(\varphi_1 + \frac{2A+2}{u}\right)R_{11} + \frac{2B}{u}R_{12} = 0, \\
 & \frac{C}{u}R_{11} + \left(\varphi_1 + \frac{A+D+2}{u}\right)R_{12} + \frac{B}{u}R_{22} = 0, \\
 & \frac{2C}{u}R_{12} + \left(\varphi_1 + \frac{2D+2}{u}\right)R_{22} = 0, \\
 & \left(\varphi_2 + \frac{2C}{u}\right)R_{11} + \frac{2D}{u}R_{12} = 0, \\
 & \frac{E}{u}R_{11} + \varphi_2R_{12} + \frac{D}{u}R_{22} = 0, \\
 & \frac{2E}{u}R_{12} + \left(\varphi_2 - \frac{2C}{u}\right)R_{22} = 0
 \end{aligned}
 \tag{10}$$

must be solvable for functions  $\varphi_1(u, v)$ ,  $\varphi_2(u, v)$ .

**Lemma 6.** *Let the Ricci tensor  $\hat{\text{Ric}}$  of a symmetric connection  $\hat{\nabla}$  of Type B in  $M_2 = \mathbb{R}^2[u, v] \setminus \{(0, v); v \in \mathbb{R}\}$  be symmetric and recurrent. Let  $R_{12} = 0$  on  $M_2$  and  $R_{11} \cdot R_{22} \neq 0$  on  $M_2$ . Then  $\hat{\text{Ric}}$  is non-degenerate,  $B = C = F = 0$ ,  $A = D \neq 0$ ,  $E \neq 0$ ,  $R_{11} = u^{-2}A$ ,  $R_{22} = -u^{-2}E$ , and  $\hat{\nabla}\hat{\text{Ric}} = \omega \otimes \hat{\text{Ric}}$ ,  $\omega = df$  where  $\omega = -2u^{-1}(A + 1)du$  and  $f = -2(A + 1)\ln|u|^2 + b$ ,  $b \in \mathbb{R}$ . The connection  $\hat{\nabla}$  is metrizable.*

**Proof.** If  $R_{12} = 0$  then the system (10) is simplified:

$$\begin{aligned}
 & \left(\varphi_1 + \frac{2A+2}{u}\right)R_{11} + \frac{2B}{u}R_{12} = 0, \quad \frac{2C}{u}R_{12} + \left(\varphi_1 + \frac{2D+2}{u}\right)R_{22} = 0, \\
 & \left(\varphi_2 + \frac{2C}{u}\right)R_{11} + \frac{2D}{u}R_{12} = 0, \quad \frac{E}{u}R_{11} + \varphi_2R_{12} + \frac{D}{u}R_{22} = 0, \\
 & \frac{C}{u}R_{11} + \left(\varphi_1 + \frac{A+D+2}{u}\right)R_{12} + \frac{B}{u}R_{22} = 0, \quad \frac{2E}{u}R_{12} + \left(\varphi_2 - \frac{2C}{u}\right)R_{22} = 0.
 \end{aligned}$$

If  $R_{11} \cdot R_{22} \neq 0$  we obtain  $\varphi_1 = -2u^{-1}(A + 1) = -2u^{-1}(D + 1)$ , hence  $A = D$ , moreover,  $\varphi_2 = 0$ ,  $C = 0$ ,  $B = 0$ ,  $R_{11} : R_{22} = -D : E$ . The rest follows.  $\square$

From Theorem 1 we deduce

**Theorem 6.** *Any choice of real constants  $B = C = F = 0$ ,  $A = D \neq 0$ ,  $E \neq 0$  defines in (a neighborhood  $U$  of)  $M_2 = \mathbb{R}^2[u, v] \setminus \{(0, v); v \in \mathbb{R}\}$  a non-flat (locally) metrizable connection.*

The choice  $A = D = -1$ ,  $E = 1$ ,  $B = C = F = 0$  gives a non-flat metrizable connection  $\hat{\nabla}$ , with (symmetric, recurrent and non-degenerate) Ricci tensor  $\text{Ric} = -u^{-2}(du \otimes du + dv \otimes dv)$  ( $\omega = 0$ ), compatible with the Riemannian metric  $g = -\text{Ric} = u^{-2}(du \otimes du + dv \otimes dv)$  of constant negative curvature  $K = -1$ . Hence (locally)  $\hat{\nabla}$  is the Levi-Civita connection of the standard hyperbolic plane. On the

other hand, the constants  $A = D = E = 1$ ,  $B = C = F = 0$  define by (4) the Levi-Civita connection of the Lorentzian metric  $g = \text{Ric} = u^{-2}(du \otimes du - dv \otimes dv)$  of constant positive curvature  $K = 1$ . The result is in accordance with [1].

**Remark 1.** As concerns the 2-dimensional sphere, it is neither of Type A, nor of Type B, see [1].

#### REFERENCES

- [1] Arias-Marco, T., Kowalski, O., *Classification of locally homogeneous affine connections with arbitrary torsion on 2-dimensional manifolds*, Monatsh. Math. **153** (1) (2008), 1–18.
- [2] Eisenhart, L.P., Veblen, O., *The Riemann geometry and its generalization*, Proc. London Math. Soc. **8** (1922), 19–23.
- [3] Jost, J., *Riemannian Geometry and Geometric Analysis*, Springer, Berlin, Heidelberg, New York, 2005.
- [4] Kobayashi, S., Nomizu, K., *Foundations of Differential Geometry I, II*, Wiley–Intersc. Publ., New York, Chichester, Brisbane, Toronto, Singapore, 1991.
- [5] Kowalski, O., *On regular curvature structures*, Math. Z. **125** (1972), 129–138.
- [6] Kowalski, O., *Metrizability of affine connections on analytic manifolds*, Note Mat. **8** (1) (1988), 1–11.
- [7] Kowalski, O., Opozda, B., Vlášek, Z., *Curvature homogeneity of affine connections on two-dimensional manifolds*, Colloq. Math. **81** (1) (1999), 123–139.
- [8] Kowalski, O., Opozda, B., Vlášek, Z., *A classification of locally homogeneous affine connections with skew-symmetric Ricci tensor on 2-dimensional manifolds*, Monatsh. Math. **130** (2000), 109–125.
- [9] Kowalski, O., Opozda, B., Vlášek, Z., *A classification of locally homogeneous connections on 2-dimensional manifolds via group-theoretical approach*, CEJM **2** (1) (2004), 87–102.
- [10] Lovelock, D., Rund, H., *Tensors, Differential Forms, and Variational Principle*, A. Wiley Intersc. Publ., New York, London, Sydney, 1975.
- [11] Nomizu, K., Sasaki, T., *Affine Differential Geometry. Geometry of Affine Immersions*, Cambridge Univ. Press, 1994.
- [12] Olver, P. J., *Equivalence, Invariants and Symmetry*, Cambridge: Univ. Press, 1995.
- [13] Opozda, B., *On curvature homogeneous and locally homogeneous affine connections*, Proc. Amer. Math. Soc. **124** (6) (1996), 1889–1893.
- [14] Opozda, B., *A classification of locally homogeneous connections on 2-dimensional manifolds*, Differential Geom. Appl. **21** (2004), 173–198.
- [15] Petrov, A. Z., *Einstein Spaces*, Moscow, 1961, in Russian.
- [16] Vanžurová, A., *Metrization of linear connections, holonomy groups and holonomy algebras*, Acta Phys. Debrecina **42** (2008), 39–48.
- [17] Vanžurová, A., *Metrization problem for linear connections and holonomy algebras*, Arch. Math. (Brno) **44** (2008), 339–348.
- [18] Vanžurová, A., *Metrization of connections with regular curvature*, Arch. Math. (Brno) **45** (4) (2009), 325–333.
- [19] Vanžurová, A., Žáčková, P., *Metrizability of connections on two-manifolds*, Acta Univ. Palack. Olomuc. Fac. Rerum Natur. Math. **48** (2009), 157–170.
- [20] Vanžurová, A., Žáčková, P., *Metrization of linear connections*, Aplimat 2009: 8th International Conference Proceedings **2** (2009), 151–163.

FACULTY OF SCIENCE, DEPARTMENT OF ALGEBRA AND GEOMETRY,  
PALACKÝ UNIVERSITY,  
TR. 17. LISTOPADU 12, 771 46 OLOMOUC, CZECH REPUBLIC  
*E-mail:* [alena.vanzurova@upol.cz](mailto:alena.vanzurova@upol.cz)

DEPARTMENT OF MATHEMATICS, FACULTY OF CIVIL ENGINEERING,  
BRNO UNIVERSITY OF TECHNOLOGY,  
VEVEŘÍ 331/95, 602 00 BRNO, CZECH REPUBLIC  
*E-mail:* [vanzurova.a@fce.vutbr.cz](mailto:vanzurova.a@fce.vutbr.cz)