

Vsevolod Yu. Gubarev; Pavel S. Kolesnikov
Operads of decorated trees and their duals

Commentationes Mathematicae Universitatis Carolinae, Vol. 55 (2014), No. 4, 421--445

Persistent URL: <http://dml.cz/dmlcz/143967>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2014

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Operads of decorated trees and their duals

VSEVOLOD YU. GUBAREV, PAVEL S. KOLESNIKOV

Abstract. This is an extended version of a talk presented by the second author on the Third Mile High Conference on Nonassociative Mathematics (August 2013, Denver, CO). The purpose of this paper is twofold. First, we would like to review the technique developed in a series of papers for various classes of di-algebras and show how the same ideas work for tri-algebras. Second, we present a general approach to the definition of pre- and post-algebras which turns out to be equivalent to the construction of dendriform splitting. However, our approach is more algebraic and thus provides simpler way to prove various properties of pre- and post-algebras in general.

Keywords: Leibniz algebra; dialgebra; dendriform algebra; pre-Lie algebra

Classification: 17A30, 17A36, 17A42, 18D50

1. Introduction

The study of a wide variety of algebraic systems that may be informally called di-algebras was initiated by J.-L. Loday and T. Pirashvili [22], who proposed the notion of an (associative) di-algebra as a tool in the cohomology theory of Lie and Leibniz algebras. A systematic study of associative di-algebras and their Koszul dual dendriform algebras was presented in [21]. Later, an algebraic approach to operads appearing in combinatorics led J.-L. Loday and M. Ronco [23] to the notions of tri-associative and tri-dendriform algebras.

In [9], F. Chapoton pointed out that the operads governing the varieties of Leibniz algebras and of di-algebras in the sense of [22] may be presented as Manin white products of the operad Perm with Lie and As, respectively. Manin products (white product \circ and black product \bullet) were originally defined for quadratic associative algebras and then for binary quadratic operads. In [29], it was proposed a conceptual approach to Manin products and Koszul duality which covers a wide range of monoids in categories with two coherent monoidal products (quadratic associative algebras and binary operads, in particular, fit this scheme). The operad Perm has an extremely simple algebraic nature, so it is obvious that the white product Perm \circ \mathfrak{M} coincides with the Hadamard product Perm \otimes \mathfrak{M} for every binary quadratic operad \mathfrak{M} (see [29]). In this way, a general definition of a di-algebra over \mathfrak{M} as an algebra governed by Perm \otimes \mathfrak{M} was considered in [18], where it was shown that di-algebras are closely related with pseudo-algebras in the sense of [3]. This relation allowed solving many algebraic problems on di-algebras [14], [19], [31]. Hence, it is interesting to find an analogous construction for tri-algebras as well. It was also shown in [29] that the operad ComTrias (introduced in [28]) has the same property

as $\text{Perm} : \text{ComTrias} \circ \mathfrak{M} = \text{ComTrias} \otimes \mathfrak{M}$. In this paper, we show how to recover an “ordinary” algebra from a given $(\text{ComTrias} \otimes \mathfrak{M})$ -algebra and apply the result to solve a series of problems on tri-algebras.

Roughly speaking, a passage from an operad \mathfrak{M} governing a variety of “ordinary” algebras (associative, Lie, Jordan, Poisson, etc.) to the operad $\text{di-}\mathfrak{M}$ or $\text{tri-}\mathfrak{M}$ may be performed by “decoration” of planar trees presenting the operad \mathfrak{M} . (For di-algebras, the procedure was proposed in [18], for tri-algebras — in [15] in the case of binary operations.) In this sense, to decorate a tree one has to emphasize one (for di-algebras) or several (for tri-algebras) leaves and assume the composition (grafting) of trees to preserve the decoration (see Section 2 for details).

A similar unified approach to the definition of dendriform algebras comes naturally from the general concept of Manin black product [29]. The class of associative dendriform di-algebras Dend [21] is known to be governed by the operad $\text{pre-Lie} \bullet \text{As}$, where pre-Lie is the operad of left-symmetric algebras. Obviously (see [13]), $\text{Dend} = (\text{Perm} \circ \text{As})^!$ since $\text{Perm}^! = \text{pre-Lie}$, $\text{As}^! = \text{As}$. The same duality between white and black Manin product holds in the general settings [29]. So, the natural way to define a dendriform version of an \mathfrak{M} -algebra is to consider the operad $\text{pre-Lie} \bullet \mathfrak{M}$ or $\text{post-Lie} \bullet \mathfrak{M}$ (the operad $\text{post-Lie} = \text{ComTrias}^!$ was introduced in [28]). The explicit description of the corresponding varieties of such systems in terms of defining identities was proposed in [2] (as di-successor and tri-successor algebras) and in [15] as (di- and tri-dendriform algebras). A generalization of the first construction has recently been published in [25]: $\mathcal{B}\text{-}(\mathcal{A}\text{-})\text{Sp}(\mathfrak{M})$ -algebras are defined for an arbitrary operad \mathfrak{M} . In this paper, we state another simple procedure of “dendriform splitting” and prove that the classes of systems obtained (called pre- or post-algebras, respectively) coincide with those already introduced in [2], [15], [25].

The paper is organized as follows. In Section 2 we recall the general definition of what is a di- or tri-algebra and explain its relation with averaging operators. Section 3 is devoted to the definition of operads $\text{pre-}\mathfrak{M}$ and $\text{post-}\mathfrak{M}$ for an arbitrary (not necessarily binary or quadratic) operad \mathfrak{M} . The varieties of $\text{pre-}\mathfrak{M}$ - and $\text{post-}\mathfrak{M}$ -algebras obtained are closely related with \mathfrak{M} -algebras equipped with Rota–Baxter operators in the very same way as $(\mathcal{A}\text{-})\text{Sp}(\mathfrak{M})$ - and $\mathcal{B}\text{Sp}(\mathfrak{M})$ -algebras in [25], thus, our approach leads to the same classes of systems. In Section 4 we observe a series of algebraic problems related with di- and tri-algebras. Most of natural problems in this area may be easily reduced to similar problems in “ordinary” algebras by means of the embedding proved in Theorem 2.8. Section 5 is devoted to analogous problems on pre- and post-algebras. In these classes, the picture is obscure: It is possible to state that many classical algebraic problems (like those stated in Section 4) make sense for pre- and post-algebras, but it is not clear how to solve them.

Throughout the paper we will use the following notations: $\mathcal{P}(n)$ is the set of all nonempty subsets of $\{1, \dots, n\}$; S_n is the group of all permutations of $\{1, \dots, n\}$. An operad \mathfrak{M} is a collection of S_n -modules $\mathfrak{M}(n)$, $n \geq 1$, equipped with associative and equivariant composition rule (see, e.g., [24]).

Given a language Σ (a set of symbols of algebraic operations f together with their arities $\nu(f)$), by a Σ -algebra we mean a linear space equipped with algebraic operations


from Σ . The class of all Σ -algebras as well as the corresponding (free) operad we denote by \mathfrak{F}_Σ . If \mathfrak{M} is a quotient operad of \mathfrak{F}_Σ and a Σ -algebra A belongs to the variety governed by \mathfrak{M} then we say A to be an \mathfrak{M} -algebra. For an operad \mathfrak{M} , we will use the same symbol \mathfrak{M} to denote the entire variety governed by \mathfrak{M} .

The free algebra in the variety of all \mathfrak{M} -algebras generated by a set X we denote by $\mathfrak{M}\langle X \rangle$.

2. Replicated algebras


2.1 Replication of a free operad. In this section we present an explanation of the idea underlying the transition from “ordinary” algebras to di- and tri-algebras and discuss why these constructions are the only possible ones in a certain context.

Let us consider the free operad $\mathfrak{F} = \mathfrak{F}_\Sigma$ generated by operations Σ . According to the natural graphical interpretation, the spaces $\mathfrak{F}(n)$, $n \geq 1$, are spanned by planar trees with enumerated leaves (variables) and labeled vertices (operations). For example, if $\Sigma = \{(\cdot * \cdot), [\cdot, \cdot]\}$ consists of two binary operations then the term $[x_1, (x_4 * x_3)] * [x_2, x_5]$ may be identified with



The general idea of replication (cf. [18]) is to set an additional feature on the trees from $\mathfrak{F}(n)$: Emphasize one or several leaves and claim that the emphasizing is preserved by composition (grafting). Let us explain the details graphically and then present an equivalent algebraic statement.


Recall the composition rule on the operad \mathfrak{F} : Given $T \in \mathfrak{F}(n)$, $T_i \in \mathfrak{F}(m_i)$, $i = 1, \dots, n$, their composition $T(T_1, \dots, T_n) \in \mathfrak{F}(m_1 + \dots + m_n)$ is a tree obtained by attaching each T_i to the i th leaf of T and by natural shift of numeration of leaves in each T_i . For example, if $\Sigma = \{(\cdot * \cdot), [\cdot, \cdot]\}$ consists of two binary operations, $T = [x_2, x_3] * x_1$, $T_1 = [x_2, [x_1, x_3]]$, $T_2 = [x_2, x_1]$, $T_3 = x_1 * x_2$, then $T(T_1, T_2, T_3)$ is presented by



Symmetric group S_n acts on $\mathfrak{F}(n)$ by permutations of leaves' numbers.

By definition, every tree in $\mathfrak{F}(n)$ may be constructed by composition and symmetric group actions from the elementary trees (generators of the operad) $f(x_1, \dots, x_n)$, $f \in \Sigma$, $\nu(f) = n$.

Now, replace the generators with “decorated” elementary trees with one or several emphasized leaves and define the composition of such trees by the same rule as in \mathfrak{F} , assuming that: (1) attaching of a tree T_i to a non-emphasized leaf of T removes decoration from T_i ; (2) attaching of a tree T_i to an emphasized leaf of T preserves decoration on T_i . An example of such a composition with emphasized leaves circled in black is stated below.



Note that if each of the trees T, T_1, \dots, T_n has only one emphasized leaf then so is their composition $T(T_1, \dots, T_n)$. However, if we are allowed to emphasize more than one leaf (say, no more than two leaves of each tree, as in example above) then the composition may contain more emphasized leaves than each of the trees T, T_1, \dots, T_n (see the example above). Hence, there are two natural cases: Either we may emphasize only one leaf of a tree (di-algebra case) or an arbitrary number of leaves (tri-algebra case). Let us denote the operads obtained by di- \mathfrak{F} or tri- \mathfrak{F} , respectively.

2.2 Operads Perm and ComTrias. Let us state definitions of two important operads.

Example 2.1 ([9]). Let Σ contains one binary operation. The operad governing the variety of associative algebras satisfying the identity $(x_1 x_2) x_3 = (x_2 x_1) x_3$ is denoted by Perm. It is easy to see that monomials $e_i^{(n)} = (x_1 \dots x_{i-1} x_{i+1} \dots x_n) x_i, i = 1, \dots, n$, form a linear basis of Perm(n), and thus $\dim \text{Perm}(n) = n$.

Example 2.2 ([28]). Given $n \geq 1$, let $C(n)$ be the formal linear span of the set of “corollas” $\{e_H^{(n)} \mid H \in \mathcal{P}(n)\}$, where $\mathcal{P}(n)$ stands for the collection of all nonempty subsets of $\{1, \dots, n\}$. For $K \in \mathcal{P}(m)$, $H_i \in \mathcal{P}(n_i)$, $i = 1, \dots, m$, define the composition of sets $K(H_1, \dots, H_m) \in \mathcal{P}(n_1 + \dots + n_m)$ as follows:

$$\begin{aligned}
 j \in K(H_1, \dots, H_m) &\iff \exists k \in K, l \in H_k : \\
 &n_1 + \dots + n_{k-1} < j \leq n_1 + \dots + n_k, \\
 &j = n_1 + \dots + n_{k-1} + l.
 \end{aligned}$$

Then

$$e_K^{(m)}(e_{H_1}^{(n_1)}, \dots, e_{H_m}^{(n_m)}) = e_{K(H_1, \dots, H_m)}^{(n)},$$

where $n = n_1 + \dots + n_m$.

With respect to the natural action of the symmetric group, the family of spaces $C(n)$, $n \geq 1$, forms a symmetric operad denoted by ComTrias .

The algebraic interpretation of ComTrias was stated in [28]. Namely, an algebra from the variety ComTrias is a linear space equipped with two binary operations \perp and \vdash satisfying the following axioms:

$$\begin{aligned}
 (x \vdash y) \vdash z &= x \vdash (y \vdash z), & (x \vdash y) \vdash z &= (y \vdash x) \vdash z, \\
 (x \perp y) \vdash z &= (x \vdash y) \vdash z, & x \vdash (y \perp z) &= (x \vdash y) \perp z, \\
 (x \perp y) \perp z &= x \perp (y \perp z).
 \end{aligned}$$

It is easy to see that $e_H^{(n)} \in \text{ComTrias}(n)$ may be identified with the monomial

$$x_{j_1} \vdash \dots \vdash x_{j_{n-k}} \vdash (x_{i_1} \perp \dots \perp x_{i_k}),$$

where $H = \{i_1, \dots, i_k\}$, $i_1 < \dots < i_k$, $j_1 < \dots < j_{n-k}$.

Example 2.3. Denote by C_2 a 2-dimensional space with a basis $\{e_1, e_2\}$ and operations

$$e_i \perp e_i = e_i, \quad e_1 \vdash e_1 = e_1, \quad e_1 \vdash e_2 = e_2,$$

other products are zero. It is easy to check that $C_2 \in \text{ComTrias}$.

Note that the composition rule in the operad Perm is completely similar to the composition in ComTrias restricted to singletons: $e_i^{(n)} \in \text{Perm}(n)$ may be identified with $e_{\{i\}}^{(n)} \in \text{ComTrias}(n)$.

Lemma 2.4. Let $m \geq 1$, $n_1, \dots, n_m \geq 1$, and let $n = n_1 + \dots + n_m$. Then

$$\begin{aligned}
 \sum_{H \in \mathcal{P}(n)} \sum_{\substack{K, H_1, \dots, H_m \\ K(H_1, \dots, H_m) = H \\ K \in \mathcal{P}(m) \\ H_i \in \mathcal{P}(n_i)}} e_K^{(m)} \otimes e_{H_1}^{(n_1)} \otimes \dots \otimes e_{H_m}^{(n_m)} \\
 = \sum_{K \in \mathcal{P}(m)} \sum_{H_1 \in \mathcal{P}(n_1)} \dots \sum_{H_m \in \mathcal{P}(n_m)} e_K^{(m)} \otimes e_{H_1}^{(n_1)} \otimes \dots \otimes e_{H_m}^{(n_m)}.
 \end{aligned}$$

A similar statement holds for Perm, if we restrict the sums to singletons only.

PROOF: For $m = 1$ the statement is obvious. It is enough to note that

$$K(H_1, \dots, H_m) = \begin{cases} (K \setminus \{m\})(H_1, \dots, H_{m-1}) \cup (n - n_m + H_m), & m \in K, \\ K(H_1, \dots, H_{m-1}), & m \notin K, \end{cases}$$

and proceed by induction on m . □

2.3 Defining identities. Let \mathfrak{M} be a variety of Σ -algebras satisfying a family of polynomial identities $\text{Id}(\mathfrak{M})$. Denote the operad governing this variety by the same symbol \mathfrak{M} . This is an image of the free operad $\mathfrak{F} = \mathfrak{F}_\Sigma$ with respect to a morphism of operads whose kernel equals $\text{Id}(\mathfrak{M})$.

Definition 2.5 ([18], [15]). Denote by $\text{di-}\mathfrak{M}$ and $\text{tri-}\mathfrak{M}$ the following Hadamard products of operads:

$$\text{di-}\mathfrak{M} = \text{Perm} \otimes \mathfrak{M}, \quad \text{tri-}\mathfrak{M} = \text{ComTrias} \otimes \mathfrak{M}.$$

As an immediate corollary of this definition, we obtain

Proposition 2.6 ([18], [19]). *Let $A \in \mathfrak{M}$, $P \in \text{Perm}$. Then $P \otimes A$ equipped with operations*

$$\begin{aligned} f_i(x_1 \otimes a_1, \dots, x_n \otimes a_n) &= e_i^{(n)}(x_1, \dots, x_n) \otimes f(a_1, \dots, a_n), \\ f \in \Sigma, \nu(f) = n, x_i \in P, a_i \in A, i = 1, \dots, n, \end{aligned}$$

belongs to the variety $\text{di-}\mathfrak{M}$.

Proposition 2.7. *Let $A \in \mathfrak{M}$, $C \in \text{ComTrias}$. Then $C \otimes A$ equipped with operations*

$$\begin{aligned} f^H(x_1 \otimes a_1, \dots, x_n \otimes a_n) &= e_H^{(n)}(x_1, \dots, x_n) \otimes f(a_1, \dots, a_n), \\ f \in \Sigma, \nu(f) = n, H \in \mathcal{P}(n), x_i \in C, a_i \in A, i = 1, \dots, n, \end{aligned}$$

belongs to the variety $\text{tri-}\mathfrak{M}$.

In general, it is not clear which operations generate a Hadamard product of two operads (even if the operads are binary). However, operads $\mathfrak{P} = \text{Perm}, \text{ComTrias}$ are good enough to allow finding generators and defining relations of $\mathfrak{P} \otimes \mathfrak{M}$. In particular, if \mathfrak{M} is a binary quadratic operad then $\mathfrak{P} \otimes \mathfrak{M} = \mathfrak{P} \circ \mathfrak{M}$, where \circ stands for the Manin white product of operads. The purpose of this section is to present explicitly defining relations of $\text{tri-}\mathfrak{M}$ (for $\text{di-}\mathfrak{M}$, the algorithm was presented in [5], see also [19]).

First, let us note that the operad $\text{tri-}\mathfrak{F}$ is generated by

$$\Sigma^{(3)} = \{f^H \mid f \in \Sigma, \nu(f) = n, H \in \mathcal{P}(n)\}.$$

Indeed, there exists a morphism of operads $\iota : \mathfrak{F}_{\Sigma^{(3)}} \rightarrow \text{tri-}\mathfrak{F}$ sending f^H to $e_H^{(n)} \otimes f$, $f \in \Sigma$, $\nu(f) = n$. Therefore, every $D \in \text{tri-}\mathfrak{M}$ may be considered as a $\Sigma^{(3)}$ -algebra. Note that for

every $f, g \in \Sigma, \nu(f) = n, \nu(g) = m$, and for every $a_k, b_j \in D$ we have

$$(1) \quad f^H(a_1, \dots, a_{i-1}, g^S(b_1, \dots, b_m), a_{i+1}, \dots, a_n) \\ = f^H(a_1, \dots, a_{i-1}, g^Q(b_1, \dots, b_m), a_{i+1}, \dots, a_n)$$

for all $H \in \mathcal{P}(n), S, Q \in \mathcal{P}(m)$ provided that $i \notin H$. Indeed, by the definition of ComTrias, the composition

$$e_H^{(n)}(\text{id}, \dots, e_S^{(m)}, \dots, \text{id})$$

does not depend on S if $i \notin H$.

Moreover, each $\iota(m) : \mathfrak{F}_{\Sigma^{(3)}}(m) \rightarrow \text{tri-}\mathfrak{F}(m), m \geq 1$, is surjective. The natural algorithm of constructing a canonical pre-image $\Phi^H \in \mathfrak{F}_{\Sigma^{(3)}}(m)$ of $e_H^{(m)} \otimes \Phi \in \text{tri-}\mathfrak{F}(m)$ with respect to $\iota(m)$ is stated in [15] for binary case. In the general case, the algorithm remains the same: Assume the pre-images are constructed for all terms of degree smaller than m . For a monomial $u = u(x_1, \dots, x_m) \in \mathfrak{F}(m)$, one may consider $e_H^{(m)} \otimes u \in \text{tri-}\mathfrak{F}(m)$ as a planar tree with emphasized leaves x_{i_1}, \dots, x_{i_k} , where $\{i_1, \dots, i_k\} = H$. If $u = f(\nu_1, \dots, \nu_n), f \in \Sigma, \nu(f) = n, \nu_i \in \mathfrak{F}(m_i)$, then choose $K = \{i \mid \nu_i \text{ contains } x_j, j \in H\}$ and set $u^H = f^K(\nu_1^{H_1}, \dots, \nu_n^{H_n})$, where

$$H_i = \begin{cases} \{j \mid j \in H, x_j \text{ appears in } \nu_i\}, & i \in K, \\ \{1\}, & i \notin K. \end{cases}$$

Next, suppose $\Phi(x_1, \dots, x_m) \in \mathfrak{F}(m)$ is a polylinear identity on all algebras of a variety \mathfrak{M} , i.e., Φ belongs to the kernel of natural morphisms of operads $\tau_{\mathfrak{M}} : \mathfrak{F} \rightarrow \mathfrak{M}$. Then $e_H^{(m)} \otimes \Phi$ belongs to the kernel of $\text{id} \otimes \tau_{\mathfrak{M}} : \text{tri-}\mathfrak{F} \rightarrow \text{tri-}\mathfrak{M}$. Hence, $\Phi^H(x_1, \dots, x_m) \in \mathfrak{F}_{\Sigma^{(3)}}(m)$ is an identity on all algebras in $\text{tri-}\mathfrak{M}$.

Suppose the variety \mathfrak{M} is defined by a set of polylinear identities $S(\mathfrak{M}) \subset \text{Id}(\mathfrak{M})$. As we have shown above, every algebra in $\text{tri-}\mathfrak{M}$ may be considered as a $\Sigma^{(3)}$ -algebra satisfying the collection of identities $S^{(3)}(\mathfrak{M})$ that consists of (1) and $\Phi^H(a_1, \dots, a_m) = 0$ for all $\Phi \in S(\mathfrak{M}) \cap \mathfrak{F}(m), H \in \mathcal{P}(m), m \geq 1$.

Let us prove that $S(\text{tri-}\mathfrak{M}) = S^{(3)}(\mathfrak{M})$, i.e., every $\Sigma^{(3)}$ -algebra satisfying $S^{(3)}(\mathfrak{M})$ is actually an algebra of the variety governed by $\text{tri-}\mathfrak{M}$.

Theorem 2.8. *Suppose $\nu(f) \geq 2$ for all $f \in \Sigma$. Then every $\Sigma^{(3)}$ -algebra satisfying $S^{(3)}(\mathfrak{M})$ may be embedded into an appropriate algebra of the form $C \otimes A \in \text{tri-}\mathfrak{M}$, where $C \in \text{ComTrias}, A \in \mathfrak{M}$.*

An analogous statement for $\text{di-}\mathfrak{M}$ was proved in [19].

PROOF: Given an algebra $T \in \text{tri-}\mathfrak{M}$, denote by $T_0 \subseteq T$ the linear span of all

$$(e_H^{(n)} \otimes f)(a_1, \dots, a_n) - (e_K^{(n)} \otimes f)(a_1, \dots, a_n),$$

$K, H \in \mathcal{P}(n), a_i \in T, f \in \Sigma, \nu(f) = n$. It follows from the definition of ComTrias that T_0 is an ideal in T , and $\bar{T} = T/T_0$ may be considered as a Σ -algebra. Moreover, the direct

sum of linear spaces

$$\widetilde{T} = \bar{T} \oplus T$$

turns into a Σ -algebra with respect to operations

$$(2) \quad f(\bar{a}_1 + b_1, \dots, \bar{a}_n + b_n) = \overline{f^K(a_1, \dots, a_n)} + \sum_{H \in \mathcal{P}(n)} f^H(c_1^H, \dots, c_n^H),$$

(K is an arbitrary set in $\mathcal{P}(n)$) $f \in \Sigma$, $\nu(f) = n$, $a_i, b_i \in T$, and

$$c_i^H = \begin{cases} a_i, & i \notin H, \\ b_i, & i \in H. \end{cases}$$

Lemma 2.9. $\widetilde{T} \in \mathfrak{M}$.

PROOF: In [15], this statement was proved in the binary case. The general case is similar.

Suppose $\Phi(x_1, \dots, x_m) \in S(\mathfrak{M})$. Then (2) and (1) imply $\Phi(\bar{a}_1 + b_1, \dots, \bar{a}_m + b_m) = 0$ for $a_i, b_j \in T$ by induction on the length of monomials. \square

Recall the algebra $C_2 \in \text{ComTrias}$ from Example 2.3. Note that the map $T \rightarrow C_2 \otimes \widetilde{T}$, given by

$$a \mapsto e_1 \otimes \bar{a} + e_2 \otimes a \in C_2 \otimes \widetilde{T}, \quad a \in T,$$

is a homomorphism of $\Sigma^{(3)}$ -algebras. Indeed, let $f \in \Sigma$, $\nu(f) = n$, $H \in \mathcal{P}(n)$, $x_i = e_1 \otimes \bar{a}_i + e_2 \otimes a_i$, $a_i \in T$, $i = 1, \dots, n$. Then

$$\begin{aligned} (e_H^{(n)} \otimes f)(x_1, \dots, x_n) &= e_H^{(n)}(e_1, \dots, e_1) \otimes f(\bar{a}_1, \dots, \bar{a}_n) \\ &= \sum_{K \in \mathcal{P}(n)} e_H^{(n)}(e_1^K, \dots, e_n^K) \otimes f(c_1^K, \dots, c_n^K), \end{aligned}$$

where

$$e_k^K = \begin{cases} e_1, & k \notin K, \\ e_2, & k \in K, \end{cases} \quad c_i^K = \begin{cases} \bar{a}_i, & i \notin K, \\ a_i, & i \in K. \end{cases}$$

It is easy to note from the definition of C_2 that $e_H^{(n)}(e_1^K, \dots, e_n^K) \neq 0$ if and only if $K = H$ (in this case, the result is equal to e_2). Hence,

$$\begin{aligned} (e_H^{(n)} \otimes f)(x_1, \dots, x_n) &= e_1 \otimes f(\bar{a}_1, \dots, \bar{a}_n) + e_2 \otimes \overline{f^H(a_1, \dots, a_n)} \\ &= e_1 \otimes f^H(a_1, \dots, a_n) + e_2 \otimes f^H(a_1, \dots, a_n). \end{aligned}$$

\square

Remark 1. Note that Theorem 2.8 remains valid for languages with unary operators $t \in \Sigma$, $\nu(t) = 1$, provided that $S(\mathfrak{M})$ includes identities stating all these t are endomorphisms or derivations with respect to all $f \in \Sigma$, $\nu(f) > 1$. In this case, T_0 is invariant with respect to t , and thus \widetilde{T} exists.

Therefore, if T satisfies $S^{(3)}(\mathfrak{M})$ then it is a subalgebra in $C_2 \otimes \widetilde{T} \in \text{tri-}\mathfrak{M}$, so $T \in \text{tri-}\mathfrak{M}$.

As it was shown in [19], the variety governed by $\text{di-}\mathfrak{M} = \text{Perm} \otimes \mathfrak{M}$ may be represented as a variety of $\Sigma^{(2)}$ -algebras defined by $S^{(2)}(\mathfrak{M})$, where $\Sigma^{(2)}$ and $S^{(2)}(\mathfrak{M})$ are obtained from Σ and $S(\mathfrak{M})$ in the same way as $\Sigma^{(3)}$ and $S^{(3)}(\mathfrak{M})$ provided that we consider only singletons $H = \{i\} \in \mathcal{P}(n)$.

Examples include Leibniz algebras (di-Lie) [20], dialgebras (di-As) [22], semi-special quasi-Jordan algebras (di-Jord) [18], [30], [4], Lie and Jordan triple di-systems (di-LTS [8] and di-JTS [5]), Malcev di-algebras (di-Mal) [7], dual pre-Poisson algebras (di-Pois) [1], triassociative algebras (tri-As) [23].

Example 2.10. Let us write down defining identities of tri-Lie-algebras. An algebra from tri-Lie is a linear space with three binary operations $[\cdot \perp \cdot]$, $[\cdot \vdash \cdot]$, and $[\cdot \dashv \cdot]$, $[a \vdash b] = -[b \dashv a]$, such that $[\cdot \perp \cdot]$ is a Lie operation, $[\cdot \dashv \cdot]$ satisfies (right) Leibniz identity, and they satisfy the following axioms:

$$(3) \quad \begin{aligned} [x_1 \perp [x_2 \dashv x_3]] &= [[x_1 \dashv x_2] \perp x_3] + [x_2 \perp [x_1 \dashv x_3]], \\ [x_1 \dashv [x_2 \perp x_3]] &= [x_1 \dashv [x_2 \dashv x_3]]. \end{aligned}$$

Let us note that the first identity of (3) appeared recently in [27].

Lemma 2.11. *If $\varphi : T \rightarrow T'$ is a homomorphism of tri- \mathfrak{M} -algebras then $\tilde{\varphi} : \widetilde{T} \rightarrow \widetilde{T}'$ defined by $\tilde{\varphi}(\bar{a}) = \overline{\varphi(a)}$, $\tilde{\varphi}(a) = \varphi(a)$, $a \in A$, is a homomorphism of \mathfrak{M} -algebras.*

PROOF: It follows from the construction (see Theorem 2.8) that $\varphi(T_0) \subseteq T'_0$. Hence, $\tilde{\varphi}$ is a well-defined map, and it is straightforward to check that φ is a homomorphism of \mathfrak{M} -algebras. □

2.4 Averaging operators. Theorem 2.8 provides a powerful tool for solving various problems for di- and tri-algebras (see Section 4). Let us state here an equivalent definition of tri- \mathfrak{M} by means of averaging operators.

Definition 2.12. Suppose A is a Σ -algebra. A linear map $t : A \rightarrow A$ is called an *averaging operator* on A if

$$f(ta_1, \dots, ta_n) = tf(ta_1, \dots, ta_{i-1}, a_i, ta_{i+1}, \dots, ta_n)$$

for all $f \in \Sigma$, $\nu(f) = n$, $a_j \in A$, $i, j = 1, \dots, n$.

Let us call t a *homomorphic averaging operator* if

$$f(ta_1, \dots, ta_n) = tf(a_1^H, \dots, a_n^H),$$

for all $H \in \mathcal{P}(n)$, where

$$(4) \quad a_i^H = \begin{cases} a_i, & i \in H, \\ ta_i, & i \notin H. \end{cases}$$

Given a Σ -algebra A equipped with a homomorphic averaging operator t , denote by $A^{(t)}$ the following $\Sigma^{(3)}$ -algebra:

$$f^H(a_1, \dots, a_n) = f(a_1^H, \dots, a_n^H),$$

where $f \in \Sigma$, $\nu(f) = n$, $H \in \mathcal{P}(n)$, $a_i \in A$, a_i^H are given by (4).

If t is an averaging operator on A then the same rule defines $\Sigma^{(2)}$ -algebra $A^{(t)}$ provided that all H are singletons.

Theorem 2.13. *Suppose $\nu(f) \geq 2$ for all $f \in \Sigma$.*

- (1) *If $A \in \mathfrak{M}$ and t is an averaging operator on A then $A^{(t)}$ is a di- \mathfrak{M} -algebra.*
- (2) *If $A \in \mathfrak{M}$ and t is a homomorphic averaging operator on A then $A^{(t)}$ is a tri- \mathfrak{M} -algebra.*
- (3) *Every $D \in \text{di-}\mathfrak{M}$ may be embedded into $A^{(t)}$ for an appropriate $A \in \mathfrak{M}$ with an averaging operator t .*
- (4) *Every $T \in \text{tri-}\mathfrak{M}$ may be embedded into $A^{(t)}$ for an appropriate $A \in \mathfrak{M}$ with a homomorphic averaging operator t .*

PROOF: Let us show (2) and (4) since (1) and (3) are in fact restrictions of the statements on tri-algebras.

To prove (2), it is enough to note (by induction on m) that for every

$$\Phi = \Phi(x_1, \dots, x_m) \in \mathfrak{F}(m)$$

and for every $H \in \mathcal{P}(m)$ we have

$$\Phi^H(a_1, \dots, a_m) = \Phi(a_1^H, \dots, a_m^H), \quad a_i \in A.$$

Moreover, (1) also hold on $A^{(t)}$ by definition of t .

Statement (4) follows from Theorem 2.8: T is a subalgebra of $C_2 \otimes \widetilde{T}$. Consider $A = \widetilde{T} = \widetilde{T} \oplus T$ and define

$$ta = \bar{a}, \quad t\bar{a} = \bar{a}, \quad a \in T.$$

It is easy to see by definition of operations on \widetilde{T} that t is indeed a homomorphic averaging operator on A , and $T \subseteq A^{(t)}$ is a $\Sigma^{(3)}$ -subalgebra. □

3. Splitted algebras

In this section, we observe an approach to the procedure of splitting of an operad [2] that leads to classes of objects in some sense dual to di- and tri-algebras.

3.1 Definition and examples. As above, let \mathfrak{M} be a variety of Σ -algebras defined by a family of polylinear identities $S(\mathfrak{M})$.

Suppose T is a $\Sigma^{(3)}$ -algebra, and let $C \in \text{ComTrias}$. Define the following Σ -algebra structure on the space $C \otimes T$:

$$(5) \quad f(a_1 \otimes u_1, \dots, a_n \otimes u_n) = \sum_{H \in \mathcal{P}(n)} e_H^{(n)}(a_1, \dots, a_n) \otimes f^H(u_1, \dots, u_n),$$

$$(6) \quad f \in \Sigma, \nu(f) = n.$$

Denote the obtained Σ -algebra by $C \boxtimes T$.

In a similar way (considering only singletons in (5)) one may define $P \boxtimes D$ for a $\Sigma^{(2)}$ -algebra D and $P \in \text{Perm}$.

Definition 3.1. A class of $\Sigma^{(2)}$ -algebras D such that $P \boxtimes D \in \mathfrak{M}$ for all $P \in \text{Perm}$ is denoted by $\text{pre-}\mathfrak{M}$.

A class of $\Sigma^{(3)}$ -algebras T such that $C \boxtimes T \in \mathfrak{M}$ for all $C \in \text{ComTrias}$ is denoted by $\text{post-}\mathfrak{M}$.

It is enough to check whether $P \boxtimes D, C \boxtimes T \in \mathfrak{M}$ for free algebras $P = \text{Perm}\langle X \rangle$ and $C = \text{ComTrias}\langle X \rangle$, where $X = \{x_1, x_2, \dots\}$ is a countable set of symbols.

It is obvious that $\text{pre-}\mathfrak{M}$ and $\text{post-}\mathfrak{M}$ are varieties of $\Sigma^{(2)}$ - and $\Sigma^{(3)}$ -algebras, respectively. Indeed, it is easy to find their defining identities by the very definition.

Example 3.2. Suppose Σ consists of one binary operation $[\cdot, \cdot]$, and let $\mathfrak{M} = \text{Lie}$. Then $\Sigma^{(2)}$ consists of two operations, say, $[\cdot \vdash \cdot]$ and $[\cdot + \cdot]$. A $\Sigma^{(2)}$ -algebra D belongs to pre-Lie if and only if $\text{Perm}\langle X \rangle \boxtimes D \in \text{Lie}$, i.e.,

$$[(x_1 \otimes a_1), (x_2 \otimes a_2)] = x_1 x_2 \otimes [a_1 \vdash a_2] + x_2 x_1 \otimes [a_1 + a_2]$$

is anti-commutative and satisfies the Jacobi identity. The anti-commutativity implies

$$[a_1 \vdash a_2] = -[a_2 + a_1], \quad a_1, a_2 \in D.$$

Denote $[a \vdash b]$ by ab . Let us check the Jacobi identity:

$$(7) \quad \begin{aligned} & [[(x_1 \otimes a_1), (x_2 \otimes a_2)], (x_3 \otimes a_3)] \\ &= x_1 x_2 x_3 \otimes (a_1 a_2) a_3 - x_3 x_1 x_2 \otimes a_3 (a_1 a_2) - x_2 x_1 x_3 \otimes (a_2 a_1) a_3 + x_3 x_2 x_1 \otimes a_3 (a_2 a_1) \\ &= e_3^{(3)} \otimes ((a_1 a_2) a_3 - (a_2 a_1) a_3) - e_2^{(3)} \otimes a_3 (a_1 a_2) + e_1^{(3)} \otimes a_3 (a_2 a_1). \end{aligned}$$

Hence,

$$(8) \quad \begin{aligned} & [[(x_1 \otimes a_1), (x_2 \otimes a_2)], (x_3 \otimes a_3)] + [[(x_2 \otimes a_2), (x_3 \otimes a_3)], (x_1 \otimes a_1)] \\ &+ [[(x_3 \otimes a_3), (x_1 \otimes a_1)], (x_2 \otimes a_2)] = e_1^{(3)} (a_3 (a_2 a_1) - (a_3 a_2) a_1 + (a_2 a_3) a_1 - a_2 (a_3 a_1)) \\ &+ e_2^{(3)} ((a_3 a_1) a_2 - a_3 (a_1 a_2) + a_1 (a_3 a_2) - (a_1 a_3) a_2) \\ &+ e_3^{(3)} ((a_1 a_2) a_3 - a_1 (a_2 a_3) + a_2 (a_1 a_3) - (a_2 a_1) a_3). \end{aligned}$$

Hence, $D \in \text{pre-Lie}$ if and only if the product ab is left-symmetric.

Other well-known examples include pre-associative (dendriform) [21], post-associative (tridendriform) [23], pre-Poisson [1], pre-Jordan [16] algebras, as well as pre-Lie triple systems [6].

3.2 Equivalent description. Suppose T is a $\Sigma^{(3)}$ -algebra. Denote by \widehat{T} the direct sum of two isomorphic copies of T as of linear space:

$$\widehat{T} = T \oplus T'.$$

Assume the isomorphism is given by the correspondence $a \leftrightarrow a', a \in T$, and define

$$(9) \quad f(a_1 + b'_1, \dots, a_n + b'_n) = \sum_{H \in \mathcal{P}(n)} f^H(a_1, \dots, a_n) + \left(\sum_{H \in \mathcal{P}(n)} f^H(c_1^H, \dots, c_n^H) \right)',$$

where $f \in \Sigma, \nu(f) = n$, and

$$c_i^H = \begin{cases} a_i, & i \notin H, \\ b_i, & i \in H. \end{cases}$$

Thus, \widehat{T} carries the structure of a Σ -algebra. For a $\Sigma^{(2)}$ -algebra D , one may define \widehat{D} in a similar way assuming $f^H(x_1, \dots, x_n) = 0$ for $|H| > 1$.

Theorem 3.3 (cf. [15]). *The following statements are equivalent:*

- (1) $T \in \text{post-}\mathfrak{M}$;
- (2) $\widehat{T} \in \mathfrak{M}$.

Similarly, a $\Sigma^{(2)}$ -algebra D belongs to pre- \mathfrak{M} if and only if $\widehat{D} \in \mathfrak{M}$.

PROOF: Let us fix $C = \text{ComTrias}(Y)$, Y is an infinite set. It is enough to prove that (2) is equivalent to $C \boxtimes T \in \mathfrak{M}$.

Suppose $\Phi = \Phi(x_1, \dots, x_n) \in \mathfrak{F}(n)$ is a polylinear term of degree n in the language Σ . Evaluate the term Φ in $C \boxtimes T$:

$$\Phi(y_1 \otimes a_1, \dots, y_n \otimes a_n) = \sum_{H \in \mathcal{P}(n)} e_H^{(n)}(y_1, \dots, y_n) \otimes \Phi_{(H)}(a_1, \dots, a_n).$$

This equation defines a family of n -linear functions $\Phi_{(H)} : T^{\otimes n} \rightarrow T, H \in \mathcal{P}(n)$.

Lemma 3.4. *In the algebra \widehat{T} , the following equations hold for $a_i \in T \subset \widehat{T} (i = 1, \dots, n)$:*

$$(10) \quad \Phi_{(H)}(a_1, \dots, a_n)' = \Phi(d_1^H, \dots, d_n^H),$$

$$(11) \quad \sum_{H \in \mathcal{P}(n)} \Phi_{(H)}(a_1, \dots, a_n) = \Phi(a_1, \dots, a_n).$$

where

$$d_i^H = \begin{cases} a'_i, & i \in H, \\ a_i, & i \notin H. \end{cases}$$

PROOF: If $n = 1$ then (10) is trivial. Proceed by induction on n . Assume

$$\Phi = f(\Psi_1, \dots, \Psi_m), \quad f \in \Sigma, \quad \nu(f) = m,$$

where $\Psi_i \in \mathfrak{F}(n_i)$, $n_1 + \dots + n_m = n$. Suppose $z_{ij} \in Y$ are pairwise different, $a_{ij} \in T$, $i = 1, \dots, m, j = 1, \dots, n_i$. To simplify notations, denote

$$\bar{z}_i = (z_{i1}, \dots, z_{in_i}), \quad \bar{a}_i = (a_{i1}, \dots, a_{in_i}), \quad i = 1, \dots, m.$$

For $H_i \in \mathcal{P}(n_i)$, denote by $\bar{a}_i^{H_i}$ the n_i -tuple $(d_{i1}^{H_i}, \dots, d_{in_i}^{H_i})$ obtained from the initial one by “adding primes” to all those components that belong to H_i .

Then

$$\begin{aligned} & f(z_{11} \otimes a_{11}, \dots, z_{mn_m} \otimes a_{mn_m}) \\ &= \sum_{\substack{K \in \mathcal{P}(m) \\ H_1 \in \mathcal{P}(n_1) \\ H_m \in \mathcal{P}(n_m)}}} e_K^{(m)}(e_{H_1}^{(n_1)}(\bar{z}_1), \dots, e_{H_m}^{(n_m)}(\bar{z}_m)) \otimes f^K(\Psi_{1(H_1)}(\bar{a}_1), \dots, \Psi_{m(H_m)}(\bar{a}_m)) \\ &= \sum_{\substack{K \in \mathcal{P}(m) \\ H_1 \in \mathcal{P}(n_1) \\ H_m \in \mathcal{P}(n_m)}}} e_{K(H_1, \dots, H_m)}^{(n)}(z_{11}, \dots, z_{mn_m}) \otimes f^K(\Psi_{1(H_1)}(\bar{a}_1), \dots, \Psi_{m(H_m)}(\bar{a}_m)), \end{aligned}$$

where $K(H_1, \dots, H_m)$ is the composition of sets from the definition of ComTrias.

Hence, for every $H \in \mathcal{P}(n)$ we have

$$(12) \quad \Phi_{(H)}(a_{11}, \dots, a_{mn_m}) = \sum_{\substack{K, H_1, \dots, H_m \\ K(H_1, \dots, H_m) = H}} f^K(\Psi_{1(H_1)}(\bar{a}_1), \dots, \Psi_{m(H_m)}(\bar{a}_m)).$$

By definition, every H uniquely determines K and H_i for $i \in K$. Other H_j (for $j \notin K$) in (12) run through the entire $\mathcal{P}(n_j)$. Therefore,

$$\Phi_{(H)}(a_{11}, \dots, a_{mn_m})' = f^K(b_1, \dots, b_m)', \quad b_i = \begin{cases} \Psi_{i(H_i)}(\bar{a}_i)', & i \in K, \\ \sum_{H_j \in \mathcal{P}(n_j)} \Psi_{i(H_j)}(\bar{a}_i), & i \notin K. \end{cases}$$

By the inductive assumption,

$$\Psi_{i(H_i)}(\bar{a}_i)' = \Psi_i(\bar{a}_i^{H_i}), \quad \sum_{H_i \in \mathcal{P}(n_i)} \Psi_{i(H_i)}(\bar{a}_i) = \Psi_i(\bar{a}_i).$$

It remains to apply the definition of operations in \widehat{D} (9) to prove (10).

To complete the proof, apply (12) and Lemma 2.4:

$$\begin{aligned} \sum_{H \in \mathcal{P}(n)} \Phi_{(H)}(a_{11}, \dots, a_{mm_m}) &= \sum_{H \in \mathcal{P}(n)} \sum_{\substack{K, H_1, \dots, H_m \\ K(H_1, \dots, H_m) = H}} f^K(\Psi_{1(H_1)}(\bar{a}_1), \dots, \Psi_{m(H_m)}(\bar{a}_m)) \\ &= \sum_{K \in \mathcal{P}(m)} \sum_{H_1 \in \mathcal{P}(n_1)} \dots \sum_{H_m \in \mathcal{P}(n_m)} f^K(\Psi_{1(H_1)}(\bar{a}_1), \dots, \Psi_{m(H_m)}(\bar{a}_m)). \end{aligned}$$

Now (11) follows from multilinearity of f^K and inductive assumption. □

Let us finish the proof of the theorem. If $\widehat{T} \in \mathfrak{M}$ then $\text{ComTrias}(Y) \boxtimes T$ satisfies all defining identities of the variety \mathfrak{M} by Lemma 3.4.

The converse is even simpler. Note that $\widehat{T} = C_2 \boxtimes T$, where C_2 is the 2-dimensional ComTrias-algebra from Example 2.3. By the very definition, $\widehat{T} \in \mathfrak{M}$. □

Remark 2. Note that the base field itself is a 1-dimensional algebra in ComTrias. Therefore, if $A \in \text{pre-}\mathfrak{M}$ or $A \in \text{post-}\mathfrak{M}$ then $\mathbb{k} \boxtimes A \in \mathfrak{M}$. This observation explains the term “splitting”: An operation $f \in \Sigma$, $\nu(f) = n$, splits into n or $2^n - 1$ operations, $f = \sum_H f^H$.

3.3 Rota-Baxter operators. Let A be a Σ -algebra.

Definition 3.5 (cf. [2]). A linear map $\tau : A \rightarrow A$ is said to be a Rota–Baxter operator of weight λ ($\lambda \in \mathbb{k}$) if

$$(13) \quad f(\tau(a_1), \dots, \tau(a_n)) = \sum_{H \in \mathcal{P}(n)} \lambda^{|H|-1} \tau(f(a_1^H, \dots, a_n^H)),$$

$$(14) \quad a_i^H = \begin{cases} a_i, & i \in H, \\ \tau(a_i), & i \notin H, \end{cases}$$

for all $f \in \Sigma$, $\nu(f) = n$, $a_i \in A$.

Obviously, if τ is a Rota–Baxter operator of nonzero weight λ then $\tau' = \frac{1}{\lambda} \tau$ is a Rota–Baxter operator of weight 1. Hence, there are two essentially different cases: $\lambda = 0$ (zero weight) and $\lambda = 1$ (unit weight).

The following statement was proved in [15] in the case of binary operations ($\nu(f) = 2$). By means of the approach presented in this paper, the proof becomes clear in the general case.

Given a Σ -algebra A equipped with a Rota–Baxter operator τ , denote by $A^{(\tau)}$ the $\Sigma^{(3)}$ -algebra defined on the space A by

$$f^H(a_1, \dots, a_n) = f(a_1^H, \dots, a_n^H),$$

where $f \in \Sigma$, $i = 1, \dots, n$, $a_1, \dots, a_n \in A$, a_i^H are given by (14).

The same relations restricted to $|H| = 1$ define a $\Sigma^{(2)}$ -algebra structure on A also denoted by $A^{(\tau)}$.

- Theorem 3.6.** (1) If $A \in \mathfrak{M}$ and τ is a Rota–Baxter operator of zero weight on A then $A^{(\tau)}$ is a pre- \mathfrak{M} -algebra.
 (2) If $A \in \mathfrak{M}$ and τ is a Rota–Baxter operator of unit weight on A then $A^{(\tau)}$ is a post- \mathfrak{M} -algebra.
 (3) Every $D \in \text{pre-}\mathfrak{M}$ may be embedded into $A^{(\tau)}$ for an appropriate $A \in \mathfrak{M}$ equipped with Rota–Baxter operator τ of zero weight.
 (4) Every $T \in \text{post-}\mathfrak{M}$ may be embedded into $A^{(\tau)}$ for an appropriate $A \in \mathfrak{M}$ equipped with Rota–Baxter operator τ of unit weight.

PROOF: As in Theorem 2.13, let us prove (2) and (4).

For (2), it is enough to consider $C \boxtimes A^{(\tau)}$ for any $C \in \text{ComTrias}$, and note (by induction on $m \geq 1$) that

$$\Phi(y_1 \otimes a_1, \dots, y_m \otimes a_m) = \sum_{H \in \mathcal{P}(m)} e^{(m)}(y_1, \dots, y_m) \otimes \Phi(a_1^H, \dots, a_m^H)$$

for every $\Phi \in \mathfrak{F}(m)$.

Hence, $C \boxtimes A^{(\tau)} \in \mathfrak{M}$.

To prove (4), consider $A = C_2 \boxtimes T \in \mathfrak{M}$, where C_2 is the algebra from Example 2.3, and define

$$(15) \quad \tau(e_1 \otimes a) = -e_1 \otimes a, \quad \tau(e_2 \otimes a) = e_1 \otimes a, \quad a \in T.$$

Let us show that (15) is a Rota–Baxter operator of unit weight on $C_2 \boxtimes T$. Indeed, suppose $f \in \Sigma$, $\nu(f) = n$, $u_i = e_{k_i} \otimes a_i$, $k_i \in \{1, 2\}$, $a_i \in T$, $i = 1, \dots, n$. Evaluate the left-hand side of (13):

$$f(\tau(u_1), \dots, \tau(u_n)) = (-1)^{|K|} f(e_1 \otimes a_1, \dots, e_1 \otimes a_n),$$

where $K = \{i \mid k_i = 1\}$. On the other hand,

$$\begin{aligned} f(u_1^H, \dots, u_n^H) &= (-1)^{|K \setminus H|} f(e_{k'_i} \otimes a_1, \dots, e_{k'_n} \otimes a_n) \\ &= (-1)^{|K \setminus H|} \sum_{M \in \mathcal{P}(n)} e_M^{(n)}(e_{k'_1}, \dots, e_{k'_n}) \otimes f^M(a_1, \dots, a_n), \end{aligned}$$

where $k'_i = \begin{cases} k_i, & i \in H, \\ 1, & i \notin H \end{cases}$. Nonzero summands appear in two cases: (1) $k'_i = 1$ for all $i = 1, \dots, n$; (2) $k'_i = 2$ if and only if $i \in M$. The first case occurs if and only if $H \subseteq K$, the second one corresponds to $M = H \setminus K$. Hence,

$$f(u_1^H, \dots, u_n^H) = \begin{cases} (-1)^{|K| - |H|} e_1 \otimes \sum_{M \in \mathcal{P}(n)} f^M(a_1, \dots, a_n), & H \subseteq K, \\ (-1)^{|K \setminus H|} e_2 \otimes f^{H \setminus K}(a_1, \dots, a_n), & H \not\subseteq K. \end{cases}$$

Let us evaluate the right-hand side of (13):

$$(16) \quad \sum_{H \in \mathcal{P}(n)} \tau(f(u_1^H, \dots, u_n^H)) = \sum_{\emptyset \neq H \subseteq K} (-1)^{|K|-|H|+1} f(e_1 \otimes a_1, \dots, e_1 \otimes a_n) + \sum_{H \not\subseteq K} (-1)^{|K \setminus H|} e_1 \otimes f^{H \setminus K}(a_1, \dots, a_n).$$

The first summand in the right-hand side of (16) is equal to $(-1)^{|K|} f(e_1 \otimes a_1, \dots, e_1 \otimes a_n)$ since

$$\sum_{H \subseteq K} (-1)^{|H|} = 1 + \sum_{\emptyset \neq H \subseteq K} (-1)^{|H|} = 0.$$

In the second summand, present $H \not\subseteq K$ as $H = U \cup M$, $U \subseteq K$, $M \neq \emptyset$, $M \cap K = \emptyset$. Then

$$\sum_{U \subseteq K} \sum_{\substack{M \neq \emptyset \\ M \cap K = \emptyset}} (-1)^{|K|-|U|} e_1 \otimes f^M(a_1, \dots, a_n) = 0$$

by the same reasons.

We have proved that (13) holds for τ ($\lambda = 1$), and thus it is a Rota–Baxter operator of unit weight. □

Remark 3. Theorem 3.6 implies that Definition 3.1 provides an equivalent description of the same class of systems as the splitting procedure described in [2]: $\text{pre-}\mathfrak{M} = \mathcal{A}\text{Sp}(\mathfrak{M})$, $\text{post-}\mathfrak{M} = \mathcal{B}\text{Sp}(\mathfrak{M})$.

In the binary case, $\text{pre-}\mathfrak{M}$ and $\text{post-}\mathfrak{M}$ coincide with operads denoted in [15] by $\text{DendDi}\mathfrak{M}$ and $\text{DendTri}\mathfrak{M}$, respectively.

Remark 4. Indeed, it was shown in [15] that if \mathfrak{M} is a binary quadratic operad then $\text{pre-}\mathfrak{M} = \text{pre-Lie} \bullet \mathfrak{M}$, $\text{post-}\mathfrak{M} = \text{post-Lie} \bullet \mathfrak{M}$, where \bullet is the Manin black product of operads [13],

$$(\text{pre-}\mathfrak{M})^! = \text{di-}(\mathfrak{M}^!), \quad (\text{post-}\mathfrak{M})^! = \text{tri-}(\mathfrak{M}^!),$$

where $!$ stands for Koszul duality of operads.

4. Problems on replicated algebras

In this section, we consider a series of problems for replicated algebras. Some of them have already been solved in particular cases. Here we will show how to solve them in general.

4.1 Codimension of varieties. Given an operad \mathfrak{M} , the number $c_n(\mathfrak{M}) = \dim \mathfrak{M}(n)$, $n \geq 1$, (if it is finite) is called *codimension* of \mathfrak{M} . The growth of codimensions, namely, of $\sqrt[n]{c_n(\mathfrak{M})}$ is intensively studied since the seminal paper [12] on associative algebras.

It follows immediately from definition that for a variety $\text{di-}\mathfrak{M}$ or $\text{tri-}\mathfrak{M}$ the codimension may be explicitly evaluated as a product of $c_n(\text{Perm})$ or $c_n(\text{ComTrias})$ with $c_n(\mathfrak{M})$.

Proposition 4.1. *For every operad \mathfrak{M} , $c_n(\text{di-}\mathfrak{M}) = n c_n(\mathfrak{M})$, $c_n(\text{tri-}\mathfrak{M}) = (2^n - 1) c_n(\mathfrak{M})$.*

In particular, if \mathfrak{M} is a variety of Lie algebras of polynomial codimension growth then so is the variety $\text{di-}\mathfrak{M}$ of Leibniz algebras.

4.2 Replication of morphisms of operads. Let $\mathfrak{M}, \mathfrak{N}$ be two operads. Suppose $\omega : \mathfrak{N} \rightarrow \mathfrak{M}$ is a morphism of operads. Then for every algebra A in \mathfrak{M} one may define $A^{(\omega)} \in \mathfrak{N}$, a new algebra structure on the same linear space A .

The well-known examples include $- : \text{Lie} \rightarrow \text{As}$, $x_1x_2 \mapsto x_1x_2 - x_2x_1$, a similar morphism $\text{Mal} \rightarrow \text{Alt}$, as well as $+ : \text{Jord} \rightarrow \text{As}$, $x_1x_2 \mapsto x_1x_2 + x_2x_1$, and many others.

For every $B \in \mathfrak{N}$ there exists unique (up to isomorphism) algebra $U_\omega(B) \in \mathfrak{M}$ such that:

- there exists a homomorphism $\iota : B \rightarrow U_\omega(B)^{(\omega)}$ of algebras in \mathfrak{N} ;
- for every algebra $A \in \mathfrak{M}$ and for every homomorphism $\psi : B \rightarrow A^{(\omega)}$ there exists unique homomorphism $\xi : U_\omega(B) \rightarrow A$ of algebras in \mathfrak{M} such that $\psi(b) = \xi(\iota(b))$ for all $b \in B$.

The algebra $U_\omega(B)$ is called the *universal enveloping* algebra of B with respect to ω . Note that ι is not necessarily injective, e.g., for the Albert algebra $H_3(\mathbb{O}) \in \text{Jord}$ the universal enveloping associative algebra (with respect to $+$) is equal to $\{0\}$.

An \mathfrak{N} -algebra B is *special* relative to ω if there exists an \mathfrak{M} -algebra A such that B is isomorphic to a subalgebra of $A^{(\omega)}$.

Definition 2.5 immediately implies

Proposition 4.2. *Given a morphism of operads $\omega : \mathfrak{N} \rightarrow \mathfrak{M}$, the map $\text{id} \otimes \omega : \text{tri-}\mathfrak{N} = \text{ComTrias} \otimes \mathfrak{N} \rightarrow \text{ComTrias} \otimes \mathfrak{M} = \text{tri-}\mathfrak{M}$ is also a morphism of operads.*

A similar statement for di-algebra case obviously holds.

4.3 PBW-type problems. The following natural problems appear each time when we consider a morphism of operads $\omega : \mathfrak{N} \rightarrow \mathfrak{M}$.

- Embedding problem: Whether every $B \in \mathfrak{N}$ is special with respect to ω ?
- Ado problem: Whether every finite-dimensional algebra $B \in \mathfrak{N}$ is a subalgebra of $A^{(\omega)}$, where $A \in \mathfrak{M}$, $\dim A < \infty$?
- Poincaré–Birkhoff–Witt (PBW) problem: Given $B \in \mathfrak{N}$, what is the structure of the universal enveloping algebra $U_\omega(B) \in \mathfrak{M}$?

Suppose \mathfrak{N} and \mathfrak{M} are varieties of Σ - and Σ' -algebras, respectively. Throughout this section, assume $\nu(f) \geq 2$ for all $f \in \Sigma \cup \Sigma'$.

The following lemma is an immediate corollary of definitions.

Lemma 4.3. *For every morphism of operads $\omega : \mathfrak{N} \rightarrow \mathfrak{M}$ and for every $A \in \text{tri-}\mathfrak{M}$, $C \in \text{ComTrias}$ we have*

$$C \otimes A^{(\omega)} = (C \otimes A)^{(\text{id} \otimes \omega)} \in \text{tri-}\mathfrak{N}.$$

A similar statement holds for di-algebras [19].

Theorem 4.4. *If the embedding problem has positive solution for $\omega : \mathfrak{N} \rightarrow \mathfrak{M}$ then it has positive solution for $\text{id} \otimes \omega : \text{di-}\mathfrak{N} \rightarrow \text{di-}\mathfrak{M}$ and for $\text{id} \otimes \omega : \text{tri-}\mathfrak{N} \rightarrow \text{tri-}\mathfrak{M}$. The same statement holds for the Ado problem.*

PROOF: Let us consider tri-algebra case. If $T \in \text{tri-}\mathfrak{M}$ then $\widetilde{T} \in \mathfrak{M}$ and $T \subseteq C_2 \otimes \widetilde{T}$. If $\widetilde{T} \subseteq A^{(\omega)}$ for some $A \in \mathfrak{M}$ then $T \subseteq C_2 \otimes \widetilde{T} \subseteq C_2 \otimes A^{(\omega)} = (C_2 \otimes A)^{(\text{id} \otimes \omega)}$, $C_2 \otimes A \in \text{tri-}\mathfrak{M}$.

Finally, if $\dim T < \infty$ then $\dim \widetilde{T} < \infty$ and $\dim(C_2 \otimes \widetilde{T}) < \infty$, the same holds for A . Hence, if \widetilde{T} has a finite-dimensional envelope then so is T . □

Let $T \in \text{tri-}\mathfrak{M}$, $\widetilde{T} \in \mathfrak{M}$, $\iota : \widetilde{T} \rightarrow U_\omega(\widetilde{T})^{(\omega)}$ as above.

Theorem 4.5. *The subalgebra generated in $C_2 \otimes U_\omega(\widetilde{T}) \in \text{tri-}\mathfrak{M}$ by the set $\{e_1 \otimes \iota(\bar{a}) + e_2 \otimes \iota(a) \mid a \in T\}$ is isomorphic to $U_{\text{id} \otimes \omega}(T) \in \text{tri-}\mathfrak{M}$.*

PROOF: Consider $\text{id} \otimes \iota : C_2 \otimes \widetilde{T} \rightarrow C_2 \otimes U_\omega(\widetilde{T})^{(\omega)} = (C_2 \otimes U_\omega(\widetilde{T}))^{(\text{id} \otimes \omega)}$. Hence, the restriction of $\text{id} \otimes \iota$ to $T \subseteq C_2 \otimes \widetilde{T}$ maps $a \in T$ to $e_1 \otimes \iota(\bar{a}) + e_2 \otimes \iota(a)$. Denote by U the subalgebra of $C_2 \otimes U_\omega(\widetilde{T}) \in \text{tri-}\mathfrak{M}$ generated by $(\text{id} \otimes \iota)(T)$. Let us check the universal property for U .

Suppose $A \in \text{tri-}\mathfrak{M}$, and $\psi : T \rightarrow A^{(\text{id} \otimes \omega)}$. By Lemma 2.11, there exists $\tilde{\psi} : \widetilde{T} \rightarrow \widetilde{A}^{(\text{id} \otimes \omega)}$. Note that $A_0^{(\text{id} \otimes \omega)} \subseteq A_0$. It follows from the construction of \tilde{A} , see the proof of Theorem 2.8. Therefore, there exists natural homomorphism $\widetilde{A}^{(\text{id} \otimes \omega)} \rightarrow \tilde{A}^{(\omega)}$, and we may consider $\tilde{\psi}$ as a homomorphism from \widetilde{T} to $\tilde{A}^{(\omega)}$, where

$$\tilde{\psi}(\bar{a}) = \overline{\psi(a)} \in A/A_0, \quad \tilde{\psi}(a) = \psi(a)$$

for $a \in T$.

By definition, there exists a homomorphism of \mathfrak{M} -algebras $\xi : U_\omega(\widetilde{T}) \rightarrow \tilde{A}$ such that $\xi(\iota(x)) = \tilde{\psi}(x)$, $x \in \widetilde{T}$. Then $\text{id} \otimes \xi : C_2 \otimes U_\omega(\widetilde{T}) \rightarrow C_2 \otimes \tilde{A}$ is a homomorphism of tri- \mathfrak{M} -algebras. Moreover, it is easy to see that $(\text{id} \otimes \xi)((\text{id} \otimes \iota)(T)) \subseteq A \subseteq C_2 \otimes \tilde{A}$. Hence, the restriction of $\text{id} \otimes \xi$ to $U \subseteq C_2 \otimes U_\omega(\widetilde{T})$ is the desired homomorphism of tri- \mathfrak{M} -algebras $U \rightarrow A$. □

The similar statement obviously holds for di-algebras (consider C_2 as an algebra in Perm). For example, the morphism $- : \text{Lie} \rightarrow \text{As}$ leads to $\text{id} \otimes - : \text{Leib} \rightarrow \text{di-As}$ considered in [22], see also [21]. The PBW Theorem for Leibniz algebras is an immediate corollary of Theorem 4.5.

Let us deduce PBW Theorem for tri-Lie algebras as an application of Theorem 4.5. Every $L \in \text{tri-Lie}$ with operations $[\cdot \vdash \cdot]$, $[\cdot \dashv \cdot]$, and $[\cdot \perp \cdot]$ gives rise to the following Lie algebras: $\bar{L} = L/L_0$ and $L_\perp = (L, [\cdot \perp \cdot])$.

Corollary 4.6. *Let $L \in \text{tri-Lie}$. Then $U_{\text{id} \otimes -}(L) \in \text{tri-As}$ as a linear space is isomorphic to $U(\bar{L}) \otimes U_0(L_\perp)$, where $U(\cdot)$ is the ordinary universal enveloping associative algebra with identity, $U_0(\cdot)$ stands for its augmentation ideal.*

PROOF: Suppose $B \subset L$ is a well-ordered linear basis of L . It is easy to see that the defining identities of $U_{\text{id} \otimes -}(L)$, namely,

$$a \vdash b - b \vdash a = [a \vdash b], \quad a \perp b - b \perp a = [a \perp b], \quad a, b \in B, a \geq b,$$

allow to present every element of $U_{\text{id} \otimes -}(L) \in \text{tri-As}$ as a linear combination of

$$(17) \quad u = (a_1 \vdash \cdots \vdash a_n \vdash b_1) \perp b_2 \perp \cdots \perp b_m,$$

$a_i, b_j \in B, a_1 \leq \dots \leq a_n, b_1 \leq \dots \leq b_m, n \geq 0, m \geq 1$, and $\bar{a}_1, \dots, \bar{a}_n$ are linearly independent in \tilde{L} .

It remains to show that the elements (17) are linearly independent in $U_{\text{id}\otimes}(L)$. By Theorem 4.5, $U_{\text{id}\otimes}(L) \subseteq C_2 \otimes U(\tilde{L})$. Identify x and $\iota(x)$ for $x \in \tilde{L}$ and evaluate

$$\tilde{u} = (\tilde{a}_1 \vdash \dots \vdash \tilde{a}_n \vdash \tilde{b}_1) \perp \tilde{b}_2 \perp \dots \perp \tilde{b}_m,$$

where $\tilde{a} = e_1 \otimes \bar{a} + e_2 \otimes a, a \in B$. By the definition of C_2 ,

$$\tilde{u} = e_1 \otimes \bar{a}_1 \dots \bar{a}_n \bar{b}_1 \dots \bar{b}_m + e_2 \otimes \bar{a}_1 \dots \bar{a}_n b_1 \dots b_m.$$

By the choice of a_i, b_j the second summands are linearly independent in $C_2 \otimes U(\tilde{L})$. \square

4.4 Special identities. Let $\omega : \mathfrak{R} \rightarrow \mathfrak{M}$ be a morphism of operads. An algebra $B \in \mathfrak{R}$ is said to be *special* with respect to ω if there exists $A \in \mathfrak{M}$ such that B is a subalgebra in $A^{(\omega)}$.

The class of all special algebras in \mathfrak{R} with respect to ω may not form a subvariety of \mathfrak{R} : It is not closed with respect to homomorphic image. The variety generated by all special algebras is denoted by $S^{(\omega)}\mathfrak{R}$. The corresponding operad is an image of \mathfrak{R} . Nonzero elements of the kernel of the corresponding morphism of operads (if they exist) are exactly all polylinear identities that hold on all special algebras in \mathfrak{R} but do not hold on the entire \mathfrak{R} . Such identities are called special (with respect to ω).

Theorem 4.7. *If $\text{char } \mathbb{k} = 0$ then the following equation holds for varieties:*

$$S^{(\text{id}\otimes\omega)}\text{tri-}\mathfrak{R} = \text{tri-}S^{(\omega)}\mathfrak{R}.$$

The proof is completely similar to di-algebra case in [19]. The only difference appears in using $C_2 \in \text{ComTrias}$ instead of $\mathbb{k}[x] \in \text{Perm}$, where the Perm-algebra structure on polynomials was given by $f(x)g(x) = f(0)g(x)$. Let us sketch the main steps of the proof in these new settings.

PROOF: (\subseteq) It is enough to prove that every $T \in \text{tri-}\mathfrak{R}$ which is special with respect to $\text{id} \otimes \omega$ satisfies replicated polylinear special identities. Indeed, if $T \subseteq A^{(\text{id}\otimes\omega)}$ for $A \in \text{tri-}\mathfrak{M}$ then $\psi : \tilde{T} \rightarrow \widetilde{A^{(\text{id}\otimes\omega)}} \rightarrow \tilde{A}^{(\omega)}$ is a homomorphism of tri- \mathfrak{R} -algebras. Then $\text{id} \otimes \psi : C_2 \otimes \tilde{T} \rightarrow C_2 \otimes \tilde{A}^{(\omega)}$ is a homomorphism of tri- \mathfrak{R} -algebras which is injective on $T \subseteq C_2 \otimes \tilde{T}$. Hence, T satisfies all identities that hold on $C_2 \otimes \tilde{A}^{(\omega)} \in \text{tri-}S^{(\omega)}\mathfrak{R}$.

(\supseteq) If $T \in \text{tri-}S^{(\omega)}\mathfrak{R}$ then $\tilde{T} \in S^{(\omega)}\mathfrak{R}$ and thus \tilde{T} is a homomorphic image of a special algebra $B \subseteq A^{(\omega)}, A \in \mathfrak{M}$. It is straightforward to deduce that $C_2 \otimes \tilde{T}$ is then a homomorphic image of a special algebra $(C_2 \otimes A)^{(\text{id}\otimes\omega)}$. Therefore, T belongs to $S^{(\text{id}\otimes\omega)}\text{tri-}\mathfrak{R}$. \square

4.5 TKK construction for tri-Jordan algebras. The classical Tits–Kantor–Koecher (TKK) construction of a Lie algebra $T(J)$ for a Jordan algebra J is known to preserve simplicity, nilpotence and strong (Penico) solvability. Moreover, $T(J)$ is a \mathbb{Z}_3 -graded Lie algebra $J^+ \oplus S(J) \oplus J^-$, where J^\pm are isomorphic copies of the space $J, S(J)$ is the structure algebra constructed by inner derivations and operators of left multiplication in J [17].

The TKK construction for Jordan dialgebras was done in [14]. There was also proved an analogue of Zhevlakov theorem [32] which for ordinary Jordan algebras states that any finitely generated solvable Jordan algebra is nilpotent.

Proposition 4.8. *A finitely generated solvable tri-Jordan algebra is nilpotent.*

PROOF: Let us consider finitely generated and solvable $J \in \text{tri-Jord}$. By Lemma 2.9, $\tilde{J} = \tilde{J} \oplus J \in \text{Jord}$, it has to be a finitely generated and solvable Jordan algebra by the construction. By the Zhevlakov theorem, \tilde{J} is nilpotent. Hence, $C_2 \otimes \tilde{J} \in \text{tri-Jord}$ is also nilpotent, and by Theorem 2.8 so is $J \subseteq C_2 \otimes \tilde{J}$. \square

The notion of strong solvability for Jordan algebras is translated to di- and tri-algebras in a straightforward way (the minimal change is due to absence of commutativity). For a tri-Jordan J , the language $\Sigma^{(3)}$ contains three operations \vdash, \dashv, \perp (note that $a \vdash b = b \dashv a$). Consider the sequence

$$J^{(1)} = J, \quad J^{(2)} = J \cdot J,$$

$$J^{(n+1)} = J^{(n)} \cdot J^{(n)} + J \cdot (J^{(n)} \cdot J^{(n)}) + (J^{(n)} \cdot J^{(n)}) \cdot J, \quad n > 1,$$

where $A \cdot B$ stands for $A \vdash B + A \perp B$. All $J^{(n)}$ are ideals of J . If there exists $N \geq 1$ such that $J^{(N)} = 0$ then J is said to be *strongly solvable* (or Penico solvable).

Let us state an analogue of the TKK construction for tri-Jordan.

Proposition 4.9. *For every $J \in \text{tri-Jord}$ there exists $T(J) \in \text{tri-Lie}$ such that the following properties hold:*

- $T(J) = J_{-1} \oplus J_0 \oplus J_1$ is \mathbb{Z}_3 -graded algebra, where the spaces J_{-1}, J_1 are copies of J ;
- $T(J)$ is nilpotent if and only if J is nilpotent;
- $T(J)$ is solvable if and only if J is strongly solvable.

PROOF: Let us consider $X(J) = C_2 \otimes T(\tilde{J})$, where $T(\tilde{J}) = \tilde{J}^- \oplus St(\tilde{J}) \oplus \tilde{J}^+$ is the TKK construction for Jordan algebra \tilde{J} , where $St(\tilde{J})$ is the structure algebra of \tilde{J} . By Lemma 2.9, $X(\tilde{J}) \in \text{tri-Lie}$. We can represent $X(J)$ as a \mathbb{Z}_3 -graded space

$$(C_2 \otimes \tilde{J}^-) \oplus (C_2 \otimes St(\tilde{J})) \oplus (C_2 \otimes \tilde{J}^+).$$

Let J^\pm be subspaces in $C_2 \otimes \tilde{J}^\pm$ spanned by isomorphic images of elements $e_1 \otimes \bar{a} + e_2 \otimes a$, $a \in J$. The subalgebra $T(J)$ generated by J^+ and J^- in $X(J) \in \text{tri-Lie}$ is the required one. Indeed, $T(J)$ is nilpotent or solvable if and only if J is nilpotent or strongly solvable, respectively, because of the definitions of $C_2 \otimes \tilde{J}$ and properties of TKK construction for ordinary algebras. \square

4.6 Tri-Jordan polynomials. Another classical question is related with Cohn’s description of Jordan polynomials in the free associative algebra [10]. Suppose $\text{char } \mathbb{k} \neq 2$. For the morphism of operads $+$: $\text{Jord} \rightarrow \text{As}$ defined by $x_1 x_2 \mapsto x_1 x_2 + x_2 x_1$, the free algebra $SJ\langle X \rangle = S^{(+)}\text{Jord}\langle X \rangle$ is a subspace of $\text{As}\langle X \rangle$, elements of $SJ\langle X \rangle$ are called Jordan polynomials. It is well-known since [10] that $SJ\langle X \rangle \subseteq H\langle X \rangle$, where $H\langle X \rangle$ is the space of

symmetric elements with respect to involution $\sigma : x_1 \dots x_n \mapsto x_n \dots x_1$. The embedding is strict if and only if $|X| > 3$.

For di-Jordan algebras a similar question was considered in [31]: It was shown that $S^{(+)}\text{di-Jord}\langle X \rangle$ lies in the space of symmetric elements (with respect to naturally defined involution), and the embedding is strict if and only $|X| > 2$. Here we use (+) for (id \otimes +) to simplify notations.

Theorem 2.8 provides a way to solve the same question for tri-Jordan algebras. Let us sketch the proof which is even simpler than the proof in di-algebra case [31].

Denote by σ the linear map $\text{tri-As}\langle X \rangle \rightarrow \text{tri-As}\langle X \rangle$ such that

$$\begin{aligned} \sigma(x) &= x, & x \in X; \\ \sigma(u \vdash v) &= \sigma(v) \dashv \sigma(u), \\ \sigma(u \dashv v) &= \sigma(v) \vdash \sigma(u), \\ \sigma(u \perp v) &= \sigma(v) \perp \sigma(u), & u, v \in \text{tri-As}\langle X \rangle. \end{aligned}$$

Denote $\text{tri-H}\langle X \rangle = \{f \in \text{tri-As}\langle X \rangle \mid \sigma(f) = f\}$.

Proposition 4.10. *For every X , $S^{(+)}\text{tri-Jord}\langle X \rangle \subseteq \text{tri-H}\langle X \rangle$. The embedding is strict if and only if $|X| > 1$.*

PROOF: Obviously, $S^{(+)}\text{tri-Jord}\langle X \rangle \subseteq \text{tri-H}\langle X \rangle$. If $|X| > 1$ then

$$f = (x_1 \vdash x_2 \vdash x_2) \perp x_1 + x_1 \perp (x_2 \dashv x_2 \dashv x_1) \in \text{tri-H}\langle X \rangle \setminus S^{(+)}\text{tri-Jord}\langle X \rangle.$$

Indeed, consider the tri-algebra analogue of the Grassmann algebra $\wedge\langle \xi_1, \dots, \xi_n \rangle$ constructed as follows. Associative algebra

$$A_n = \text{As}\langle \bar{\xi}_1, \dots, \bar{\xi}_n, \xi_1, \dots, \xi_n \mid ab = -ba, a^2 = 0, a, b \in \{\xi_i, \bar{\xi}_i \mid i = 1, \dots, n\} \rangle$$

is equipped with homomorphic averaging operator τ given by $\xi_i \mapsto \bar{\xi}_i, \bar{\xi}_i \mapsto \xi_i$. Therefore, $A_n^{(\tau)} \in \text{tri-As}$ by 2.13.

The epimorphism $\theta : \text{tri-As}\langle x_1, x_2 \rangle \rightarrow A_2^{(\tau)}$ defined by $x_1 \mapsto \xi_1, x_2 \mapsto \xi_2$ annihilates $S^{(+)}\text{tri-Jord}\langle x_1, x_2 \rangle$, but does not annihilate f :

$$\theta(f) = \bar{\xi}_1 \bar{\xi}_2 \xi_2 \xi_1 + \xi_1 \xi_2 \bar{\xi}_2 \bar{\xi}_1 = 2\bar{\xi}_1 \bar{\xi}_2 \xi_2 \xi_1 \neq 0.$$

If $|X| = 1, X = \{x\}$, then the equality $S^{(+)}\text{tri-Jord}\langle x \rangle \subseteq \text{tri-H}\langle x \rangle$ may be derived from Theorem 2.8 and the Cohn Theorem for ordinary algebras. The involution σ of $\text{tri-As}\langle X \rangle$ may be extended to $\widetilde{\text{tri-As}}\langle X \rangle$ and $C_2 \otimes \widetilde{\text{tri-As}}\langle X \rangle$ in the natural ways. Note that $\widetilde{\text{tri-As}}\langle x \rangle$ is a homomorphic image of $\text{As}\langle \bar{x}, x \rangle$, and so $C_2 \otimes \text{As}\langle \bar{x}, x \rangle$ maps onto $\text{tri-As}\langle x \rangle \subseteq C_2 \otimes \widetilde{\text{tri-As}}\langle x \rangle$. If $\sigma(f) = f$ for $f \in \text{tri-As}\langle x \rangle$ then f has a preimage in $C_2 \otimes H\langle \bar{x}, x \rangle$. The latter coincides with $C_2 \otimes SJ\langle \bar{x}, x \rangle$ and thus f belongs to $S^{(+)}\text{tri-Jord}\langle x \rangle$. \square

5. Problems on splitted algebras

Less is known about relations between operads of pre- and post-algebras that are (in quadratic binary case) Koszul dual to di- and tri-algebras, respectively. Apart from already considered relations with Rota–Baxter operators, we may prove analogues of some results from the previous section.

5.1 Splitting morphisms of operads. Let us show how a morphism of operads $\omega : \mathfrak{R} \rightarrow \mathfrak{M}$ induces a functor on the corresponding varieties of pre- and post-algebras. We will consider the case of post-algebras since all constructions for pre-algebras may be obtained by restriction.

As above, assume Σ and Σ' are the languages of \mathfrak{M} and \mathfrak{R} , respectively.

Let $A \in \text{post-}\mathfrak{M}$. Define a structure of a $\Sigma'^{(3)}$ -algebra on the space A as follows. Given $f \in \Sigma'$, $\nu(f) = n$, $H \in \mathcal{P}(n)$, we have to define $f^H(a_1, \dots, a_n)$, $a_i \in A$. Consider $(\text{ComTrias}\langle y_1, y_2, \dots \rangle \boxtimes A)^{(\omega)} \in \mathfrak{R}$, and evaluate

$$f(y_1 \otimes a_1, \dots, y_n \otimes a_n) = \sum_{H \in \mathcal{P}(n)} e_H^{(n)}(y_1, \dots, y_n) \otimes b_H.$$

Here $b_H \in A$ are uniquely defined. Finally, set

$$f^H(a_1, \dots, a_n) = b_H.$$

Denote the Σ' -algebra obtained by $A^{(\text{post-}\omega)}$.

In a similar way $(|H| = 1)$, $A^{(\text{pre-}\omega)} \in \text{pre-}\mathfrak{R}$ may be defined for $A \in \text{pre-}\mathfrak{M}$.

Proposition 5.1. *If $\omega : \mathfrak{R} \rightarrow \mathfrak{M}$ is a morphism of operads and $A \in \text{post-}\mathfrak{M}$ then $A^{(\text{post-}\omega)} \in \text{post-}\mathfrak{R}$.*

PROOF: Immediately follows from the definition since

$$(\text{ComTrias}\langle y_1, y_2, \dots \rangle \boxtimes A)^{(\omega)} = \text{ComTrias}\langle y_1, y_2, \dots \rangle \boxtimes A^{(\text{post-}\omega)}.$$

□

Example 5.2. Consider the following morphism from the operad of Lie triple systems (LTS) to the operad of Lie algebras:

$$\begin{aligned} \omega : LTS &\rightarrow \text{Lie}, \\ [x_1, x_2, x_3] &\mapsto [[x_1, x_2], x_3] \end{aligned}$$

Then for every $L \in \text{pre-Lie}$ the following operations define $L^{(\text{pre-}\omega)} \in \text{pre-LTS}$:

$$\begin{aligned} [x_1, x_2, x_3]_1 &= x_3(x_2x_1), & [x_1, x_2, x_3]_2 &= -[x_2, x_1, x_3]_1 = -x_3(x_1x_2), \\ [x_1, x_2, x_3]_3 &= (x_1x_2)x_3 - (x_2x_1)x_3. \end{aligned}$$

Indeed, consider $P = \text{Perm}\langle y_1, y_2, y_3 \rangle$, and evaluate

$$[y_1 \otimes x_1, y_2 \otimes x_2, y_3 \otimes x_3] = [[y_1 \otimes x_1, y_2 \otimes x_2], y_3 \otimes x_3]$$

in $P \boxtimes \text{pre-Lie}\langle x_1, x_2, x_3 \rangle$ assuming $ab = [a, b]_2 = -[b, a]_1$ in pre-Lie:

$$\begin{aligned} [[y_1 \otimes x_1, y_2 \otimes x_2], y_3 \otimes x_3] &= [y_1 y_2 \otimes x_1 x_2 - y_2 y_1 \otimes x_2 x_1, y_3 \otimes x_3] \\ &= y_1 y_2 y_3 \otimes (x_1 x_2) x_3 - y_3 y_1 y_2 \otimes x_3 (x_1 x_2) - y_2 y_1 y_3 \otimes (x_2 x_1) x_3 + y_3 y_2 y_1 \otimes x_3 (x_2 x_1) \end{aligned}$$

It remains to collect similar terms to get the desired expressions.

5.2 On the special identities for pre- and post-algebras. It remains unclear how to solve in general the analogues of PBW-type problems for pre- and post-algebras. For special identities, however, we may state a partial result and show by example that an analogue of Theorem 4.7 does not hold.

Given a morphism of operads $\omega : \mathfrak{N} \rightarrow \mathfrak{M}$, one may consider the induced morphisms pre- ω , post- ω , and define varieties $S^{(\text{pre-}\omega)} \text{pre-}\mathfrak{N}$ and $S^{(\text{post-}\omega)} \text{post-}\mathfrak{N}$ generated by all special algebras in pre- \mathfrak{N} and post- \mathfrak{N} , respectively.

Proposition 5.3. *Over a field of zero characteristic, we have the following relations: $S^{(\text{pre-}\omega)} \text{pre-}\mathfrak{N} \subseteq \text{pre-}S^{(\omega)} \mathfrak{N}$, $S^{(\text{post-}\omega)} \text{post-}\mathfrak{N} \subseteq \text{post-}S^{(\omega)} \mathfrak{N}$.*

PROOF: Let us consider the case of post-algebras. It is enough to show that every special algebra in post- \mathfrak{N} belongs to post- $S^{(\omega)} \mathfrak{N}$. Suppose $T \in S^{(\text{post-}\omega)} \text{post-}\mathfrak{N}$, $T \subseteq A^{(\text{post-}\omega)}$. Fix $C = \text{ComTrias}\langle y_1, y_2, \dots \rangle$ and note that $C \boxtimes T \subseteq C \boxtimes A^{(\text{post-}\omega)} = (C \boxtimes A)^{(\omega)}$, i.e., $C \boxtimes T \in S^{(\omega)} \mathfrak{N}$. By definition, $T \in \text{post-}S^{(\omega)} \mathfrak{N}$. \square

Let us state an example to show that the converse embedding may not hold. Although the language in the example below contains unary operation, it is a derivation with respect to the binary product. Hence, Theorem 4.7 for di- or tri-algebras would remain valid in these settings (see Remark 1). Thus, the example stated below shows an essential difference between di-, tri-algebras and pre-, post-algebras.

Example 5.4. Let $\mathfrak{N} = \text{Perm}$, and let \mathfrak{M} govern the variety of associative commutative algebras with a derivation (unary operation) ∂ such that $\partial^2 = 0$. Consider $\omega : \mathfrak{N} \rightarrow \mathfrak{M}$ given by $x_1 x_2 \mapsto \partial(x_1) x_2$.

Here $\Sigma' = \{\cdot\}$, one binary operation; $\Sigma = \{\cdot, \partial\}$.

It is well-known that ω determines a functor from the variety \mathfrak{M} to $\mathfrak{N} = \text{Perm}$ [21]. Moreover, every algebra of the form $A^{(\omega)}$, $A \in \mathfrak{M}$, is 3-nilpotent. Since there are no identities of smaller degree, the variety $S^{(\omega)} \mathfrak{N}$ coincides with N_3 , the variety of algebras satisfying $x(yz) = (xy)z = 0$.

It is straightforward to find the defining identities of pre- N_3 :

$$(18) \quad \begin{aligned} (x < y) < z = 0, \quad (x > y) < z = 0, \quad (x < y + x > y) > z = 0, \\ x < (y < z + y > z) = 0, \quad x > (y < z) = 0, \quad x > (y > z) = 0. \end{aligned}$$

Here $\Sigma'^{(3)} = \{>, <\}$, two binary operations.

On the other hand, pre- \mathfrak{M} consists of Perm-algebras equipped with a derivation ∂ such that $\partial^2 = 0$. If $A \in \text{pre-}\mathfrak{M}$ then the operations on $A^{(\text{pre-}\omega)} \in \text{pre-}\mathfrak{N}$ are given by

$$a > b = \partial(a)b, \quad a < b = b\partial(a).$$

Note that $a \succ b + b \prec a = \partial(a)b + a\partial(b) = \partial(ab)$, and $\partial(ab) \succ c = 0$ for all $a, b, c \in A$. Hence, every algebra in $S^{(\text{pre-}\omega)}\text{pre-}\mathfrak{R}$ satisfies an identity

$$(x \succ y + y \prec x) \succ z = 0$$

which does not follow from (18).

Remark 5. It is interesting to mention two similar classes of “replicated” and “splitted” operads: totally compatible and linearly compatible ones [11], [26]. These operads may also be obtained as white and black Manin products with operads denoted by ${}^2\text{Com}$ and Lie^2 , respectively. All problems mentioned in Section 4 make sense for these classes.

Acknowledgments. The authors are very grateful to Bruno Vallette and to the anonymous referee for valuable remarks and comments. The second author is pleased to acknowledge hospitality of the University of Denver during the Third Mile High Conference on Nonassociative Mathematics. This work was partially supported by RFBR 12-01-00329 and 12-01-33031.

REFERENCES

- [1] Aguiar M., *Pre-Poisson algebras*, Lett. Math. Phys. **54** (2000), 263–277.
- [2] Bai C., Bellier O., Guo L., Ni X., *Splitting of operations, Manin products, and Rota–Baxter operators*, Int. Math. Res. Notes **3** (2013), 485–524.
- [3] Bakalov B., D’Andrea A., Kac V.G., *Theory of finite pseudoalgebras*, Adv. Math. **162** (2001), no. 1, 1–140.
- [4] Bremner M.R., *On the definition of quasi-Jordan algebra*, Comm. Algebra **38** (2010), 4695–4704.
- [5] Bremner M.R., Felipe R., Sánchez-Ortega J., *Jordan triple disystems*, Comput. Math. Appl. **63** (2012), 1039–1055.
- [6] Bremner M.R., Madariaga S., *Dendriform analogues of Lie and Jordan triple systems*, Comm. Algebra, to appear, arXiv:1305.1389 [math.RA].
- [7] Bremner M.R., Peresi L.A., Sánchez-Ortega J., *Malcev dialgebras*, Linear Multilinear Algebra **60** (2012), 1125–1141.
- [8] Bremner M.R., Sánchez-Ortega J., *Leibniz triple systems*, Commun. Contemp. Math. **16** (2014), 1350051 (19 pages); DOI: 10.1142/S021919971350051X.
- [9] Chapoton F., *Un endofoncteur de la catégorie des opérades*, in: Loday J.-L., Frabetti A., Chapoton F., Goichot F. (Eds), Dialgebras and related operads, Lectures Notes in Mathematics, 1763, Springer, Berlin, 2001, pp. 105–110.
- [10] Cohn P.M., *On homomorphic images of special Jordan algebras*, Canadian J. Math. **6** (1954), 253–264.
- [11] Dotsenko V., Khoroshkin A., *Character formulas for the operad of two compatible brackets and for the bihamiltonian operad*, Funktsional. Anal. i Prilozhen. **41** (2007), no. 1, 1–17.
- [12] Giambruno A., Zaicev M., *On codimension growth of finitely generated associative algebras*, Adv. Math. **140** (1998), no. 2, 145–155.
- [13] Ginzburg V., Kapranov M., *Koszul duality for operads*, Duke Math. J. **76** (1994), no. 1, 203–272.
- [14] Gubarev V.Yu., Kolesnikov P.S., *The Tits–Kantor–Koecher construction for Jordan dialgebras*, Comm. Algebra **39** (2011), no. 2, 497–520.
- [15] Gubarev V.Yu., Kolesnikov P.S., *Embedding of dendriform algebras into Rota–Baxter algebras*, Cent. Eur. J. Math. **11** (2013), no. 2, 226–245.
- [16] Hou D.P., Ni X., Bai C., *Pre-Jordan algebras*, Math. Scand. **112** (2013), no. 1, 19–48.
- [17] Jacobson N., *Structure and Representations of Jordan Algebras*, American Mathematical Society, Providence, 1968.
- [18] Kolesnikov P.S., *Varieties of dialgebras and conformal algebras*, Sib. Math. J. **49** (2008), no. 2, 257–272.

- [19] Kolesnikov P.S., Voronin V.Yu., *On special identities for dialgebras*, Linear Multilinear Algebra **61** (2013), no. 3, 377–391.
- [20] Loday J.-L., *Une version non commutative des algèbres de Lie: les algèbres de Leibniz*, Enseign. Math. **39** (1993), 269–293.
- [21] Loday J.-L., *Dialgebras*, in: Loday J.-L., Frabetti A., Chapoton F., Goichot F. (Eds.), *Dialgebras and related operads*, Lectures Notes in Mathematics, 1763, Springer, Berlin, 2001, pp. 105–110.
- [22] Loday J.-L., Pirashvili T., *Universal enveloping algebras of Leibniz algebras and (co)homology*, Math. Ann. **296** (1993), 139–158.
- [23] Loday J.-L., Ronco M., *Trialgebras and families of polytopes*, in: Goerss P.G., Priddy S. (Eds.), *Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-theory*, Contemp. Math. **346** (2004), 369–398.
- [24] Loday J.-L., Vallette B., *Algebraic Operads*, Grundlehren der mathematischen Wissenschaften, 346, Springer, Heidelberg, 2012.
- [25] Pei J., Bai C., Guo L., *Splitting of operads and Rota–Baxter operators on operads*, arXiv:1306.3046 [math.CT].
- [26] Strohmayr H., *Operads of compatible structures and weighted partitions*, J. Pure Appl. Algebra **212** (2008), 2522–2534.
- [27] Uchino K., *On distributive laws in derived bracket construction and homotopy theory of derived bracket Leibniz algebras*, arXiv:1110.4188v5 [math.QA].
- [28] Vallette B., *Homology of generalized partition posets*, J. Pure Appl. Algebra **208** (2007), no. 2, 699–725.
- [29] Vallette B., *Manin products, Koszul duality, Loday algebras and Deligne conjecture*, J. Reine Angew. Math. **620** (2008), 105–164.
- [30] Velasquez R., Felipe R., *Quasi-Jordan algebras*, Comm. Algebra **36** (2008), no. 4, 1580–1602.
- [31] Voronin V., *Special and exceptional Jordan dialgebras*, J. Algebra Appl. **11** (2012), no. 2, 23 p.
- [32] Zhevlakov K.A., Slin'ko A.M., Shestakov I.P., Shirshov A.I., *Rings That Are Nearly Associative*, Academic Press, New York, 1982.

SOBOLEV INSTITUTE OF MATHEMATICS, AKAD. KOPTYUG PROSP. 4, 630090 NOVOSIBIRSK, RUSSIA

E-mail: pavelsk@math.nsc.ru
vsevolodgu@mail.ru

(Received October 28, 2013)