

Andrea Caggegi

Some Additive $2 - (v, 5, \lambda)$ Designs

Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica, Vol. 54 (2015),
No. 1, 65–80

Persistent URL: <http://dml.cz/dmlcz/144368>

Terms of use:

© Palacký University Olomouc, Faculty of Science, 2015

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Some Additive $2 - (v, 5, \lambda)$ Designs

Andrea CAGGEGI

DEIM, Viale delle Scienze Ed. 8, I-90128 Palermo, Italy
e-mail: andrea.caggegi@unipa.it

(Received March 17, 2014)

Abstract

Given a finite additive abelian group G and an integer k , with $3 \leq k \leq |G|$, denote by $\mathcal{D}_k(G)$ the simple incidence structure whose point-set is G and whose blocks are the k -subsets $C = \{c_1, c_2, \dots, c_k\}$ of G such that $c_1 + c_2 + \dots + c_k = 0$. It is known (see [2]) that $\mathcal{D}_k(G)$ is a 2-design, if G is an elementary abelian p -group with p a prime divisor of k . From [3] we know that $\mathcal{D}_3(G)$ is a 2-design if and only if G is an elementary abelian 3-group. It is also known (see [4]) that G is necessarily an elementary abelian 2-group, if $\mathcal{D}_4(G)$ is a 2-design. Here we shall prove that $\mathcal{D}_5(G)$ is a 2-design if and only if G is an elementary abelian 5-group.

Key words: Conformal mapping, geodesic mapping, conformal-geodesic mapping, initial conditions, (pseudo-) Riemannian space.

2010 Mathematics Subject Classification: 53B20, 53B30, 53C21

1 Introduction and preliminary results

Let v, k, t, λ be positive integers with $v > k > t$. By a t -design with parameters v, k, λ (or shortly: a $t - (v, k, \lambda)$ design) one understands a pair $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ where \mathcal{P} is a finite set with v elements (called points) and \mathcal{B} is a set of subsets of \mathcal{P} called blocks such that each block contains k points and any t distinct points are contained in exactly λ common blocks (cf. [1], [5]). We say that a $t - (v, k, \lambda)$ design $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ is an additive design, if there are a finite abelian group G , written additively, and an injective mapping $\chi: \mathcal{P} \rightarrow G$ with the property that $\chi(c_1) + \chi(c_2) + \dots + \chi(c_k) = 0$ whenever $C = \{c_1, c_2, \dots, c_k\} \in \mathcal{B}$ is a block of $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ (cf. [2]). For every finite additive abelian group G and for any integer $k \in \{3, 4, \dots, |G| - 1\}$ we denote by $\mathcal{D}_k(G)$ the simple incidence structure the point-set of which is G and the blocks of which are the k -subsets $C = \{c_1, c_2, \dots, c_k\}$ of G such that $c_1 + c_2 + \dots + c_k = 0$. Note that each 2-design of the form $\mathcal{D}_k(G)$ is an additive 2-design.

Throughout this paper we shall be concerned only with finite abelian groups, written additively. If G is such a group, the notation that follows will remain fixed: $|G|$ is the order of G ; $\langle a \rangle$ is the subgroup of G generated by $a \in G$; if m is a positive integer, mG and G_m are the subgroups of G given by $mG = \{mg \mid g \in G\}$ and $G_m = \{g \in G \mid mg = 0\}$; if $|G| > 4$ and if x, y are distinct elements of G , $N_{x,y}$ denotes the number of pairs $\{c, C\}$ where $c \in G \setminus \{x, y\}$ and C is a block of $\mathcal{D}_5(G)$ through $\{x, y, c\}$.

We state now some preliminary results.

Lemma 1 *If $\mathcal{D}_5(G)$ is a $2 - (|G|, 5, \lambda)$ design for some λ , then $N_{x,y}$ is a constant (equal to 3λ).*

Proof Suppose $\mathcal{D}_5(G)$ is a $2 - (|G|, 5, \lambda)$ design for some λ . Then there are λ blocks of $\mathcal{D}_5(G)$ through any given two distinct elements $x, y \in G$; on the other hand, each block of $\mathcal{D}_5(G)$ through $\{x, y\}$ contains exactly 3 points distinct from x, y . Therefore $N_{x,y} = 3\lambda$ and the Lemma 1 is proved. \square

Proposition 1 *$\mathcal{D}_5(G)$ is not a 2-design if one of the statements below is true:*

- 1) G is an elementary abelian 2-group;
- 2) G is direct sum of cyclic groups of order 4;
- 3) G is direct sum of groups of order 2 and cyclic groups of order 4;
- 4) G contains just one involution and $2G$ is an elementary abelian 3-group.

Proof We may assume that G has order greater than 4.

1) Suppose G is an elementary abelian 2-group of order $n = 2^\nu \geq 8$. Let $g \in G$, $g \neq 0 \in G$ and let $x \in G \setminus \{0, g\}$. We show that $N_{0,g} \neq N_{x,g}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design. There are no blocks of $\mathcal{D}_5(G)$ through $\{0, g, x, g+x\}$, however $\{0, g, x\}$ may be extended to a block $\{0, g, x, y, g+x+y\}$ for every $y \in G \setminus \{0, g, x, g+x\}$. Therefore

$$N_{0,g} = (n-2) \frac{n-4}{2}.$$

There are no blocks of $\mathcal{D}_5(G)$ through $\{g, x, g+x\}$, however there are $\frac{n-4}{2}$ blocks through $\{0, g, x\}$ and $\frac{n-6}{2}$ blocks through $\{x, g, z\}$ for any given $z \in G \setminus \{0, g, x, g+x\}$. Therefore

$$N_{x,g} = \frac{n-4}{2} + (n-4) \frac{n-6}{2}.$$

From $n \neq 4$ it follows $N_{0,g} \neq N_{x,g}$ and hence $\mathcal{D}_5(G)$ is not a 2-design.

2) Suppose G is direct sum of $\nu \geq 2$ cyclic groups of order 4. So G is a finite abelian group of order $n = 4^\nu \geq 16$ and $2G = G_2$ is an elementary abelian 2-group of order $2^\nu \geq 4$.

Let $a \in G_2$, $a \neq 0$ and let $b \in G_4 \setminus G_2$. We show that $N_{0,a} \neq N_{0,b}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design.

If $x \in G_2 \setminus \langle a \rangle$, there are no blocks of $\mathcal{D}_5(G)$ through $\{0, a, x, a + x\}$; if $y \in G \setminus G_2$ with $2y \neq a$, any block of $\mathcal{D}_5(G)$ through $\{0, a, y\}$ does not intersect $\{a - y, -y, a + 2y\}$. These facts imply:

if $g \in G$ with $2g = a$, then ($g \in G \setminus G_2$ and) there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$;

if $g \in G \setminus G_2$ with $2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$;

if $g \in G_2 \setminus \langle a \rangle$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$.

Therefore

$$N_{0,a} = |G_2| \cdot \frac{n-4}{2} + (n-2|G_2|) \cdot \frac{n-6}{2} + (|G_2|-2) \cdot \frac{n-4-|G_2|}{2}$$

can be written as

$$N_{0,a} = 3|G_2| - \frac{1}{2}|G_2|^2 + \frac{n^2 - 8n + 8}{2}. \quad (1.1)$$

There are no blocks of $\mathcal{D}_5(G)$ containing the group $\langle b \rangle = \{0, b, 2b, -b\}$;

if $g \in b + G_2$ with $b \neq g \neq -b$, there are no blocks of $\mathcal{D}_5(G)$ through $\{0, b, g, -b - g\}$;

if $2b \neq g \in G \setminus b + G_2$, any block of $\mathcal{D}_5(G)$ through $\{0, b, g\}$ does not meet $\{3b - g, 2b - g, 2g - b\}$.

These facts guarantee that:

$\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{0, b, -b\}$;

there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, b, 2b\}$;

if $g \in b + G_2$ with $b \neq g \neq -b$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$;

if $g \in G$ with $g \neq 2b \neq 2g$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$.

Therefore

$$N_{0,b} = \frac{n-2-|G_2|}{2} + \frac{n-4}{2} + (|G_2|-2) \cdot \frac{n-4-|G_2|}{2} + (n-|G_2|-2) \cdot \frac{n-6}{2}$$

can be written as

$$N_{0,b} = \frac{3}{2} \cdot |G_2| - \frac{1}{2} \cdot |G_2|^2 + \frac{n^2 - 8n + 14}{2} \quad (1.2)$$

Since $|G_2| \neq 2$, (1.1) and (1.2) yield $N_{0,a} \neq N_{0,b}$ and hence $\mathcal{D}_5(G)$ is not a 2-design.

3) Suppose G is direct sum of $h \geq 1$ groups of order 2 and $\nu \geq 1$ cyclic groups of order 4. So G is a finite abelian group of order $n = |G| = 2^h \cdot 4^\nu \geq 8$; $2G$ is an elementary abelian 2-group of order 2^ν ; G_2 is an elementary abelian 2-group of order $2^{h+\nu} \geq 4$ which admits $2G$ as a proper subgroup.

Let $a \in G_2 \setminus 2G$ and let $b \in 2G$, $b \neq 0$. We show now that $N_{0,a} \neq N_{0,b}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design.

If $a \neq g \in a + 2G$, then $a + g \in 2G$ and there are no blocks of $\mathcal{D}_5(G)$ through $\{0, a, g, a + g\}$;

if $0 \neq g \in G_2 \setminus a + 2G$, then $a + g \notin 2G$ and there are no blocks of $\mathcal{D}_5(G)$ through $\{0, a, g, a + g\}$;

if $g \in G \setminus G_2$, then $a + g \notin 2G$ and any block of $\mathcal{D}_5(G)$ through $\{0, a, g\}$ does not intersect $\{a - g, -g, a + 2g\}$.

From these facts we deduce that:

if $a \neq g \in a + 2G$, then $\frac{n-4-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$;

if $g \in G_2$ with $0 \neq g \notin a + 2G$, there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$;

if $g \in G \setminus G_2$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$.

Therefore

$$N_{0,a} = (|2G| - 1) \cdot \frac{n-4-|G_2|}{2} + (|G_2| - |2G| - 1) \cdot \frac{n-4}{2} + (n - |G_2|) \cdot \frac{n-6}{2}$$

which, since $|2G| \cdot |G_2| = |G| = n$, simplifies to

$$N_{0,a} = \frac{3}{2} \cdot |G_2| + \frac{n^2 - 9n + 8}{2}. \quad (1.3)$$

If $b = 2g$ with $g \in G$, then $b + g \notin 2G$ and there are no blocks of $\mathcal{D}_5(G)$ through $\{0, b, g, -g\}$;

if $g \in 2G \setminus \{0, b\}$, then $b + g \in 2G$ and there are no blocks of $\mathcal{D}_5(G)$ through $\{0, b, g, b + g\}$;

if $g \in G_2 \setminus 2G$, then $b + g \notin 2G$ and there are no blocks of $\mathcal{D}_5(G)$ through $\{0, b, g, b + g\}$;

if $g \in G \setminus G_2$ and $2g \neq b$, then $b + g \notin 2G$ and any block of $\mathcal{D}_5(G)$ through $\{0, b, g\}$ does not meet $\{b - g, -g, b + 2g\}$.

These facts enable us to conclude that:

if $g \in G$ has the property that $2g = b$, there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$;

if $g \in 2G \setminus \{0, b\}$, then $\frac{n-4-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$;

if $g \in G_2 \setminus 2G$, there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$;

if $g \in G \setminus G_2$ with $2g \neq b$, then $\frac{n-6}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$.

Therefore

$$\begin{aligned} N_{0,b} &= \\ &= |G_2| \cdot \frac{n-4}{2} + (|2G| - 2) \cdot \frac{n-4-|G_2|}{2} + (|G_2| - |2G|) \cdot \frac{n-4}{2} + (n - 2|G_2|) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, can be rewritten as

$$N_{0,b} = 3|G_2| + \frac{n^2 - 9n + 8}{2}. \quad (1.4)$$

Since $|G_2| \neq 0$, (1.3) and (1.4) give $N_{0,a} \neq N_{0,b}$ and hence $\mathcal{D}_5(G)$ is not a 2-design.

4) In this case $G_2 = \{0, a\}$ is a group of order two and a is the unique involution of G ; G can be written as direct sum $G = G_2 \oplus 2G$ and $2G = G_3$ is an elementary abelian 3-group. If $2G = G_3$ has order 3, then G is cyclic of order 6 and clearly $\mathcal{D}_5(G)$ is not a 2-design. Thus we may assume that $|2G| = 3^m$ for some integer $m > 1$. Then G has order $n = |G| = 2|2G| \geq 18$ and we have:

if $a \neq g \in G \setminus 2G$ and $x \in \{2g, -g\}$, there are no blocks of $\mathcal{D}_5(G)$ through $\{0, a, g, x\}$;

if $0 \neq g \in 2G$, any block of $\mathcal{D}_5(G)$ through $\{0, a, g\}$ does not intersect $\{a - g, -g, a - 2g\}$.

These facts imply:

if $g \in G \setminus 2G$ with $g \neq a$, there are $\frac{n-4-|G_2|}{2} = \frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$;

if $g \in 2G$ is not equal to $0 \in G$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$.

Therefore

$$N_{0,a} = (|2G| - 1) \cdot \frac{n-6}{2} + (|2G| - 1) \cdot \frac{n-6}{2}. \quad (1.5)$$

Let $b \in G_3$, $b \neq 0$. Clearly ($b \neq a$ and) we have:

there are no blocks of $\mathcal{D}_5(G)$ containing $\{0, b, -b\}$;

if $g \in G \setminus 2G$, any block of $\mathcal{D}_5(G)$ through $\{0, b, g\}$ does not intersect $\{2b - g, b - g, 2b - 2g\}$;

if $g \in 2G \setminus \langle b \rangle$, then $b + g \in 2G$ and any block of $\mathcal{D}_5(G)$ through $\{0, b, g\}$ does not intersect $\{2b - g, b - g, 2b - 2g\}$.

These facts imply:

if $g \in G \setminus 2G$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$;

if $g \in 2G \setminus \langle b \rangle$, there are $\frac{n-6-|G_2|}{2} = \frac{n-8}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, b, g\}$.

Therefore

$$N_{0,b} = (n - |2G|) \cdot \frac{n-6}{2} + (|2G| - 3) \cdot \frac{n-8}{2}$$

which, since $2|2G| = |G| = n$, simplifies to

$$N_{0,b} = |2G| \cdot (n - 6) + 12 - 2n. \quad (1.6)$$

Since $n \neq 6$, (1.5) and (1.6) yield $N_{0,a} \neq N_{0,b}$ and hence $\mathcal{D}_5(G)$ is not a 2-design. This last result completes the proof. \square

Lemma 2 *Let G be a finite additive abelian group of even order $n > 4$. If there is $a \in G$ such that $a \notin 2G$ and $2a \neq 0$, then*

$$N_{a,-a} = |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Proof We first note that there are $\frac{n-2-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 0\}$. We now discuss five cases.

Case (L. 1. 1): $4a = 0$ and $|G_3| = 1$. In this case we have:

$\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 2a\}$;
 if $g \in 2G$ with $0 \neq g \neq 2a$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in G - 2G$ with $a \neq g \neq -a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$N_{a,-a} = 2 \cdot \frac{n-2-|G_2|}{2} + (|2G|-2) \cdot \frac{n-6-|G_2|}{2} + (n-|2G|-2) \cdot \frac{n-6}{2}$$

which, since $|2G| \cdot |G_2| = |G| = n$ and $|G_3| = 1$, can be written as

$$N_{a,-a} = |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 1. 2): $4a = 0$ and $|G_3| \neq 1$. In this case we get:

$\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 2a\}$;
 if $g \in G_3$ is distinct from 0, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in 2G \setminus G_3$ with $g \neq 2a$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in G \setminus 2G$ with $a \neq g \neq -a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= 2 \cdot \frac{n-2-|G_2|}{2} + (|G_3|-1) \cdot \frac{n-4-|G_2|}{2} \\ &+ (|2G|-|G_3|-1) \cdot \frac{n-6-|G_2|}{2} + (n-|2G|-2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, simplifies to

$$N_{a,-a} = |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 1. 3): a has order 6. In this case we have:

if $g \in \{-2a, 2a\}$, then $\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in G_3 \setminus \{0, -2a, 2a\}$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in 2G \setminus G_3$, then $\frac{n-6-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ containing $\{a, -a, g\}$;
 if $g \in G \setminus 2G$ with $a \neq g \neq -a$, there are $\frac{n-6}{2}$ blocks $\mathcal{D}_5(G)$ including $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= 3 \cdot \frac{n-2-|G_2|}{2} + (|G_3|-3) \cdot \frac{n-4-|G_2|}{2} \\ &+ (|2G|-|G_3|) \cdot \frac{n-6-|G_2|}{2} + (n-|2G|-2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, gives

$$N_{a,-a} = |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 1. 4): $4a \neq 0 \neq 6a$ and $|G_3| = 1$. In this case we get:
 if $g \in \{-2a, 2a\}$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in 2G \setminus \{0, -2a, 2a\}$, $\frac{n-6-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ including $\{a, -a, g\}$;
 if $g \in G \setminus 2G$ with $a \neq g \neq -a$, $\frac{n-6}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= \frac{n-2-|G_2|}{2} + 2 \cdot \frac{n-4-|G_2|}{2} \\ &+ (|2G| - 3) \cdot \frac{n-6-|G_2|}{2} + (n - |2G| - 2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$ and $|G_3| = 1$, can be rewritten as

$$N_{a,-a} = |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 1. 5): $4a \neq 0 \neq 6a$ and $|G_3| \neq 1$. In this case we obtain:
 there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$ if $g \in \{-2a, 2a\}$ or $0 \neq g \in G_3$;
 if $g \in 2G \setminus G_3$ with $2a \neq g \neq -2a$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in G \setminus 2G$ with $a \neq g \neq -a$, $\frac{n-6}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= \frac{n-2-|G_2|}{2} + (|G_3| + 1) \cdot \frac{n-4-|G_2|}{2} \\ &+ (|2G| - |G_3| - 2) \cdot \frac{n-6-|G_2|}{2} + (n - |2G| - 2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, simplifies to

$$N_{a,-a} = |G_3| + \frac{n^2 - 9n + 18}{2}.$$

The Lemma 2 is proved. □

Proposition 2 $\mathcal{D}_5(G)$ is not a 2-design if G is a finite abelian group of even order $n > 4$ with the property that $2G = 4G$.

Proof From $2G = 4G$ it follows $G_2 = G_4$ and this requires that the Sylow 2-subgroup of G is an elementary abelian 2-group. Therefore G can be written as direct sum $G = G_2 \oplus 2G$ and, by Proposition 1, we may assume that $2G$ is a finite abelian group of odd order $|2G| > 1$. Then any $z \in G$ of the form $z = x + y$, with $x \in G_2$ and $y \in 2G$ both distinct from 0, is not equal to $-z$ and does not belong to $2G$. Thus, using Lemma 2 we see that

$$N_{z,-z} = |G_3| + \frac{n^2 - 9n + 18}{2}. \quad (1.7)$$

Choose $a \in 2G$, $a \neq 0$ and let α be the unique element in $2G$ such that $a = 2\alpha$. We shall prove that $N_{a,-a} \neq N_{z,-z}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design. We first note that $\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 0\}$. We now discuss five cases.

Case (P. 2. 1): $|G_3| = 1$ and $5a \neq 0$. In this case we have:

if $g \in \{-2a, 2a, -\alpha, \alpha\}$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $-\alpha \neq g \in G$ with $2g = -a$, then ($g \in -\alpha + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $\alpha \neq g \in G$ with $2g = a$, then ($g \in \alpha + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in 2G \setminus \{a, -a, 0, \alpha, -\alpha, 2a, -2a\}$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= \frac{n-2-|G_2|}{2} + 4 \cdot \frac{n-4-|G_2|}{2} + 2(|G_2| - 1) \cdot \frac{n-4}{2} \\ &+ (|2G| - 7) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G| + 2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$ and $|G_3| = 1$, can be rewritten as

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}$$

Because $|G_2| \neq 0$, this equality together with (1.7) gives $N_{a,-a} \neq N_{z,-z}$ and hence $\mathcal{D}_5(G)$ is not a 2-design.

Case (P. 2. 2): $|G_3| = 1$ and a has order 5. In this case we have ($\alpha = -2a$ and):

if $g \in \{2a, -2a\}$, there are $\frac{n-2-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $2a \neq g \in G$ with $2g = -a$, then ($g \in 2a + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $-2a \neq g \in G$ with $2g = a$, then ($g \in -2a + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in 2G \setminus \langle a \rangle$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= 3 \cdot \frac{n-2-|G_2|}{2} + 2 \cdot (|G_2| - 1) \frac{n-4}{2} \\ &+ (|2G| - 5) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G| + 2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$ and $|G_3| = 1$, simplifies to

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}$$

Since $|G_2| \neq 0$, this equality together with (1.7) gives $N_{a,-a} \neq N_{z,-z}$ and hence $\mathcal{D}_5(G)$ is not a 2-design.

Case (P. 2. 3): $|G_3| \neq 1$ and a has order 5. In this case we have ($\alpha = -2a$ and):

if $g \in \{2a, -2a\}$, there are $\frac{n-2-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $2a \neq g \in G$ with $2g = -a$, then ($g \in 2a + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $-2a \neq g \in G$ with $2g = a$, then ($g \in -2a + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $0 \neq g \in G_3$, then $\frac{n-4-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in 2G \setminus G_3$ with $2a \neq g \neq -2a$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= 3 \cdot \frac{n-2-|G_2|}{2} + 2 \cdot (|G_2| - 1) \cdot \frac{n-4}{2} + (|G_3| - 1) \cdot \frac{n-4-|G_2|}{2} \\ &+ (|2G| - |G_3| - 2) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G|) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, simplifies to

$$N_{a,-a} = 2|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}$$

Since $|G_2| \neq 0$, this equality together with (1.7) gives $N_{a,-a} \neq N_{z,-z}$ and hence $\mathcal{D}_5(G)$ is not a 2-design.

Case (P. 2. 4): $|G_3| \neq 1$ and $3a \neq 0 \neq 5a$. In this case we have:

if $g \in \{2a, -2a, \alpha, -\alpha\}$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $-\alpha \neq g \in G$ with $2g = -a$, then ($g \in -\alpha + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $\alpha \neq g \in G$ with $2g = a$, then ($g \in \alpha + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $0 \neq g \in G_3$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in 2G \setminus G_3$ and $g \notin \{a, -a, \alpha, -\alpha, 2a, -2a\}$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= \frac{n-2-|G_2|}{2} + 4 \cdot \frac{n-4-|G_2|}{2} + 2 \cdot (|G_2| - 1) \cdot \frac{n-4}{2} \\ &+ (|G_3| - 1) \cdot \frac{n-4-|G_2|}{2} + (|2G| - |G_3| - 6) \cdot \frac{n-6-|G_2|}{2} \\ &+ (n-2 \cdot |G_2| - |2G| + 2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, can be rewritten as

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}$$

Since $|G_2| \neq 0$, this equality together with (1.7) gives $N_{a,-a} \neq N_{z,-z}$ and hence $\mathcal{D}_5(G)$ is not a 2-design.

Case (P. 2. 5): $a \in G_3$. In this case we obtain ($\alpha = -a$ and):

if $a \neq g \in G$ with $2g = -a$, then ($g \in a + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $-a \neq g \in G$ with $2g = a$, then ($g \in -a + G_2$ hence) $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in G_3 \setminus \langle a \rangle$, then $\frac{n-4-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in 2G \setminus G_3$, then $\frac{n-6-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
 if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= \frac{n-2-|G_2|}{2} + 2(|G_2| - 1) \cdot \frac{n-4}{2} + (|G_3| - 3) \cdot \frac{n-4-|G_2|}{2} \\ &+ (|2G| - |G_3|) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G| + 2) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, can be rewritten as

$$N_{a,-a} = 3|G_2| - 6 + |G_3| + \frac{n^2 - 9n + 18}{2}$$

This equality together with (1.7) yields $|G_2| = 2$. Such a result and those obtained from the above cases allow us to conclude that: G has just one involution and $2G$ must be an elementary abelian 3-group. Now using Proposition 1 we see that $\mathcal{D}_5(G)$ is not a 2-design, the Proposition is proved. \square

Lemma 3 *Suppose G is a finite abelian group of even order $n > 4$ in which $G_4 \neq G \neq G_2 + 2G$ and choose $\alpha \in G$ in such a way that $\alpha \notin G_2 + 2G$, $4\alpha \neq 0$. Then $a = 2\alpha$ and $-a$ are distinct elements of G and*

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Proof Clearly, from $a = 2\alpha$ it follows $a \in 2G$, $2a \neq 0$, $a \notin 4G$, $3a \neq 0$, $5a \neq 0$. We first note that: $\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 0\}$; if $g \in G \setminus \{a, -a, 0\}$, any block of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$ does not intersect $\{-a - g, a - g, -2g\}$. We now discuss five cases.

Case (L. 2. 1): $4a = 0$ and $|G_3| = 1$. In this case we have:

$\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 2a\}$;

if $g \in G$ and $2g \in \{a, -a\}$, then $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in 2G \setminus \langle a \rangle$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= 2 \cdot \frac{n-2-|G_2|}{2} + 2 \cdot |G_2| \cdot \frac{n-4}{2} \\ &+ (|2G| - 4) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G|) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$ and $|G_3| = 1$, yields

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 2. 2): $4a = 0$ and $|G_3| \neq 1$. In this case we have:

$\frac{n-2-|G_2|}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 2a\}$;

if $g \in G$ and $2g \in \{a, -a\}$, then $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in G_3$ is distinct from 0, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in 2G \setminus G_3$ does not belong to $\langle a \rangle$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;

if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$; there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$N_{a,-a} = 2 \cdot \frac{n-2-|G_2|}{2} + 2 \cdot |G_2| \cdot \frac{n-4}{2} + (|G_3| - 1) \cdot \frac{n-4-|G_2|}{2} \\ + (|2G| - |G_3| - 3) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G|) \cdot \frac{n-6}{2}$$

which, since $|2G| \cdot |G_2| = |G| = n$, simplifies to

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 2. 3): a has order 6. In this case we obtain:

- if $g \in \{2a, -2a\}$, there are $\frac{n-2-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G$ and $2g \in \{a, -a\}$, then $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G_3 \setminus \{0, 2a, -2a\}$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in 2G \setminus G_3$ with $-a \neq g \neq a$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$N_{a,-a} = 3 \cdot \frac{n-2-|G_2|}{2} + 2 \cdot |G_2| \cdot \frac{n-4}{2} + (|G_3| - 3) \cdot \frac{n-4-|G_2|}{2} \\ + (|2G| - |G_3| - 2) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G|) \cdot \frac{n-6}{2}$$

which, since $|2G| \cdot |G_2| = |G| = n$, yields

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 2. 4): $4a \neq 0 \neq 6a$ and $|G_3| = 1$. In this case we get:

- if $g \in \{2a, -2a\}$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G$ and $2g \in \{-a, a\}$, then $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in 2G \setminus \{a, -a, 0, -2a, 2a\}$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$N_{a,-a} = \frac{n-2-|G_2|}{2} + 2 \cdot \frac{n-4-|G_2|}{2} + 2 \cdot |G_2| \cdot \frac{n-4}{2} \\ + (|2G| - 5) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G|) \cdot \frac{n-6}{2}$$

which, since $|2G| \cdot |G_2| = |G| = n$ and $|G_3| = 1$, yields

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}.$$

Case (L. 2. 5): $4a \neq 0 \neq 6a$ and $|G_3| \neq 1$. In this case we deduce:

- if $g \in \{2a, -2a\}$, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G$ and $2g \in \{a, -a\}$, then $g \notin 2G$ and there are $\frac{n-4}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G_3$ is distinct from 0, there are $\frac{n-4-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in 2G \setminus G_3$ and $g \notin \{a, -a, 2a, -2a\}$, there are $\frac{n-6-|G_2|}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
- if $g \in G \setminus 2G$ with $-a \neq 2g \neq a$, there are $\frac{n-6}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.

Therefore

$$\begin{aligned} N_{a,-a} &= \frac{n-2-|G_2|}{2} + (|G_3| + 1) \cdot \frac{n-4-|G_2|}{2} + 2 \cdot |G_2| \cdot \frac{n-4}{2} \\ &\quad + (|2G| - |G_3| - 4) \cdot \frac{n-6-|G_2|}{2} + (n-2 \cdot |G_2| - |2G|) \cdot \frac{n-6}{2} \end{aligned}$$

which, since $|2G| \cdot |G_2| = |G| = n$, yields

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}.$$

The Lemma 3 is proved. \square

Theorem 1 *If $\mathcal{D}_5(G)$ is a 2-design, then $n = |G|$ must be odd integer.*

Proof We may assume that G is a finite additive abelian group of order $n = |G| > 4$. Suppose n is an even integer: so we must show that $\mathcal{D}_5(G)$ is not a 2-design. We discuss five cases.

Case (T. 1): $2g = 0$ whenever $g \in G \setminus 2G$. In this case G must be an elementary abelian 2-group and hence, by Proposition 1, $\mathcal{D}_5(G)$ is not a 2-design.

Case (T. 2): G is an abelian group of exponent 4. In this case either G is direct sum of cyclic groups of order 4 or G is direct sum of groups of order 2 and cyclic groups of order 4. Then, by Proposition 1, $\mathcal{D}_5(G)$ is not a 2-design.

Case (T. 3): $2G = 4G$. Then Proposition 2 asserts that $\mathcal{D}_5(G)$ is not a 2-design.

Case (T. 4): $2G \neq 4G$ and $4x = 0$ for every $x \notin G_2 + 2G$. Then G must be an abelian group of exponent 4 and hence, by statements 2) and 3) of Proposition 1, $\mathcal{D}_5(G)$ is not a 2-design.

Case (T. 5): $2G \neq 4G$ and $G \neq G_4$. Then $G_2 + 2G$ is a proper subgroup of G and there is $\alpha \in G$ such that $\alpha \notin G_2 + 2G$ and $4\alpha \neq 0$. Thus $a = 2\alpha \neq -a$ and, by Lemma 3, we obtain

$$N_{a,-a} = 3|G_2| + |G_3| + \frac{n^2 - 9n + 18}{2}. \quad (1.8)$$

On the other hand, since $2G \neq 4G$ implies that G is not an elementary abelian 2-group, there is $z \in G$ such that $z \notin 2G$ and $2z \neq 0$. Then using Lemma 2 we deduce that

$$N_{z,-z} = |G_3| + \frac{n^2 - 9n + 18}{2}. \quad (1.9)$$

Since $|G_2| \neq 0$, combining (1.8) and (1.9) we deduce that $N_{a,-a} \neq N_{z,-z}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design. Now the proof of the theorem is complete. \square

2 Main result

Proposition 3 $\mathcal{D}_5(G)$ is not a 2-design if one of the statements below is true:

1. G is a finite abelian group of odd order n divisible by 3;
2. G is a finite abelian group of odd order n not divisible by 5.

Proof

1. Choose $a \in G_3$, $a \neq 0$. Then clearly we have:
 $\frac{n-3}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{a, -a, 0\}$;
if $g \in G_3 \setminus \langle a \rangle$, there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$;
if $g \in G \setminus G_3$, there are $\frac{n-7}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{a, -a, g\}$.
Therefore

$$N_{a,-a} = \frac{n-3}{2} + (|G_3| - 3) \cdot \frac{n-5}{2} + (n - |G_3|) \cdot \frac{n-7}{2}. \quad (2.1)$$

Note that if G is an elementary abelian 3-group, then $G = G_3$ and (2.1) can be rewritten as

$$N_{a,-a} = \frac{n-3}{2} + (n-3) \cdot \frac{n-5}{2}. \quad (2.2)$$

Suppose $|G_5| \neq 1$ and choose $\alpha \in G_5$, $\alpha \neq 0$. Then we obtain:
if $g \in \{0, 2\alpha, -2\alpha\}$, there are $\frac{n-3}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{\alpha, -\alpha, g\}$;
if $0 \neq g \in G_3$, there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{\alpha, -\alpha, g\}$;
if $g \in G \setminus G_3$ and $g \notin \{\alpha, -\alpha, -2\alpha, 2\alpha\}$, there are $\frac{n-7}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{\alpha, -\alpha, g\}$.
Therefore

$$N_{\alpha,-\alpha} = 3 \cdot \frac{n-3}{2} + (|G_3| - 1) \cdot \frac{n-5}{2} + (n-4 - |G_3|) \cdot \frac{n-7}{2}. \quad (2.3)$$

Combining (2.1) and (2.3) we deduce that $N_{a,-a} \neq N_{\alpha,-\alpha}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design.

Suppose $|G_5| = 1$, $G_3 \neq G$ and choose $\beta \in G \setminus G_3$. Then we find:
 $\frac{n-3}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{\beta, -\beta, 0\}$;
 if $0 \neq g \in \{2\beta, -2\beta\}$, there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{\beta, -\beta, g\}$;
 if $\gamma \in G$ with $2\gamma = -\beta$, there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{\beta, -\beta, \gamma\}$;
 there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{\beta, -\beta, -\gamma\}$;
 if $0 \neq g \in G_3$, there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{\beta, -\beta, g\}$;
 if $g \in G \setminus G_3$ and $g \notin \{\beta, -\beta, 2\beta, -2\beta, \gamma, -\gamma\}$, there are $\frac{n-7}{2}$ blocks $\mathcal{D}_5(G)$ through $\{\beta, -\beta, g\}$.

Therefore

$$N_{\beta, -\beta} = \frac{n-3}{2} + (|G_3| + 3) \cdot \frac{n-5}{2} + (n-6 - |G_3|) \cdot \frac{n-7}{2}. \quad (2.4)$$

Combining (2.1) and (2.4) we find $N_{a, -a} \neq N_{\beta, -\beta}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design.

We can now assume that $G = G_3$. Then for any $g \in G \setminus \langle a \rangle$ there are $\frac{n-7}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$. Therefore

$$N_{0, a} = \frac{n-3}{2} + (n-3) \cdot \frac{n-7}{2}. \quad (2.5)$$

Combining (2.2) and (2.5) we find $N_{a, -a} \neq N_{0, a}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design.

2. By **1** we may assume that n and 15 are (odd integers) relatively prime. Choose $x \in G$, $x \neq 0$ and let y, z be elements of G such that $2y = x$, $2z = 7x$. Then clearly we have:

$\frac{n-3}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through the 3-set $\{x, -x, 0\}$;
 if $g \in \{2x, -2x, y, -y\}$, there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{x, -x, g\}$;
 if $0 \neq g \in G \setminus \{x, -x, 2x, -2x, y, -y\}$, there are $\frac{n-7}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{x, -x, g\}$.

Therefore

$$N_{x, -x} = \frac{n-3}{2} + 4 \cdot \frac{n-5}{2} + (n-7) \cdot \frac{n-7}{2}. \quad (2.6)$$

On the other hand we have:

if $g \in \{6x, 11x, z\}$, there are $\frac{n-5}{2}$ blocks of $\mathcal{D}_5(G)$ through the 3-set $\{x, -4x, g\}$;
 if $g \in G \setminus \{x, -4x, 6x, 11x, z\}$, there are $\frac{n-7}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{x, -4x, g\}$.

Therefore

$$N_{x, -4x} = 3 \cdot \frac{n-5}{2} + (n-5) \cdot \frac{n-7}{2}. \quad (2.7)$$

Combining (2.6) and (2.7) we obtain $N_{x, -x} \neq N_{x, -4x}$ and hence, by Lemma 1, $\mathcal{D}_5(G)$ is not a 2-design. Now the Proposition 3 is proved. \square

We can now state our main result.

Theorem 2 $\mathcal{D}_5(G)$ is a 2-design if and only if G is an elementary abelian 5-group. When this is so, there are

$$\lambda = \frac{|G| - 3}{2} + \frac{(|G| - 5) \cdot (|G| - 7)}{6}$$

blocks of $\mathcal{D}_5(G)$ through any given 2-set $\{x, y\} \subset G$.

Proof Suppose $\mathcal{D}_5(G)$ is a 2-design. By Theorem 1 and Proposition 3, $n = |G|$ must be an odd integer multiple of 5 not divisible by 3. Let $a \in G_5$, $a \neq 0$. Then we find: if $g \in \langle a \rangle$ with $0 \neq g \neq a$, then $\frac{n-3}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$; if $g \in G \setminus \langle a \rangle$, there are $\frac{n-7}{2}$ blocks of $\mathcal{D}_5(G)$ through $\{0, a, g\}$.

Therefore

$$N_{0,a} = 3 \cdot \frac{n-3}{2} + (n-5) \cdot \frac{n-7}{2} \quad (2.8)$$

Assume that $5b \neq 0$ for some $b \in G$ and let β be the unique element in G such that $2\beta = 7b$. Then $(\beta \in G \setminus \{b, -4b, 6b, 11b\})$ and we obtain: if $g \in \{6b, 11b, \beta\}$, then $\frac{n-5}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{b, -4b, g\}$; if $g \in G \setminus \{b, -4b, 6b, 11b, \beta\}$, then $\frac{n-7}{2}$ is the number of blocks of $\mathcal{D}_5(G)$ through $\{b, -4b, g\}$.

Therefore

$$N_{b,-4b} = 3 \cdot \frac{n-5}{2} + (n-5) \cdot \frac{n-7}{2}$$

and thus, since $\mathcal{D}_5(G)$ is a 2 design, we find (by Lemma 1)

$$3 \cdot \frac{n-3}{2} + (n-5) \cdot \frac{n-7}{2} = N_{0,a} = N_{b,-4b} = 3 \cdot \frac{n-5}{2} + (n-5) \cdot \frac{n-7}{2}$$

and this gives $n-3 = n-5$ a contradiction. Such a contradiction shows that $5g = 0$ for all $g \in G$: in other words, G is an elementary abelian 5-group. Furthermore, from Lemma 1 and equation (2.8) we know that

$$3 \cdot \frac{n-3}{2} + (n-5) \cdot \frac{n-7}{2} = N_{0,a} = 3\lambda$$

from which it follows that $\lambda = \frac{|G|-3}{2} + (|G|-5) \cdot \frac{|G|-7}{6}$ is the number of blocks of $\mathcal{D}_5(G)$ through any given two distinct elements $x, y \in G$.

To finish, assume that G is an elementary abelian 5-group. If we regard G as a vector space over the field with five elements, then we see that the affine group $\text{Aff}(G)$ acts 2-homogeneously on G and the block-set \mathcal{B} of $\mathcal{D}_5(G)$ may be written as $\mathcal{B} = C^{\text{Aff}(G)}$ (i.e. $\mathcal{B} = \{C^\gamma \mid \gamma \in \text{Aff}(G)\}$) is the $\text{Aff}(G)$ -orbit of a fixed block $C \in \mathcal{B}$. Hence, by [1, Proposition 4.6], $\mathcal{D}_5(G)$ is a 2-design. The Theorem is proved. \square

References

- [1] Beth, T., Jungnickel, D., Lenz, H.: Design Theory. 2nd ed., *Cambridge University Press*, Cambridge, 1999.
- [2] Caggegi, A., Di Bartolo, A., Falcone, G.: *Boolean 2-designs and the embedding of a 2-design in a group*. arxiv 0806.3433v2, (2008), 1–8.
- [3] Caggegi, A., Falcone, G., Pavone, M.: *On the additivity of block design*. submitted.
- [4] Caggegi, A.: *Some additive 2 – (v, 4, λ) designs*. *Boll. Mat. Pura e Appl.* **2** (2009), 1–3.
- [5] Colbourn, C. J., Dinitz, J. H.: *The CRC Handbook of Combinatorial Designs*. *Discrete Mathematics and Its Applications*, 2nd ed., *Chapman & Hall/CRC Press*, 2007.