

Luisa Di Piazza; Valeria Marraffa; Kazimierz Musiał  
Variational Henstock integrability of Banach space valued functions

*Mathematica Bohemica*, Vol. 141 (2016), No. 2, 287–296

Persistent URL: <http://dml.cz/dmlcz/145716>

## Terms of use:

© Institute of Mathematics AS CR, 2016

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

VARIATIONAL HENSTOCK INTEGRABILITY OF  
BANACH SPACE VALUED FUNCTIONSLUISA DI PIAZZA, Palermo, VALERIA MARRAFFA, Palermo,  
KAZIMIERZ MUSIAŁ, WrocławReceived March 6, 2016  
Communicated by Dagmar Medková

*Cordially dedicated to Professor Jaroslav Kurzweil on the occasion  
of his 90th birthday*

*Abstract.* We study the integrability of Banach space valued strongly measurable functions defined on  $[0, 1]$ . In the case of functions  $f$  given by  $\sum_{n=1}^{\infty} x_n \chi_{E_n}$ , where  $x_n$  are points of a Banach space and the sets  $E_n$  are Lebesgue measurable and pairwise disjoint subsets of  $[0, 1]$ , there are well known characterizations for Bochner and Pettis integrability of  $f$ . The function  $f$  is Bochner integrable if and only if the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is absolutely convergent. Unconditional convergence of the series is equivalent to Pettis integrability of  $f$ . In this paper we give some conditions for variational Henstock integrability of a certain class of such functions.

*Keywords:* Kurzweil-Henstock integral; variational Henstock integral; Pettis integral

*MSC 2010:* 26A39

## 1. INTRODUCTION

In this paper we study the variational Henstock integrability of strongly measurable functions. It is well known (cf. [5], Lemma 5.1) that each strongly measurable Banach valued function, defined on a measurable space, can be written as  $f = g + \sum_{n=1}^{\infty} x_n \chi_{E_n}$ , where  $g$  is a bounded strongly measurable function,  $x_n$  are vectors of the given Banach space and  $E_n$  are measurable and pairwise disjoint sets. As each bounded strongly measurable function is Bochner integrable, it is enough to study

---

The research has been supported by the grant GNAMPA 2016-Di Piazza.

integrability only for functions of the form  $\sum_{n=1}^{\infty} x_n \chi_{E_n}$ . In the case of Bochner and Pettis integrals, a necessary and sufficient condition for integrability of a function given by  $\sum_{n=1}^{\infty} x_n \chi_{E_n}$  is, respectively, the absolute and the unconditional convergence of the series  $\sum_{n=1}^{\infty} x_n |E_n|$  (see Theorem A). In the case of Kurzweil-Henstock or variational Henstock integrals, in general the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is only conditionally convergent. So the conditions for integrability depend on the order of the terms  $x_n |E_n|$ . In [1], [3] and [4] conditions for the Kurzweil-Henstock integrability of functions of the form  $\sum_{n=1}^{\infty} x_n \chi_{E_n}$  are given. Here we go a bit further in this investigation. We give another characterization of the Kurzweil-Henstock integrability (see Theorem 3.1). The main results are Proposition 4.1 and Theorem 4.1. In the latter, a necessary and sufficient condition for the variational Henstock integrability of a special type of such functions is given. It needs a particular order of the sets  $E_n$ .

## 2. BASIC FACTS

Let  $[0, 1]$  be the unit interval of the real line equipped with the usual topology and Lebesgue measure. If a set  $E \subset [0, 1]$  is Lebesgue measurable, then  $|E|$  denotes its Lebesgue measure.  $\mathcal{I}$  denotes the family of all closed subintervals of  $[0, 1]$ .

A *partition* in  $[0, 1]$  is a finite collection of pairs  $\mathcal{P} = \{(I_1, t_1), \dots, (I_p, t_p)\}$ , where  $I_1, \dots, I_p$  are nonoverlapping subintervals of  $[0, 1]$  and  $t_i \in I_i$ ,  $i = 1, \dots, p$ . If  $\bigcup_{i=1}^p I_i = [0, 1]$ , we say that  $\mathcal{P}$  is a *partition* of  $[0, 1]$ . A *gauge* on  $E \subset [0, 1]$  is a positive function on  $E$ . For a given gauge  $\delta$ , we say that a partition  $\{(I_1, t_1), \dots, (I_p, t_p)\}$  is  $\delta$ -*fine* if  $I_i \subset (t_i - \delta(t_i), t_i + \delta(t_i))$ ,  $i = 1, \dots, p$ .

Throughout this paper,  $X$  is a Banach space with dual  $X^*$ . We recall the following definitions:

**Definition 2.1.** A function  $f: [0, 1] \rightarrow X$  is said to be *Kurzweil-Henstock integrable* (or simply *KH-integrable*) on  $[0, 1]$  if there exists  $w \in X$  with the following property:

For every  $\varepsilon > 0$  there exists a gauge  $\delta$  on  $[0, 1]$  such that

$$\left\| \sum_{i=1}^p f(t_i) |I_i| - w \right\| < \varepsilon,$$

for each  $\delta$ -fine partition  $\{(I_1, t_1), \dots, (I_p, t_p)\}$  of  $[0, 1]$ . We set  $(\text{KH}) \int_0^1 f := w$ .

**Definition 2.2.** A function  $f: [0, 1] \rightarrow X$  is said to be *variationally Henstock integrable* (briefly *vH-integrable*) on  $[0, 1]$ , if there exists an additive function  $F: \mathcal{I} \rightarrow X$ , satisfying the following condition:

Given  $\varepsilon > 0$  there exists a gauge  $\delta$  such that if  $\mathcal{P} = \{(I_i, t_i): i = 1, \dots, p\}$  is a  $\delta$ -fine partition in  $[0, 1]$ , then

$$\sum_{i=1}^p \|f(t_i)|I_i| - F(I_i)\| < \varepsilon.$$

It is obvious that each vH-integrable function is KH-integrable. It is also well known that in the case of real-valued functions the variational Henstock and the Kurzweil-Henstock integrals are equivalent.

We recall the following classical result for the Bochner and Pettis integrals:

**Theorem A** ([2], page 55). *Let  $f = \sum_{n=1}^{\infty} x_n \chi_{E_n}$ , where  $x_n \in X$  and the sets  $E_n$  are Lebesgue measurable and pairwise disjoint subsets of  $[0, 1]$ . Then*

- (1)  *$f$  is Pettis integrable if and only if the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is unconditionally convergent;*
- (2)  *$f$  is Bochner integrable if and only if the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is absolutely convergent.*

*In both cases  $\int_E f = \sum_{n=1}^{\infty} x_n |E_n \cap E|$ , for every measurable set  $E$ .*

### 3. KURZWEIL-HENSTOCK INTEGRABILITY

In [1], Theorem 1, a necessary condition for the Kurzweil-Henstock integrability of the function  $f = \sum_{n=1}^{\infty} x_n \chi_{E_n}$  is given. Here we prove that the condition is also sufficient.

**Theorem 3.1.** *Let  $f: [0, 1] \rightarrow X$  be defined by  $f = \sum_{n=1}^{\infty} x_n \chi_{E_n}$ , where  $x_n \in X$  and the sets  $E_n$  are Lebesgue measurable and pairwise disjoint. Then the following conditions are equivalent:*

- (A)  *$f$  is Kurzweil-Henstock integrable with*

$$(KH) \int_I f(t) dt = \sum_{n=1}^{\infty} x_n |E_n \cap I|,$$

*for every interval  $I \in \mathcal{I}$ ;*

(B) for every  $\varepsilon > 0$  there exist a gauge  $\delta$  and  $k_0 \in \mathbb{N}$  such that given a  $\delta$ -fine partition  $\{(I_1, t_1), \dots, (I_p, t_p)\}$  of  $[0, 1]$  and given  $s > r > k_0$  we have

$$\left\| \sum_{k=r}^s x_k \left| \bigcup_{t_j \in E_k} I_j \right| \right\| < \varepsilon.$$

*Proof.* (B)  $\Rightarrow$  (A) was proved in [1].

(A)  $\Rightarrow$  (B) We assume that  $f$  is Kurzweil-Henstock integrable with

$$(KH) \int_0^1 f(t) dt = \sum_{n=1}^{\infty} x_n |E_n|.$$

According to [3], Theorem 2, for every  $\varepsilon > 0$  there exists a gauge  $\delta$  on  $[0, 1]$  such that if  $\mathcal{P} := \{(i_1, t_1), \dots, (I_p, t_p)\}$  is a  $\delta$ -fine partition of  $[0, 1]$ , then there exists  $n_{\mathcal{P}} \in \mathbb{N}$  such that

$$\left\| \sum_{n=1}^n x_k \left( \left| \bigcup_{t_i \in E_k} I_i \right| - |E_k| \right) \right\| < \frac{\varepsilon}{3} \quad \text{for all } n > n_{\mathcal{P}}.$$

Since the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is convergent, there is  $n_1 > n_{\mathcal{P}}$  such that if  $s > r > n_1$ , then

$$\left\| \sum_{i=r}^s x_i |E_i| \right\| < \frac{\varepsilon}{3}.$$

Hence, if  $s > r > n_1$ , then

$$\begin{aligned} \left\| \sum_{k=r}^s x_k \left| \bigcup_{t_j \in E_k} I_j \right| \right\| &\leq \left\| \sum_{k=1}^s x_k \left| \bigcup_{t_j \in E_k} I_j \right| - \sum_{k=1}^s x_k |E_k| \right\| \\ &\quad + \left\| \sum_{k=1}^{r-1} x_k \left| \bigcup_{t_j \in E_k} I_j \right| - \sum_{k=1}^{r-1} x_k |E_k| \right\| + \left\| \sum_{i=r}^s x_i |E_i| \right\| < \varepsilon. \end{aligned}$$

□

#### 4. VARIATIONAL HENSTOCK INTEGRABILITY

The aim of this section is to formulate conditions for the variational Henstock integrability of a certain class of strongly measurable functions.

**Proposition 4.1.** Let  $\{a_n\}$  be a decreasing sequence converging to zero such that  $a_1 = 1$ . Let  $\{x_n\} \subset X$  be arbitrary and define  $f: [0, 1] \rightarrow X$  by  $f = \sum_{n=1}^{\infty} x_n \chi_{E_n}$ , where each  $E_n \subseteq [a_{n+1}, a_n]$  is Lebesgue measurable. Then the following conditions are equivalent:

- (i) the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is convergent;
- (ii)  $f$  is vH-integrable;
- (iii)  $f$  is KH-integrable.

In each case

$$(4.1) \quad (\text{vH}) \quad \int_I f = \sum_{n=1}^{\infty} x_n |E_n \cap I| \quad \text{for every } I \in \mathcal{I}$$

and the series  $\sum_{n=1}^{\infty} x_n |E_n \cap I|$  is uniformly convergent on  $\mathcal{I}$ .

*Proof.* (i)  $\Rightarrow$  (ii) Assume that the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is convergent. Notice then that the series  $\sum_{n=1}^{\infty} x_n |E_n \cap I|$  is convergent for every  $I \in \mathcal{I}$ . Let  $F(I) = \sum_{n=1}^{\infty} x_n |E_n \cap I|$ . Now we show that  $f$  is vH-integrable. Without loss of generality we may assume that  $f(0) = 0$ .

Let  $\varepsilon > 0$ . Since the series  $\sum_{n=1}^{\infty} x_n |E_n|$  is convergent, there is  $K \in \mathbb{N}$  such that for  $s \geq n \geq K$ ,

$$\left\| \sum_{k=n}^s x_k |E_k| \right\| < \frac{\varepsilon}{4}.$$

Moreover, for each  $n \in \mathbb{N}$ , let  $\delta_n: [a_{n+1}, a_n] \rightarrow (0, \infty)$  be a gauge such that if  $\mathcal{P} = \{(I_i, t_i), i = 1, \dots, p\}$  is a  $\delta_n$ -fine partition of  $[a_{n+1}, a_n]$ , then

$$\sum_{i=1}^p \|f(t_i) |I_i| - F(I_i)\| < \frac{\varepsilon}{2^{n+1}}.$$

We may assume that  $\delta_{n+1}(a_{n+1}) = \delta_n(a_{n+1})$ .

Define  $\delta(t)$  on  $[0, 1]$  as follows:

$$\delta(t) = \begin{cases} \delta_n(t) & \text{if } t \in (a_{n+1}, a_n), \\ \min\{\delta_n(a_n), \delta_{n-1}(a_n)\} & \text{if } t = a_n, \\ a_K & \text{if } t = 0. \end{cases}$$

Let us consider now a  $\delta$ -fine partition  $\mathcal{P} = \{(I_i, t_i), i = 1, \dots, p\}$  of  $[0, 1]$  and the corresponding sum

$$\sum_{i=1}^p \|f(t_i)|I_i| - F(I_i)\|.$$

If  $q \geq K$  is the largest integer such that  $I_1 \subset [0, a_q]$ , then

$$\begin{aligned} (4.2) \quad \|f(t_1)|I_1| - F(I_1)\| &= \left\| \sum_{n=1}^{\infty} x_n |E_n \cap I_1| \right\| \\ &= \left\| \sum_{k=q}^{\infty} x_k |E_k \cap I_1| \right\| = \left\| x_q |E_q \cap I_1| + \sum_{k=q+1}^{\infty} x_k |E_k| \right\| \\ &\leq \|x_q\| |E_q \cap I_1| + \left\| \sum_{k=q+1}^{\infty} x_k |E_k| \right\| < \frac{\varepsilon}{2}. \end{aligned}$$

Hence

$$\begin{aligned} \sum_{i=1}^p \|f(t_i)|I_i| - F(I_i)\| &= \|f(t_1)|I_1| - F(I_1)\| \\ &\quad + \sum_{n=1}^{\infty} \sum_{t_i \in (a_{n+1}, a_n]} \|f(t_i)|I_i \cap [a_{n+1}, a_n]| - F(I_i \cap [a_{n+1}, a_n])\| \\ &\leq \frac{\varepsilon}{2} + \sum_{n=1}^{\infty} \frac{\varepsilon}{2^{n+1}} = \varepsilon, \end{aligned}$$

which proves the vH-integrability of  $f$  and equality (4.1) for  $I = [0, 1]$ .

(iii)  $\Rightarrow$  (i) If  $f$  is KH-integrable, its primitive  $F(t) = (\text{vH}) \int_0^t f$  is continuous on  $[0, 1]$ . Let  $F(I)$  be the additive interval function associated to  $F(t)$ . We have

$$F([0, 1]) = \sum_{k=1}^n F([a_{k+1}, a_k]) + F([0, a_{n+1}]) = \sum_{k=1}^n x_k |E_k| + F([0, a_{n+1}]).$$

Letting  $n \rightarrow \infty$ , the convergence of the series  $\sum_{n=1}^{\infty} x_n |E_n|$  follows.

In the same way, setting  $F_I(t) := (\text{vH}) \int_{\alpha}^t f$  if  $t \in I = [\alpha, \beta]$ , we obtain (4.1).

Now we are going to prove that the series  $\sum_{n=1}^{\infty} x_n |E_n \cap I|$  is uniformly convergent on  $\mathcal{I}$ .

Since  $F$  is uniformly continuous, there is  $n_0 \in \mathbb{N}$  such that if  $I \subset [0, a_{n_0}]$ , then

$$(4.3) \quad \left\| \sum_{n=1}^{\infty} x_n |E_n \cap I| \right\| = \|F(I)\| \leq \varepsilon.$$

Now, if  $I \in \mathcal{I}$  and  $m > n_0$ , then applying (4.1) and (4.3), we have the following inequalities:

$$\begin{aligned}
& \left\| F(I) - \sum_{n=1}^m x_n |E_n \cap I| \right\| \\
& \leq \left\| F(I \cap [0, a_m]) - \sum_{n=1}^m x_n |E_n \cap I \cap [0, a_m]| \right\| \\
& \quad + \left\| F(I \cap [a_m, 1]) - \sum_{n=1}^m x_n |E_n \cap I \cap [a_m, 1]| \right\| \\
& \leq \left\| F(I \cap [0, a_m]) - \sum_{n=1}^{\infty} x_n |E_n \cap I \cap [0, a_m]| \right\| \\
& \quad + \left\| \sum_{n=m+1}^{\infty} x_n |E_n \cap I \cap [0, a_m]| \right\| \\
& \quad + \left\| F(I \cap [a_m, 1]) - \sum_{n=1}^{\infty} x_n |E_n \cap I \cap [a_m, 1]| \right\| \\
& \quad + \left\| \sum_{n=m+1}^{\infty} x_n |E_n \cap I \cap [a_m, 1]| \right\| \\
& \stackrel{(4.1)}{=} \left\| \sum_{n=m+1}^{\infty} x_n |E_n \cap I \cap [0, a_m]| \right\| + \left\| \sum_{n=m+1}^{\infty} x_n |E_n \cap I \cap [a_m, 1]| \right\| \\
& = \left\| \sum_{n=1}^{\infty} x_n |E_n \cap I \cap [0, a_m]| \right\| \stackrel{(4.3)}{\leq} \varepsilon \quad \text{for every } I \in \mathcal{I}.
\end{aligned}$$

The last equality follows from the fact that  $E_n \cap [a_m, 1] = \emptyset$  if  $n > m$ . □

Reordering the sets  $E_n$  in a suitable way, we obtain the following more general result:

**Theorem 4.1.** *Let  $\{a_n\}$  and  $\{b_n\}$  be decreasing sequences converging to zero such that  $a_1 = 1$  and  $a_{n+1} \leq b_n \leq a_n$ , for every  $n \in \mathbb{N}$ . Let  $\{x_n\} \subset X$  be arbitrary and define  $f: [0, 1] \rightarrow X$  by  $f = \sum_{k=1}^{\infty} x_k \chi_{E_k}$ , where  $\{E_k: k \in \mathbb{N}\}$  is a sequence of pairwise disjoint Lebesgue measurable sets of positive measure with the following properties:*

- (j)  $\lim_k \text{diam}(E_k) = 0$ ;
- (jj) *for each  $n \in \mathbb{N}$ , the set  $\{E_k: E_k \subset [a_{n+1}, a_n]\}$  is split into two disjoint collections (one of them may be empty):*

$$\{E_{2n-1, p_i}: \forall i \in \mathbb{N} \sup E_{2n-1, p_{i+1}} \leq \inf E_{2n-1, p_i}\} \subset [a_{n+1}, b_n]$$



and

$$\{E_{2n,q_i} : \forall i \in \mathbb{N} \inf E_{2n,q_{i+1}} \geq \sup E_{2n,q_i}\} \subset [b_n, a_n];$$

(jjj) for each  $n \in \mathbb{N}$ ,  $\lim_i d_H(\{a_{n+1}\}, E_{2n-1,p_i}) = 0 = \lim_i d_H(\{a_n\}, E_{2n,q_i})$ , where  $d_H(\cdot, \cdot)$  is the Hausdorff distance between two sets.

Let  $c_{2n-1,i}(I) := x_n |E_{2n-1,p_i} \cap I|$  and  $c_{2n,i}(I) := x_n |E_{2n,q_{i+1}} \cap I|$ ,  $n \in \mathbb{N}$ . We order the series  $\sum_{k=1}^{\infty} x_k |E_k \cap I|$  in the following way:

$$(4.4) \quad \sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I).$$

Then, the following conditions are equivalent:

- (a) the series (4.4) is uniformly convergent on the family  $\mathcal{I}$ ;
- (b)  $f$  is vH-integrable;
- (c)  $f$  is KH-integrable.

In each case

$$(4.5) \quad (\text{vH}) \int_I f = \sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I) \quad \text{for every } I \in \mathcal{I}.$$

**Proof.** Without loss of generality, we may assume that if for some  $n \in \mathbb{N}$  one has  $\{E_{n,p_i} : i \in \mathbb{N}, \text{ for all } i \in \mathbb{N}\} = \emptyset$ , then  $a_{n+1} = b_n$ , and if  $\{E_{n,q_i} : i \in \mathbb{N}, \text{ for all } i \in \mathbb{N}\} = \emptyset$ , then  $a_n = b_n$ . We may assume also that each interval  $[a_{n+1}, a_n]$  contains infinitely many sets  $E_k$  and  $f(0) = 0$ .

(a)  $\Rightarrow$  (b) It follows from Proposition 4.1 that if for every  $I \in \mathcal{I}$  the series  $\sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I)$  is convergent, then  $f$  is vH-integrable on every interval  $[a_{n+1}, b_n]$  and  $[b_n, a_n]$ . Consequently,  $f$  is vH-integrable on  $[a_{n+1}, a_n]$  and

$$(\text{vH}) \int_{a_{n+1}}^{a_n} f = \sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}([a_{n+1}, a_n]).$$

Now let  $F(I) = \sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I)$  for every  $I \in \mathcal{I}$  and let  $\varepsilon > 0$ . For each  $n \in \mathbb{N}$  there exists a gauge  $\delta_n : [a_{n+1}, a_n] \rightarrow (0, \infty)$  with the property that for each  $\delta_n$ -partition  $\{(J_1, s_1), \dots, (J_p, s_p)\}$  of  $[a_{n+1}, a_n]$  one has

$$\sum_{j=1}^p \left\| f(s_j) |J_j| - (\text{vH}) \int_{J_j} f \right\| \leq \frac{\varepsilon}{2^{n+2}}.$$

Taking  $\min\{\delta_{n+1}(a_{n+1}), \delta_n(a_{n+1})\}$ , one may assume that  $\delta_{n+1}(a_{n+1}) = \delta_n(a_{n+1})$ .

Let  $k_0 \in \mathbb{N}$  be such that  $k \geq k_0$  yields

$$\left\| \sum_{n=k}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I) \right\| < \frac{\varepsilon}{4} \quad \text{for every } I \in \mathcal{I}.$$

Then, let  $n_0 \in \mathbb{N}$  be such that all sets  $E_j$  built into some  $c_{i,n+1-i}(I)$  with  $n \leq k_0$  are contained in  $(a_{n_0}, 1]$ .

Define  $\delta(t)$  on  $[0, 1]$  as follows:

$$\delta(t) = \begin{cases} \delta_n(t) & \text{if } t \in [a_{n+1}, a_n], \quad n \in \mathbb{N}, \\ a_{n_0} & \text{if } t = 0. \end{cases}$$

Let  $\mathcal{P} = \{(I_i, t_i), i = 1, \dots, p\}$  be a  $\delta$ -fine partition of  $[0, 1]$  and let us consider the sum

$$\sum_{i=1}^p \|f(t_i)|I_i| - F(I_i)\|.$$

Without loss of generality, one may assume that the right end point of  $I_1$  is equal to a point  $a_m$  with  $m > n_0$ .

It follows that

$$\sum_{j=2}^p \left\| f(s_j)|J_j| - (\text{vH}) \int_{J_j} f \right\| \leq \frac{\varepsilon}{2}.$$

Then,

$$\begin{aligned} \left\| f(0)|J_1| - \sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(J_1) \right\| \\ = \left\| \sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(J_1) \right\| = \left\| \sum_{k=k_0}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(J_1) \right\| < \frac{\varepsilon}{4} \end{aligned}$$

and so  $f$  is vH-integrable.

(c)  $\Rightarrow$  (a) Assume the KH-integrability of  $f$  and let  $\varepsilon > 0$  and  $n_0 \in \mathbb{N}$  be such that  $I \subset [0, a_{n_0}]$  yields  $\|F(I)\| < \varepsilon/2$ . In virtue of Proposition 4.1, the series  $\sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(J)$  is uniformly convergent to  $F(J)$  on the family  $\mathcal{I} \cap [a_{n_0}, 1]$ . Let  $k_0 > n_0$  be such that if  $m > k_0$ , then  $E_m \cap (a_{n_0}, 1] = \emptyset$  and

$$\left\| F(J \cap [a_{n_0}, 1]) - \sum_{n=1}^m \sum_{i=1}^n c_{i,n+1-i}(J \cap [a_{n_0}, 1]) \right\| \leq \frac{\varepsilon}{2} \quad \text{for every } J \in \mathcal{I}.$$

If  $I \in \mathcal{I}$  is fixed and  $m > k_0$ , then

$$\left\| \sum_{n=m+1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I \cap [0, a_{n_0}]) \right\| = \|F(I \cap [0, a_{n_0}])\| \leq \frac{\varepsilon}{2}$$

and so, taking into account (4.5), we have

$$\begin{aligned} & \left\| F(I \cap [0, a_{n_0}]) - \sum_{n=1}^m \sum_{i=1}^n c_{i,n+1-i}(I \cap [0, a_{n_0}]) \right\| \\ & \leq \left\| F(I \cap [0, a_{n_0}]) - \sum_{n=1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I \cap [0, a_{n_0}]) \right\| \\ & \quad + \left\| \sum_{n=m+1}^{\infty} \sum_{i=1}^n c_{i,n+1-i}(I \cap [0, a_{n_0}]) \right\| \leq \frac{\varepsilon}{2}. \end{aligned}$$

As a result, if  $m > k_0$ , then

$$\left\| F(I) - \sum_{n=1}^m \sum_{i=1}^n c_{i,n+1-i}(I) \right\| \leq \varepsilon \quad \text{for every } I \in \mathcal{I},$$

which proves the uniform convergence of the series (4.5) on  $\mathcal{I}$ .  $\square$

**Remark 4.1.** In the same way as Theorem 4.1 was deduced from Proposition 4.1, one can obtain subsequent generalizations of Theorem 4.1.

#### References

- [1] *B. Bongiorno, L. Di Piazza, K. Musiał*: Kurzweil-Henstock and Kurzweil-Henstock-Pettis integrability of strongly measurable functions. *Math. Bohem.* 131 (2006), 211–223.
- [2] *J. Diestel, J. J. Uhl, Jr.*: *Vector Measures*. Mathematical Surveys 15, American Mathematical Society 13, Providence, 1977.
- [3] *V. Marraffa*: A characterization of strongly measurable Kurzweil-Henstock integrable functions and weakly continuous operators. *J. Math. Anal. Appl.* 340 (2008), 1171–1179.
- [4] *V. Marraffa*: Strongly measurable Kurzweil-Henstock type integrable functions and series. *Quaest. Math.* 31 (2008), 379–386.
- [5] *K. Musiał*: Topics in the theory of Pettis integration. School on Measure Theory and Real Analysis, Grado, 1991. *Rend. Ist. Mat. Univ. Trieste* 23, 1993, pp. 177–262.

*Authors' addresses:* *Luisa Di Piazza, Valeria Marraffa*, Department of Mathematics, University of Palermo, via Archirafi 34, 90123 Palermo, Italy, e-mail: [luisa.dipiazza@unipa.it](mailto:luisa.dipiazza@unipa.it), [valeria.marraffa@unipa.it](mailto:valeria.marraffa@unipa.it); *Kazimierz Musiał*, Institute of Mathematics, Wrocław University, Plac Grunwaldzki 2/4, 50-384 Wrocław, Poland, e-mail: [kazimierz.musial@math.uni.wroc.pl](mailto:kazimierz.musial@math.uni.wroc.pl).