

Mehdi Badie

Comaximal graph of $C(X)$

Commentationes Mathematicae Universitatis Carolinae, Vol. 57 (2016), No. 3, 353–364

Persistent URL: <http://dml.cz/dmlcz/145840>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2016

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Comaximal graph of $C(X)$

MEHDI BADIE

Abstract. In this article we study the comaximal graph $\Gamma'_2 C(X)$ of the ring $C(X)$. We have tried to associate the graph properties of $\Gamma'_2 C(X)$, the ring properties of $C(X)$ and the topological properties of X . Radius, girth, dominating number and clique number of the $\Gamma'_2 C(X)$ are investigated. We have shown that $2 \leq \text{Rad} \Gamma'_2 C(X) \leq 3$ and if $|X| > 2$ then $\text{girth} \Gamma'_2 C(X) = 3$. We give some topological properties of X equivalent to graph properties of $\Gamma'_2 C(X)$. Finally we have proved that X is an almost P -space which does not have isolated points if and only if $C(X)$ is an almost regular ring which does not have any principal maximal ideals if and only if $\text{Rad} \Gamma'_2 C(X) = 3$.

Keywords: rings of continuous functions; comaximal graph; radius; girth; dominating number; clique number; zero cellularity; P -space; almost P -space; connected space; regular ring

Classification: 54C40

1. Introduction

Throughout this paper, G stands for an undirected graph. Distance between two vertices u and v is defined as the length of shortest path between u and v , and is denoted by $d(u, v)$, then the *diameter* of G is denoted by $\text{diam}(G)$, and is defined to be the supremum of $\{d(u, v) : u, v \in G\}$. If u is a vertex of a graph G , then *eccentricity* of u , denoted by $\text{ecc}(u)$, is defined $\max\{d(u, v) : v \in G\}$. The set of all vertices with the smallest eccentricity is called *center* of G and $\min\{\text{ecc}(u) : u \in G\}$ is called the *radius* of G and is denoted by $\text{Rad}(G)$. The minimum length of cycles in a graph G is called the *girth* of G and is denoted by $\text{girth}(G)$. For every $u, v \in G$, let us denote by $\text{gi}(u, v)$ the length of the shortest cycle containing u and v . It is clear that $\text{girth}(G) = \min\{\text{gi}(u, v) : u, v \in G\}$. G is called *triangulated* (*hypertriangulated*) if each vertex (edge) of G is a vertex (edge) of a triangle. A subset A of G is called a *dominating set* if for each $u \in G \setminus A$, there exists v in A such that u is adjacent to v . The *dominating number* of G , denoted by $\text{dt}(G)$, is the smallest cardinal number of the form $|A|$, where A is a dominating set of G . It is said that two vertices u and v of G are *orthogonal*, written $u \perp v$, if u and v are adjacent and there is no a vertex w of G which is adjacent to both u and v . A graph G is called *complemented* if for each vertex u of G , there is a vertex v of G such that $u \perp v$. A clique of a graph G is defined as

a maximal complete subgraph of G and the supremum of $|A|$, where A is clique of G , is called the clique number of G , and is denoted by $\kappa(G)$.

Let R be a commutative ring with unity. R is called an almost ring if each non-unit element of R is a zero-divisor element of R . *Comaximal graph* $\Gamma(R)$ is defined as a graph with vertices of elements of R , where two distinct vertices a and b are adjacent if and only if $Ra + Rb = R$. Also consider a subgraph $\Gamma_2(R)$ of $\Gamma(R)$ which consists of all non-unit elements of R . If $J(R)$ is Jacobson radical of R , then $\Gamma_2(R) \setminus J(R)$ is denoted by $\Gamma'_2(R)$.

We assume throughout the paper that $C(X)$ is the ring of all real valued continuous functions on a Tychonoff space X . The *density (weight)* of X , denoted by $d(X)$ ($w(X)$), is the infimum of the cardinalities of dense subsets (bases) of X . The character of X at a point p , denoted by $\chi(p, X)$, is the infimum of the cardinalities of neighborhood bases at x and the *character* of space X , denoted by $\chi(X)$, is the supremum of $\chi(p, X)$, where $p \in X$. A space X is called *first (second) countable* if $w(X)$ ($\chi(X)$) is countable. The *cellularity* of X , denoted by $c(X)$, is defined by

$$\sup\{|\mathcal{U}| : \mathcal{U} \text{ is a family of mutually disjoint nonempty open subsets of } X\}.$$

For any $f \in C(X)$, we denote $f^{-1}\{0\}$ and $X \setminus f^{-1}\{0\}$ by $Z(f)$ and $\text{Coz}(f)$, respectively. Every set of the form $Z(f)$ ($\text{Coz}(f)$) is called *zeroset (cozeroset)*. A subset S of X is C -embedded in X if for every f in $C(S)$, there exists g in $C(X)$ such that $g|_S = f$. It is clear that every clopen subset of X is C -embedded in X . Suppose $p \in \beta X$, then by M^p we mean the set $\{f \in C(X) : p \in \text{cl}_{\beta X} Z(f)\}$. By [18, Theorem 7.3 (Gelfand-Kolmogoroff)], $\{M^p : p \in \beta X\}$ is the family of all maximal ideal of $C(X)$. X is a P -space if every prime ideal of X is maximal and we say that X is an *almost P -space* if the interior of every nonempty zeroset of X is nonempty. It is easy to check that X is an almost P -space if and only if $C(X)$ is an almost regular ring. By [18, Theorem 14.28], X is a P -space if and only if every zeroset of X is open. For more details we refer the reader to [15], [18], [11] and [29].

The study of translating graph properties to algebraic properties is an interesting subject for mathematicians. In [14], linear algebra and some properties of polynomials were used to describe properties of graphs. In [13], the studying of zero-divisor graph of commutative rings has been started. The investigation on zero-divisor graph of commutative rings was then continued in [7], [10], [20], [25], [4], [9], [6], and [8].

In [27], comaximal graph of a commutative ring was defined. On later, in [21], [26], [16], [28], [24], [22], [19], [2], [3], [1], [30], and [23], this investigation was continued.

In [12] and in a section of [5] the zero-divisor graph and the comaximal ideal graph of $C(X)$ were studied, respectively. These investigations tried to associate the ring properties of $C(X)$, the graph properties of graphs on $C(X)$ and the topological properties of X .

In this article we study the $\Gamma'_2 C(X)$. Since $J(C(X)) = 0$, so $\Gamma'_2 C(X) = \Gamma_2(C(X)) - \{0\}$. If X is singleton, then $\Gamma'_2 C(X)$ is empty. Thus, subsequently we assume $|X| > 1$.

By [21, Theorem 3.1, Lemma 3.2 and Proposition 3.3] and [24, Corollary 3.4], we can conclude the following.

Proposition 1.1. *For each Tychonoff space X ,*

- (a) $\Gamma'_2 C(X)$ is connected;
- (b) $\text{diam } \Gamma'_2 C(X) = 3$;
- (c) if X is infinite, then $\text{girth } \Gamma'_2 C(X) = 3$.

Lemma 1.2. *Suppose $f, g \in \Gamma'_2 C(X)$. Then f is adjacent to g if and only if $Z(f) \cap Z(g) = \emptyset$.*

PROOF: f is not adjacent to g if and only if both f and g are contained in a maximal ideal, that is

$$\begin{aligned} \exists p \in \beta X \quad f, g \in M^p &\Leftrightarrow \exists p \in \beta X \quad p \in \text{cl}_{\beta X} Z(f) \wedge p \in \text{cl}_{\beta X} Z(g) \\ &\Leftrightarrow \exists p \in \text{cl}_{\beta X} Z(f) \cap \text{cl}_{\beta X} Z(g) = \text{cl}_{\beta X} (Z(f) \cap Z(g)) \\ &\Leftrightarrow Z(f) \cap Z(g) \neq \emptyset. \quad \square \end{aligned}$$

In Section 2 we investigate the radius of $\Gamma'_2 C(X)$ and show that $2 \leq \text{Rad } \Gamma'_2 C(X) \leq 3$. The girth of this graph is investigated in Section 3 and we show that if $|X| > 2$, then $\text{girth } \Gamma'_2 C(X) = 3$. In Section 4 we study the dominating number and the clique number of the graph $\Gamma'_2 C(X)$. We prove that $d(X) \leq \text{dt } \Gamma'_2 C(X) \leq w(X)$, introduce zeroset cellularity of X and show that it is equal to $\text{clique } \Gamma'_2 C(X)$. In Section 5 we use the notions of the previous sections to associate the topological properties of X , the ring properties $C(X)$ and the graph properties of $\Gamma'_2 C(X)$. In this section we observe that $\Gamma'_2 C(X)$ is triangulated (hypertriangulated, complemented) if and only if X does not have any isolated points (X is connected, X is a P -space), and finally we conclude that X is an almost P -space which does not have isolated points if and only if $C(X)$ is regular ring which does not have any principal maximal ideals if and only if $\text{Rad } \Gamma'_2 C(X) = 3$.

Similar results to Theorem 4.4, Proposition 4.7 and Corollary 4.8 devoted to zero divisor graphs may be found in [12]. Here we prove them for comaximal graphs.

2. Radius of the graph

Lemma 2.1. *For any f and g in $\Gamma'_2 C(X)$*

- (a) $d(f, g) = 1$ if and only if $Z(f) \cap Z(g) = \emptyset$;
- (b) $d(f, g) = 2$ if and only if $Z(f) \cap Z(g) \neq \emptyset$ and $Z(f) \cup Z(g) \neq X$;
- (c) $d(f, g) = 3$ if and only if $Z(f) \cap Z(g) \neq \emptyset$ and $Z(f) \cup Z(g) = X$.

PROOF: (a) By Lemma 1.2, it is clear.

(b) \Rightarrow Since $d(f, g) = 2$, f is not adjacent to g , thus $Z(f) \cap Z(g) \neq \emptyset$, by Lemma 1.2. We now show that $Z(f) \cup Z(g) \neq X$. Suppose that, on the contrary $Z(f) \cup Z(g) = X$. From $d(f, g) = 2$ it follows that h in $\Gamma'_2 C(X)$ exists such that h is adjacent to both f and g , hence by Lemma 1.2,

$$\begin{cases} Z(h) \cap Z(f) = \emptyset \\ Z(h) \cap Z(g) = \emptyset \end{cases} \Rightarrow Z(h) = Z(h) \cap X = Z(h) \cap [Z(f) \cup Z(g)] = \emptyset$$

which is a contradiction.

(b) \Leftarrow Since $Z(f) \cap Z(g) \neq \emptyset$, $d(f, g) > 1$, by Lemma 1.2. Since $Z(fg) = Z(f) \cup Z(g) \neq X$, $p \in X \setminus Z(fg)$ exists, hence there is some h in $C(X)$ such that $p \in Z(h)$ and $Z(fg) \cap Z(h) = \emptyset$, thus $h \in \Gamma'_2 C(X)$ and

$$\begin{aligned} \emptyset &= [Z(f) \cup Z(g)] \cap Z(h) = [Z(f) \cap Z(h)] \cup [Z(g) \cap Z(h)] \\ &\Rightarrow \begin{cases} Z(f) \cap Z(h) = \emptyset \\ Z(g) \cap Z(h) = \emptyset. \end{cases} \end{aligned}$$

Hence h is adjacent to both f and g , thus $d(f, g) = 2$.

(c) By Proposition 1.1(b), it is clear. □

Lemma 2.2. For every $f \in \Gamma'_2 C(X)$, $\text{ecc}(f) \geq 2$.

PROOF: Since $Z(f) \cap Z(2f) = Z(f) \neq \emptyset$ and $Z(f) \cup Z(2f) = Z(f) \neq X$, $d(f, 2f) = 2$, by Lemma 2.1. This implies that $\text{ecc}(f) \geq 2$. □

Proposition 2.3. Suppose $f \in \Gamma'_2 C(X)$. Then $\text{ecc}(f) = 2$ if and only if either $\text{int}Z(f) = \emptyset$ or $Z(f) = \{p\}$, in which p is an isolated point.

PROOF: \Rightarrow By Lemma 2.2, $d(f, g) \neq 3$, for every $g \in \Gamma'_2 C(X)$. From Lemma 2.1, it follows that

$$\begin{aligned} &\forall g \in \Gamma'_2 C(X) \quad Z(f) \cup Z(g) \neq X \vee Z(f) \cap Z(g) = \emptyset \\ \equiv &\forall g \in \Gamma'_2 C(X) \quad Z(f) \cup Z(g) = X \Rightarrow Z(f) \cap Z(g) = \emptyset \\ \equiv &\forall g \in \Gamma'_2 C(X) \quad \text{Coz}(g) \subseteq Z(f) \Rightarrow Z(f) = \text{Coz}(g) \quad (1) \end{aligned}$$

If $\text{int}Z(f) \neq \emptyset$, then $Z(f)$ is open. It is sufficient to show that $Z(f)$ is singleton. Suppose, on the contrary, there are two distinct points p and q in $Z(f)$, thus there is a function $h : Z(f) \rightarrow \mathbb{R}$, such that $h(p) = 0$ and $h(q) = 1$. Since $Z(f)$ is clopen, $Z(f)$ is C -embedded in X , thus k in $C(X)$ exists such that $k|_{Z(f)} = h$.

Let $g : X \rightarrow \mathbb{R}$ be given by

$$g(x) = \begin{cases} 1 & x \in Z(f) \\ 0 & x \notin Z(f). \end{cases}$$

Since $Z(f)$ is clopen, $g \in \Gamma'_2 C(X)$ and therefore $gk \in \Gamma'_2 C(X)$. $\text{Coz}(gk) = \text{Coz}(g) \cap \text{Coz}(k) \subseteq Z(f)$, but $Z(f) \neq \text{Coz}(gk)$ since $p \in Z(f) \setminus \text{Coz}(gk)$. This contradicts the fact (1).

\Leftarrow By Proposition 1.1, it suffices to prove that

$$\forall g \in \Gamma'_2 C(X) \quad d(f, g) \neq 3.$$

According to the first part of the proof, the above statement is equivalent to

$$\forall g \in \Gamma'_2 C(X) \quad \text{Coz}(g) \subseteq Z(f) \Rightarrow Z(f) = \text{Coz}(g).$$

By the assumption, the above statement is clear. □

An immediate conclusion of Proposition 1.1, and Lemma 2.2, is the following corollary.

Corollary 2.4. $2 \leq \text{Rad } \Gamma'_2 C(X) \leq 3.$

3. Girth of the graph

Lemma 3.1. *Let $f \in \Gamma'_2 C(X)$. Then $\text{Coz}(f)$ is not singleton if and only if f is a vertex of a triangle.*

PROOF: \Rightarrow Let p and q be distinct elements of $\text{Coz}(f)$. There are two disjoint zerosets Z_1 and Z_2 containing p and q , respectively. Since $p, q \notin Z(f)$, there are two zerosets Z_3 and Z_4 containing p and q , respectively, such that $Z_3 \cap Z(f) = Z_4 \cap Z(f) = \emptyset$. Put $Z(g) = Z_3 \cap Z_1$ and $Z(h) = Z_4 \cap Z_2$. Consequently, $g, h \in \Gamma'_2 C(X)$, $Z(f) \cap Z(g) = \emptyset$, $Z(g) \cap Z(h) = \emptyset$ and $Z(h) \cap Z(f) = \emptyset$. Lemma 1.2 now shows that f is adjacent to g , g is adjacent to h and h is adjacent to f , thus f is vertex of a triangle.

\Leftarrow There are vertices g and h in $\Gamma'_2 C(X)$ such that f is adjacent to g , g is adjacent to h and h is adjacent to f . By Lemma 1.2

$$\begin{cases} Z(f) \cap Z(g) = \emptyset \\ Z(f) \cap Z(h) = \emptyset \\ Z(g) \cap Z(h) = \emptyset \end{cases} \Rightarrow \begin{cases} \emptyset \neq Z(g) \subseteq \text{Coz}(f) \\ \emptyset \neq Z(h) \subseteq \text{Coz}(f) \\ Z(g) \cap Z(h) = \emptyset. \end{cases}$$

Hence $\text{Coz}(f)$ is not singleton. □

Theorem 3.2. *If $|X| > 2$, then $\text{girth } \Gamma'_2 C(X) = 3.$*

PROOF: Since X has some non-singleton cozeroset, $\text{girth } \Gamma'_2 C(X) = 3$, by Lemma 3.1. □

Example 3.3. If $|X| > 2$ and finite, then $C(X)$ has finitely many maximal ideal and $\text{girth } \Gamma'_2 C(X) = 3$, by Theorem 3.2. This is a counterexample to the converse of [24, Corollary 3.4].

4. Dominating and clique number

Lemma 4.1. *Let $f, g \in \Gamma'_2 C(X)$.*

- (a) *If $Z(f) \cap Z(g) = \emptyset$ and $Z(f) \cup Z(g) = X$, then $\text{gi}(f, g) = 4.$*
- (b) *If $Z(f) \cap Z(g) = \emptyset$ and $Z(f) \cup Z(g) \neq X$, then $\text{gi}(f, g) = 3.$*

- (c) If $Z(f) \cap Z(g) \neq \emptyset$ and $Z(f) \cup Z(g) \neq X$, then $\text{gi}(f, g) = 4$.
- (d) If $Z(f) \cap Z(g) \neq \emptyset$ and $Z(f) \cup Z(g) = X$, then $\text{gi}(f, g) = 6$.

PROOF: (a) $Z(f) \cap Z(g) = \emptyset$, $Z(g) \cap Z(2f) = \emptyset$, $Z(2f) \cap Z(2g) = \emptyset$ and $Z(2g) \cap Z(f) = \emptyset$. By Lemma 1.2, f is adjacent to g , g is adjacent to $2f$, $2f$ is adjacent to $2g$ and $2g$ is adjacent to f , it follows that $\text{gi}(f, g) \leq 4$. We claim that $\text{gi}(f, g) \neq 3$ and therefore $\text{gi}(f, g) = 4$. On the contrary, suppose $\text{gi}(f, g) = 3$, then h in $\Gamma'_2 C(X)$ exists such that h is adjacent to both f and g , by Lemma 1.2

$$\begin{cases} Z(h) \cap Z(f) = \emptyset \\ Z(h) \cap Z(g) = \emptyset \end{cases} \Rightarrow Z(h) = Z(h) \cap X = Z(h) \cap [Z(f) \cup Z(g)] = \emptyset$$

which is impossible.

(b) Suppose $x \in X \setminus [Z(f) \cup Z(g)]$. There is some h in $\Gamma'_2 C(X)$ such that $x \in Z(h)$ and

$$Z(h) \cap [Z(f) \cup Z(g)] = \emptyset \Rightarrow Z(h) \cap Z(f) = \emptyset \text{ and } Z(h) \cap Z(g) = \emptyset$$


thus $h \in \Gamma'_2 C(X)$ and h is adjacent to both f and g , by Lemma 1.2, hence $\text{gi}(f, g) = 3$.

(c) Suppose $x \in X \setminus [Z(f) \cup Z(g)]$, then there is some $h \in \Gamma_2 C(X)$, such that $x \in Z(h)$ and


$$Z(h) \cap [Z(f) \cup Z(g)] = \emptyset \Rightarrow Z(h) \cap Z(f) = \emptyset \text{ and } Z(h) \cap Z(g) = \emptyset$$

thus $Z(f) \cap Z(2h) = \emptyset$ and $Z(2h) \cap Z(g) = \emptyset$. From Lemma 4.1, we deduce that f is adjacent to h , h is adjacent to g , g is adjacent to $2h$ and $2h$ is adjacent to f , this gives $\text{gi}(f, g) \leq 4$. Since $Z(f) \cap Z(g) \neq \emptyset$, so f is not adjacent to g and therefore $\text{gi}(f, g) \neq 3$, and so $\text{gi}(f, g) = 4$.


(d) By Lemma 2.1, $d(f, g) = 3$, thus $\text{gi}(f, g) \leq 6$ and there are h and k in $\Gamma'_2 C(X)$ such that f is adjacent to h , h is adjacent to k and k is adjacent to g . It is easily seen that g is adjacent to $2k$, $2k$ is adjacent to $2h$ and $2h$ is adjacent to f . This clearly forces $\text{gi}(f, g) = 6$. □




$$\begin{aligned} Z(f) \cap Z(g) &= \emptyset \\ Z(f) \cup Z(g) &= X \\ \text{gi}(f, g) &= 4 \end{aligned}$$



$$\begin{aligned} Z(f) \cap Z(g) &= \emptyset \\ Z(f) \cup Z(g) &\neq X \\ \text{gi}(f, g) &= 3 \end{aligned}$$



$$\begin{aligned} Z(f) \cap Z(g) &\neq \emptyset \\ Z(f) \cup Z(g) &\neq X \\ \text{gi}(f, g) &= 4 \end{aligned}$$



$$\begin{aligned} Z(f) \cap Z(g) &\neq \emptyset \\ Z(f) \cup Z(g) &= X \\ \text{gi}(f, g) &= 6 \end{aligned}$$

- Corollary 4.2.** (a) Every cycle in $\Gamma'_2 C(X)$ has length 3 or 4.
 (b) Every edge of $\Gamma'_2 C(X)$ is edge of a cycle with length 3 or 4.
 (c) Every vertex of $\Gamma'_2 C(X)$ is vertex of a square.

PROOF: (a) and (b) are immediate conclusions of Lemma 4.1.

(c) For each $f \in \Gamma'_2 C(X)$, we have

$$\begin{cases} Z(f) \cap Z(2f) \neq \emptyset \\ Z(f) \cup Z(2f) \neq X. \end{cases}$$

By Lemma 4.1, $\text{gi}(f, 2f) = 4$, and therefore f is vertex of a square. □

Lemma 4.3. *If X is an infinite space, then every dominating set of $\Gamma'_2 C(X)$ is infinite.*

PROOF: We show that none of the finite subsets of $\Gamma'_2 C(X)$ is a dominating set. Suppose $A = \{f_1, f_2, \dots, f_n\}$ is a finite subset of $\Gamma'_2 C(X)$. Each $Z(f_i)$ is nonempty, thus p_i in $Z(f_i)$ exists. Since X is infinite, p_0 in X distinct from p_i 's exists. Thus there are zerosets Z_0, Z_1, \dots, Z_n in $\Gamma'_2 C(X)$ such that $p_i \in Z_i$, for every $0 \leq i \leq n$, and $i \neq j$ implies $Z_i \cap Z_j = \emptyset$. Set $Z(g) = Z_1 \cup Z_2 \cup \dots \cup Z_n$. Then $p_0 \notin Z(g) \neq X$ and $p_i \in Z(g) \cap Z(f_i) \neq \emptyset$, thus $g \in \Gamma'_2 C(X)$ and not adjacent to any f_i . This follows that A is not a dominating set. □

Theorem 4.4. $d(X) \leq \text{dt} \Gamma'_2 C(X) \leq w(X)$. In particular, whenever $d(X) = w(X)$, then $\text{dt} \Gamma'_2 C(X) = w(X)$.

PROOF: If X is finite, then it is easy to check $d(X) = \text{dt} \Gamma'_2 C(X) = w(X)$, thus we assume X is infinite. Let A be a dominating set in $\Gamma'_2 C(X)$. For each $f \in A$, we pick $x_f \in Z(f)$ and $y_f \in \text{Coz}(f)$. Set $D = \{x_f : f \in A\} \cup \{y_f : f \in A\}$. For every cozeroset $\text{Coz}(g)$, if $g \in A$, then $y_g \in D \cap \text{Coz}(g)$, if $g \notin A$, then $f \in A$ exists such that

$$Z(f) \cap Z(g) = \emptyset \Rightarrow Z(f) \subseteq \text{Coz}(g) \Rightarrow x_f \in Z(f) \cap D \subseteq \text{Coz}(g) \cap D.$$

Hence D is dense in X . Since D is infinite, $d(X) \leq \text{dt} \Gamma'_2 C(X)$.

We now suppose that \mathcal{B} is a base for X . Without loss of generality we can assume that \mathcal{B} does not have any empty members. Then for every $B \in \mathcal{B}$, there

is some f_B in $\Gamma'_2 C(X)$ such that $Z(f_B) \subseteq B$. For each f in $\Gamma'_2 C(X)$ there is some B in \mathcal{B} such that

$$Z(f_B) \subseteq B \subseteq \text{Coz}(f) \Rightarrow Z(f_B) \cap Z(f) = \emptyset.$$

By Lemma 1.2, f is adjacent to f_B . Therefore $\{f_B : B \in \mathcal{B}\}$ is a dominating set and finally that $\text{dt } \Gamma'_2 C(X) \leq w(X)$. \square

An immediate conclusion of the above theorem is the following corollary.

Corollary 4.5. *If X is an infinite second countable space, then $\text{dt } \Gamma'_2 C(X) = \omega$.*

Example 4.6. Let X be Moore plane. For every (x_o, y_o) in X , set $f_{x_o, y_o} : X \rightarrow \mathbb{R}$ as $f_{x_o, y_o}(x, y) = \sqrt{(x - x_o)^2 + (y - y_o)^2}$. It is clear that $f \in \Gamma'_2 C(X)$ and $Z(f_{x_o, y_o}) = \{(x_o, y_o)\}$. Suppose $A = \{f_{x,y} : x, y \in \mathbb{Q} \text{ and } y > 0\}$. If $Z(f) \cap Z(f_{x,y}) \neq \emptyset$, for each $f_{x,y} \in A$, then $\mathbb{Q} \times \mathbb{Q}^{>0} \subseteq Z(f)$ and therefore $XZ(f)$. This implies that A is a dominating set and therefore $\text{dt } \Gamma'_2 C(X) = \omega \neq \mathbf{c} = \omega(X)$.

Proposition 4.7. *Suppose $\Gamma C(X)$ is the zero divisor graph of $C(X)$. If $\chi(X) \leq d(X)$, then $\text{dt } \Gamma'_2 C(X) = d(X) = \text{dt } \Gamma C(X)$.*

PROOF: According to Theorem 4.4, we only need to show that $d(X) \geq \text{dt } \Gamma'_2 C(X)$. Clearly, if X is finite, then $\text{dt } \Gamma'_2 C(X) = d(X)$.

Now suppose X is infinite, then every dominating set is infinite, by Lemma 4.3. Let D be a dense subset of X and \mathfrak{B}_x is a neighborhood base at x , for each x in D . For every $x \in D$ and $B \in \mathfrak{B}_x$, there is some $f_{x,B} \in \Gamma'_2 C(X)$ such that $x \in Z(f_{x,B}) \subseteq B$. Put $A = \{f_{x,B} : x \in D \text{ and } B \in \mathfrak{B}_x\}$. If $g \in \Gamma'_2 C(X)$, then $\text{Coz}(g) \neq \emptyset$, and it follows that $x \in D \cap \text{Coz}(g)$ exists. Hence there is a $B \in \mathfrak{B}_x$ such that $Z(f_{B,x}) \subseteq B \subseteq \text{Coz}(g)$, thus $Z(f_{B,x}) \cap Z(g) = \emptyset$, and, in consequence, $f_{B,x}$ is adjacent to g . This implies that A is dominating set. Since $|A| \leq \chi(X)|D| \leq d(X)d(X) = d(X)$, $d(X) \geq \text{dt } \Gamma'_2 C(X)$. The equality $\text{dt } \Gamma C(X) = d(X)$ was shown in [12, Proposition 3.4]. \square

By [17, Thorem 1.5.7], $w(X) \leq \exp d(X)$, hence the following corollary is immediate.

Corollary 4.8. $d(X) \leq \text{dt } \Gamma'_2 C(X) \leq \exp d(X)$.

Definition 4.9. We define zero cellularity of X , denoted by $zc(X)$, by the supremum of $\{|\mathcal{Z}| : \mathcal{Z} \text{ is a family of pairwise disjoint nonempty zero subsets of } X\}$.

Theorem 4.10. *We have $\text{clique } \Gamma'_2 C(X) = zc(X) \leq |X|$. In particular if X is first countable, then $\text{clique } \Gamma'_2 C(X) = zc(X) = |X|$.*

PROOF: By Lemma 1.2, $A \subseteq \Gamma'_2 C(X)$ is a complete subgraph of $\Gamma'_2 C(X)$ if and only if $Z(A) = \{Z(f) : f \in A\}$ is a family of pairwise disjoint zerosets, thus clique number of $\Gamma'_2 C(X)$ is the supremum of

$$\{|\mathcal{Z}| : \mathcal{Z} \text{ is a family of pairwise disjoint zero sets of } X\},$$

hence $\text{clique } \Gamma'_2 C(X) = zc(X)$. It is clear that $zc(X) \leq |X|$.

If X is first countable, then for every p in X , $\{p\}$ is a zeroset and thus $zc(X) = |X|$. \square

5. Some applications

Theorem 5.1. $\Gamma'_2 C(X)$ is triangulated if and only if X does not have any isolated points.

PROOF: \Rightarrow If p is an isolated point of X , then there is some f in $\Gamma'_2 C(X)$ such that $\text{Coz}(f) = \{p\}$, thus f is not a vertex of a triangle, by Lemma 3.1. Consequently, $\Gamma'_2 C(X)$ is not triangulated.

\Leftarrow Suppose $\Gamma'_2 C(X)$ is not triangulated, hence there is some f in $\Gamma'_2 C(X)$ such that f is not a vertex of any triangle. By Lemma 3.1, $\text{Coz}(f) = \{p\}$ is singleton, hence p is an isolated point of X . \square

Theorem 5.2. $\Gamma'_2 C(X)$ is hypertriangulated if and only if X is connected.

PROOF: \Rightarrow If X is disconnected, then there are zerosets $Z(f)$ and $Z(g)$ such that $Z(f) \cap Z(g) = \emptyset$ and $Z(f) \cup Z(g) = X$. From Lemma 4.1, $\text{gi}(f, g) = 4$, which yields $\{f, g\}$ is not edge of any triangle, and therefore $\Gamma'_2 C(X)$ is not hypertriangulated.

\Leftarrow Suppose $\Gamma'_2 C(X)$ is not hypertriangulated. Then there is an edge $\{f, g\}$ of $\Gamma'_2 C(X)$, which is not an edge of any triangle, thus $\text{gi}(f, g) = 4$. Lemma 4.1 now shows that $Z(f) \cup Z(g) = X$ and $Z(f) \cap Z(g) = \emptyset$, this implies X is disconnected. \square

Theorem 5.3. $\Gamma'_2 C(X)$ is complemented if and only if X is a P -space.

PROOF: \Rightarrow For every zeroset $Z(f)$ there is a zeroset $Z(g)$ such that $f \perp g$, thus $\text{gi}(f, g) = 3$. We conclude from Lemma 4.1 that $\text{Coz}(f) \cap \text{Coz}(g) = \emptyset$ and $\text{Coz}(f) \cup \text{Coz}(g) = X$, therefore $Z(f)$ is open. This follows that X is a P -space.

\Leftarrow For every vertex f in $\Gamma'_2 C(X)$, $Z(f)$ is open, thus g in $\Gamma'_2 C(X)$ exists such that $Z(f) \cap Z(g) = \emptyset$ and $Z(f) \cup Z(g) = X$. Now Lemma 4.1 becomes $\text{gi}(f, g) = 3$, thus $f \perp g$ and consequently $\Gamma'_2 C(X)$ is complemented. \square

Lemma 5.4. Suppose M^p is a maximal ideal of $C(X)$, for some $p \in \beta X$. Then M^p is principal if and only if p is an isolated point.

PROOF: \Rightarrow Let $M^p = \langle f \rangle$, for some $f \in C(X)$. $Z(f) = \{p\}$, since $\bigcap_{Z \in Z(M^p)} Z = Z(f)$. If p is not an isolated point, then $p \in \overline{\text{Coz}(f)}$, and therefore there is a net (x_λ) in $\text{Coz}(f)$, which converges to p . We conclude from $Z(f^{\frac{1}{3}}) = Z(f)$ that there is some g in $C(X)$ such that $f^{\frac{1}{3}} = gf$, hence that $g(x) = 1/f^{\frac{2}{3}}(x)$, for each x in $\text{Coz}(f)$, therefore $g(p) = \lim g(x_\lambda) = \infty$, which is a contradiction.

\Leftarrow It is straightforward. \square

Proposition 5.5. Let p be a G_δ -point of X . If $Z(f) = \{p\}$, then

$$M^p = \{g \in \Gamma'_2 C(X) : d(f, g) = 2\} \cup \{0, f\}.$$

PROOF: Set $I = \{g \in \Gamma'_2 C(X) : d(f, g) = 2\} \cup \{0, f\}$. If $g \in M^p$, then $p \in Z(g)$, hence $Z(f) \cup Z(g) \neq X$ and $Z(f) \cap Z(g) \neq \emptyset$. Lemma 2.1 shows that $d(f, g) = 2$, and therefore $M^p \subseteq I$ (1).

Now suppose $g \in I$. Since $d(f, g) = 2$, $\{p\} \cap Z(g) = Z(f) \cap Z(g) \neq \emptyset$. This implies $p \in Z(g)$, and thus $g \in M^p$. Hence $I \subseteq M^p$ (2). By (1) and (2), $M^p = I$. □

Theorem 5.6. *The following are equivalent.*

- (a) X is an almost P -space which does not have any isolated points.
- (b) $C(X)$ is almost regular ring which does not have any principal maximal ideals.
- (c) $\text{Rad } \Gamma'_2 C(X) = 3$.
- (d) For each $f \in \Gamma'_2 C(X)$, there is some $g \in \Gamma'_2 C(X)$ such that $\text{gi}(f, g) = 6$.

PROOF: (a) \Leftrightarrow (b) By Lemma 5.4 it is obvious.

(a) \Rightarrow (c) For each $f \in \Gamma'_2 C(X)$, $\text{int}Z(f) \neq \emptyset$, thus $\text{ecc}(f) = 3$. Hence $\text{Rad } \Gamma'_2 C(X) = 3$.

(c) \Rightarrow (a) Since $\text{Rad } \Gamma'_2 C(X) = 3$, for every $f \in \Gamma'_2 C(X)$ we have $\text{ecc}(f) = 3$. We conclude from Proposition 2.3, that $\text{int}Z(f) \neq \emptyset$ and X does not have any isolated points, hence that X is an almost P -space without any isolated points.

(d) \Rightarrow (b) For each $f \in \Gamma'_2 C(X)$, there is some $g \in \Gamma'_2 C(X)$ such that $\text{gi}(f, g) = 6$, thus $Z(f) \cap Z(g) \neq \emptyset$ and $Z(f) \cup Z(g) = X$. We conclude that f is a zero divisor and consequently $C(X)$ is an almost regular ring.

(c) \Rightarrow (d) Since $\text{Rad } \Gamma'_2 C(X) = 3$, for each $f \in \Gamma'_2 C(X)$ we have $\text{ecc}(f) = 3$, thus g in $\Gamma'_2 C(X)$ exists such that $d(f, g) = 3$ and therefore $\text{gi}(f, g) = 6$. □

Proposition 5.7. *The following statements are equivalent.*

- (a) $C(X)$ is almost regular ring which has some principal maximal ideal.
- (b) X is an almost P -space which has some isolated point.
- (c) $\text{Rad } \Gamma'_2 C(X) = 2$ and for each f in the center of $\Gamma'_2 C(X)$ there is $g \in \Gamma'_2 C(X)$ such that $\{f, g\}$ is an edge of a square.

PROOF: (a) \Leftrightarrow (b) By Lemma 5.4, it is evident.

(b) \Rightarrow (c) By Theorem 5.6, $\text{Rad } \Gamma'_2 C(X) = 2$, thus for every f in center of $\Gamma'_2 C(X)$, $\text{ecc}(f) = 2$. Theorem 5.4 shows that $Z(f) = \{p\}$, for some isolated point p . Hence $g \in \Gamma'_2 C(X)$ exists such that $\text{Coz}(g) = \{p\}$. Since $Z(f) \cap Z(g) = \emptyset$ and $Z(f) \cup Z(g) = X$, by Lemma 4.1, $\{f, g\}$ is an edge of a square.

(c) \Rightarrow (b) If $Z(f) = \emptyset$, for some $f \in \Gamma'_2 C(X)$, then f belongs to the center of $\Gamma'_2 C(X)$. This implies that there is a $g \in \Gamma'_2 C(X)$ such that $\{f, g\}$ is an edge of some square, thus $Z(f)$ is open, by Lemma 4.1, a contradiction. Since $\text{Rad } \Gamma'_2 C(X) = 2$, from Theorem 5.6, X has some isolated point. □

Acknowledgments. We are very grateful to the referee whose valuable suggestions and comments improved an earlier version of this article.

REFERENCES

- [1] Afkhami M., Barati Z., Khashyarmanesh K., *When the comaximal and zero-divisor graphs are ring graphs and outerplanar*, Rocky Mountain J. Math. **44** (2014), no. 6, 1745–1761.
- [2] Afkhami M., Khashyarmanesh K., *On the cozero-divisor graphs and comaximal graphs of commutative rings*, J. Algebra Appl. **12** (2013), no. 3, 1250173, 9pp.
- [3] Akbari S., Habibi M., Majidinya A., Manaviyat R., *A note on comaximal graph of non-commutative rings*, Algebr. Represent. Theory **16** (2013), no. 2, 303–307.
- [4] Akbari S., Maimani H.R., Yassemi S., *When a zero-divisor graph is planar or a complete r -partite graph*, J. Algebra **270** (2003), no. 1, 169–180.
- [5] Amini A., Amini B., Momtahan E., Shirdareh Haghighi M.H., *On a graph of ideals*, Acta Math. Hungar. **134** (2011), no. 3, 369–384.
- [6] Anderson D.F., Mulay S.B., *On the diameter and girth of a zero-divisor graph*, J. Pure Appl. Algebra **210** (2007), no. 2, 543–550.
- [7] Anderson D.D., Naseer M., *Beck's coloring of a commutative ring*, J. Algebra **159** (1993), no. 2, 500–514.
- [8] Anderson D.F., Badawi A., *On the zero-divisor graph of a ring*, Comm. Algebra **36** (2008), no. 8, 3073–3092.
- [9] Anderson D.F., Levy R., Shapiro J., *Zero-divisor graphs, von Neumann regular rings, and Boolean algebras*, J. Pure Appl. Algebra **180** (2003), no. 3, 221–241.
- [10] Anderson D.F., Livingston P.S., *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), no. 2, 434–447.
- [11] Atiyah M.F., Macdonald I.G., *Introduction to Commutative Algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [12] Azarpanah F., Motamedi M., *Zero-divisor graph of $C(X)$* , Acta Math. Hungar. **108** (2005), no. 1–2, 25–36.
- [13] Beck I., *Coloring of commutative rings*, J. Algebra **116** (1988), no. 1, 208–226.
- [14] Biggs N., *Algebraic Graph Theory*, Cambridge University Press, Cambridge, 1993.
- [15] Bondy J.A., Murty U.S.R., *Graph Theory with Application*, The Macmillan Press, New York, 1976.
- [16] Dheena P., Elavarasan B., *On comaximal graphs of near-rings*, Kyungpook Math. J. **49** (2009), no. 2, 283–288.
- [17] Engelking R., *General Topology*, Heldermann-Verlag, Berlin, 1989.
- [18] Gillman L., Jerison M., *Rings of Continuous Functions*, Transactions of the New York Academy of Sciences **27** (1964), no. 1 Series II, 5–6.
- [19] Jinnah M.I., Mathew Sh.C., *When is the comaximal graph split?*, Comm. Algebra **40** (2012), no. 7, 2400–2404.
- [20] Levy R., Shapiro J., *The zero-divisor graph of von Neumann regular rings*, Comm. Algebra **30** (2002), no. 2, 745–750.
- [21] Maimani H.R., Salimi M., Sattari A., Yassemi S., *Comaximal graph of commutative rings*, J. Algebra **319** (2008), no. 4, 1801–1808.
- [22] Maimani H.R., Pournaki M.R., Tehrani A., Yassemi S., *Graphs attached to rings revisited*, Arab. J. Sci. Eng. **36** (2011), no. 6, 997–1011.
- [23] Mehdi-Nezhad E., Rahimi A.M., *Dominating sets of the comaximal and ideal-based zero-divisor graphs of commutative rings*, Quaest. Math. **38** (2015), 1–17.
- [24] Moconja S.M., Petrović Z., *On the structure of comaximal graphs of commutative rings with identity*, Bull. Aust. Math. Soc. **83** (2011), no. 1, 11–21.
- [25] Mulay Sh.B., *Cycles and symmetries of zero-divisors*, Comm. Algebra **30** (2002), no. 7, 3533–3558.
- [26] Petrovic Z.Z., Moconja S.M., *On graphs associated to rings*, Novi Sad J. Math. **38** (2008), no. 3, 33–38.
- [27] Sharma P.K., Bhatwadekar S.M., *A note on graphical representation of rings*, J. Algebra **176** (1995), no. 1, 124–127.

- [28] Wang H.-J., *Co-maximal graph of non-commutative rings*, Linear Algebra Appl. **430** (2009), no. 2, 633–641.
- [29] Willard S., *General Topology*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1970.
- [30] Ye M., Wu T., Liu Q., Yu H., *Implements of graph blow-up in co-maximal ideal graphs*, Comm. Algebra **42** (2014), no. 6, 2476–2483.

DEPARTMENT OF BASIC SCIENCES, JUNDI-SHAPUR UNIVERSITY OF TECHNOLOGY, DEZFUL, IRAN

E-mail: Badie@jsu.ac.ir

(Received December 6, 2015, revised March 22, 2016)