

Applications of Mathematics

Yusaku Yamamoto

On the optimality and sharpness of Laguerre's lower bound on the smallest eigenvalue of a symmetric positive definite matrix

Applications of Mathematics, Vol. 62 (2017), No. 4, 319–331

Persistent URL: <http://dml.cz/dmlcz/146832>

Terms of use:

© Institute of Mathematics AS CR, 2017

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE OPTIMALITY AND SHARPNESS OF LAGUERRE'S
LOWER BOUND ON THE SMALLEST EIGENVALUE
OF A SYMMETRIC POSITIVE DEFINITE MATRIX

YUSAKU YAMAMOTO, Tokyo

Received January 31, 2017. First published June 29, 2017.

Abstract. Lower bounds on the smallest eigenvalue of a symmetric positive definite matrix $A \in \mathbb{R}^{m \times m}$ play an important role in condition number estimation and in iterative methods for singular value computation. In particular, the bounds based on $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ have attracted attention recently, because they can be computed in $O(m)$ operations when A is tridiagonal. In this paper, we focus on these bounds and investigate their properties in detail. First, we consider the problem of finding the optimal bound that can be computed solely from $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ and show that the so called Laguerre's lower bound is the optimal one in terms of sharpness. Next, we study the gap between the Laguerre bound and the smallest eigenvalue. We characterize the situation in which the gap becomes largest in terms of the eigenvalue distribution of A and show that the gap becomes smallest when $\{\text{Tr}(A^{-1})\}^2/\text{Tr}(A^{-2})$ approaches 1 or m . These results will be useful, for example, in designing efficient shift strategies for singular value computation algorithms.

Keywords: eigenvalue bound; symmetric positive definite matrix; Laguerre bound; singular value computation; dqds algorithm

MSC 2010: 15A18, 15A42

1. INTRODUCTION

Let $A \in \mathbb{R}^{m \times m}$ be a symmetric positive definite matrix and denote its smallest eigenvalue by $\lambda_m(A)$. In this paper, we are interested in a lower bound on $\lambda_m(A)$. For the Cholesky factorization $A = BB^T$, where $B \in \mathbb{R}^{m \times m}$ is a nonsingular lower triangular matrix, the smallest singular value of B can be written as $\sigma_m(B) = \sqrt{\lambda_m(A)}$. Hence, finding a lower bound on $\lambda_m(A)$ is equivalent to finding a lower bound on $\sigma_m(B)$.

This study has been supported by JSPS KAKENHI Grant Numbers JP26286087, JP15H02708, JP15H02709, JP16KT0016, and 17H02828.

A lower bound on $\lambda_m(A)$ or $\sigma_m(B)$ plays an important role in various scientific computations. For example, when combined with an upper bound on $\|A\|_2$, a lower bound on $\lambda_m(A)$ can be used to give an upper bound on the condition number of A . In singular value computation algorithms such as the dqds algorithm [3], the orthogonal qd algorithm [10], and the mdLVs algorithm [6], a lower bound on $\sigma_m(B)$ is used as a shift to accelerate the convergence. In the latter case, the matrix B is usually a lower bidiagonal matrix as a result of preprocessing by the Householder method [4].

Several types of lower bounds on $\lambda_m(A)$ or $\sigma_m(B)$ have been proposed so far. There are bounds based on eigenvalue inclusion theorems such as Gershgorin's circle theorem [7] or Brauer's oval of Cassini [8]. The norm of the inverse, $\|A^{-1}\|_\infty$, can also be used to bound the maximum eigenvalue of A^{-1} from above, and therefore to bound $\lambda_m(A)$ from below. There are also bounds based on the traces of the inverses, namely, $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$. Among these, the latter class of bounds is attractive in the context of singular value computation, because they always give a valid (positive) lower bound, as opposed to the bounds based on the eigenvalue inclusion theorems, and they can be computed in $O(m)$ operations using efficient algorithms [9], [11], [13]. Examples of lower bounds of this type include the Newton bound [10], the generalized Newton bound [9], [1], and the Laguerre bound [10].

In this paper, we focus on the lower bounds of $\lambda_m(A)$ derived from $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ and investigate their properties. In particular, we will address the following two questions. The first is to identify an *optimal* formula for a lower bound on $\lambda_m(A)$ that is based solely on $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$. Here, the word "optimal" means that the formula always gives a sharper (that is, larger) bound than any other formula using only $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$. As a result of our analysis, we show that the Laguerre bound mentioned above is the optimal formula in this sense. The second question is to evaluate the *gap* between the Laguerre bound and $\lambda_m(A)$. Unlike the Laguerre bound, $\lambda_m(A)$ is not determined uniquely only from $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$. Hence, for some of the matrices, there must be a gap between the bound and $\lambda_m(A)$. Our problem is to quantify the maximum possible gap and identify the conditions under which the maximum gap is attained. These results will be useful, for example, in designing efficient shift strategies for singular value computation algorithms, which combine the Laguerre bound with other bounds with complementary characteristics [12].

The rest of this paper is structured as follows. In Section 2, we investigate the lower bounds on $\lambda_m(A)$ derived from $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ and show that the Laguerre bound is an optimal one in terms of sharpness. Section 3 deals with the gap between the Laguerre bound and $\lambda_m(A)$. In particular, we characterize the situation in which the gap becomes largest in terms of the eigenvalue distribution of A . Section 4 gives some concluding remarks.

2. AN OPTIMAL LOWER BOUND BASED ON $\text{Tr}(A^{-1})$ AND $\text{Tr}(A^{-2})$

2.1. Lower bounds based on $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$. Let A be an $m \times m$ real symmetric positive definite matrix. We denote the k th largest eigenvalue of A by $\lambda_k(A)$, or λ_k for short. Let $f(\lambda) = \det(\lambda I - A)$ be the characteristic polynomial of A . To find a lower bound on the smallest eigenvalue λ_m , we consider applying a root finding method for the algebraic equation to $f(\lambda) = 0$ starting from the initial value $\lambda^{(0)} = 0$. There are several root finding methods, such as the Bailey's (Halley's) method [2], Householder's method [5], and Laguerre's method [14], [10], for which the iteration formulas can be written as follows:

$$(2.1) \quad \lambda_B^{(n+1)} = \lambda^{(n)} - \frac{f(\lambda^{(n)})}{f'(\lambda^{(n)})} \left(1 - \frac{f(\lambda^{(n)})f''(\lambda^{(n)})}{2f'(\lambda^{(n)})^2} \right)^{-1},$$

$$(2.2) \quad \lambda_H^{(n+1)} = \lambda^{(n)} - \frac{f(\lambda^{(n)})}{f'(\lambda^{(n)})} \left(1 + \frac{f(\lambda^{(n)})f''(\lambda^{(n)})}{2f'(\lambda^{(n)})^2} \right),$$

$$(2.3) \quad \lambda_L^{(n+1)} = \lambda^{(n)} - \frac{f(\lambda^{(n)})}{f'(\lambda^{(n)})} \\ \times m \left(1 + \sqrt{(m-1) \left\{ m \cdot \frac{f'(\lambda^{(n)})^2 - f(\lambda^{(n)})f''(\lambda^{(n)})}{f'(\lambda^{(n)})^2} - 1 \right\}} \right)^{-1}.$$

Equations (2.1), (2.2), and (2.3) represent the iteration formulas of Bailey's method, Householder's method and Laguerre's method, respectively. When applied to $f(\lambda) = \det(\lambda I - A)$ starting from $\lambda^{(0)} = 0$, these formulas produce a sequence that increases monotonically and converges to λ_m . Hence, all of $\lambda_B^{(1)}$, $\lambda_H^{(1)}$, and $\lambda_L^{(1)}$ can be used as a lower bound on λ_m .

Noting that $f(\lambda) = \prod_{k=1}^m (\lambda - \lambda_k)$, we have

$$(2.4) \quad f'(\lambda) = - \sum_{k=1}^m \prod_{j \neq k} (\lambda_j - \lambda) \\ = - \prod_{j=1}^m (\lambda_j - \lambda) \sum_{k=1}^m \frac{1}{\lambda_k - \lambda} = -f(\lambda) \text{Tr}((A - \lambda I)^{-1}),$$

$$(2.5) \quad f''(\lambda) = -f'(\lambda) \text{Tr}((A - \lambda I)^{-1}) - f(\lambda) \sum_{k=1}^m \frac{1}{(\lambda_k - \lambda)^2} \\ = -f'(\lambda) \text{Tr}((A - \lambda I)^{-1}) - f(\lambda) \text{Tr}((A - \lambda I)^{-2}).$$

Hence,

$$(2.6) \quad \frac{f(\lambda)}{f'(\lambda)} = -\frac{1}{\text{Tr}((A - \lambda I)^{-1})},$$

$$(2.7) \quad \frac{f(\lambda)f''(\lambda)}{f'(\lambda)^2} = 1 - \frac{\text{Tr}((A - \lambda I)^{-2})}{\{\text{Tr}((A - \lambda I)^{-1})\}^2}.$$

Inserting these into equations (2.1), (2.2), and (2.3) with $\lambda^{(0)} = 0$, we obtain the following lower bounds on $\lambda_m(A)$:

$$(2.8) \quad L_B(A) = \frac{2\text{Tr}(A^{-1})}{\{\text{Tr}(A^{-1})\}^2 + \text{Tr}(A^{-2})},$$

$$(2.9) \quad L_H(A) = \frac{1}{\text{Tr}(A^{-1})} \left[\frac{3}{2} - \frac{1}{2} \frac{\text{Tr}(A^{-2})}{\{\text{Tr}(A^{-1})\}^2} \right],$$

$$(2.10) \quad L_L(A) = \frac{1}{\text{Tr}(A^{-1})} \cdot m \left(1 + \sqrt{(m-1) \left[m \cdot \frac{\text{Tr}(A^{-2})}{\{\text{Tr}(A^{-1})\}^2} - 1 \right]} \right)^{-1}$$

We call $L_B(A)$, $L_H(A)$, and $L_L(A)$ the Bailey bound, the Householder bound, and the Laguerre bound, respectively. In addition to these, we also have a simple bound:

$$(2.11) \quad L_N(A) = \{\text{Tr}(A^{-2})\}^{-1/2} \leq \left(\sum_{k=1}^m \frac{1}{\lambda_k^2} \right)^{-1/2} < \lambda_m,$$

which is called the Newton bound of order 2 (see [10], [9], [1]). In the case, where A is a tridiagonal matrix, both $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ can be computed in $O(m)$ operations from its Cholesky factor B (see [9], [11], [13]). Accordingly, any of these bounds can be employed in a practical shift strategy for singular value computation algorithms. The problem then is which of the four lower bounds, or possibly another bound derived from $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$, is optimal in terms of sharpness.

2.2. The optimal lower bound. To answer the question, we reformulate the problem as follows. Assume that $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ are specified for a symmetric positive definite matrix A . Then, how small can the smallest eigenvalue $\lambda_m(A)$ be? If this bound can be obtained explicitly as a function of $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$, then it will be the optimal formula for the lower bound of $\lambda_m(A)$.

Now, let $a \equiv \text{Tr}(A^{-1})$, $b \equiv \text{Tr}(A^{-2})$, and $x_k \equiv 1/\lambda_k$ ($k = 1, 2, \dots, m$). Then the upper bound on x_m (the reciprocal of the lower bound on λ_m) can be obtained by

solving the following constrained optimization problem:

$$(2.12) \quad \text{maximize } x_m$$

$$(2.13) \quad \text{s.t. } \sum_{k=1}^m x_k = a,$$

$$(2.14) \quad \sum_{k=1}^m x_k^2 = b,$$

$$(2.15) \quad x_k > 0 \quad (k = 1, 2, \dots, m),$$

$$(2.16) \quad x_1 \leq x_2 \leq \dots \leq x_m.$$

Actually, the constraint (2.16) is redundant, because if $(x_1^*, x_2^*, \dots, x_m^*)$ is a solution of the optimization problem without constraint (2.16), then from symmetry, $(x_{\sigma(1)}^*, x_{\sigma(2)}^*, \dots, x_{\sigma(m)}^*)$ is also a solution for any permutation σ of $\{1, 2, \dots, m\}$, and therefore we can choose a solution that satisfies (2.16). Hence, we omit (2.16) in the sequel.

To solve the optimization problem (2.12)–(2.15), we remove the constraint (2.15) and consider a relaxed problem described by (2.12)–(2.14). By introducing the Lagrange multipliers μ and ν , we can write the Lagrangian as

$$(2.17) \quad L = x_m - \mu \left(\sum_{k=1}^m x_k - a \right) - \nu \left(\sum_{k=1}^m x_k^2 - b \right).$$

Then the solution to (2.12)–(2.14) must satisfy

$$(2.18) \quad \frac{\partial L}{\partial x_m} = 1 - \mu - 2\nu x_m = 0,$$

$$(2.19) \quad \frac{\partial L}{\partial x_k} = -\mu - 2\nu x_k = 0 \quad (k = 1, 2, \dots, m-1),$$

$$(2.20) \quad \frac{\partial L}{\partial \mu} = \sum_{k=1}^m x_k - a = 0,$$

$$(2.21) \quad \frac{\partial L}{\partial \nu} = \sum_{k=1}^m x_k^2 - b = 0.$$

From (2.19) we have either $\nu = 0$ or $x_1 = x_2 = \dots = x_{m-1}$. However, when $\nu = 0$, we have $\mu = 0$ from (2.19) and $\mu = 1$ from (2.18), which is a contradiction. Thus $x_1 = x_2 = \dots = x_{m-1}$ must hold. Inserting this into (2.20) and (2.21) leads to

$$(2.22) \quad x_m + (m-1)x_1 - a = 0,$$

$$(2.23) \quad x_m^2 + (m-1)x_1^2 - b = 0.$$

Solving these simultaneous equations with respect to x_m gives

$$(2.24) \quad x_m^\pm = \frac{a \pm \sqrt{m(m-1)b - (m-1)a^2}}{m}.$$

Note that the x_m given by (2.24) is real, since

$$(2.25) \quad \begin{aligned} m(m-1)b - (m-1)a^2 &= (m-1) \left\{ m \sum_{k=1}^m x_k^2 - \left(\sum_{k=1}^m x_k \right)^2 \right\} \\ &= (m-1) \sum_{k=1}^m \sum_{l=1}^{k-1} (x_k - x_l)^2 \geq 0. \end{aligned}$$

Now we return to the relaxed optimization problem (2.12)–(2.14). Since the feasible set of this problem is compact and both the objective function and the constraints are differentiable, it must have a minimum and a maximum at points, where the gradient of the Lagrangian is zero. Furthermore, since the objective function is x_m itself, the maximum is attained when $x_m = x_m^+$. Then from (2.22) we have

$$(2.26) \quad x_1 = x_2 = \dots = x_{m-1} = \frac{(m-1)a - \sqrt{m(m-1)b - (m-1)a^2}}{m(m-1)}.$$

Hence, (2.26) and $x_m = x_m^+$ are the solution of the relaxed optimization problem.

Finally, we consider the positivity constraint (2.15). It is clear from (2.24) that $x_m^+ > 0$. To investigate the positivity of the other variables, note that

$$(2.27) \quad a^2 - b = \left(\sum_{k=1}^m \frac{1}{\lambda_k} \right)^2 - \sum_{k=1}^m \frac{1}{\lambda_k^2} = 2 \sum_{k=1}^m \sum_{l=1}^{k-1} \frac{1}{\lambda_k} \cdot \frac{1}{\lambda_l} > 0,$$

where we have used the fact that a and b are the traces of the inverse of a matrix with positive eigenvalues. Then (2.26) can be rewritten as

$$(2.28) \quad \begin{aligned} x_1 = x_2 = \dots = x_{m-1} &= \frac{(m-1)^2 a^2 - \{m(m-1)b - (m-1)a^2\}}{m(m-1)\{(m-1)a + \sqrt{m(m-1)b - (m-1)a^2}\}} \\ &= \frac{m(m-1)(a^2 - b)}{m(m-1)\{(m-1)a + \sqrt{m(m-1)b - (m-1)a^2}\}} > 0. \end{aligned}$$

This shows that the solution to the relaxed problem (2.12)–(2.14) automatically satisfies the constraint (2.15). Hence, it is also a solution to the original problem

(2.12)–(2.15). Returning to the original variables $\lambda_k = 1/x_k$, we know that the smallest value that λ_m can take is

$$(2.29) \quad \frac{1}{\text{Tr}(A^{-1})} \cdot m \left(1 + \sqrt{(m-1) \left[m \cdot \frac{\text{Tr}(A^{-2})}{\{\text{Tr}(A^{-1})\}^2} - 1 \right]} \right)^{-1}.$$

This gives the optimal lower bound on $\lambda_m(A)$ in terms of $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$. Since equation (2.29) is exactly the Laguerre bound (2.10), we arrive at the following theorem.

Theorem 2.1. *Among the lower bounds on $\lambda_m(A)$ computed from $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$, the Laguerre bound (2.10) is optimal in terms of sharpness.*

3. THE GAP BETWEEN THE LAGUERRE BOUND AND THE SMALLEST EIGENVALUE

Now that we have established that the Laguerre bound is the optimal lower bound, we next study the gap between the bound and the minimum eigenvalue. We begin with a lemma that holds for a 3×3 matrix and then proceed to the general case. In the course of discussion, we also allow infinite eigenvalues to make the arguments simpler.

Assume that $A \in \mathbb{R}^{3 \times 3}$ is a symmetric positive definite matrix with $\text{Tr}(A^{-1}) = a$ and $\text{Tr}(A^{-2}) = b$. Let the eigenvalues of A be $\lambda_1 \geq \lambda_2 \geq \lambda_3 > 0$. To evaluate the gap, we consider how large λ_3 can be under the fixed values of a and b . First, we show the following lemma.

Lemma 3.1. *For fixed $a = \text{Tr}(A^{-1})$ and $b = \text{Tr}(A^{-2})$, if λ_3 attains maximum value, then necessarily either $\lambda_2 = \lambda_3$ or $\lambda_1 = \infty$ holds.*

Proof. Let $x = 1/\lambda_3$, $y = 1/\lambda_2$, and $z = 1/\lambda_1$. Since we allow infinite eigenvalues, the point (x, y, z) lies in a region D of the xyz space specified by $x + y + z = a$, $x^2 + y^2 + z^2 = b$, and $x \geq y \geq z \geq 0$. Since D is a compact set, the continuous function x attains a minimum somewhere in D . Hence, if we can show that x does not attain a minimum when $x > y$ and $z > 0$, it means that x attains a minimum when $x = y$ or $z = 0$.

Assume that the point (x, y, z) is in D and both $x > y$ and $z > 0$ hold. Then, let $\varepsilon > 0$ be some small quantity and $t \in \mathbb{R}$ and consider changing (x, y, z) to (x', y', z') as follows:

$$(3.1) \quad x' = x - \varepsilon,$$

$$(3.2) \quad y' = y + t\varepsilon,$$

$$(3.3) \quad z' = z + (1 - t)\varepsilon.$$

Clearly, the new point (x', y', z') lies on the plane $x + y + z = a$. We determine t so that it is also on the sphere $x^2 + y^2 + z^2 = b$. The condition can be written as

$$(3.4) \quad (x - \varepsilon)^2 + (y + t\varepsilon)^2 + \{z + (1 - t)\varepsilon\}^2 = x^2 + y^2 + z^2,$$

or

$$(3.5) \quad \varepsilon t^2 + (y - z - \varepsilon)t + (-x + z + \varepsilon) = 0.$$

Solving this with respect to t gives

$$(3.6) \quad t_{\pm} = \frac{-(y - z - \varepsilon) \pm \sqrt{(y - z - \varepsilon)^2 + 4\varepsilon(x - z - \varepsilon)}}{2\varepsilon}.$$

In the following, we adopt the solution $t = t_+$. Now we consider two cases. First, for the case of $y = z$, we have from (3.6)

$$(3.7) \quad t_{+\varepsilon} = \frac{\varepsilon + \sqrt{\varepsilon^2 + 4\varepsilon(x - z - \varepsilon)}}{2} = O(\sqrt{\varepsilon}).$$

Inserting this into (3.1) through (3.3), we know that the changes in x , y , and z are at most $O(\sqrt{\varepsilon})$ when ε is small.

Next, consider the case of $y > z$. In this case, we can rewrite (3.6) as

$$(3.8) \quad t_+ = \frac{2(x - z - \varepsilon)}{(y - z - \varepsilon) + \sqrt{(y - z - \varepsilon)^2 + 4\varepsilon(x - z - \varepsilon)}}.$$

Since $x - z - \varepsilon > 0$ and $y - z - \varepsilon > 0$ for sufficiently small ε , we have

$$(3.9) \quad 0 < t_+ < \frac{x - z - \varepsilon}{y - z - \varepsilon} = 1 + \frac{x - y}{y - z - \varepsilon}.$$

The right-hand side is smaller than $1 + 2(x - y)/(y - z)$, which is a constant that does not depend on ε , when $\varepsilon < \frac{1}{2}(y - z)$. Hence, $t_{+\varepsilon} = O(\varepsilon)$ when ε is small and therefore, the changes in x , y , and z are at most $O(\varepsilon)$ in this case.

In summary, in both cases, the changes of x , y , and z can be made arbitrarily small. Thus, by choosing ε sufficiently small, we can make x' smaller than x while keeping the relation $x' > y' > 0$ and $x' > z' > 0$ (Fig. 1). The relation $y' \geq z'$ may not hold, but in that case, we can interchange y' and z' . In this way, we can obtain another point $(x', y', z') \in D$ which attains a smaller value of x . Hence, x cannot attain a minimum when both $x > y$ and $z > 0$ hold and the lemma is proved. \square




Figure 1. The values of x , y and z before and after the perturbation.

Using this lemma, we can prove the following theorem for any $m \times m$ real symmetric positive definite matrix A , where $m \geq 3$.

Theorem 3.2. *Let $a = \text{Tr}(A^{-1})$ and $b = \text{Tr}(A^{-2})$ be fixed. Since $1 < a^2/b \leq m$ by virtue of (2.25) and (2.27), a positive integer q satisfying $q < a^2/b \leq q + 1$ is determined uniquely. Then, $\lambda_m(A)$ takes a maximum when*

$$\lambda_1(A) = \dots = \lambda_{m-q-1}(A) = \infty \quad \text{and} \quad \lambda_{m-q+1} = \dots = \lambda_m(A).$$

The maximum is given as

$$(3.10) \quad \lambda_m^*(A) = \frac{1}{\text{Tr}(A^{-1})} \cdot q(q+1) \left(q + \sqrt{q \left[(q+1) \cdot \frac{\text{Tr}(A^{-2})}{\{\text{Tr}(A^{-1})\}^2} - 1 \right]} \right)^{-1}.$$

Proof. Let $x_k = 1/\lambda_k$. First, assume that there are two or more eigenvalues which are neither infinite nor equal to $\lambda_m(A)$. In this case, as we will show in the sequel, we can make λ_m smaller by adding appropriate perturbations. We divide the proof into two cases depending on the multiplicity q of the smallest eigenvalue.

When $q = 1$, from the assumption, both λ_{m-2} and λ_{m-1} are neither infinite nor equal to $\lambda_m(A)$. Thus, we have $0 < x_{m-2} \leq x_{m-1} < x_m$. Then, by picking up these three variables and adding the same perturbations as in Lemma 3.1, we can make x_m smaller while keeping the condition $0 < x_{m-2} \leq x_{m-1} < x_m$ (Fig. 2). Clearly, the values of $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ are unchanged by this perturbation. Hence, x_m cannot take a minimum in this case.




Figure 2. The values of x_m , x_{m-1} , and x_{m-2} before and after the perturbation.

When $q > 1$, $0 < x_{m-q-1} \leq x_{m-q} < x_{m-q+1} = \dots = x_m$ holds from the assumption. Then, by picking up the three variables x_{m-q-1} , x_{m-q} and x_{m-q+1} and adding the perturbations as in Lemma 3.1, we can make x_{m-q+1} smaller while keeping $0 < x_{m-q-1} \leq x_{m-q} < x_{m-q+1}$. This does not change the smallest eigenvalue, but reduces its multiplicity from q to $q - 1$ (Fig. 3). Moreover, the condition that there are two or more eigenvalues which are neither infinite nor equal to $\lambda_m(A)$ still holds. Hence, we can repeat this procedure and reduce q to 1, while keeping the value of the smallest eigenvalue unchanged. But in this last situation, x_m cannot take a minimum, as concluded in the analysis of the $q = 1$ case.




Figure 3. The values of $x_{m-q-1}, x_{m-q}, \dots, x_m$ before and after the perturbation.

From the above analysis, we can conclude that x_m cannot attain a minimum when there are two or more eigenvalues which are neither infinite nor equal to $\lambda_m(A)$. Thus, the only possible case is when $x_1 = \dots = x_{m-q-1} = 0$ and $x_{m-q+1} = \dots = x_m$ holds for some q . In this case, we have

$$(3.11) \quad x_{m-q} + qx_m = a,$$

$$(3.12) \quad x_{m-q}^2 + qx_m^2 = b,$$

or

$$(3.13) \quad x_m^\pm = \frac{aq \pm \sqrt{q\{(q+1)b - a^2\}}}{q(q+1)},$$

$$(3.14) \quad x_{m-q}^\pm = \frac{a \mp \sqrt{q\{(q+1)b - a^2\}}}{q+1}.$$

For x_m and x_{m-q} to be real, q must satisfy $q+1 \geq a^2/b$. Then, for $x_{m-q} \leq x_m$ to hold, we have to choose x_m^+ and x_{m-q}^+ . In addition, for $x_{m-q}^+ > 0$ to hold, we must have $q < a^2/b$. From the condition $q < a^2/b \leq q+1$, q is determined uniquely. Hence, there is only one set of q , x_m , and x_{m-q} that satisfy the condition for minimum x_m . Since the feasible region of (x_1, x_2, \dots, x_m) , specified by $\sum_{k=1}^m x_k = a$, $\sum_{k=1}^m x_k^2 = b$ and $0 \leq x_1 \leq x_2 \leq \dots \leq x_m$, is compact, x_m must attain a minimum somewhere in

this region. Accordingly, we conclude that x_m takes a minimum when $q < a^2/b \leq q + 1$, $x_1 = \dots = x_{m-q-1} = 0$, $x_{m-q} = x_{m-q}^+$ and $x_{m-q+1} = \dots = x_m = x_m^+$. Equation (3.10) is obtained from $\lambda_m = 1/x_m$. \square

To measure the gap between the Laguerre bound and the smallest eigenvalue, we use the quantity $L_L(A)/\lambda_m^*(A)$, which becomes one when there is no gap and zero when the gap is maximal. Let $\alpha \equiv \{\text{Tr}(A^{-1})\}^2/\text{Tr}(A^{-2})$ and let q be an integer specified in Theorem 3.2. Then from (3.10) and (2.10) we have

$$(3.15) \quad \frac{L_L(A)}{\lambda_m^*(A)} = \frac{m}{q(q+1)} \cdot \frac{q + \sqrt{q\{(q+1)\alpha^{-1} - 1\}}}{1 + \sqrt{(m-1)(m\alpha^{-1} - 1)}}.$$

Thus, we have obtained an expression for the maximum possible gap as a function of m and α (note that q is determined from α uniquely).

So far, we have allowed infinite eigenvalues. However, of course, actual matrices have only finite eigenvalues. Accordingly, except for the case of $q = m - 1$, for which no infinite eigenvalues are required for $\lambda_m(A)$ to take a maximum, the right-hand side of (3.15) is a lower bound that can be approached arbitrarily closely.

Finally, we investigate the behavior of the right-hand side of (3.15) as a function of α . Note that $1 < \alpha \leq m$ from (2.25) and (2.27). We consider three extreme cases, namely, $1 < \alpha \leq 2$, $1 \ll \alpha \ll m$, and $m - 1 < \alpha \leq m$.

▷ When $1 < \alpha \leq 2$, we have $q = 1$ and therefore,

$$(3.16) \quad \frac{L_L(A)}{\lambda_m^*(A)} = \frac{m}{2} \cdot \frac{1 + \sqrt{2\alpha^{-1} - 1}}{1 + \sqrt{(m-1)(m\alpha^{-1} - 1)}}.$$

For $1 < \alpha \leq 2$, this is a decreasing function in α that attains the minimum value

$$(3.17) \quad \frac{1}{\sqrt{2}} \cdot \frac{1}{\sqrt{2}m^{-1} + \sqrt{(1-m^{-1})(1-2m^{-1})}}$$

at $\alpha = 2$ and approaches 1 as $\alpha \rightarrow 1$. Since the minimum value (3.17) is larger than $1/\sqrt{2}$ when $m \geq 3$, $L_L(A)/\lambda_m^*(A)$ is always larger than $1/\sqrt{2}$ when $m \geq 3$.

▷ When $1 \ll \alpha \ll m$, we have $1 \ll q \ll m$ and therefore, $L_L(A)/\lambda_m^*(A) \simeq 1/\sqrt{q}$.

▷ When $m - 1 < \alpha \leq m$, we have $q = m - 1$ and therefore,

$$(3.18) \quad \frac{L_L(A)}{\lambda_m^*(A)} = \frac{1}{m-1} \cdot \left\{ 1 + \frac{m-2}{1 + \sqrt{(m-1)(m\alpha^{-1} - 1)}} \right\}.$$

This is an increasing function in α and takes the maximum value 1 at $\alpha = m$ and approaches $\frac{1}{2}m/(m-1)$ as $\alpha \rightarrow m - 1$. Hence, $L_L(A)/\lambda_m^*(A) > \frac{1}{2}$ all over the region.

In summary, we can conclude that the Laguerre bound is fairly tight when α is smaller than 2 or close to m and can be loose when α is in the intermediate region.

In Fig. 4, we plot the smallest eigenvalues of randomly generated 5×5 symmetric positive definite matrices. Any such matrix can be written as $A = Q\Lambda Q^T$, where Q is an orthogonal matrix and Λ is a diagonal matrix with positive diagonal elements. However, its traces $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$ depend only on Λ and not on Q . We therefore set Q to be the identity matrix and generate the diagonal elements of Λ randomly. These matrices are normalized so that $\text{Tr}(A^{-1}) = 1$ and the horizontal axis is $\alpha = \{\text{Tr}(A^{-1})\}^2 / \text{Tr}(A^{-2})$. The Laguerre bound (2.10) and the upper bound (3.10) on the smallest eigenvalue are also shown in the graph. From the graph, we can confirm the optimality of the Laguerre bound, since it actually constitutes the lower boundary of the region, where the smallest eigenvalues exist. We can also confirm the upper bound given by (3.10) numerically. Finally, it is clear that the Laguerre bound is tight when $\alpha \leq 2$ or $\alpha \simeq m$ and loose in the intermediate region.




Figure 4. The smallest eigenvalues of randomly generated 5×5 symmetric positive definite matrices as a function of α .

4. CONCLUSION

In this paper, we investigated the properties of lower bounds on the smallest eigenvalue of a symmetric positive definite matrix A computed from $\text{Tr}(A^{-1})$ and $\text{Tr}(A^{-2})$. We studied two problems, namely, finding the optimal bound and evaluating its sharpness. As for the former question, we found that the Laguerre bound is the optimal one in terms of sharpness. As for the latter question, we characterized the situation in which the gap becomes largest in terms of the eigenvalue

distribution of A . Furthermore, we showed that the gap becomes smallest when $\{\text{Tr}(A^{-1})\}^2/\text{Tr}(A^{-2})$ approaches 1 or m . These results will help designing efficient shift strategies for singular value computation methods such as the dqds algorithm and the mdLVs algorithm.

Acknowledgment. The author would like to thank Prof. Kinji Kimura, Dr. Yuji Nakatsukasa, and Dr. Takumi Yamashita for fruitful discussion. The author is also grateful to the anonymous referees whose comments helped improving the quality of this paper.

References

- [1] *K. Aishima, T. Matsuo, K. Murota, M. Sugihara*: A survey on convergence theorems of the dqds algorithm for computing singular values. *J. Math-for-Ind.* *2* (2010), 1–11. [zbl](#) [MR](#)
- [2] *G. Alefeld*: On the convergence of Halley’s method. *Am. Math. Monthly* *88* (1981), 530–536. [zbl](#) [MR](#) [doi](#)
- [3] *K. V. Fernando, B. N. Parlett*: Accurate singular values and differential qd algorithms. *Numer. Math.* *67* (1994), 191–229. [zbl](#) [MR](#) [doi](#)
- [4] *G. H. Golub, C. F. Van Loan*: *Matrix Computations*. Johns Hopkins Studies in the Mathematical Sciences, Johns Hopkins University Press, Baltimore, 2013. [zbl](#) [MR](#)
- [5] *A. S. Householder*: *The Numerical Treatment of a Single Nonlinear Equation*. International Series in Pure and Applied Mathematics, McGraw-Hill Book, New York, 1970. [zbl](#) [MR](#)
- [6] *M. Iwasaki, Y. Nakamura*: Accurate computation of singular values in terms of shifted integrable schemes. *Japan J. Ind. Appl. Math.* *23* (2006), 239–259. [zbl](#) [MR](#) [doi](#)
- [7] *C. R. Johnson*: A Gersgorin-type lower bound for the smallest singular value. *Linear Algebra Appl.* *112* (1989), 1–7. [zbl](#) [MR](#) [doi](#)
- [8] *C. R. Johnson, T. Szulc*: Further lower bounds for the smallest singular value. *Linear Algebra Appl.* *272* (1998), 169–179. [zbl](#) [MR](#) [doi](#)
- [9] *K. Kimura, T. Yamashita, Y. Nakamura*: Conserved quantities of the discrete finite Toda equation and lower bounds of the minimal singular value of upper bidiagonal matrices. *J. Phys. A, Math. Theor.* *44* (2011), Article ID 285207, 12 pages. [zbl](#) [MR](#) [doi](#)
- [10] *U. von Matt*: The orthogonal qd-algorithm. *SIAM J. Sci. Comput.* *18* (1997), 1163–1186. [zbl](#) [MR](#) [doi](#)
- [11] *T. Yamashita, K. Kimura, Y. Nakamura*: Subtraction-free recurrence relations for lower bounds of the minimal singular value of an upper bidiagonal matrix. *J. Math-for-Ind.* *4* (2012), 55–71. [zbl](#) [MR](#)
- [12] *T. Yamashita, K. Kimura, M. Takata, Y. Nakamura*: An application of the Kato-Temple inequality on matrix eigenvalues to the dqds algorithm for singular values. *JSIAM Lett.* *5* (2013), 21–24. [MR](#) [doi](#)
- [13] *T. Yamashita, K. Kimura, Y. Yamamoto*: A new subtraction-free formula for lower bounds of the minimal singular value of an upper bidiagonal matrix. *Numer. Algorithms* *69* (2015), 893–912. [zbl](#) [MR](#) [doi](#)
- [14] *J. H. Wilkinson*: *The Algebraic Eigenvalue Problem*. Monographs on Numerical Analysis, Oxford Science Publications, Clarendon Press, Oxford, 1988. [zbl](#) [MR](#)

Author’s address: Yusaku Yamamoto, The University of Electro-Communications, 1-5-1 Chofugaoka, Chofu, Tokyo, 182-8585, Japan, e-mail: yusaku.yamamoto@uec.ac.jp.