

Yahya Talebi; Atefeh Darzi

On graph associated to co-ideals of commutative semirings

Commentationes Mathematicae Universitatis Carolinae, Vol. 58 (2017), No. 3, 293–305

Persistent URL: <http://dml.cz/dmlcz/146913>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2017

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

On graph associated to co-ideals of commutative semirings

YAHYA TALEBI, ATEFEH DARZI

Abstract. Let R be a commutative semiring with non-zero identity. In this paper, we introduce and study the graph $\Omega(R)$ whose vertices are all elements of R and two distinct vertices x and y are adjacent if and only if the product of the co-ideals generated by x and y is R . Also, we study the interplay between the graph-theoretic properties of this graph and some algebraic properties of semirings. Finally, we present some relationships between the zero-divisor graph $\Gamma(R)$ and $\Omega(R)$.

Keywords: semiring; co-ideal; maximal co-ideal

Classification: 16Y60, 05C75

1. Introduction

The concept of the *zero-divisor graph* of a commutative ring R was first introduced by Beck [3]. He defined this graph as a simple graph where all elements of the ring R are the vertex-set of this graph and two distinct elements x and y are adjacent if and only if $xy = 0$. Beck conjectured that $\chi(R) = \omega(R)$ for every ring R . In [2], Anderson and Livingston introduced the zero-divisor graph with vertices $Z(R)^* = Z(R) \setminus \{0\}$, the set of non-zero zero-divisors of R . Some other investigations into properties of zero-divisor graph over commutative semiring may be found in [5], [6]. In [11], Sharma and Bhatwadekar defined another graph on a ring R with vertices as elements of R and there is an edge between two distinct vertices x and y in R if and only if $Rx + Ry = R$. Further, in [10], Maimani et al. studied the graph defined by Sharma and Bhatwadekar and called it *comaximal graph*. Also, in [1], Akbari et al. studied the comaximal graph over non-commutative ring.

Note that throughout this paper all semirings are considered to be commutative semirings with non-zero identity. First, we introduce the concept of *product* of co-ideals in the semiring R . Next, we define an undirected graph over commutative semiring in which vertices are all elements of R and two distinct vertices x and y are adjacent if and only if the product of the co-ideals generated by x and y is R (i.e. $F(x)F(y) = R$). We denote this graph by $\Omega(R)$. In Section 2, we recall some notions of semirings which will be used in this paper. In other sections, we study some graph-theoretic properties of $\Omega(R)$ and its subgraphs such as diameter, radius, girth, clique number and chromatic number.

In a graph G , we denote the vertex-set of G by $V(G)$ and the edge-set by $E(G)$. A graph G is said to be *connected*, if there is a path between every two distinct vertices and we say that G is *totally disconnected*, if no two vertices of G are adjacent. For a given vertex x , the number of all vertices adjacent to it, is called *degree* of the vertex x , denoted by $deg(x)$. For distinct vertices x and y of G , let $d(x, y)$ be the length of the shortest path from x to y ($d(x, x) = 0$ and $d(x, y) = \infty$ if there is no such path). The *diameter* of G is $diam(G) = \sup\{d(x, y) : x \text{ and } y \text{ are distinct vertices of } G\}$. The *girth* of G , denoted by $gr(G)$, is defined as the length of the shortest cycle in G . If G has no cycles, then $gr(G) = \infty$ and G is called a *forest*. Also, G is called a *tree* if G is connected and has no cycles. A *clique* in a graph G is a complete subgraph of G . The *clique number* of G , denoted by $\omega(G)$, is the number of vertices in a largest clique of G . An *independent set* in a graph G is a set of pairwise non-adjacent vertices. A graph in which each pair of distinct vertices is joined by an edge is called a *complete graph*. We denote the complete graph on n vertices by K_n . For a positive integer k , a *k-partite* graph is one whose vertex-set can be partitioned into k independent sets. A *k-partite* graph G is said to be a *complete k-partite* graph, if each vertex is joined to every vertex that is not in the same partition. The *complete bipartite* graph (2-partite graph) with parts of sizes m and n is denoted by $K_{m,n}$. We will sometimes call a $K_{1,n}$ a *star graph*. We write $G \setminus \{x\}$ or $G \setminus S$ for the subgraph of G obtained by deleting a vertex x or set of vertices S . An *induced subgraph* is a subgraph obtained by deleting a set of vertices. Also, a *spanning subgraph* of G is a subgraph with vertex-set $V(G)$. A general reference for graph theory is [12].

2. Preliminaries

In this section, we recall various notions about semirings which will be used throughout the paper. A *semiring* R is an algebraic system $(R, +, \cdot)$ such that $(R, +)$ is a commutative monoid with identity element 0 and (R, \cdot) is a semigroup. In addition, operations $+$ and \cdot are connected by distributivity and 0 annihilates R (i.e. $x0 = 0x = 0$ for each $x \in R$). A semiring R is said to be *commutative* if (R, \cdot) is a commutative semigroup and R is said to have an *identity* if there exists $1 \in R$ such that $1x = x1 = x$.

Recall that, throughout this paper, all semirings are commutative with non-zero identity. The following definitions are given in [7], [9].

2.1 Definition. Let R be a semiring.

(1) A non-empty subset I of R is called a *co-ideal* of R if and only if it is closed under multiplication and satisfies the condition that $a + r \in I$ for all $a \in I$ and $r \in R$. According to this definition, $0 \in I$ if and only if $I = R$. Also, a co-ideal I of R is called *strong*, if $1 \in I$.

(2) A co-ideal I of semiring R is called *subtractive* if $x \in I$ and $xy \in I$, implies $y \in I$ for all $x, y \in R$. So every subtractive co-ideal is a strong co-ideal.

(3) A proper co-ideal P of R is called *prime* if $a + b \in P$, implies $a \in P$ or $b \in P$ for all $a, b \in R$.

(4) A proper co-ideal I of R is called *maximal* if there is no co-ideal J such that $I \subset J \subset R$.

(5) An element a of a semiring R is *multiplicatively idempotent* if and only if $a^2 = a$ and a is called *additively idempotent* if and only if $a + a = a$. A semiring R is said to be idempotent if it is both additively and multiplicatively idempotent.

(6) An element x of a semiring R is called a *zero-sum* of R , if there exists an element $y \in R$ such that $x + y = 0$. It is clear that, y is the unique element which satisfies $x + y = 0$. We will denote the set of all zero-sums of R by $ZS(R)$. It is easy to see that $ZS(R)$ is an ideal of R . Also, a semiring R is a ring if and only if $ZS(R) = R$ and R is called *zero-sumfree* if and only if $ZS(R) = 0$.

(7) If A is a non-empty subset of a semiring R , then the set $F(A)$ of all elements of R of the form $a_1 a_2 \dots a_n + r$, where $a_i \in A$ for all $1 \leq i \leq n$ and $r \in R$, is a co-ideal of R containing A . In fact, $F(A)$ is the unique smallest co-ideal of R containing A .

By the above definition, we can consider the co-ideal generated by a single element $x \in R$ as follows: $F(x) = \{x^n + r : r \in R \text{ and } n \in \mathbf{N}\}$. It is obvious that, if $x \in I$ for some co-ideal I , then $F(x) \subseteq I$.

By definition of co-ideal, if R is a ring, then R has no proper co-ideals and so throughout this paper we consider semirings which are not rings. For a semiring R , we denote the set of maximal co-ideals, the union of all the maximal co-ideals and the intersection of all the maximal co-ideals of R by $Co - Max(R)$, $UM(R)$ and $IM(R)$, respectively. Also, if the semiring R has exactly one maximal co-ideal, then we say that the semiring R is *c-local* and R is said to be a *c-semilocal* semiring, if R has only a finite number of maximal co-ideals.

2.2 Lemma ([7]). *Let I_1, \dots, I_n be co-ideals of a semiring R and P be a prime co-ideal containing $\bigcap_{i=1}^n I_i$. Then $I_i \subseteq P$ for some $i = 1, \dots, n$. Moreover, if $P = \bigcap_{i=1}^n I_i$, then $P = I_i$ for some i .*

2.3 Lemma. *Let R be a semiring. Then $x \in \sqrt{ZS(R)}$ if and only if $F(x) = R$.*

PROOF: Let $x \in \sqrt{ZS(R)}$. Thus $x^n \in ZS(R)$ for some positive integer n . This implies $x^n + r = 0$ for some $r \in R$. Hence $0 \in F(x)$, since $x^n + r \in F(x)$ and so $F(x) = R$.

The converse follows, since all conclusions are reversible. □

2.4 Proposition. *Let R be a semiring. Then $R \setminus \sqrt{ZS(R)} = UM(R)$.*

PROOF: Assume that $x \in R \setminus \sqrt{ZS(R)}$. Thus $F(x) \neq R$ and by [7, Proposition 2.1], there exists $m \in Co - Max(R)$ such that $x \in F(x) \subseteq m$. Hence $R \setminus \sqrt{ZS(R)} \subseteq UM(R)$.

Conversely, suppose that $x \in UM(R)$. Thus there is a maximal co-ideal m such that $x \in m$. Now, if $x \in \sqrt{ZS(R)}$, then $F(x) = R$ by Lemma 2.3 and so $R = F(x) \subseteq m$, that is impossible. Hence $UM(R) \subseteq R \setminus \sqrt{ZS(R)}$. This implies $R \setminus \sqrt{ZS(R)} = UM(R)$. □

2.5 Remark. Note that the Prime Avoidance Theorem is explained for subtractive prime co-ideals of a commutative semiring R in [4, Theorem 3.8]. Also, by [8, Proposition 2.5] and [7, Theorem 3.10], every maximal co-ideal is a subtractive and prime co-ideal, so we can conclude that the Prime Avoidance Theorem and Lemma 2.2 also hold for the case where co-ideals are maximal.

In the following, we define the product of co-ideals of a semiring R . It is straightforward to verify that the product of co-ideals with this definition is a co-ideal.

2.6 Definition. Let I and J be two co-ideals of a semiring R . We define the product of I and J as follows:

$$IJ = \{xy + r : x \in I, y \in J \text{ and } r \in R\}.$$

Similarly, we define the product of any finite family of co-ideals. Moreover, I^n is defined for any co-ideal I and $I^n = \{a_1 \dots a_n + r : a_i \in I \text{ and } r \in R\}$.

Let I and J be co-ideals of R such that $x \in I$ and $y \in J$. Note that with this definition, if I and J are strong co-ideals, then $x, y \in IJ$ because $x = x1 + 0$ and $y = 1y + 0$ but this may not be true in general.

3. Some basic properties of $\Omega(R)$

As mentioned in the introduction, the graph $\Omega(R)$ is a graph with all the elements of R as its vertex-set and two distinct vertices x and y are adjacent if and only if $F(x)F(y) = R$. Let $\Omega_1(R)$ be the subgraph of $\Omega(R)$ with vertex-set $\sqrt{ZS(R)}$ and $\Omega_2(R)$ be the subgraph of $\Omega(R)$ with vertex-set $UM(R)$. If $x \in \sqrt{ZS(R)}$, then by Lemma 2.3, $F(x) = R$ and this implies x is adjacent to any other vertex of R . With this comment, we can say that $\Omega_1(R)$ is a complete graph. Also, if $x, y \in m$ for some maximal co-ideal m of R , then x and y cannot be adjacent because $F(x)F(y) \subseteq m$. Hence, if the semiring R has one maximal co-ideal, then $\Omega_2(R)$ is a totally disconnected graph.

3.1 Lemma. Let m be a maximal co-ideal of a semiring R and $x \in R$. If $x \notin m$, then $mF(x) = R$.

PROOF: Suppose that $x \notin m$. Thus $F(m \cup \{x\}) = R$ since $m \subsetneq F(m \cup \{x\})$ and m is a maximal co-ideal. Now, since $0 \in R$, we split the proof into three cases for $F(m \cup \{x\})$:

Case 1: There exist $a_1, \dots, a_k \in m$ and $r \in R$ for some positive integer k such that $a_1 \dots a_k + r = 0$. This implies $0 \in m$ since m is co-ideal. This is a contradiction because m is a maximal co-ideal.

Case 2: $x^t + r = 0$ for some $r \in R$ and a positive integer t . In this case, $F(x) = R$ because $0 = x^t + r \in F(x)$ and so $mF(x) = R$.

Case 3: $yx^t + r = 0$ for some $y \in m$, $r \in R$ and a positive integer t . Hence $mF(x) = R$ since $0 = yx^t + r \in mF(x)$. \square

As an immediate consequence of Lemma 3.1, we have the next proposition:

3.2 Proposition. *Let m be a maximal co-ideal of a semiring R and $x \in R$. If $x \notin m$, then there is an element $y \in m$ such that x is adjacent to y in $\Omega(R)$.*

PROOF: Suppose that m is a maximal co-ideal and $x \notin m$. By Lemma 3.1, we have $mF(x) = R$. This implies $y(x^t + r) + k = 0$ for some $r, k \in R, y \in m$ and a positive integer t . Hence $yx^t + s = 0$ for some $s \in R$ and so $F(x)F(y) = R$ since $0 = yx^t + s \in F(x)F(y)$. Therefore, x and y are adjacent in $\Omega(R)$. \square

3.3 Proposition. *Let R be a semiring and $x \in R$. Then $x \in IM(R)$ if and only if x is adjacent to no vertex of $\Omega_2(R)$.*

PROOF: Let $x \in IM(R)$. Assume contrary that $y \in UM(R)$ is adjacent to x in $\Omega_2(R)$. Thus there exists $m \in Co - Max(R)$ such that $y \in m$ and $F(x)F(y) = R$. On the other hand, $x \in IM(R)$ gives $x \in m$. Hence $F(x)F(y) \subseteq m$, that is a contradiction.

Conversely, assume that x is not adjacent to any vertex of $\Omega_2(R)$. If $x \notin IM(R)$, there exists $m \in Co - Max(R)$ such that $x \notin m$. By Proposition 3.2, there is an element $y \in m$ such that x is adjacent to y , which is contrary to our assumption. \square

By Proposition 3.3, for each $x \in IM(R)$, $deg_{\Omega_2(R)}(x) = 0$. So it will be interesting to study the properties of the graph $\Omega_2(R) \setminus IM(R)$ with vertex-set $UM(R) \setminus IM(R)$. Note that if R is a c-local semiring, then $\Omega_2(R) \setminus IM(R)$ is an empty graph.

3.4 Theorem. *Let R be a semiring which is not c-local. Then $\Omega_2(R) \setminus IM(R)$ is a complete bipartite graph if and only if R has exactly two maximal co-ideals.*

PROOF: First, assume that $\Omega_2(R) \setminus IM(R)$ is a complete bipartite graph with vertex-sets V_1 and V_2 . Clearly, m is contained in one of the partitions for any maximal co-ideal m . Thus, suppose that $m_i \setminus IM(R) \subseteq V_i$ for $i = 1, 2$. If R has another maximal co-ideal such as m_3 , then $m_3 \setminus IM(R) \subseteq V_i$ for some $i = 1, 2$, which is impossible, since $m_1m_3 = m_2m_3 = R$. Hence R can have only two maximal co-ideals.

Conversely, suppose that $Co - Max(R) = \{m_1, m_2\}$. Then the vertex-set of $\Omega_2(R) \setminus IM(R)$ is $(m_1 \setminus m_2) \cup (m_2 \setminus m_1)$. Clearly, the subgraphs $m_1 \setminus m_2$ and $m_2 \setminus m_1$ are totally disconnected. Let $x \in m_1 \setminus m_2$ and $y \in m_2 \setminus m_1$. Now to complete the proof, it suffices to show that $F(x)F(y) \not\subseteq m_1$ and $F(x)F(y) \not\subseteq m_2$. If $F(x)F(y) \subseteq m_1$, then $xy \in m_1$. This implies that $y \in m_1$, since m_1 is subtractive, a contradiction. Similarly, it can be shown that $F(x)F(y) \not\subseteq m_2$. Therefore we have $F(x)F(y) = R$. Hence $\Omega_2(R) \setminus IM(R)$ is complete bipartite graph with vertex-set $m_1 \setminus m_2$ and $m_2 \setminus m_1$. \square

In the following, we give an example of semiring R in which R has two maximal co-ideals and show that $\Omega_2(R) \setminus IM(R)$ is complete bipartite graph.

3.5 Example. Let $S = \{0, 1, a\}$ be an idempotent semiring in which $a + 1 = 1 + a = a$ and let $R = S \times S$. The maximal co-ideals of R are as follows:

$$m_1 = \{(0, 1), (0, a), (1, a), (a, 1), (1, 1), (a, a)\},$$

$$m_2 = \{(1, 0), (a, 0), (1, a), (a, 1), (1, 1), (a, a)\}.$$

It can be shown that $\Omega_2(R) \setminus IM(R)$ is complete bipartite with vertex-sets $\{(0, 1), (0, a)\}$ and $\{(1, 0), (a, 0)\}$.

In the next theorem, we study the clique number of the graph $\Omega_2(R) \setminus IM(R)$ for a c-semilocal semiring. Also, with this theorem, we give a result about the girth of $\Omega_2(R) \setminus IM(R)$.

3.6 Theorem. *Let R be a c-semilocal semiring and $|Co - Max(R)| \geq n$ with $n \geq 2$. Then $\Omega_2(R) \setminus IM(R)$ has a clique of order n . In particular, if $|Co - Max(R)| = n$, then $\omega(\Omega_2(R) \setminus IM(R)) = n$.*

PROOF: Let $\{m_1, \dots, m_n\}$ be a subset of $Co - Max(R)$. We claim that for any $x_1 \in m_1 \setminus \bigcup_{j=2}^n m_j$, there exists a clique with vertex-set $\{x_1, \dots, x_n\}$ in $\Omega_2(R) \setminus IM(R)$, where $x_i \in m_i \setminus \bigcup_{j \neq i}^n m_j$ for $i = 1, \dots, n$. We prove this claim

by induction on n . For $n = 2$, the proof is similar to the proof of Theorem 3.4. Now, suppose that $n \geq 3$. By Remark 2.5, $m_1 \cap m_n \not\subseteq \bigcup_{j=2}^{n-1} m_j$. Thus there exists $y \in (m_1 \cap m_n) \setminus \bigcup_{j=2}^{n-1} m_j$ and so $x_1 + y \in (m_1 \cap m_n) \setminus \bigcup_{j=2}^{n-1} m_j$. By induction hypothesis, there is a clique with vertex-set $\{x_1 + y, x_2, \dots, x_{n-1}\}$, where $x_i \in m_i \setminus \bigcup_{j \neq i}^{n-1} m_j$ for $2 \leq i \leq n-1$. Indeed, $x_2, \dots, x_{n-1} \notin m_n$ since $x_1 + y \in m_n$.

On the other hand, since $x_1 + y$ is adjacent to x_2, \dots, x_{n-1} , hence x_1 is adjacent to x_2, \dots, x_{n-1} because $F(x_1 + y) \subseteq F(x_1)$. Now, since $x_1 + \dots + x_{n-1} \notin m_n$ (m_n is prime), so by Proposition 3.2, there exists $x_n \in m_n$ which is adjacent to $x_1 + \dots + x_{n-1}$. This implies that x_n is adjacent to x_1, \dots, x_{n-1} and we can conclude $\{x_1, \dots, x_n\}$ is a clique of order n in $\Omega_2(R) \setminus IM(R)$.

Now, suppose that $|Co - Max(R)| = n$. Thus we have $\omega(\Omega_2(R) \setminus IM(R)) \geq n$. If $\Omega_2(R) \setminus IM(R)$ has a clique of order k in which $k \geq n$, then by the Pigeon Hole Principal, two elements of the clique should belong to one maximal co-ideal, which is a contradiction. Hence $\omega(\Omega_2(R) \setminus IM(R)) = n$. □

Theorem 3.6 leads to the following corollary:

3.7 Corollary. *Let R be a c-semilocal semiring with $|Co - Max(R)| \geq 3$. Then $gr(\Omega_2(R) \setminus IM(R)) = 3$.*

PROOF: Let $|Co - Max(R)| \geq 3$. By Theorem 3.6, $\Omega_2(R) \setminus IM(R)$ has a clique of order 3, so $gr(\Omega_2(R) \setminus IM(R)) = 3$. □

In the next theorem, we will compute the girth of $\Omega_2(R) \setminus IM(R)$ when R is a c-semilocal semiring.

3.8 Theorem. *Let R be a c-semilocal semiring with $|Co - Max(R)| \geq 2$. If $\Omega_2(R) \setminus IM(R)$ contains a cycle, then $gr(\Omega_2(R) \setminus IM(R)) \leq 4$.*

PROOF: Assume that $\Omega_2(R) \setminus IM(R)$ contains a cycle and $gr(\Omega_2(R) \setminus IM(R)) \neq 3$. So Corollary 3.7 implies that $|Co - Max(R)| = 2$. Hence by Theorem 3.4, $\Omega_2(R) \setminus IM(R)$ is complete bipartite graph and so $gr(\Omega_2(R) \setminus IM(R)) = 4$. \square

3.9 Example. Let $X = \{a, b, c\}$ and $R = (P(X), \cup, \cap)$ be a semiring, where $P(X)$ is the power set of X . For this semiring we have $1_R = X$ and $0_R = \emptyset$. In this case, the maximal co-ideals of semiring R are as follows:

$$\begin{aligned} m_1 &= \{\{a\}, \{a, b\}, \{a, c\}, X\}, \\ m_2 &= \{\{b\}, \{a, b\}, \{b, c\}, X\}, \\ m_3 &= \{\{c\}, \{a, c\}, \{b, c\}, X\}. \end{aligned}$$

For the graph $\Omega_2(R) \setminus IM(R)$ the vertex-set is $P(X) \setminus \{\emptyset, X\}$ and $\{\{a\}, \{b\}, \{c\}\}$ is a maximal clique. This implies that $\omega(\Omega_2(R) \setminus IM(R)) = 3$ and so $gr(\Omega_2(R) \setminus IM(R)) = 3$.

3.10 Proposition. *Let R be a c -semilocal semiring with $|Co - Max(R)| \geq 2$. Then $\Omega_2(R) \setminus IM(R)$ is star graph if and only if there is a vertex of $\Omega_2(R) \setminus IM(R)$ which is adjacent to every other vertex.*

PROOF: The necessity is obvious by definition, thus we need to prove the sufficiency. Assume that there exists $x \in \Omega_2(R) \setminus IM(R)$ that is adjacent to every other vertex. Let $x \in m$ for some $m \in Co - Max(R)$. We must have $|m \setminus IM(R)| = 1$, because if x and y are distinct vertices of $m \setminus IM(R)$, then by assumption x and y are adjacent, which is impossible. Now, if $|Co - Max(R)| \geq 3$, then $|m \setminus IM(R)| \geq 3$ for any maximal co-ideal m of R . Hence R cannot contain more than two maximal co-ideals. It is straightforward to verify that $\Omega_2(R) \setminus IM(R)$ is a star graph by Theorem 3.4. \square

3.11 Theorem. *Let R be a c -semilocal semiring with $|Co - Max(R)| \geq 2$. Then the following statements are equivalent:*

- (1) $\Omega_2(R) \setminus IM(R)$ is a tree;
- (2) $\Omega_2(R) \setminus IM(R)$ is a forest;
- (3) $|Co - Max(R)| = 2$ and $|m \setminus IM(R)| = 1$ for some $m \in Co - Max(R)$;
- (4) $\Omega_2(R) \setminus IM(R)$ is a star graph.

PROOF: (1) \Rightarrow (2), (3) \Rightarrow (4) and (4) \Rightarrow (1) are clear.

(2) \Rightarrow (3) Let $\Omega_2(R) \setminus IM(R)$ be a forest. Thus by Corollary 3.7, we have $|Co - Max(R)| = 2$. Now, if $|m \setminus IM(R)| \geq 2$ for each maximal co-ideal m , then $\Omega_2(R) \setminus IM(R)$ contains a cycle of order 4, because by Theorem 3.4, $\Omega_2(R) \setminus IM(R)$ is a complete bipartite graph, a contradiction. Hence $|m \setminus IM(R)| = 1$ for some $m \in Co - Max(R)$. \square

3.12 Proposition. *Let R be a c -semilocal semiring. Then $\Omega_2(R) \setminus IM(R)$ is a complete graph if and only if it is in the form $K_{1,1}$.*

PROOF: Let $\Omega_2(R) \setminus IM(R)$ be a complete graph. So we can say that there is a vertex of $\Omega_2(R) \setminus IM(R)$ that is adjacent to every other vertex. Hence by

Proposition 3.10, $\Omega_2(R) \setminus IM(R)$ is a star graph and Theorem 3.11 implies that R has exactly two maximal co-ideals m_1 and m_2 so that $|m_i \setminus IM(R)| = 1$ for some i . Now, since for each maximal co-ideal m_i , the vertex-set $m_i \setminus IM(R)$ is a partition of $\Omega_2(R) \setminus IM(R)$, we must have $|m_i \setminus IM(R)| = 1$ for any i , because the elements of $m_i \setminus IM(R)$ are not adjacent to each other. In this case, $\Omega_2(R) \setminus IM(R)$ is in the form $K_{1,1}$.

The converse is obvious. □

3.13 Example. Let $X = \{a, b\}$ and $R = (P(X), \cup, \cap)$ be a semiring, where $P(X)$ is power set of X and $1_R = X$ and $0_R = \emptyset$. The maximal co-ideals of semiring R are as follows:

$$m_1 = \{\{a\}, X\},$$

$$m_2 = \{\{b\}, X\}.$$

Thus by Theorem 3.4, $\Omega_2(R) \setminus IM(R)$ is a complete bipartite graph with vertex-sets $V_1 = \{\{a\}\}$ and $V_2 = \{\{b\}\}$. Indeed, $\Omega_2(R) \setminus IM(R)$ forms $K_{1,1}$. Hence $\Omega_2(R) \setminus IM(R)$ is complete graph that is a star graph and a tree. Also, since $\Omega_2(R) \setminus IM(R)$ does not contain any cycle, so it is a forest and $gr(\Omega_2(R) \setminus IM(R)) = \infty$.

3.14 Theorem. *Let R be a c -semilocal semiring which is not a c -local. Then the following hold.*

- (i) *If $|Co - Max(R)| = n$, then $\Omega_2(R) \setminus IM(R)$ is n -partite.*
- (ii) *If $\Omega_2(R) \setminus IM(R)$ is n -partite, then $|Co - Max(R)| \leq n$. In this case, if $\Omega_2(R) \setminus IM(R)$ is not $(n - 1)$ -partite, then $|Co - Max(R)| = n$.*

PROOF: (i) Suppose that $Co - Max(R) = \{m_1, \dots, m_n\}$. Let $V_1 = m_1 \setminus IM(R)$ and $V_i = m_i \setminus \bigcup_{j=1}^{i-1} m_j$ for $2 \leq i \leq n$. By Remark 2.5, $V_i \neq \emptyset$ for each i . Also, clearly that $\bigcup_{i=1}^n V_i = UM(R) \setminus IM(R)$ and for every $x, y \in V_i$, they are not adjacent in $\Omega_2(R) \setminus IM(R)$. Hence $\Omega_2(R) \setminus IM(R)$ is n -partite graph.

(ii) Assume contrary that $|Co - Max(R)| \geq n + 1$. By Theorem 3.6, $\Omega_2(R) \setminus IM(R)$ has a clique with cardinality $n + 1$. Thus by the Pigeon Hole Principal, two elements of this clique should belong to one part of $\Omega_2(R) \setminus IM(R)$, which is a contradiction.

Now, if $\Omega_2(R) \setminus IM(R)$ is not $(n - 1)$ -partite and $|Co - Max(R)| = k < n$, then by part (i), $\Omega_2(R) \setminus IM(R)$ can be a k -partite graph, a contradiction. □

3.15 Proposition. *Let R be a semiring with $|Co - Max(R)| \geq 2$. If $\Omega_2(R) \setminus IM(R)$ is complete n -partite graph, then $n = 2$.*

PROOF: Let $\{m_1, m_2\} \subseteq Co - Max(R)$. By Proposition 3.2, it is clear that there exists at least one element of $m_1 \setminus IM(R)$ which is adjacent to one element of $m_2 \setminus IM(R)$. Also, $m_i \setminus IM(R)$ is totally disconnected for any $m_i \in Co - Max(R)$, so $m_1 \setminus IM(R)$ and $m_2 \setminus IM(R)$ are entirely contained in one of partitions of $\Omega_2(R) \setminus IM(R)$. This implies that $(m_1 \setminus IM(R)) \cap (m_2 \setminus IM(R)) = \emptyset$ and hence

$m_1 \cap m_2 \subseteq IM(R)$. Therefore we have $m_1 \cap m_2 = IM(R)$. Thus $|Co-Max(R)| = 2$ and by Theorem 3.4, $\Omega_2(R) \setminus IM(R)$ is a complete bipartite graph. \square

As mentioned in the introduction, Beck conjectured that $\chi(R) = \omega(R)$ for every ring R . In the following theorem we want to establish Beck's conjecture for the graph $\Omega_2(R) \setminus IM(R)$ of c-semilocal semiring.

We recall that the *chromatic number* of the graph G , denoted by $\chi(G)$, is the minimal number of colors which can be assigned to the vertices of G in such a way that any two adjacent vertices have different colors.

3.16 Theorem. *Let R be a c-semilocal semiring with $|Co-Max(R)| = n$. Then $\chi(\Omega_2(R) \setminus IM(R)) = \omega(\Omega_2(R) \setminus IM(R)) = n$.*

PROOF: Let $Co - Max(R) = \{m_1, \dots, m_n\}$. By Theorem 3.6, we know that $\omega(\Omega_2(R) \setminus IM(R)) = n$. Also, it is obvious that $\chi(G) \geq \omega(G)$ for any graph G , so $\chi(\Omega_2(R) \setminus IM(R)) \geq n$. On the other hand, $\Omega_2(R) \setminus IM(R)$ is n -partite by Theorem 3.14, thus the elements of each part can be colored by an identical color because these elements are not adjacent. Hence $\chi(\Omega_2(R) \setminus IM(R)) = n$. \square

4. Diameter and radius of $\Omega(R)$

In this section, we show that $\Omega_2(R) \setminus IM(R)$ is a connected graph and $\text{diam}(\Omega_2(R) \setminus IM(R)) \leq 3$. Also, we compute the eccentricity of the vertices of $\Omega_2(R) \setminus IM(R)$.

4.1 Theorem. *Let R be a semiring. The graph $\Omega_2(R) \setminus IM(R)$ is connected with $\text{diam}(\Omega_2(R) \setminus IM(R)) \leq 3$.*

PROOF: Let $x, y \in \Omega_2(R) \setminus IM(R)$ that are not adjacent. We consider two cases:

Case 1: Suppose that $x + y \notin IM(R)$. By Proposition 3.3, $F(x + y)F(a) = R$, for some $a \in \Omega_2(R) \setminus IM(R)$. This implies that $F(x)F(a) = F(y)F(a) = R$ since $F(x + y) \subseteq F(x), F(y)$. Hence $x - a - y$ is a path in $\Omega_2(R) \setminus IM(R)$ and $d(x, y) = 2$.

Case 2: Suppose that $x + y \in IM(R)$. Thus for each $m \in Co - Max(R)$, we have $x \in m$ or $y \in m$. Since $x \notin IM(R)$, by Proposition 3.3, there exists $a \in \Omega_2(R) \setminus IM(R)$ such that x is adjacent to a in $\Omega_2(R) \setminus IM(R)$. Hence if $x \in m$ for maximal co-ideal m , then $a \notin m$. Now, there exists $n \in Co - Max(R)$ in which $y \notin n$, since $y \notin IM(R)$. This implies that $x \in n$ and $a \notin n$. As n is prime co-ideal, we have $a + y \notin IM(R)$. So by Case 1, $d(a, y) \leq 2$ and hence $d(x, y) \leq 3$. \square

We recall that for a graph G , the *eccentricity* of a vertex x is $e(x) = \text{Max}\{d(y, x); y \in V(G)\}$. A vertex x with smallest eccentricity is called a *center* of G and its eccentricity is called the *radius* of G and is denoted by $\text{rad}(G)$.

4.2 Proposition. *Let R be a c-semilocal semiring with $|Co - Max(R)| \geq 3$. If $x \in \Omega_2(R) \setminus IM(R)$ belongs to at least two maximal co-ideals, then $e(x) = 3$.*

PROOF: Suppose that for $x \in \Omega_2(R) \setminus IM(R)$ there exist at least two maximal co-ideals m_i and m_j so that x is contained in $m_i \cap m_j$. By Theorem 4.1, $d(x, y) \leq 3$ for any $y \in \Omega_2(R) \setminus IM(R)$. Now to complete the proof, it suffices to show that, there is an element y in $\Omega_2(R) \setminus IM(R)$ such that $d(x, y) = 3$. Let $y \in \bigcap_{\substack{k=1 \\ k \neq i}}^n m_k \setminus IM(R)$. Clearly that $d(x, y) \neq 1$, since $x, y \in m_j$. If $d(x, y) = 2$, then $x - a - y$ is a path for some $a \in \Omega_2(R) \setminus IM(R)$. Now, as $x \in m_i \cap m_j$, thus $a \notin m_i, m_j$. Also, $y \in \bigcap_{\substack{k=1 \\ k \neq i}}^n m_k \setminus IM(R)$ implies that $a \notin m_k$, for $1 \leq k \leq n$ and $k \neq i$. Indeed, this implies that $a \notin m$ for any $m \in Co - Max(R)$, that is impossible. So we can conclude that $d(x, y) = 3$ and hence $e(x) = 3$. \square

4.3 Corollary. *Let R be a c -semilocal semiring with $|Co - Max(R)| \geq 3$. Then $\text{diam}(\Omega_2(R) \setminus IM(R)) = 3$.*

PROOF: We know that $\text{diam}(\Omega_2(R) \setminus IM(R)) \leq 3$, by Theorem 4.1. On the other hand, $|Co - Max(R)| \geq 3$ implies that there is an element x in $\Omega_2(R) \setminus IM(R)$ that belongs to at least two maximal co-ideals. Now, the proof is immediate from Proposition 4.2. \square

4.4 Proposition. *Let R be a semiring with $|Co - Max(R)| = 2$. If $|m_i \setminus IM(R)| \geq 2$ for some i , then $\text{diam}(\Omega_2(R) \setminus IM(R)) = 2$.*

PROOF: Assume that $|Co - Max(R)| = 2$. By Theorem 3.4, $\Omega_2(R) \setminus IM(R)$ is complete bipartite graph and thus $\text{diam}(\Omega_2(R) \setminus IM(R)) \leq 2$. On the other hand, $\text{diam}(\Omega_2(R) \setminus IM(R)) \neq 1$ because $|m_i \setminus IM(R)| \geq 2$ for some i . Hence $\text{diam}(\Omega_2(R) \setminus IM(R)) = 2$. \square

4.5 Theorem. *Let R be a semiring. If $\text{diam}(\Omega_2(R) \setminus IM(R)) = 2$, then R has an infinite number of maximal co-ideals or $|Co - Max(R)| = 2$ such that $|m_i \setminus IM(R)| \geq 2$ for some $i = 1, 2$.*

PROOF: Assume that $\text{diam}(\Omega_2(R) \setminus IM(R)) = 2$ and $|Co - Max(R)|$ is finite. If $n \geq 3$, then by Corollary 4.3, $\text{diam}(\Omega_2(R) \setminus IM(R)) = 3$, which is a contradiction. Thus we must have $|Co - Max(R)| = 2$. Now, if $|m_i \setminus IM(R)| = 1$ for each i , then $\text{diam}(\Omega_2(R) \setminus IM(R)) = 1$ because $\Omega_2(R) \setminus IM(R)$ is a complete bipartite graph, this is a contradiction. Hence $|m_i \setminus IM(R)| \geq 2$ for some i . \square

4.6 Theorem. *Let R be a c -semilocal semiring with $|Co - Max(R)| = n \geq 2$. If $\Omega_2(R) \setminus IM(R)$ is not a star graph, then we have:*

$$e(x) = \begin{cases} 2 & \text{if } x \in m_i \setminus \bigcup_{\substack{j=1 \\ j \neq i}}^n m_j \\ 3 & \text{otherwise.} \end{cases}$$

PROOF: First, we claim that for any $a \in \Omega_2(R) \setminus IM(R)$, $e(a) \neq 1$. Suppose that there is an element x of $\Omega_2(R) \setminus IM(R)$ such that $e(x) = 1$. This means that x is adjacent to any vertex of $\Omega_2(R) \setminus IM(R)$ and so $\Omega_2(R) \setminus IM(R)$ is a star graph by Proposition 3.10, which is a contradiction. Now, suppose that $x \in m_i \setminus \bigcup_{\substack{j=1 \\ j \neq i}}^n m_j$.

For any $y \in \bigcup_{j \neq i}^n m_j \setminus m_i$, if $F(x)F(y) \neq R$, then $F(x)F(y) \subseteq m_k$ for some $m_k \in Co - Max(R)$. Hence $x, y \in m_k$, that is a contradiction. Therefore, in this case $d(x, y) = 1$. But, if $y \in m_i \setminus IM(R)$ and $y \neq x$, then by proof of Theorem 4.1, $d(x, y) \leq 2$ since $x + y \notin IM(R)$. Clearly x and y are not adjacent and so $d(x, y) = 2$. According to the assumption, since $\Omega_2(R) \setminus IM(R)$ is not star graph thus by Theorem 3.11 ((4) \Rightarrow (3)) $|Co - Max(R)| \geq 2$ and $|m \setminus IM(R)| \geq 2$ for each $m \in Co - Max(R)$. Hence $e(x) = 2$ for any $x \in m_i \setminus \bigcup_{j \neq i}^n m_j$.

Now, suppose that $x \notin m_i \setminus \bigcup_{j \neq i}^n m_j$ for any maximal co-ideal m_i . Hence there are at least two maximal co-ideals m_k and m_j so that x is contained in $m_k \cap m_j$. This implies that $|Co - Max(R)| \geq 3$, thus by Proposition 4.2 we have $e(x) = 3$. □

4.7 Corollary. *Let R be a c-semilocal semiring with $|Co - Max(R)| = n \geq 2$. If $\Omega_2(R) \setminus IM(R)$ is not a star graph, then the elements of $m_i \setminus \bigcup_{j \neq i}^n m_j$ are center of $\Omega_2(R) \setminus IM(R)$ for each $m_i \in Co - Max(R)$ and $rad(\Omega_2(R) \setminus IM(R)) = 2$.*

PROOF: This is an immediate consequence of Theorem 4.6. □

4.8 Proposition. *Let R be a semiring with $|Co - Max(R)| = 2$. Then $rad(\Omega_2(R) \setminus IM(R)) = 1$ or 2 .*

PROOF: We know by Theorem 3.4, $\Omega_2(R) \setminus IM(R)$ is a complete bipartite graph when $|Co - Max(R)| = 2$. Now, if $\Omega_2(R) \setminus IM(R)$ is a star graph, clearly $rad(\Omega_2(R) \setminus IM(R)) = 1$. Otherwise, $rad(\Omega_2(R) \setminus IM(R)) = 2$ and all elements of $\Omega_2(R) \setminus IM(R)$ are center. □

5. The relations between $\Omega(R)$ and $\Gamma(R)$

In this section, we will investigate the relations between the zero-divisor graph $\Gamma(R)$ and $\Omega(R)$. We show that $\Gamma(R)$ is a subgraph of the $\Omega(R)$. Also, we determine a family of commutative semirings whose zero-divisor graph $\Gamma(R)$ and $\Omega_2(R)$ are isomorphic.

We recall that an *isomorphism* from a simple graph G to a simple graph H is a bijection $f : V(G) \rightarrow V(H)$ such that x and y are adjacent in G if and only if $f(x)$ and $f(y)$ are adjacent in H . We say G is isomorphic to H , if there is an isomorphism from G to H , denoted by $G \cong H$.

5.1 Theorem. *The zero-divisor graph $\Gamma(R)$ is a subgraph of the graph $\Omega(R)$.*

PROOF: Suppose that x and y are two distinct adjacent vertices in $\Gamma(R)$. Thus $xy = 0$ and this implies $F(x)F(y) = R$, since $0 = xy \in F(x)F(y)$. Hence x and y are adjacent in $\Omega(R)$. Now, since the vertex-set of zero-divisor graph is $Z(R)^*$, thus we can conclude that $\Gamma(R)$ is a subgraph of $\Omega(R)$. □

5.2 Theorem. *Let R be a multiplicatively idempotent and zero-sumfree semiring. Then the zero-divisor graph $\Gamma(R)$ is an induced subgraph of the graph $\Omega(R)$.*

PROOF: By Theorem 5.1, $\Gamma(R)$ is a subgraph of $\Omega(R)$. Thus it is enough to show that if $x, y \in Z(R)^*$ and they are adjacent in $\Omega(R)$, then x and y are adjacent in $\Gamma(R)$. Assume that $x, y \in Z(R)^*$ and $F(x)F(y) = R$. So we have $(x^n + r)(y^m + s) + k = 0$ for some positive integers n, m and $r, s, k \in R$. Since R is a multiplicatively idempotent, then we have $xy + a = 0$ for some $a \in R$. Hence $xy = 0$ because R is a zero-sumfree semiring. This implies x and y are adjacent in $\Gamma(R)$. \square

Note that if $UM(R) = Z(R)^*$, then $\Gamma(R)$ is a spanning subgraph of $\Omega_2(R)$ by Theorem 5.1. Thus, if R is a multiplicatively idempotent and zero-sumfree semiring, then we have the following result:

5.3 Corollary. *Let R be a multiplicatively idempotent and zero-sumfree semiring. If $Z(R)^* = UM(R)$, then the zero-divisor graph $\Gamma(R)$ and $\Omega_2(R)$ are isomorphic. In particular, if $Z(R)^* = UM(R) \setminus IM(R)$, then $\Gamma(R)$ and $\Omega_2(R) \setminus IM(R)$ are isomorphic.*

PROOF: This is an immediate consequence of Theorems 5.1 and 5.2. \square

To this end, we give an example that clarifies the previous results:

5.4 Example. Let $S = \{0, 1, a\}$ and $R = (S \times S, +, \cdot)$ be a semiring as defined in Example 3.5. We know that R is a multiplicatively idempotent. For this semiring, the vertex-set of $\Gamma(R)$ is

$$Z(R)^* = \{(0, 1), (1, 0), (0, a), (a, 0)\}$$

and the vertex-set of $\Omega_2(R)$ is $UM(R) = R \setminus \{(0, 0)\}$. Clearly $\Gamma(R)$ is an induced subgraph of $\Omega(R)$ and $\Omega_2(R)$. On the other hand, $(0, 0)$ is only zero-sum of R , thus R is zero-sumfree semiring. We see that $UM(R) \setminus IM(R) = Z(R)^*$, so we can conclude that $\Gamma(R)$ and $\Omega_2(R) \setminus IM(R)$ are isomorphic by Corollary 5.3.

REFERENCES

- [1] Akbari S., Habibi M., Majidinya A., Manaviyat R., *A note on co-maximal graph of non-commutative rings*, Algebr. Represent. Theory **16** (2013), 303–307.
- [2] Anderson D.F., Livingston P.S., *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), 434–447.
- [3] Beck I., *Coloring of commutative rings*, J. Algebra **116** (1988), 208–226.
- [4] Chaudhari J.N., Ingale K.J., *Prime avoidance theorem for co-ideals in semirings*, Research J. Pure Algebra **1(9)** (2011), 213–216.
- [5] Ebrahimi Atani S., *The zero-divisor graph with respect to ideals of a commutative semiring*, Glas. Mat. **43(63)** (2008), 309–320.
- [6] Ebrahimi Atani S., *An ideal-based zero-divisor graph of a commutative semiring*, Glas. Mat. **44(64)** (2009), 141–153.
- [7] Ebrahimi Atani S., Dolati Pish Hesari S., Khoramdel M., *Strong co-ideal theory in quotients of semirings*, J. Adv. Res. Pure Math. **5** (2013), no. 3, 19–32.
- [8] Ebrahimi Atani S., Dolati Pish Hesari S., Khoramdel M., *A fundamental theorem of co-homomorphisms for semirings*, Thai J. Math. **12** (2014), no. 2, 491–497.

- [9] Golan J.S., *Semirings and Their Applications*, Kluwer Academic Publishers, Dordrecht, 1999.
- [10] Maimani H.R., Salimi M., Sattari A., Yassemi S., *Comaximal graph of commutative rings*, *J. Algebra* **319** (2008), 1801–1808.
- [11] Sharma P.K., Bhatwadekar S.M., *A note on graphical representation of rings*, *J. Algebra* **176** (1995), 124–127.
- [12] West D.B., *Introduction to Graph Theory*, Prentice-Hall of India Pvt. Ltd, 2003.

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, UNIVERSITY OF MAZANDARAN, BABOLSAR, IRAN

E-mail: talebi@umz.ac.ir

a.darzi@stu.umz.ac.ir

(Received December 7, 2016, revised February 2, 2017)