

Suying Liu; Minghua Yang

The weighted Hardy spaces associated to self-adjoint operators and their duality on product spaces

Czechoslovak Mathematical Journal, Vol. 68 (2018), No. 2, 415–431

Persistent URL: <http://dml.cz/dmlcz/147226>

Terms of use:

© Institute of Mathematics AS CR, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE WEIGHTED HARDY SPACES ASSOCIATED TO
SELF-ADJOINT OPERATORS AND THEIR DUALITY
ON PRODUCT SPACES

SUYING LIU, Xian, MINGHUA YANG, Nanchang

Received September 7, 2016. First published February 7, 2018.

Abstract. Let L be a non-negative self-adjoint operator acting on $L^2(\mathbb{R}^n)$ satisfying a pointwise Gaussian estimate for its heat kernel. Let w be an A_r weight on $\mathbb{R}^n \times \mathbb{R}^n$, $1 < r < \infty$. In this article we obtain a weighted atomic decomposition for the weighted Hardy space $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$, $0 < p \leq 1$ associated to L . Based on the atomic decomposition, we show the dual relationship between $H_{L,w}^1(\mathbb{R}^n \times \mathbb{R}^n)$ and $\text{BMO}_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$.

Keywords: weighted Hardy space; operator; Gaussian estimate; duality; product space

MSC 2010: 42B35, 42B30, 47F05

1. INTRODUCTION

The theory of Hardy spaces has been a central part of modern harmonic analysis. The theory of Hardy spaces on product domains was initiated by Gundy and Stein in [23]. The atomic decompositions for Hardy spaces on product domains were obtained by Chang and Fefferman in [11], [13]. Later, Fefferman in [21], Krug in [29], Sato in [36] and others established the weighted theory of the classical Hardy spaces on product domains. See [42], [12], [8] for more results on product domain.

The classical Hardy spaces on \mathbb{R}^n can be characterized by certain estimates via the Laplacian, but there are some important situations in which the theory of classical Hardy spaces is not applicable. Some interesting operators were found to be out of the Calderón-Zygmund class. See [17], [2], [25], [4], [5], [6]. In order to handle such beyond Calderón-Zygmund operators, many researchers study Hardy spaces that

S. Y. Liu is supported in part by NSFC (No. 11701453) and Postdoctoral Science Foundation of China (No. 2016M600811).

are adapted to a linear operator L . For some results on the one-parameter Hardy spaces adapted to an operator, we refer the reader to [3], [20], [7], [25], [19], [26], [24], [9], [16], [38], [28] and their references. For the theory of weighted Hardy spaces associated with an operator, we refer the reader to a series of papers [1], [39], [31], [33], [34].

In [32], the authors studied the atomic decomposition of weighted Hardy spaces $H_{L,w}^1(\mathbb{R}^n \times \mathbb{R}^n)$ associated to self-adjoint operators on product spaces. Inspired by the work [32], it is natural to study the weighted Hardy space $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$, $0 < p < 1$. Let L be a non-negative self-adjoint operator acting on $L^2(\mathbb{R}^n)$ satisfying a pointwise Gaussian estimate for its heat kernel. Let w be an A_r weight on $\mathbb{R}^n \times \mathbb{R}^n$ (see Section 2 for its definition), where $1 < r < \infty$. The main purpose of this paper is to introduce a weighted Hardy space $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$ and prove a weighted atomic decomposition for $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$. See Theorem 4.1. And then, we introduce weighted BMO spaces $\text{BMO}_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$ on product spaces. Using the atomic decomposition for $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$, we show that the dual space of the weighted Hardy space $H_{L,w}^1(\mathbb{R}^n \times \mathbb{R}^n)$ is $\text{BMO}_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$. See Theorem 5.2. When L is the Laplacian of \mathbb{R}^n , the weighted atomic decompositions in product spaces were obtained in [29], [30]. Here, we generalize the results of [29], [30].

Since there is no Whitney decomposition in product domains, the situation becomes considerably complicated. As pointed out by [10], the product atoms should be supported in open sets rather than rectangles, which leads to many difficulties. The important tool is a weighted version of Journé's covering lemma, which is a good substitute in product domains for Whitney decomposition. It will be used to prove (i) of Theorem 4.1.

The layout of the article is as follows. In Section 2, we introduce some basic assumptions and state some preliminary results. In Section 3, we define the weighted Hardy space $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$, $0 < p \leq 1$ associated to a non-negative self-adjoint operator with Gaussian upper bounds on its heat kernel. We give an atomic characterization of the weighted Hardy spaces on the product domain associated to the operator. In Section 4, we state and prove the relationship between $H_{L,w}^1(\mathbb{R}^n \times \mathbb{R}^n)$ and $\text{BMO}_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$, which is the main result of this paper.

Throughout this article, the letter “ C ” or “ c ” will denote (possibly different) constants that are independent of the essential variables.

2. THE PRELIMINARIES

2.1. Assumption (H). Assume that L is a non-negative self-adjoint operator on $L^2(\mathbb{R}^n)$ and that each of the semigroups e^{-tL} , generated by $-L$ on $L^2(\mathbb{R}^n)$, has the kernel $p_t(x, y)$ which satisfies the Gaussian upper bound. That is, there exist constants $C, c > 0$ such that

$$(GE) \quad |p_t(x, y)| \leq \frac{C}{t^{n/2}} \exp\left(-\frac{|x-y|^2}{ct}\right).$$

We note that such estimates are typical for elliptic or sub-elliptic differential operators of second order (see for example, [14] and [18]).

Lemma 2.1. *Let L be an operator satisfying the assumption (H). For every $k = 0, 1, \dots$, there exist two positive constants C_k, c_k such that the kernel $p_{t,k}(x, y)$ of the operator $(t^2L)^k e^{-t^2L}$ satisfies*

$$(2.1) \quad |p_{t,k}(x, y)| \leq \frac{C_k}{(4\pi t)^n} \exp\left(-\frac{|x-y|^2}{c_k t^2}\right)$$

for all $t > 0$ and almost every $x, y \in \mathbb{R}^n$.

Proof. For the proof, we refer the reader to [14] and [35], Theorem 6.17. □

2.2. Muckenhoupt weights. We review some needed background on Muckenhoupt weights.

A weight w is a non-negative locally integrable function on \mathbb{R}^n . We say that $w \in A_p$, $1 < p < \infty$, if there exists a constant C such that for every ball $B \subseteq \mathbb{R}^n$,

$$\left(\frac{1}{|B|} \int_B w(x) \, dx\right) \left(\frac{1}{|B|} \int_B w^{-1/(p-1)}(x) \, dx\right)^{p-1} \leq C.$$

For $p = 1$, we say that $w \in A_1$ if there is a constant C such that for every ball $B \subseteq \mathbb{R}^n$,

$$\frac{1}{|B|} \int_B w(y) \, dy \leq Cw(x) \quad \text{for a.e. } x \in B.$$

We denote the Hardy-Littlewood maximal function by

$$\mathcal{M}(f)(x) := \sup_{x \in B} |B|^{-1} \int_B |f(y)| \, dy.$$

We sum up some of the properties of the above classes.

Proposition 2.2. *The following assertions hold:*

- (i) $A_1 \subseteq A_p \subseteq A_q$ for $1 \leq p \leq q < \infty$.
- (ii) If $w \in A_p$, $1 < p < \infty$, then there exists $1 < q < p$ such that $w \in A_q$.
- (iii) $A_\infty = \bigcup_{1 \leq p < \infty} A_p$.
- (iv) If $1 < p < \infty$, $w \in A_p$ if and only if $w^{1-p'} \in A_{p'}$.
- (v) Let $w \in A_p$, $p \geq 1$. Then for any ball B and $\lambda > 1$, there exists a constant C (independent of B and λ) such that

$$w(\lambda B) \leq C \lambda^{np} w(B).$$

- (vi) If $1 < p < \infty$, $w \in A_p$, then

$$\int_{\mathbb{R}^n} (\mathcal{M}(f)(x))^p w(x) \, dx \leq C \int_{\mathbb{R}^n} |f(x)|^p w(x) \, dx.$$

P r o o f. Properties (i)–(vi) are standard, see for instance [22] and [40]. □

2.3. Muckenhoupt weights on product spaces. In this section, we recall some basic facts on the product $A_p(\mathbb{R}^n \times \mathbb{R}^n)$.

A non-negative locally integrable function $w(x, y)$ on $\mathbb{R}^n \times \mathbb{R}^n$ is said to belong to $A_p(\mathbb{R}^n \times \mathbb{R}^n)$, $1 < p < \infty$, if there exists a constant C such that

$$\left(\frac{1}{|R|} \iint_R w(x, y) \, dx \, dy \right) \left(\frac{1}{|R|} \iint_R w(x, y)^{-1/(p-1)} \, dx \, dy \right)^{p-1} \leq C,$$

where R runs over all rectangles in $\mathbb{R}^n \times \mathbb{R}^n$. When $p = 1$, we say that $w \in A_1(\mathbb{R}^n \times \mathbb{R}^n)$ if there exists a constant C such that

$$\frac{1}{|R|} \iint_R w(x, y) \, dx \, dy \leq C \operatorname{ess\,inf}_{(x,y) \in R} w(x, y) \quad \text{for all } R.$$

We also define $A_\infty(\mathbb{R}^n \times \mathbb{R}^n) := \bigcup_{r \geq 1} A_r(\mathbb{R}^n \times \mathbb{R}^n)$. For any $1 \leq p < \infty$, the weighted Lebesgue spaces $L_w^p(\mathbb{R}^n \times \mathbb{R}^n)$ can be defined by

$$\left\{ f: \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f(x, y)|^p w(x, y) \, dx \, dy < \infty \right\}$$

with the norm

$$\|f\|_{L_w^p} := \left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f(x, y)|^p w(x, y) \, dx \, dy \right)^{1/p}.$$

We denote the strong maximal function by

$$\mathcal{M}_s(f)(x, y) := \sup_{(x, y) \in R} |R|^{-1} \iint_R |f(u, v)| \, du \, dv,$$

where R is any rectangle in $\mathbb{R}^n \times \mathbb{R}^n$.

The following properties of product A_p weights are well known, see [36], [30].

Proposition 2.3. *The following assertions hold:*

- (i) *If $w(x, y) \in A_r(\mathbb{R}^n \times \mathbb{R}^n)$, then $w(x, \cdot)$ satisfies $A_r(\mathbb{R}^n)$ condition uniformly for a.e. x ; similarly for $w(\cdot, y)$.*
- (ii) *Product weights satisfy $A_1 \subseteq A_r \subseteq A_t \subseteq A_\infty$ for $1 \leq r \leq t < \infty$.*
- (iii) *If $w \in A_r(\mathbb{R}^n \times \mathbb{R}^n)$, $1 < r < \infty$, there exists $1 < s < r$ such that $w \in A_s(\mathbb{R}^n \times \mathbb{R}^n)$.*
- (iv) *If $1 < r < \infty$, $w \in A_r(\mathbb{R}^n \times \mathbb{R}^n)$ if and only if $w^{1-r'} \in A_{r'}(\mathbb{R}^n \times \mathbb{R}^n)$.*
- (v) *If $1 < r < \infty$, $w \in A_r(\mathbb{R}^n \times \mathbb{R}^n)$, then*

$$\left\| \left(\sum_k \mathcal{M}_s^2(f_k) \right)^{1/2} \right\|_{L_w^r} \leq C \left\| \left(\sum_k |f_k|^2 \right)^{1/2} \right\|_{L_w^r}.$$

3. WEIGHTED PRODUCT HARDY SPACES AND ATOMS ASSOCIATED TO OPERATORS

3.1. Weighted product Hardy spaces and weighted product atoms. Given a function f on $\mathbb{R}^n \times \mathbb{R}^n$, $K = [n/2] + 1$, let us use Q_{t_1, t_2} to denote the operator $(t_1^2 L)^K e^{-t_1^2 L} \otimes (t_2^2 L)^K e^{-t_2^2 L}$. The area function associated with operators L is defined by

$$(3.1) \quad S_{\alpha_1, \alpha_2} f(x) := \left(\iint_{\substack{|x_1 - y_1| < \alpha_1 t_1 \\ |x_2 - y_2| < \alpha_2 t_2}} |Q_{t_1, t_2} f(y)|^2 \frac{dy \, dt}{t_1^{n+1} t_2^{n+1}} \right)^{1/2},$$

where $\alpha_i > 0$, $x = (x_1, x_2)$, $y = (y_1, y_2)$, $t = (t_1, t_2)$ and $x_i, y_i \in \mathbb{R}^n$, $t_i \in (0, \infty)$ for $i = 1, 2$. Here, we replace $(t_i^2 L) e^{-t_i^2 L}$ by $(t_i^2 L)^K e^{-t_i^2 L}$ for technical reasons. For simplicity, we write $S(f) := S_{1,1}(f)$. For the theory of unweighted Hardy spaces associated to the operator L on product domains, we refer to [15], [37].

Definition 3.1. Let L be an operator satisfying the assumption (H) and $0 < p \leq 1$. Suppose that $w \in A_\infty(\mathbb{R}^n \times \mathbb{R}^n)$. The weighted product Hardy space $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$ associated to L is defined as the completion of

$$\{f \in L^2(\mathbb{R}^n \times \mathbb{R}^n) : \|S(f)\|_{L_w^p(\mathbb{R}^n \times \mathbb{R}^n)} < \infty\}$$

with respect to the norm

$$\|f\|_{H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)} = \|S(f)\|_{L_w^p(\mathbb{R}^n \times \mathbb{R}^n)}.$$

Remark 3.2. An argument similar to Theorem 1 of Chapter IV in [41] implies the following result. Suppose $w \in A_r(\mathbb{R}^n \times \mathbb{R}^n)$, $1 \leq r < \infty$. Then for all $\alpha_1 \geq 1$, $\alpha_2 \geq 1$, there exist positive constants $c_{\alpha_1, \alpha_2, p}$ and $C_{\alpha_1, \alpha_2, p}$ such that

$$c_{\alpha_1, \alpha_2, p} \|S(f)\|_{L_w^1(\mathbb{R}^n \times \mathbb{R}^n)} \leq \|S_{\alpha_1, \alpha_2}(f)\|_{L_w^1(\mathbb{R}^n \times \mathbb{R}^n)} \leq C_{\alpha_1, \alpha_2, p} \|S(f)\|_{L_w^1(\mathbb{R}^n \times \mathbb{R}^n)}.$$

We now introduce the product (p, r, M, w) -atom associated to operators.

Definition 3.3. Suppose that $M \in \mathbb{N}$, $0 < p \leq 1$ and $w \in A_r$, $1 \leq r < \infty$. A function $a \in L^2(\mathbb{R}^{2n})$ is called a product (p, r, M, w) -atom associated to L , if it satisfies

- (1) $\text{supp } a \subseteq \Omega$, where Ω is an open set with finite measure;
- (2) a can be further decomposed into $a = \sum_{R \in m(\Omega)} a_R$, where for each $R \in m(\Omega)$

there exists a function $b_R \in \mathcal{D}(L^M \otimes L^M)$ such that

- (i) $a_R = (L^M \otimes L^M)b_R$;
- (ii) $\text{supp}(L^{k_1} \otimes L^{k_2})b_R \subseteq 10R$, $k_1, k_2 = 0, 1, \dots, M$;
- (iii) $\|a\|_{L_w^r(\mathbb{R}^n \times \mathbb{R}^n)} \leq w(\Omega)^{1/r-1/p}$ and for $k_1, k_2 = 0, 1, \dots, M$,

$$\sum_{R \in m(\Omega)} \ell(I)^{-2rM} \ell(J)^{-2rM} \|(\ell(I)^2 L)^{k_1} \otimes (\ell(J)^2 L)^{k_2} b_R\|_{L_w^r(\mathbb{R}^n \times \mathbb{R}^n)} \leq w(\Omega)^{1-r/p},$$

where $m(\Omega)$ denotes the set of maximal dyadic subrectangles of Ω ; $L^0 := \mathbb{I}$ denotes the identity operator on \mathbb{R}^n ; $R = I \times J$ denotes the dyadic rectangle of $\mathbb{R}^n \times \mathbb{R}^n$, I, J denote the dyadic cube of \mathbb{R}^n ; $10R$ denotes the 10-fold dilate of R concentric with R .

Now we can define a weighted product Hardy space $H_{L,w}^{p,r,M}$ by atoms.

Definition 3.4. Let M and w be the same as in Definition 3.3. The weighted product Hardy space $H_{L,w}^{p,r,M}(\mathbb{R}^n \times \mathbb{R}^n)$ is defined as follows. We say that $f = \sum_j \lambda_j a_j$ is a product atomic (p, r, M, w) -representation of f if $\{\lambda_j\}_{j=0}^\infty \in \ell^p$, each a_j is a product (p, r, M, w) -atom, and the sum converges in $L^2(\mathbb{R}^{2n})$. Set

$$\mathbb{H}_{L,w}^{p,r,M}(\mathbb{R}^n \times \mathbb{R}^n) = \{f: f \text{ has a product atomic } (p, r, M, w)\text{-representation}\},$$

with the norm given by

$$\begin{aligned} & \|f\|_{\mathbb{H}_{L,w}^{p,r,M}(\mathbb{R}^n \times \mathbb{R}^n)} \\ &= \inf \left\{ \left(\sum_{j=0}^\infty |\lambda_j|^p \right)^{1/p} : f = \sum_j \lambda_j a_j \text{ has a product atomic } (p, r, M, w)\text{-representation} \right\}. \end{aligned}$$

The space $H_{L,w}^{p,r,M}(\mathbb{R}^n \times \mathbb{R}^n)$ is then defined as the completion of $\mathbb{H}_{L,w}^{p,r,M}(\mathbb{R}^n \times \mathbb{R}^n)$ with respect to this norm.

3.2. Some auxiliary lemmas. For any $\alpha > 0$, let us introduce the product cone $\Gamma(x) := \Gamma(x_1) \times \Gamma(x_2)$, where $\Gamma(x_i) := \{(y_i, t_i) \in \mathbb{R}_+^{n+1} : |x_i - y_i| < t_i\}$, $i = 1, 2$. Suppose $\Omega \subseteq \mathbb{R}^n \times \mathbb{R}^n$ is an open set of finite measure. Denote by $m(\Omega)$ the maximal dyadic subrectangles of Ω . Let $m_1(\Omega)$ denote those dyadic subrectangles $R \subseteq \Omega$, $R = I \times J$ which are maximal in the x_1 direction. In other words, if $S = I' \times J \supseteq R$ is a dyadic subrectangle of Ω , then $I = I'$. Define $m_2(\Omega)$ similarly.

In order to prove our main result, we need some auxiliary results.

Lemma 3.5. *Let $w \in A_\infty$ and let Ω be an open set of $\mathbb{R}^n \times \mathbb{R}^n$. For any $R = I \times J \in m_2(\Omega)$, we set*

$$\gamma_1(R) := \sup_{\substack{I' \subseteq I \\ I' \times J \subseteq \Omega^*}} |I'|/|I|,$$

where $\Omega^* = \{x \in \mathbb{R}^n \times \mathbb{R}^n : \mathcal{M}_s(\chi_\Omega)(x) > 1/2\}$. Then for any $\delta > 0$,

$$\sum_{R \in m_2(\Omega)} w(R) \gamma_1^{-\delta}(R) \leq c_\delta w(\Omega),$$

where c_δ is a constant depending only on δ , not on Ω .

Similarly, for any $R = I \times J \in m_1(\Omega)$, we set

$$\gamma_2(R) := \sup_{\substack{J' \subseteq J \\ I \times J' \subseteq \Omega^*}} |J'|/|J|.$$

Then for any $\delta > 0$,

$$\sum_{R \in m_1(\Omega)} w(R) \gamma_2^{-\delta}(R) \leq c_\delta w(\Omega),$$

where c_δ is a constant depending only on δ , not on Ω .

Proof. For the proof, we refer to [21], Lemma 2, and [27]. In fact, this is a weighted version of Journé's lemma. \square

We also need the following results.

Lemma 3.6. *Let L be an operator satisfying the assumption (H). Then for any $1 < r < \infty$ and $w \in A_r(\mathbb{R}^n \times \mathbb{R}^n)$, there exist constants C_1, C_2 such that*

$$(3.2) \quad C_1 \|f\|_{L_w^r(\mathbb{R}^n \times \mathbb{R}^n)} \leq \|S(f)\|_{L_w^r(\mathbb{R}^n \times \mathbb{R}^n)} \leq C_2 \|f\|_{L_w^r(\mathbb{R}^n \times \mathbb{R}^n)}$$

holds for all $f \in L_w^r(\mathbb{R}^n \times \mathbb{R}^n)$.

Proof. For the details of the proof, we refer the reader to [32]. □

Lemma 3.7. Fix $M \in \mathbb{N}$ and $w \in A_\infty(\mathbb{R}^n \times \mathbb{R}^n)$, $0 < p \leq 1$. Suppose that T is a non-negative sublinear operator satisfying the weak-type $(2, 2)$ bound

$$\omega\{x \in \mathbb{R}^n \times \mathbb{R}^n : |Tf(x)| > \eta\} \leq C_T \eta^{-2} \|f\|_{L^2(\mathbb{R}^n \times \mathbb{R}^n)}^2, \quad \eta > 0,$$

and for every product (p, r, M, w) -atom a we have

$$\|Ta\|_{L_w^p(\mathbb{R}^n \times \mathbb{R}^n)} \leq C,$$

with constant C independent of a . Then T is bounded from $H_{L,w}^{p,r,M}(\mathbb{R}^n \times \mathbb{R}^n)$ to $L_w^p(\mathbb{R}^{2n})$, and

$$\|Tf\|_{L_w^p(\mathbb{R}^{2n})} \leq C \|f\|_{H_{L,w}^{p,r,M}(\mathbb{R}^n \times \mathbb{R}^n)}.$$

Proof. The proof is similar to that of Lemma 4.3 in [24] and so we skip it here. □

4. THE ATOMIC CHARACTERIZATIONS OF WEIGHTED PRODUCT HARDY SPACES

In this section we will state and prove the atomic characterizations of weighted product Hardy spaces $H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)$, $0 < p \leq 1$.

Theorem 4.1. Suppose that $w \in A_s(\mathbb{R}^n \times \mathbb{R}^n)$, $1 < s < \infty$ and $0 < p \leq 1$.

- (i) Suppose that $M \in \mathbb{N}$, $M > (r/p - 1)n/2$ and $r \geq s$. Let $f = \sum_{i=0}^{\infty} \lambda_i a_i$, where $\{\lambda_i\}_{i=0}^{\infty} \in l^p$, the a_i are product (p, r, M, w) -atoms, and the sum converges in $L^2(\mathbb{R}^{2n})$. Then $f \in H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n) \cap L^2(\mathbb{R}^{2n})$ and

$$\left\| \sum_{i=0}^{\infty} \lambda_i a_i \right\|_{H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)} \leq C \left(\sum_{i=0}^{\infty} |\lambda_i|^p \right)^{1/p}.$$

- (ii) Let $M \in \mathbb{N}$ and $1 < r < \infty$. If $f \in H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n) \cap L^2(\mathbb{R}^{2n})$, then there exist a family of product (p, r, M, w) -atoms $\{a_i\}_{i=0}^{\infty}$ and a sequence of numbers $\{\lambda_i\}_{i=0}^{\infty}$ such that f can be represented in the form $f = \sum_{i=0}^{\infty} \lambda_i a_i$, and the sum converges in the sense of $L^2(\mathbb{R}^{2n})$ -norm. Moreover,

$$\left(\sum_{i=0}^{\infty} |\lambda_i|^p \right)^{1/p} \leq C \|f\|_{H_{L,w}^p(\mathbb{R}^n \times \mathbb{R}^n)}.$$

The proof of (ii) in Theorem 4.1 is very similar to the case $p = 1$ considered by [32] (in fact, one just needs to set $\lambda_i := c_{\Psi} 2^i w(\Omega_i)^{1/p}$ to get the result), we do not discuss it here. In this paper, we give the details of the proof of (i).

P r o o f of (i) of Theorem 4.1. Let $f(x) = \sum_j \lambda_j a_j(x)$, where the a_j are product (p, r, M, w) -atoms and $\left(\sum_{i=0}^{\infty} |\lambda_i|^p\right)^{1/p} < \infty$. By applying Lemma 3.7 and the fact that a (p, s, M, w) -atom is also a (p, r, M, w) -atom for $s \geq r > 1$, it shall be enough to show that for every product (p, r, M, w) -atom a there exists a constant C (independent of a) such that

$$\|S(a)\|_{L_w^p(\mathbb{R}^n \times \mathbb{R}^n)} \leq C.$$

By the assumption on M , we can choose N such that $n(r/p - 1) < N < 2M$. We assume that $a = \sum_{R \in m(\Omega)} a_R$ is supported in some open set Ω with finite measure. For any $R = I \times J \subseteq \Omega$, let I' be the biggest dyadic cube containing I such that

$$I' \times J \subseteq \tilde{\Omega} = \{x \in \mathbb{R}^n \times \mathbb{R}^n : \mathcal{M}_s(\chi_{\Omega})(x) > 1/2\}.$$

Then $I' \times J$ is in $m_1(\tilde{\Omega})$ and let S be the biggest dyadic cube such that $S \supseteq J$ and $I' \times S \subseteq \tilde{\tilde{\Omega}}$, where $\tilde{\tilde{\Omega}} = \{x \in \mathbb{R}^n \times \mathbb{R}^n : \mathcal{M}_s(\chi_{\tilde{\Omega}})(x) > 1/2\}$. Let \check{R} be the 10-fold dilate of $I' \times S$ concentric with $I' \times S$. By (v) of Proposition 2.3, we have $w(\bigcup \check{R}) \leq Cw(\Omega)$.

Making use of Hölder's inequality and the definition of product atoms, we have

$$\begin{aligned} \int_{\bigcup \check{R}} S(a)^p(x)w(x) dx &\leq \int_{\bigcup \check{R}} S(a)(x)w^{p/r}(x)w^{1-p/r}(x) dx \\ &\leq C \left(\int_{\bigcup \check{R}} w dx \right)^{1-p/r} \|S(a)\|_{L_w^r}^p \\ &\leq Cw(\Omega)^{1-p/r} \|a\|_{L_w^r}^p \\ &\leq Cw(\Omega)^{1-1/p} w(\Omega)^{(1/r-1/p)p} \\ &\leq C, \end{aligned}$$

where in the third inequality we have used Lemma 3.6.

It remains to prove

$$(4.1) \quad \int_{(\bigcup \check{R})^c} S(a)^p(x)w(x) dx \leq C.$$

By the definition of product atoms, one can write

$$\begin{aligned}
 (4.2) \quad \int_{(\cup \check{R})^c} S(a)^p(x)w(x) dx &\leq \sum_{R \in m(\Omega)} \int_{\check{R}^c} S(a_R)^p(x)w(x) dx \\
 &\leq \sum_{R \in m(\Omega)} \int_{(10I')^c \times \mathbb{R}^n} S(a_R)^p(x)w(x) dx \\
 &\quad + \sum_{R \in m(\Omega)} \int_{\mathbb{R}^n \times (10S)^c} S(a_R)^p(x)w(x) dx \\
 &=: \mathbf{D} + \mathbf{E}.
 \end{aligned}$$

We only estimate the term \mathbf{D} since the term \mathbf{E} can be estimated similarly.

For the term \mathbf{D} , we have

$$\begin{aligned}
 \int_{(10I')^c \times \mathbb{R}^n} S(a_R)^p(x)w(x) dx &= \int_{(10I')^c \times (10J)^c} S(a_R)^p(x_1, x_2)w(x_1, x_2) dx_1 dx_2 \\
 &\quad + \int_{(10I')^c \times 10J} S(a_R)^p(x_1, x_2)w(x_1, x_2) dx_1 dx_2 \\
 &=: \mathbf{D}_1 + \mathbf{D}_2.
 \end{aligned}$$

By the same analysis as that of Theorem 4.1 in [32], we know that

$$\begin{aligned}
 (4.3) \quad \mathbf{D}_1 &\leq C \left(\frac{|R|}{w(R)^{1/p}} \|c_R\|_{L_w^p} \right)^p \\
 &\quad \times \int_{(10I')^c \times (10J)^c} w(x) \frac{\ell(I)^{pN}}{|x_1 - x_I|^{p(n+N)}} \frac{\ell(J)^{pN}}{|x_2 - x_J|^{p(n+N)}} dx
 \end{aligned}$$

where $c_R := |a_R| + \ell(J)^{-2M}|a_{R,1}| + \ell(I)^{-2M}|a_{R,2}| + \ell(I)^{-2M}\ell(J)^{-2M}|b_R|$, and we write $a_{R,1} := (L^M \otimes \mathbb{1})b_R$ and $a_{R,2} := (\mathbb{1} \otimes L^M)b_R$.

Noting that $N > n(r/p - 1)$, one can compute

$$\begin{aligned}
 (4.4) \quad &\int_{(10I')^c \times (10J)^c} w(x) \frac{\ell(I)^{pN}}{|x_1 - x_I|^{p(n+N)}} \frac{\ell(J)^{pN}}{|x_2 - x_J|^{p(n+N)}} dx \\
 &\leq C \sum_{l_1=1}^{\infty} \sum_{l_2=1}^{\infty} \int_{2^{l_1} I' \setminus 2^{l_1-1} I'} \int_{2^{l_2} J \setminus 2^{l_2-1} J} w(x_1, x_2) \frac{\ell(I)^{pN}}{|x_1 - x_I|^{p(n+N)}} \\
 &\quad \times \frac{\ell(J)^{pN}}{|x_2 - x_J|^{p(n+N)}} dx_2 dx_1
 \end{aligned}$$

$$\begin{aligned}
&\leq C \sum_{l_1=1}^{\infty} \sum_{l_2=1}^{\infty} \frac{\ell(I)^{pN}}{(2^{l_1} \ell(I'))^{p(n+N)}} \frac{\ell(J)^{pN}}{(2^{l_2} \ell(J))^{p(n+N)}} w(2^{l_1} I' \times 2^{l_2} J) \\
&\leq C \sum_{l_1=1}^{\infty} \sum_{l_2=1}^{\infty} \frac{1}{2^{l_1(p(n+N)-rn)}} \frac{1}{2^{l_2(p(n+N)-rn)}} \frac{\ell(I)^N}{\ell(I')^{n+N}} w(R) |J|^{-p} \\
&\leq C \left(\frac{\ell(I)}{\ell(I')} \right)^{pN} w(R) |R|^{-p},
\end{aligned}$$

where in the third inequality above we have used (v) of Proposition 2.2.

Combining (4.3) and (4.4), we write

$$(4.5) \quad D_1 \leq C \gamma_1(R)^{-pN} w(R)^{1-p/r} \|c_R\|_{L_w^r}^p,$$

where $\gamma_1(R) = \ell(I')/\ell(I)$.

We turn to estimating the term D_2 . By Hölder's inequality, we have

$$\begin{aligned}
D_2 &\leq \sum_{k=3}^{\infty} \int_{2^{k+1} I' \setminus 2^k I'} \int_{10J} |S(a_R)(x_1, x_2)|^p w(x_1, x_2) dx_2 dx_1 \\
&\leq \sum_{k=3}^{\infty} \left(\int_{2^{k+1} I' \setminus 2^k I'} \int_{10J} |S(a_R)(x_1, x_2)|^r w(x_1, x_2) dx_2 dx_1 \right)^{p/r} \\
&\quad \times \left(\int_{2^{k+1} I'} \int_{10J} w(x_1, x_2) dx_2 dx_1 \right)^{1-p/r}.
\end{aligned}$$

We have

$$\begin{aligned}
(4.6) \quad &\int_{2^{k+1} I' \setminus 2^k I'} \int_{10J} |S(a_R)(x_1, x_2)|^r w(x_1, x_2) dx_2 dx_1 \\
&\leq C \int_{2^{k+1} I' \setminus 2^k I'} \int \left(\iint_{\Gamma(x_1)} |(t_1^2 L)^K e^{-t_1^2 L} a_R(y_1, x_2)|^2 \frac{dy_1 dt_1}{t_1^{n+1}} \right)^{r/2} \\
&\quad \times w(x_1, x_2) dx_2 dx_1.
\end{aligned}$$

By an argument similar to that of $D_{1,i}$, $i = 1, \dots, 4$ we obtain

$$\begin{aligned}
&\iint_{\Gamma(x_1)} |(t_1^2 L)^K e^{-t_1^2 L} a_R(y_1, x_2)|^2 \frac{dy_1 dt_1}{t_1^{n+1}} \\
&\leq C \frac{\ell(I)^{2N}}{|x_1 - x_I|^{2(n+N)}} \left[\left(\int |a_R(z_1, x_2)| dz_1 \right)^2 + \left(\ell(I)^{-2M} \int |a_{R,2}(z_1, x_2)| dz_1 \right)^2 \right],
\end{aligned}$$

since $2M > N$.

Let $\mathcal{M}^{(1)}$ denote the Hardy–Littlewood maximal operator in the first variable. Then we have

$$\begin{aligned} \int |c_R(z_1, x_2)| dz_1 &= \frac{|2^{k+1}I'|}{|2^{k+1}I'|} \int_{2^{k+1}I'} |c_R(z_1, x_2)| dz_1 \\ &\leq |2^{k+1}I'| \inf_{z_1 \in 2^{k+1}I'} \mathcal{M}^{(1)} c_R(z_1, x_2). \end{aligned}$$

Thus, one can write

$$\begin{aligned} (4.7) \quad &\left(\int_{2^{k+1}I' \setminus 2^k I'} \int_{10J} |S(a_R)(x)|^r w(x) dx \right)^{1/r} \\ &\leq C \frac{\ell(I)^N 2^{kn} |I'|}{(2^k \ell(I'))^{n+N}} \left(\int_{2^{k+1}I'} \int \left[\inf_{z_1 \in 2^{k+1}I'} \mathcal{M}^{(1)} c_R(z_1, x_2) \right]^r \right. \\ &\quad \left. \times w(x_1, x_2) dx_2 dx_1 \right)^{1/r} \\ &\leq C \gamma_1(R)^{-N} 2^{-kN} \left(\iint [\mathcal{M}^{(1)} c_R(x_1, x_2)]^r w(x_1, x_2) dx_2 dx_1 \right)^{1/r} \\ &\leq C \gamma_1(R)^{-N} 2^{-kN} \|c_R\|_{L_w^r}, \end{aligned}$$

where in the third inequality above we have used (vi) of Proposition 2.2 and (i) of Proposition 2.3.

By virtue of (v) of Proposition 2.2 we have

$$(4.8) \quad \left(\int_{2^{k+1}I'} \int_{10J} w(x) dx \right)^{1-p/r} \leq C 2^{(k+1)n(r-p)} w(R)^{1-p/r}.$$

Combining (4.7) and (4.8), we have

$$\begin{aligned} D_2 &\leq C \sum_{k=3}^{\infty} 2^{-k(pN-n(r-p))} \|c_R\|_{L_w^p}^p \gamma_1(R)^{-pN} w(R)^{1-p/r} \\ &\leq C \|c_R\|_{L_w^p}^p \gamma_1(R)^{-pN} w(R)^{1-p/r}, \end{aligned}$$

since $N > n(r/p - 1)$.

Estimates of D_1 and D_2 , together with Hölder's inequality and weighted Journé's covering lemma (Lemma 3.5), imply that

$$\begin{aligned} D &\leq C \sum_{R \in m(\Omega)} \|c_R\|_{L_w^p}^p \gamma_1(R)^{-pN} w(R)^{1-p/r} \\ &\leq C \left(\sum_{R \in m(\Omega)} \|c_R\|_{L_w^r}^r \right)^{p/r} \left(\sum_{R \in m(\Omega)} w(R) \gamma_1(R)^{-Np(r-p)/r} \right)^{1-p/r} \\ &\leq C w(\Omega)^{p/r-1} w(\Omega)^{1-p/r} \leq C. \end{aligned}$$

The desired estimate of (4.1) follows readily. This completes the proof of (i) of Theorem 4.1. \square

5. $BMO_{L,w}$: DUALITY WITH $H^1_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$ SPACES

In this section, we introduce and study the duality of the weighted Hardy spaces $H^1_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$. Given a real-valued function f on $\mathbb{R}^n \times \mathbb{R}^n$ and a weight $w \in A_s(\mathbb{R}^n \times \mathbb{R}^n)$, $1 \leq s < \infty$, we consider the following situation: if Ω is a bounded open set in $\mathbb{R}^n \times \mathbb{R}^n$, and $R = I \times J$ runs over the collection of the maximal dyadic rectangles contained in Ω , then we introduce the following definition.

Definition 5.1. Let L be a non-negative self-adjoint operator such that the corresponding heat kernel satisfies conditions (GE). For $w \in A_s$, $1 \leq s < \infty$ and $1 \leq p < \infty$, an element $f \in L^2$ is said to belong to $BMO^p_{L,w}$ if

$$\begin{aligned} & \|f\|_{BMO^p_{L,w}} \\ & =: \sup_{\Omega} \left(\frac{1}{w(\Omega)} \sum_{R \subset \Omega} \iint_{I \times J} |(\mathbb{1} - (1 + \ell(I)^2 L \otimes \ell(J)^2 L)^{-1})^M f|^p w^{1-p} dx \right)^{1/p} < \infty, \end{aligned}$$

where the supremum ranges over the whole open set Ω of $\mathbb{R}^n \times \mathbb{R}^n$ with finite measure, and $\mathbb{1}$ denotes the identity operator on $\mathbb{R}^n \times \mathbb{R}^n$. In particular, for $p = 1$ we denote $BMO^1_{L,w} =: BMO_{L,w}$.

We have the following theorem. In fact, it expresses the dual relationship of $BMO_{L,w}$ and $H^1_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$.

Theorem 5.2. $H^{1,r,M}_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)^* = BMO^{r'}_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$, $r \geq 1$.

Proof. First, we show that each $f \in BMO^{r'}_{L,w}$ induces a bounded linear functional on $H^{1,r,M}_{L,w}(\mathbb{R}^n \times \mathbb{R}^n)$. Suppose that a is a $(1, r, M, w)$ -atom in $H^{1,r,M}_{L,w}(\mathbb{R}^n)$, and let $f \in BMO^{r'}_{L,w}(\mathbb{R}^n)$. For simplicity, denote $\mathbb{1} - (1 + \ell(I)^2 L \otimes \ell(J)^2 L)^{-1})^M =: P_M$, then by the atomic definition we have

$$\begin{aligned} & \left| \iint_{\mathbb{R}^n \times \mathbb{R}^n} a(x, y) f(x, y) dx dy \right| \leq \sum_R \left| \iint_{10R} a_R(x, y) f(x, y) dx dy \right| \\ & \leq \sum_R \left| \iint_{10R} P_M a_R(x, y) f(x, y) dx dy \right| \\ & \quad + \sum_R \left| \iint_{10R} (\mathbb{1} - P_M) a_R(x, y) f(x, y) dx dy \right| =: J_1 + J_2. \end{aligned}$$

For the term J_1 , by Hölder's inequality and the definitions of atom and $f \in \text{BMO}_{L,w}^{r'}$ we obtain

$$\begin{aligned} J_1 &\leq \left(\sum_R \|a_R\|_{L_w^{r'}}^r \right)^{1/r} \sum_R \left(\iint_{10R} |P_M f(x,y)|^{r'} w^{1-r'} dx dy \right)^{1/r'} \\ &\leq C \|f\|_{\text{BMO}_{L,1/w}^{r'}} w(B)^{1/r-1} w(B)^{1/r'} \\ &\leq C \|f\|_{\text{BMO}_{L,1/w}^{r'}}. \end{aligned}$$

To analyse J_2 , by the condition $a_R = (L^M \otimes L^M)b_R$ and the fact that L is self-adjoint, we write

$$\begin{aligned} (\mathbb{1} - P_M)a_R &= (\mathbb{1} - P_M)(L^M \otimes L^M)b_R = (L^M \otimes L^M)(\mathbb{1} - P_M)b_R \\ &= \sum_{k_1=1}^M \sum_{k_2=1}^M \frac{M!}{(M-k_1)k_1!} \frac{M!}{(M-k_2)k_2!} (\ell(I)^{-2k_1} L^{M-k_1} \otimes \ell(J)^{-2k_2} L^{M-k_2}) P_M b_R \\ &=: \sum_{k_1=1}^M \sum_{k_2=1}^M C_{k_1} C_{k_2} (\ell(I)^{-2k_1} L^{M-k_1} \otimes \ell(J)^{-2k_2} L^{M-k_2}) P_M b_R. \end{aligned}$$

Thus, by the definitions of an atom and $f \in \text{BMO}_{L,w}^{q'}$ and Hölder's inequality, we have

$$\begin{aligned} J_2 &\leq \sum_{k_1=1}^M \sum_{k_2=1}^M C_{k_1} C_{k_2} \sum_R \left| \ell(I)^{-2M} \ell(J)^{-2M} \right. \\ &\quad \times \left. \iint_{I \times J} (\ell(I)^2 L)^{M-k_1} \otimes (\ell(J)^2 L)^{M-k_2} b_R P_M f(x,y) dx dy \right| \\ &\leq \sum_{k_1=1}^M \sum_{k_2=1}^M C_{k_1} C_{k_2} \\ &\quad \times \left(\sum_R |\ell(I)^{-2rM} \ell(J)^{-2rM}| (\ell(I)^2 L)^{M-k_1} \otimes (\ell(J)^2 L)^{M-k_2} b_R \|_{L_w^{r'}} \right)^{1/r} \\ &\quad \times \sum_R \iint_{I \times J} |P_M f(x,y)|^{r'} w^{1-r'} dx dy \leq C w(\Omega)^{1/r-1} w(\Omega)^{1/r'} \leq C. \end{aligned}$$

Therefore, for every $h = \sum_j \lambda_j a_j \in H_{L,w}^{1,q,M}(\mathbb{R}^n \times \mathbb{R}^n)$, where a_j are weighted atoms, we have

$$\begin{aligned} \left| \iint_{\mathbb{R}^n \times \mathbb{R}^n} f(x,y) h(x,y) dx dy \right| &\leq \sum_j |\lambda_j| \left| \iint_{\mathbb{R}^n \times \mathbb{R}^n} f(x,y) a_j(x,y) dx dy \right| \\ &\leq C \sum_j |\lambda_j| \|f\|_{\text{BMO}_{L,1/w}^{q'}} \leq C \|h\|_{H_{L,w}^1} \|f\|_{\text{BMO}_{L,w}^{q'}}. \end{aligned}$$

Conversely, suppose that $l \in H_{L,w}^{1,r,M}(\mathbb{R}^n \times \mathbb{R}^n)^*$ can be represented by $f(x, y)$ in the form

$$l(g) = \iint_{\mathbb{R}^n \times \mathbb{R}^n} f(x, y)g(x, y) \, dx \, dy.$$

Let Ω be a bounded open set in $\mathbb{R}^n \times \mathbb{R}^n$, and suppose that $\Omega = \bigcup_R R$, where the R 's are the maximal dyadic rectangles contained in Ω , and that

$$\left(\sum_R \|\varphi\|_{L_w^r(R)}^r \right)^{1/r} = 1.$$

Set

$$a(x, y) = \frac{1}{w(\Omega)^{1-1/r}} \sum_R (\mathbb{1} - (1 + \ell(I)^2 L \otimes \ell(J)^2 L)^{-1})^M \varphi.$$

Then it is not difficult to check that a is a $(1, r, M, w)$ -atom (see Theorem 6.4 in [24]).

Consequently,

$$\begin{aligned} \|l\| &\geq \|l(a)\| \\ &= \frac{1}{w(\Omega)^{1-1/r}} \sum_R \iint_R (\mathbb{1} - (1 + \ell(I)^2 L \otimes \ell(J)^2 L)^{-1})^M \varphi(x, y) f(x, y) \, dx \, dy \\ &= \frac{1}{w(\Omega)^{1-1/r}} \sum_R \iint_R \varphi(x, y) (\mathbb{1} - (1 + \ell(I)^2 L \otimes \ell(J)^2 L)^{-1})^M f(x, y) \, dx \, dy. \end{aligned}$$

Then by duality it readily follows that

$$\left(\frac{1}{\omega(\Omega)} \sum_R \iint_{I \times J} |(\mathbb{1} - (1 + \ell(I)^2 L \otimes \ell(J)^2 L)^{-1})^M f|^{r'} w^{1-r'} \, dx \right)^{1/r'} \leq \|l\|,$$

which is what we want to show. \square

Acknowledgment. The authors would like to thank the referee for carefully reading the manuscript and for her/his offering valuable suggestions to improve the paper. The authors also thank Lixin Yan and Liang Song for helpful suggestions and discussions.

References

- [1] *B. T. Anh, X. T. Duong*: Weighted Hardy spaces associated to operators and boundedness of singular integrals. Available at <https://arxiv.org/abs/1202.2063>.
- [2] *P. Auscher*: On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates. *Mem. Am. Math. Soc.* *186* (2007), 75 pages. [zbl](#) [MR](#) [doi](#)
- [3] *P. Auscher, X. T. Duong, A. McIntosh*: Boundedness of Banach space valued singular integral operators and Hardy spaces. Unpublished preprint (2005).
- [4] *P. Auscher, J. M. Martell*: Weighted norm inequalities, off-diagonal estimates and elliptic operators I: General operator theory and weights. *Adv. Math.* *212* (2007), 225–276. [zbl](#) [MR](#) [doi](#)
- [5] *P. Auscher, J. M. Martell*: Weighted norm inequalities, off-diagonal estimates and elliptic operators II: Off-diagonal estimates on spaces of homogeneous type. *J. Evol. Equ.* *7* (2007), 265–316. [zbl](#) [MR](#) [doi](#)
- [6] *P. Auscher, J. M. Martell*: Weighted norm inequalities, off-diagonal estimates and elliptic operators III: Harmonic analysis of elliptic operators. *J. Funct. Anal.* *241* (2006), 703–746. [zbl](#) [MR](#) [doi](#)
- [7] *P. Auscher, A. McIntosh, E. Russ*: Hardy spaces of differential forms on Riemannian manifolds. *J. Geom. Anal.* *18* (2008), 192–248. [zbl](#) [MR](#) [doi](#)
- [8] *M. Bownik, B. Li, D. Yang, Y. Zhou*: Weighted anisotropic product Hardy spaces and boundedness of sublinear operators. *Math. Nachr.* *283* (2010), 392–442. [zbl](#) [MR](#) [doi](#)
- [9] *H.-Q. Bui, X. T. Duong, L. Yan*: Calderón reproducing formulas and new Besov spaces associated with operators. *Adv. Math.* *229* (2012), 2449–2502. [zbl](#) [MR](#) [doi](#)
- [10] *L. Carleson*: A counterexample for measures bounded on H^p for the bi-disc. Mittag Leffler Report 7 (1974).
- [11] *S.-Y. A. Chang, R. Fefferman*: A continuous version of duality H^1 with BMO on the bidisc. *Ann. Math. (2)* *112* (1980), 179–201. [zbl](#) [MR](#) [doi](#)
- [12] *S.-Y. A. Chang, R. Fefferman*: The Calderón-Zygmund decomposition on product domains. *Am. J. Math.* *104* (1982), 445–468. [zbl](#) [MR](#) [doi](#)
- [13] *S.-Y. A. Chang, R. Fefferman*: Some recent developments in Fourier analysis and H^p -theory on product domains. *Bull. Am. Math. Soc., New Ser.* *12* (1985), 1–43. [zbl](#) [MR](#) [doi](#)
- [14] *E. B. Davies*: Heat Kernels and Spectral Theory. Cambridge Tracts in Mathematics 92, Cambridge University Press, Cambridge, 1989. [zbl](#) [MR](#) [doi](#)
- [15] *D. Deng, L. Song, C. Tan, L. Yan*: Duality of Hardy and BMO spaces associated with operators with heat kernel bounds on product domains. *J. Geom. Anal.* *17* (2007), 455–483. [zbl](#) [MR](#) [doi](#)
- [16] *X. T. Duong, J. Li*: Hardy spaces associated to operators satisfying Davies-Gaffney estimates and bounded holomorphic functional calculus. *J. Funct. Anal.* *264* (2013), 1409–1437. [zbl](#) [MR](#) [doi](#)
- [17] *X. T. Duong, A. MacIntosh*: Singular integral operators with non-smooth kernels on irregular domains. *Rev. Mat. Iberoam.* *15* (1999), 233–265. [zbl](#) [MR](#) [doi](#)
- [18] *X. T. Duong, E. M. Ouhabaz, A. Sikora*: Plancherel-type estimates and sharp spectral multipliers. *J. Funct. Anal.* *196* (2002), 443–485. [zbl](#) [MR](#) [doi](#)
- [19] *X. T. Duong, J. Xiao, L. Yan*: Old and new Morrey spaces with heat kernel bounds. *J. Fourier Anal. Appl.* *13* (2007), 87–111. [zbl](#) [MR](#) [doi](#)
- [20] *X. T. Duong, L. Yan*: Duality of Hardy and BMO spaces associated with operators with heat kernel bounds. *J. Am. Math. Soc.* *18* (2005), 943–973. [zbl](#) [MR](#) [doi](#)
- [21] *R. Fefferman*: A_p weights and singular integrals. *Am. J. Math.* *110* (1988), 975–987. [zbl](#) [MR](#) [doi](#)
- [22] *J. García-Cuerva, J. Rubio de Francia*: Weighted Norm Inequalities and Related Topics. North-Holland Mathematics Studies 116, North-Holland, Amsterdam, 1985. [zbl](#) [MR](#) [doi](#)

- [23] *R. F. Gundy, E. M. Stein*: H^p theory for the poly-disc. Proc. Natl. Acad. Sci. USA 76 (1979), 1026–1029. [zbl](#) [MR](#) [doi](#)
- [24] *S. Hofmann, G. Lu, D. Mitrea, M. Mitrea, L. Yan*: Hardy spaces associated to non-negative self-adjoint operators satisfying Davies-Gaffney estimates. Mem. Am. Math. Soc. 214 (2011), 78 pages. [zbl](#) [MR](#) [doi](#)
- [25] *S. Hofmann, S. Mayboroda*: Hardy and BMO spaces associated to divergence form elliptic operators. Math. Ann. 344 (2009), 37–116. [zbl](#) [MR](#) [doi](#)
- [26] *R. Jiang, D. Yang*: New Orlicz-Hardy spaces associated with divergence form elliptic operators. J. Funct. Anal. 258 (2010), 1167–1224. [zbl](#) [MR](#) [doi](#)
- [27] *J.-L. Journé*: Calderón-Zygmund operators on product spaces. Rev. Mat. Iberoam. 1 (1985), 55–91. [zbl](#) [MR](#) [doi](#)
- [28] *G. Kerkycharian, P. Petrushev*: Heat kernel based decomposition of spaces of distributions in the framework of Dirichlet spaces. Trans. Am. Math. Soc. 367 (2015), 121–189. [zbl](#) [MR](#) [doi](#)
- [29] *D. Krug*: A weighted version of the atomic decomposition for H^p (bi-halfspace). Indian Univ. Math. J. 37 (1988), 277–300. [zbl](#) [MR](#) [doi](#)
- [30] *D. Krug, A. Torchinsky*: A weighted version of Journé’s lemma. Rev. Math. Iberoam. 10 (1994), 363–378. [zbl](#) [MR](#) [doi](#)
- [31] *S. Liu, L. Song*: An atomic decomposition of weighted Hardy spaces associated to self-adjoint operators. J. Funct. Anal. 265 (2013), 2709–2723. [zbl](#) [MR](#) [doi](#)
- [32] *S. Liu, L. Song*: The atomic decomposition of weighted Hardy spaces associated to self-adjoint operators on product spaces. J. Math. Anal. Appl. 443 (2016), 92–115. [zbl](#) [MR](#) [doi](#)
- [33] *J. M. Martell, C. Prisuelos-Arribas*: Weighted Hardy spaces associated with elliptic operators I: Weighted norm inequalities for conical square functions. Trans. Am. Math. Soc. 369 (2017), 4193–4233. [zbl](#) [MR](#) [doi](#)
- [34] *J. M. Martell, C. Prisuelos-Arribas*: Weighted Hardy spaces associated with elliptic operators. Part II: Characterizations of $H_L^1(w)$. Available at <https://arxiv.org/abs/1701.00920>.
- [35] *E. M. Ouhabaz*: Analysis of Heat Equations on Domains. London Mathematical Society Monographs Series 31, Princeton University Press, Princeton, 2005. [zbl](#) [MR](#) [doi](#)
- [36] *S. Sato*: Weighted inequalities on product domains. Stud. Math. 92 (1989), 59–72. [zbl](#) [MR](#)
- [37] *L. Song, C. Tan, L. Yan*: An atomic decomposition for Hardy spaces associated to Schrödinger operators. J. Aust. Math. Soc. 91 (2011), 125–144. [zbl](#) [MR](#) [doi](#)
- [38] *L. Song, J. Xiao, L. Yan*: Preduals of quadratic Campanato spaces associated to operators with heat kernel bounds. Potential Anal. 41 (2014), 849–867. [zbl](#) [MR](#) [doi](#)
- [39] *L. Song, L. Yan*: Riesz transforms associated to Schrödinger operators on weighted Hardy spaces. J. Funct. Anal. 259 (2010), 1466–1490. [zbl](#) [MR](#) [doi](#)
- [40] *E. M. Stein*: Harmonic Analysis: Real Variable Methods, Orthogonality and Oscillatory Integrals. Princeton Mathematical Series 43, Princeton University Press, Princeton, 1993. [zbl](#) [MR](#)
- [41] *J.-O. Strömberg, A. Torchinsky*: Weighted Hardy Spaces. Lecture Notes in Mathematics 1381, Springer, Berlin, 1989. [zbl](#) [MR](#) [doi](#)
- [42] *X. X. Zhu*: Atomic decomposition for weighted H^p spaces on product domains. Sci. China, Ser. A 35 (1992), 158–168. [zbl](#) [MR](#)

Authors’ addresses: Suying Liu, School of Science, Northwestern Polytechnical University, 127 Youyi W Rd, Xian, 710000, Shaanxi, P.R. China, e-mail: liusuying0319@126.com; Minghua Yang, School of Information Technology, Jiangxi University of Finance and Economics, Jupu Rd, Nanchang, 330032, Jiangxi, P.R. China, e-mail: yhmh20062007@163.com.