

Enrico Jabara

Every 2-group with all subgroups normal-by-finite is locally finite

*Czechoslovak Mathematical Journal*, Vol. 68 (2018), No. 2, 491–496

Persistent URL: <http://dml.cz/dmlcz/147231>

## Terms of use:

© Institute of Mathematics AS CR, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

EVERY 2-GROUP WITH ALL SUBGROUPS NORMAL-BY-FINITE  
IS LOCALLY FINITE

ENRICO JABARA, Venezia

Received September 30, 2016. First published February 14, 2018.

*Abstract.* A group  $G$  has all of its subgroups normal-by-finite if  $H/H_G$  is finite for all subgroups  $H$  of  $G$ . The Tarski-groups provide examples of  $p$ -groups ( $p$  a “large” prime) of nonlocally finite groups in which every subgroup is normal-by-finite. The aim of this paper is to prove that a 2-group with every subgroup normal-by-finite is locally finite. We also prove that if  $|H/H_G| \leq 2$  for every subgroup  $H$  of  $G$ , then  $G$  contains an Abelian subgroup of index at most 8.

*Keywords:* 2-group; locally finite group; normal-by-finite subgroup; core-finite group

*MSC 2010:* 20F50, 20F14, 20D15

## 1. INTRODUCTION

A group  $G$  is a **CF**-group (core-finite) if each of its subgroups is normal-by-finite, that is,  $H/H_G$  is finite for all  $H \leq G$ . The subgroup  $H_G$ , the core of  $H$  in  $G$ , defined by

$$H_G = \bigcap_{g \in G} H^g,$$

is the biggest normal subgroup of  $G$  contained in  $H$ .

If  $G$  is a **CF**-group, we denote by  $\sigma(G)$  the  $\sup_{H \leq G} |H : H_G|$  and we say that  $G$  is **BCF** (boundedly core-finite) if  $\sigma(G) < \infty$ .

The main Theorem in [1] states that every locally finite **CF**-group is Abelian-by-finite and **BCF**. In order to prove the previous result, the hypothesis that  $G$  is locally finite is essential, as indicated by the existence of so-called Tarski-groups, for instance the examples due to Rips and Ol’shanskii (see [5]) of infinite groups all of whose nontrivial subgroups have prime order.

The aim of this paper is to provide an elementary proof of the following results.

**Theorem A.** *Let  $G$  be a 2-group. If  $G$  is a **CF**-group, then  $G$  is locally finite.*

As a consequence of Theorem A and the main result of [1] we have that every **CF** 2-group is Abelian-by-finite and **BCF**.

In the particular case of 2-groups  $G$  with  $\sigma(G) \leq 2$  we can provide a proof that  $G$  is locally finite and hypercentral in a direct way (see Lemma 8). We also extend to all groups a result proved in [7] for finite groups.

**Theorem B.** *Let  $G$  be a group. If  $\sigma(G) \leq 2$ , then  $G$  contains an Abelian subgroup of index at most 8.*

## 2. PROOFS

A quasicyclic 2-group is a group of type  $C_{2^\infty}$ , that is, an Abelian group with generators  $x_1, x_2, \dots, x_n, \dots$  and defining relations  $x_1^2 = 1$  and  $x_{i+1}^2 = x_i$  for all  $i > 1$ . A group of type  $C_{2^\infty}$  is not finitely generated and has every proper subgroup cyclic of infinite index; in particular, if a **CF**-group  $G$  contains a quasicyclic 2-group  $C$ , then  $C \trianglelefteq G$ .

Let  $G$  be a group and let  $\Lambda$  be a well-ordered set; an indexed set  $(G_\lambda)_{\lambda \in \Lambda}$  of subgroups of  $G$  is an ascending series if whenever  $\lambda, \mu \in \Lambda$  and  $\lambda \leq \mu$ , then  $G_\lambda \leq G_\mu$ .

**Lemma 1.** *If a group  $G$  is the union of an ascending series  $(G_\lambda)_{\lambda \in \Lambda}$  of locally finite groups, then  $G$  is locally finite.*

*Proof.* Trivial (see the proof of Lemma 1.A.2 in [3]). □

**Lemma 2.** *Every periodic hyperabelian group is locally finite.*

*Proof.* It is clear that a periodic Abelian group is locally finite. Since extensions of locally finite groups by locally finite groups are locally finite (Lemma 1.A.2 of [3] or 14.3.1 of [6]), the conclusion follows by Lemma 1. □

The following result is the key to prove our Theorem A.

**Lemma 3.** *Let  $G$  be an infinite 2-group containing no subgroups of type  $C_{2^\infty}$ . Then the centralizer of every finite subgroup of  $G$  is infinite.*

*Proof.* This is Theorem C of [4]. □

**Proof of Theorem A.** Let  $G$  be an infinite 2-group which is **CF**. Since every quotient of a **CF**-group is trivially a **CF**-group, in order to prove that  $G$  is locally finite, by Lemma 2 it suffices to prove that  $G$  contains a nontrivial normal Abelian subgroup. We must distinguish between two cases.

(I)  $G$  contains a subgroup  $A$  of type  $C_{2^\infty}$ .

Since  $|A : A_G|$  is finite and  $A$  does not contain proper subgroups of finite index, we have  $A_G = A$  and hence  $A$  is a nontrivial normal Abelian subgroup of  $G$ .

(II)  $G$  contains no subgroups of type  $C_{2^\infty}$ .

By the Zorn's Lemma in  $G$  there is a maximal Abelian subgroup  $A$ . If  $A$  is finite, then by Lemma 3,  $C_G(A)$  is infinite. Let  $g \in C_G(A) \setminus A$ , then clearly  $\langle A, g \rangle$  is Abelian, against the hypothesis that  $A$  is maximal. Hence  $A$  is infinite and since  $|A : A_G| < \infty$ , we have that  $A_G$  is infinite. So  $A_G$  is a nontrivial normal Abelian subgroup of  $G$ .

This proves Theorem A. □

**Remark 4.** Let  $G$  be a finite 2-group.

- (a) If  $\sigma(G) = 1$ , then  $G$  is Abelian or Hamiltonian (for the structure of Hamiltonian groups see 5.3.7 of [6]), in particular,  $G$  contains an Abelian normal subgroup of index 2.
- (b) In [7] it is proved that if  $\sigma(G) \leq 2$ , then  $G$  contains an Abelian normal subgroup of index at most 4 and this bound is sharp.
- (c) In [2] it is proved that if  $\sigma(G) \leq 2^s$ , then  $G$  contains an Abelian subgroup of index at most  $2^t$  with

$$t \leq \frac{1}{2}(s+1)(2s^3 + 7s^2 + 9s + 2)(s^3 + 2s^2 + 3s + 3).$$

Using an inverse limit argument (see Proposition 1.K.2 of [3] and the Remark that follows) it is not difficult to check that all previous results are valid if  $G$  is locally finite, and hence, by our Theorem A, for all 2-groups.

By proving Theorem B we extend the Corollary of [7] to infinite groups. In the proof we freely make use of the fact that if  $G$  is a **BCF**-group and  $H \leq G$ , then

$$\sigma(H) \leq \sigma(G),$$

moreover if  $H \trianglelefteq G$ , then

$$\sigma(G/H) \leq \sigma(G).$$

**Lemma 5.** *Let  $G$  be a group with  $\sigma(G) \leq 2$  and let  $A$  be the subgroup of  $G$  generated by all infinite cyclic normal subgroups and by all cyclic subgroups of odd order. Then  $A$  is Abelian.*

*Proof.* Finite odd-order subgroups of  $G$  are normal and also Dedekind groups, which implies that the elements of  $G$  of odd order generate an abelian normal subgroup of  $G$ .

Let  $x \in G$  with  $o(x) = \infty$  and  $\langle x \rangle \trianglelefteq G$ . The only nonidentity element  $\alpha$  of  $\text{Aut}(\langle x \rangle)$  inverts  $x$ ; if  $a \in G$  and  $x^a = x^{-1}$ , then  $\langle x \rangle \cap \langle a \rangle \leq C_{\langle x \rangle}(\alpha) = 1$ . Thus,  $x$  is centralized by every odd order element and by every infinite cyclic normal subgroup of  $G$ . This shows that  $A$  is abelian.  $\square$

**Lemma 6.** *Let  $G$  be a nonperiodic group with  $\sigma(G) \leq 2$  and let  $A$  be the subgroup generated by all normal infinite cyclic subgroups of  $G$  and by all cyclic subgroups of odd order of  $G$ . Then for each  $g \in G$  either  $a^g = a$  for all  $a \in A$  or  $a^g = a^{-1}$  for all  $a \in A$ . Therefore  $|G : C_G(A)| \leq 2$  and moreover  $g \in C_G(A)$  for every  $g \in G$  of infinite order.*

*Proof.* By Lemma 5,  $A$  is Abelian and hence if  $g \in G$  induces by conjugation a nontrivial automorphism on  $A$ , then  $g^2 \in A$ .

Let  $x \in G$  with  $o(x) = \infty$  and let  $u \in A$  of odd order. We have  $\langle x^2 \rangle \trianglelefteq G$ , hence  $[u, x^2] = 1$  and we can write  $(x^2u)^2 = x^4u^2$ , so  $o(x^2u) = \infty$  and  $\langle (x^2u)^2 \rangle \trianglelefteq G$ . Now

$$\langle u \rangle = \langle u^2 \rangle \leq \langle x^2, (x^2u)^2 \rangle.$$

Accordingly, if  $G$  is a nonperiodic group, then the subgroup  $A$  is the one generated by the normal infinite cyclic subgroups of  $G$ . Let  $\langle x \rangle \trianglelefteq G$  and  $\langle y \rangle \trianglelefteq G$  with  $o(x) = \infty = o(y)$ . If there is  $g \in G$  such that  $x^g = x^{-1}$  and  $y^g = y$ , then  $\langle x, y \rangle = \langle x \rangle \times \langle y \rangle$ , so  $o(xy) = \infty$ , but  $\langle (xy)^2 \rangle$  is not normal in  $G$ . This shows that  $a^g = a^{-1}$  for every  $a \in A$  and hence  $|G : C_G(A)| \leq 2$ .  $\square$

**Remark 7.** From Lemmas 5 and 6 it follows that every group  $G$  with  $\sigma(G) \leq 2$  and without elements of order 2 is Abelian.

**Lemma 8.** *Let  $G$  be a 2-group. If  $\sigma(G) \leq 2$ , then  $G$  is hypercentral.*

*Proof.* Suppose  $G \neq 1$ . If  $G$  has no elements of order 4, then  $G$  is elementary Abelian and the claim is proved. If there is an element  $g \in G$  with  $o(g) = 4$ , then  $\langle g^2 \rangle$  is a normal subgroup of order 2 of  $G$  and hence  $g^2 \in Z(G)$ , so  $Z(G) \neq 1$ .  $\square$

**Lemma 9.** *Let  $G$  be a hypercentral periodic group. If  $\sigma(G) \leq 2$ , then  $G$  contains an Abelian normal subgroup of index at most 4.*

**Proof.** Since  $G$  is hypercentral, it is locally finite and, by Lemma 5, we can write  $G = R_1 \times R_2$  with  $R_1$  an Abelian 2'-group and  $R_2$  a locally finite (hypercentral) group. By Remark 4(b),  $R_2$  contains a normal Abelian subgroup  $S$  of index at most 4. We can conclude that  $R_1 \times S$  is a normal Abelian subgroup of index at most 4 in  $G$ .  $\square$

**Proof of Theorem B.** Suppose that  $G$  is a periodic group and let  $A$  be the subgroup of  $G$  generated by all elements of odd order. By Lemma 5,  $A$  is Abelian, and hence locally finite. Since  $G/A$  is a 2-group with  $\sigma(G/A) \leq 2$ , it is locally finite and hence we can deduce that  $G$  is locally finite. By Corollary of [7], every finite group  $X$  with  $\sigma(X) \leq 2$  contains an Abelian subgroup of index at most 8, applying Proposition 1.K.2 of [3] we obtain that also  $G$  has an Abelian subgroup of index at most 8. So in this case the claim is proved.

From now we suppose that  $G$  contains elements of infinite order.

Let  $A$  be the subgroup of  $G$  generated by all normal infinite cyclic subgroups of  $G$  and by all cyclic subgroups of odd order of  $G$  and let  $C = C_G(A)$  (by the proof of Lemma 6,  $A$  is actually generated by the normal infinite cyclic subgroups of  $G$ ). By construction we have that  $A$  is normal and  $G/A$  is a 2-group, by Lemma 5 we have that  $A$  is Abelian and by Lemma 6 that  $|G : C| \leq 2$ .

First of all we prove that if  $G = C$ , then  $G$  contains a normal Abelian subgroup of index at most 4. In this case  $G$  is hypercentral because  $G/A$  is a 2-group, which is hypercentral by Lemma 8, and  $A \leq Z(G)$  by hypothesis.

Let  $a \in A$  be such that  $\langle a \rangle \trianglelefteq G$  and  $o(a) = \infty$ .

Let  $h \in G$ , then  $[h, a] = 1$ , moreover if  $o(h) < \infty$ , then  $o(ah) = \infty$  and  $a^2h^2 = (ah)^2 \in A$ , so  $h^2 \in A$ . Hence for every  $g \in G$  we have  $g^2 \in A$  and  $G/A$  is an elementary Abelian 2-group. Since  $A \leq Z(G)$  for  $g, h \in G$ , we can write

$$[g, h]^2 = [g^2, h] = 1$$

and this shows  $G'$  to be an elementary Abelian 2-subgroup of  $A$ . By the Zorn's Lemma in  $A$  there is a free Abelian subgroup of maximal rank  $Q$ , moreover we can suppose  $Q \trianglelefteq G$  because since  $|Q : Q_G| \leq 2$ , we can consider  $Q_G$  instead of  $Q$ . Then  $A/Q$  is an Abelian periodic group. Let  $T$  be the full preimage in  $G$  of the 2'-Hall subgroup of  $A/Q$ , then  $T$  does not have elements of finite even order and  $G/T$  is a 2-group. Moreover,  $G' \cap T = 1$  and hence the full preimage of any Abelian subgroup of  $G/T$  is an Abelian subgroup of  $G$ . Since  $G/T$  is periodic and hypercentral, then, by Lemma 9, it contains a normal Abelian subgroup  $S/T$  of index at most 4. The full preimage  $S$  of  $S/T$  in  $G$  is a normal Abelian subgroup of index at most 4 of  $G$ .

Suppose now  $|G : C| = 2$ . For the previous argument we know that in  $C$  there is an Abelian (normal) subgroup  $S$  of index at most 4 and hence  $|G : S| \leq 8$ .  $\square$

**Remark 10.** The bound found in Theorem B is best possible, as has been shown in the proof of Corollary of [7].

**Example (i).** Let  $G = \langle x_i, y; x_i^y = x_i^{-1} \rangle$  be the semidirect product of the quasicyclic 2-group  $\langle x_1, x_2, \dots, x_n, \dots \rangle$  by the infinite cyclic group  $\langle y \rangle$ , then  $G$  is a nonperiodic **CF**-group. The element  $x_n y^2 \in G$  has infinite order and it is easy to prove that  $|\langle x_n y^2 \rangle : \langle x_n y^2 \rangle_G| = 2^{n-1}$ , so  $G$  is not a **BCF**-group. This shows that the main theorem of [1] cannot be extended to nonperiodic groups.

**Remark 11.** Let  $G$  be a group and suppose that for every  $g \in G$  we have  $|\langle g \rangle : \langle g \rangle_G| \leq 2$ . Then the subgroup  $N = \langle g^2; g \in G \rangle$  is Abelian or Hamiltonian and  $G/N$  has exponent 2, so it is elementary Abelian. Hence  $G' \leq N$  and  $|G''| \leq 2$  (in particular if  $G$  is periodic, then it is locally finite), but it can happen that  $\sigma(G) = \infty$ , as shown in Example (ii).

**Example (ii).** Let

$$G = \bigoplus_{i \in \mathbb{N}} D_i$$

with  $D_i = \langle x_i, y_i; x_i^A, y_i^2, x_i^{y_i} x_i \rangle$  isomorphic to the dihedral group of order 8. It is clear that if  $g \in G$ , then  $|\langle g \rangle : \langle g \rangle_G| \leq 2$ , but  $G$  does not possess Abelian subgroups of finite index. Moreover, if  $n \in \mathbb{N}$ , the elementary Abelian subgroup  $H = \langle y_1, y_2, \dots, y_n \rangle$  of  $G$  is such that  $|H| = 2^n$  and  $H_G = 1$ , in particular  $\sigma(G) = \infty$ .

**Acknowledgments.** The author would like to thank the referee for many helpful suggestions.

### References

- [1] J. T. Buckley, J. C. Lennox, B. H. Neumann, H. Smith, J. Wiegold: Groups with all subgroups normal-by-finite. *J. Aust. Math. Soc., Ser. A* 59 (1995), 384–398. [zbl](#) [MR](#)
- [2] G. Cutolo, E. I. Khukhro, J. C. Lennox, S. Rinauro, H. Smith, J. Wiegold: Locally finite groups all of whose subgroups are boundedly finite over their cores. *Bull. Lond. Math. Soc.* 29 (1997), 563–570. [zbl](#) [MR](#) [doi](#)
- [3] O. H. Kegel, B. A. F. Wehrfritz: *Locally Finite Groups*. North-Holland Mathematical Library 3, North-Holland Publishing, Amsterdam, 1973. [zbl](#) [MR](#)
- [4] J. C. Lennox, A. Mohammadi Hassanabadi, A. G. R. Stewart, J. Wiegold: Nilpotent extensibility and centralizers in infinite 2-groups. *Proceedings of the Second International Group Theory Conference (Bressanone, 1989)*. *Rend. Circ. Mat. Palermo (2) Suppl.* No. 23, 1990, pp. 209–219. [zbl](#) [MR](#)
- [5] A. Yu. Ol’shanskii: *Geometry of Defining Relations in Groups*. Mathematics and Its Applications. Soviet Series 70, Kluwer Academic Publishers, Dordrecht, 1991. [zbl](#) [MR](#) [doi](#)
- [6] D. J. S. Robinson: *A Course in the Theory of Groups*. Graduate Texts in Mathematics 80, Springer, New York, 1996. [zbl](#) [MR](#) [doi](#)
- [7] B. Wilkens: More on core-2 2-groups. *J. Group Theory* 20 (2017), 193–225. [zbl](#) [MR](#) [doi](#)

*Author’s address:* Enrico Jabara, DFBC—Università di Venezia, Dorsoduro 3484/D—30123 Venezia, Italy, e-mail: [jabara@unive.it](mailto:jabara@unive.it).