

Xiao-Hui Hua; Li Chen

Valency seven symmetric graphs of order $2pq$

Czechoslovak Mathematical Journal, Vol. 68 (2018), No. 3, 581–599

Persistent URL: <http://dml.cz/dmlcz/147353>

Terms of use:

© Institute of Mathematics AS CR, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

VALENCY SEVEN SYMMETRIC GRAPHS OF ORDER $2pq$

XIAO-HUI HUA, LI CHEN, Xinxiang

Received October 6, 2015. Published online June 5, 2018.

Abstract. A graph is said to be symmetric if its automorphism group acts transitively on its arcs. In this paper, all connected valency seven symmetric graphs of order $2pq$ are classified, where p, q are distinct primes. It follows from the classification that there is a unique connected valency seven symmetric graph of order $4p$, and that for odd primes p and q , there is an infinite family of connected valency seven one-regular graphs of order $2pq$ with solvable automorphism groups, and there are four sporadic ones with nonsolvable automorphism groups, which is 1, 2, 3-arc transitive, respectively. In particular, one of the four sporadic ones is primitive, and the other two of the four sporadic ones are bi-primitive.

Keywords: arc-transitive graph; symmetric graph; s -regular graph

MSC 2010: 05C25, 20B25

1. INTRODUCTION

For a finite, simple and undirected graph X , let $V(X)$, $E(X)$, $A(X)$ and $\text{Aut}(X)$ denote the vertex set, edge set, arc set and full automorphism group of X , respectively. Note that an *arc* is an ordered edge, that is, an ordered pair of adjacent vertices. For $u, v \in V(X)$, $\{u, v\}$ denotes the edge incident to u and v in X . An s -arc in a graph X for some nonnegative integer s is an ordered $(s + 1)$ -tuple (v_0, v_1, \dots, v_s) of $s + 1$ vertices such that $(v_{i-1}, v_i) \in A(X)$ for $1 \leq i \leq s$ and $v_{i-1} \neq v_{i+1}$ for $1 \leq i \leq s - 1$. For a subgroup G of the automorphism group $\text{Aut}(X)$ of a graph X , the graph X is said to be (G, s) -arc-transitive or (G, s) -regular if G acts transitively or regularly on the set of s -arcs of X , and (G, s) -transitive if G acts transitively on the set of s -arcs but not on the set of $(s + 1)$ -arcs of X . A graph X is said to be s -arc-transitive, s -regular or s -transitive if it is $(\text{Aut}(X), s)$ -arc-transitive,

This work was supported by the National Natural Science Foundation of China (11301159, 11671030, 11601132, 11501176, 11526082), the Education Department of Henan Science and Technology Research Key Project (13A110543).

$(\text{Aut}(X), s)$ -regular or $(\text{Aut}(X), s)$ -transitive. In particular, 0-arc-transitive means *vertex-transitive*, and 1-arc-transitive means *arc-transitive* or *symmetric*. A graph is said to be *primitive* if its automorphism group is primitive on the vertex set, and a graph is said to be *bi-primitive* if it is a bipartite graph with bi-parts Δ_1, Δ_2 , and the setwise stabilizer of its automorphism group is primitive on both Δ_1 and Δ_2 . Throughout this paper, we will denote by \mathbb{Z}_n the cyclic group of order n , by \mathbb{Z}_n^* the multiplicative group of units modulo n , by D_{2n} the dihedral group of order $2n$, by F_n the Frobenius group of order n , and by A_n and S_n the alternating group and the symmetric group of degree n , respectively.

It is well known that a graph Γ is G -arc-transitive if and only if G is vertex-transitive and the vertex stabilizer G_v of $v \in V(\Gamma)$ in G is transitive on $N_\Gamma(v)$. Hence the structure of the vertex stabilizer of G_v plays an important role in the study of (G, s) -transitive graphs. For example, benefitted from Djoković and Miller [4] result about the vertex stabilizer of cubic symmetric graphs, lots of works about classifications of cubic symmetric graphs were obtained by many authors (see [7], [8], [9], [23], [24]). Due to the vertex stabilizers given in [27], symmetric tetravalent graphs have also been studied extensively in the literature (see [11], [12], [22], [32], [34]). Similarly, Guo and Feng [14] determined structure of vertex stabilizers of pentavalent symmetric graphs, some works about classifications of pentavalent symmetric graphs were also obtained (see [6], [14], [17], [18], [26]). Naturally, the next step is to characterize valency seven symmetric graphs. Recently, Guo et al. [15] gave the structure of vertex stabilizers of valency seven symmetric graphs, and this encourages us to consider some work on valency seven symmetric graphs. In [16], Guo et al. classified valency seven symmetric graphs of order $4p$, and in [25], Pan et al. classified prime-valent symmetric graphs of square-free order. But, we obtain this result for valency seven symmetric graphs of order $2pq$ independently. Let p, q be two distinct primes. In this paper, we classify valency seven symmetric graphs of order $2pq$.

2. PRELIMINARIES

Let X be a graph, and N a subgroup of $\text{Aut}(X)$. Denote by X_N the quotient graph corresponding to the orbits of N , that is the graph having the orbits of N as vertices with two orbits adjacent in X_N if there is an edge in X between those orbits. In view of [20], Theorem 9, we have the following proposition.

Proposition 2.1. *Let X be a connected symmetric graph of prime valency p and G an s -arc-transitive subgroup of $\text{Aut}(X)$ for some $s \geq 1$. If a normal subgroup N of G has more than two orbits on $V(X)$ then X_N is also a symmetric graph*

of valency p and N is the kernel of the action of G on the set of orbits of N . Moreover, N is semiregular on $V(X)$ and G/N is an s -arc-transitive subgroup of $\text{Aut}(X_N)$.

By Guo [15], we have the following statement.

Proposition 2.2. *Let X be a connected (G, s) -transitive graph of valency seven for some $G \leq \text{Aut}(X)$ and $s \geq 1$. Let $v \in V(X)$. Then $s \leq 3$ and one of the following holds:*

- (1) *If G_v is soluble, then $|G_v| \mid 2^2 \cdot 3^2 \cdot 7$. Further, the triple $(s, G_v, |G_v|)$ lies in the following table:*

$s = 1$		$s = 2$		$s = 3$	
G_v	Order	G_v	Order	G_v	Order
\mathbb{Z}_7	7	F_{42}	$2 \cdot 3 \cdot 7$	$F_{42} \times \mathbb{Z}_6$	$2^2 \cdot 3^2 \cdot 7$
D_{14}	$2 \cdot 7$	$F_{42} \times \mathbb{Z}_2$	$2^2 \cdot 3 \cdot 7$		
F_{21}	$3 \cdot 7$	$F_{42} \times \mathbb{Z}_3$	$2 \cdot 3^2 \cdot 7$		
D_{28}	$2^2 \cdot 7$				
$F_{21} \times \mathbb{Z}_3$	$3^2 \cdot 7$				

- (2) *If G_v is insoluble, then $s \geq 2$ and $|G_v| \mid 2^{24} \cdot 3^4 \cdot 5^2 \cdot 7$. Further, the triple $(s, G_v, |G_v|)$ lies in the following table:*

$s = 2$		$s = 3$	
G_v	Order	G_v	Order
$\text{PSL}(3, 2)$	$2^3 \cdot 3 \cdot 7$	$\text{PSL}(3, 2) \times S_4$	$2^6 \cdot 3^2 \cdot 7$
A_7	$2^3 \cdot 3^2 \cdot 5 \cdot 7$	$A_7 \times A_6$	$2^6 \cdot 3^4 \cdot 5^2 \cdot 7$
S_7	$2^4 \cdot 3^2 \cdot 5 \cdot 7$	$S_7 \times S_6$	$2^8 \cdot 3^4 \cdot 5^2 \cdot 7$
$\mathbb{Z}_2^3 \times \text{SL}(3, 2)$	$2^6 \cdot 3 \cdot 7$	$(A_7 \times A_6) \rtimes \mathbb{Z}_2$	$2^7 \cdot 3^4 \cdot 5^2 \cdot 7$
$\mathbb{Z}_2^4 \times \text{SL}(3, 2)$	$2^7 \cdot 3 \cdot 7$	$\mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$	$2^{10} \cdot 3^2 \cdot 7$
		$([2^{20}] \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2)))$	$2^{24} \cdot 3^2 \cdot 7$

From [3], pages 12–14, 3-prime factor simple groups can be found. And by [13], pages 134–136, one can obtain the following proposition by checking the orders of nonabelian simple groups.

Proposition 2.3. *Let p, q be distinct odd primes, and let G be a nonabelian simple group of order $|G| = 2^i \cdot 3^j \cdot 5^k \cdot 7 \cdot p \cdot q$ with $1 \leq i \leq 26$, $0 \leq j \leq 4$, $0 \leq k \leq 2$ and $7 \mid |G|$. Then G has 3-prime factor, 4-prime factor, 5-prime factor or 6-prime factor, and is one of the groups in Table 1.*

G	Order	G	Order
PSL(2, 7)	$2^3 \cdot 3 \cdot 7$	A_{11}	$2^7 \cdot 3^4 \cdot 5^2 \cdot 7 \cdot 11$
PSL(2, 8)	$2^3 \cdot 3^2 \cdot 7$	A_{12}	$2^9 \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$
PSU(3, 3)	$2^5 \cdot 3^3 \cdot 7$	M_{22}	$2^7 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$
A_7	$2^3 \cdot 3^2 \cdot 5 \cdot 7$	HS	$2^9 \cdot 3^2 \cdot 5^3 \cdot 7 \cdot 11$
A_8	$2^6 \cdot 3^2 \cdot 5 \cdot 7$	PSL(2, 2^6)	$2^6 \cdot 3^2 \cdot 5 \cdot 7 \cdot 13$
A_9	$2^6 \cdot 3^4 \cdot 5 \cdot 7$	PSL(2, 2^9)	$2^9 \cdot 3^3 \cdot 7 \cdot 19 \cdot 73$
A_{10}	$2^7 \cdot 3^4 \cdot 5^2 \cdot 7$	PSL(2, 5^3)	$2^2 \cdot 3^2 \cdot 5^3 \cdot 7 \cdot 31$
PSL(2, 13)	$2^2 \cdot 3 \cdot 7 \cdot 13$	PSL(4, 4)	$2^{12} \cdot 3^4 \cdot 5^2 \cdot 7 \cdot 17$
PSL(2, 27)	$2^2 \cdot 3^3 \cdot 7 \cdot 13$	PSL(5, 2)	$2^{10} \cdot 3^2 \cdot 5 \cdot 7 \cdot 31$
PSL(3, 4)	$2^6 \cdot 3^2 \cdot 5 \cdot 7$	PSp(4, 8)	$2^{12} \cdot 3^4 \cdot 5 \cdot 7^2 \cdot 13$
PSL(3, 8)	$2^9 \cdot 3^2 \cdot 7^2 \cdot 73$	${}^2D_4(2)$	$2^{12} \cdot 3^4 \cdot 5 \cdot 7 \cdot 17$
PSU(3, 5)	$2^4 \cdot 3^2 \cdot 5^3 \cdot 7$	$G_2(4)$	$2^{12} \cdot 3^3 \cdot 5^2 \cdot 7 \cdot 13$
PSU(3, 8)	$2^9 \cdot 3^4 \cdot 7 \cdot 19$	$G_2(8)$	$2^{18} \cdot 3^5 \cdot 7^2 \cdot 19 \cdot 73$
J_2	$2^7 \cdot 3^3 \cdot 5^2 \cdot 7$	M_{23}	$2^7 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 23$
$Sz(8)$	$2^6 \cdot 5 \cdot 7 \cdot 13$	M_{24}	$2^{10} \cdot 3^3 \cdot 5 \cdot 7 \cdot 11 \cdot 23$
$D_4(2)$	$2^{12} \cdot 3^5 \cdot 5^2 \cdot 7$	J_1	$2^3 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 19$
${}^3D_4(2)$	$2^{12} \cdot 3^4 \cdot 7^2 \cdot 13$	PSL(3, 16)	$2^{12} \cdot 3^2 \cdot 5^2 \cdot 7 \cdot 13 \cdot 17$
PSp(6, 2)	$2^9 \cdot 3^4 \cdot 5 \cdot 7$	PSL(2, t)	$t = \pm 1 \pmod{7}$ and $t > 13$
PSp(8, 2)	$2^{16} \cdot 3^5 \cdot 5^2 \cdot 7$		
PSL(2, 49)	$2^4 \cdot 3 \cdot 5^2 \cdot 7^2$		

Table 1. Nonabelian simple $\{2, 3, 5, 7, p, q\}$ -groups.

Proof. Clearly, we have

$$(2.1) \quad 2^{27} \nmid |G|, \quad 3^6 \nmid |G|, \quad 5^4 \nmid |G|, \quad 7^3 \nmid |G|, \quad 7 \mid |G|, \quad t^2 \nmid |G|$$

where $t \in \{q, p\}$ and $t \geq 11$.

From [3], pages 12–14, 3-prime factor simple groups can be found. If $7 \mid |G|$, one has $G \cong \text{PSL}(2, 7)$, $\text{PSL}(2, 8)$ or $\text{PSU}(3, 3)$. Specially, if $7^2 \mid |G|$, $3^5 \mid |G|$ or $5^3 \mid |G|$, then $|G|$ has at most five prime divisors. By [31], page 3, each finite nonabelian simple group is isomorphic to A_n with $n \geq 5$, one of 26 sporadic simple groups, or a classical group or an exceptional group of Lie type. For the orders of these simple groups, one can see [13], Table 2.4, pages 134–136, and for more details, see [31], Sections 3, 4, 5.

For A_n with $n \geq 5$, since $3^6 \nmid |G|$ and $7 \mid |G|$, we have $G \cong A_7, A_8, A_9, A_{10}, A_{11}$ or A_{12} . For the 26 sporadic simple groups, by equation (2.1) we have $G \cong M_{22}, M_{23}, M_{24}, J_1, J_2, HS$.

For the groups of Lie type, since each odd prime divisor of $|G|$ has power at most 5, by [13], Table 2.4, pages 134–136, $G \cong D_4(2), {}^2D_4(2), {}^3D_4(2)$, $\text{PSL}(n, t)$ with $n \geq 3$, $\text{PSU}(n, t)$ with $n \geq 3$, $\text{PSp}(2n, t)$ with $n \geq 2$, or $\text{Sz}(2^{2n+1})$ with $n \geq 1$, where t is a prime power.

Let $G \cong \text{PSL}(n, t)$. Then $|G| = (n, t - 1)^{-1} t^{n(n-1)/2} \prod_{i=2}^n (t^i - 1)$. First assume $n \geq 3$. Then $n(n-1)/2 \geq 3$, and by equation (2.1), we have $n = 3$ and $t = 3, 5$ or 2^i with $i < 9$, $n = 4$ and $t = 2^i$ with $i < 5$, or $n = 5$ and $t = 2^i$ with $i < 3$. For each case, by checking orders with equation (2.1) again, we have $G \cong \text{PSL}(3, 4)$, $\text{PSL}(3, 8)$, $\text{PSL}(3, 16)$, $\text{PSL}(4, 2) (\cong A_8)$, $\text{PSL}(4, 4)$ or $\text{PSL}(5, 2)$. Now assume $n = 2$. Then $|G| = (2, t - 1)^{-1} t(t^2 - 1)$. If $t = 2^i$ then $i \leq 26$ by equation (2.1). Similarly, if $t = 3^i$ then $i \leq 5$; if $t = 5^i$ then $i \leq 3$; if $t = 7^i$ then $i \leq 2$; if $t = s^i$ with $s > 7$ and $s \in \{q, p\}$ then $i = 1$. For each case, checking the orders of $\text{PSL}(2, t)$ again, we have $G \cong \text{PSL}(2, 2^6)$, $\text{PSL}(2, 2^9)$, $\text{PSL}(2, 27)$, $\text{PSL}(2, 125)$, $\text{PSL}(2, 49)$, or $\text{PSL}(2, t)$ with some prime $t \geq 13$ and $t \in \{q, p\}$.

Let $G \cong \text{PSU}(n, t)$ with $n \geq 3$. Then

$$|G| = (n, t + 1)^{-1} t^{n(n-1)/2} \prod_{i=2}^n (t^i - (-1)^i).$$

Since $n(n-1)/2 \geq 3$, we have $n = 3$ and $t = 3, 5$ or $t = 2^i$ with $i < 9$, $n = 4$ and $t = 2^i$ with $i < 5$, or $n = 5$ and $t = 2^i$ with $i < 3$ by equation (2.1). Hence, $G \cong \text{PSU}(3, 8)$, $\text{PSU}(3, 5)$. For the other two infinite families $\text{PSp}(2n, t)$ of order $(n, t - 1)^{-1} t^{n^2} \prod_{i=2}^n (t^{2i} - 1)$ with $n \geq 2$ and $\text{Sz}(2^{2n+1})$ of order $2^{4n+2}(2^{4n+2} + 1) \times (2^{2n+1} - 1)$ with $n \geq 1$, one can similarly obtain that $G \cong \text{PSp}(4, 8), \text{Sz}(8)$. \square

From [30], page 417, we have the following proposition.

Proposition 2.4. *Let p be a prime, and $q = p^n \geq 5$. Then a maximal subgroup of $\text{PSL}(2, q)$ is isomorphic to one of the following groups:*

- (1) $D_{2(q-1)/d}$, where $d = (2, q - 1)$ and $q \neq 5, 7, 9, 11$;
- (2) $D_{2(q+1)/d}$, where $d = (2, q - 1)$ and $q \neq 7, 9$;
- (3) $\mathbb{Z}_q \rtimes \mathbb{Z}_{(q-1)/d}$;
- (4) A_4 , when $q = p = 5$, or $q = p \equiv 3, 13, 27, 37 \pmod{40}$;
- (5) S_4 , when $q = p \equiv \pm 1 \pmod{8}$;
- (6) A_5 , when $q = p \equiv \pm 1 \pmod{5}$, or $q = p^2 \equiv -1 \pmod{5}$ with p an odd prime;
- (7) $\text{PSL}(2, r)$, when $q = r^m$ with m an odd prime;
- (8) $\text{PGL}(2, r)$, when $q = r^2$.

To extract a classification of connected valency seven symmetric graphs of order $2p$ for a prime p from Cheng and Oxley [2], we introduce the graphs $G(2p, r)$. Let V and V' be two disjoint copies of \mathbb{Z}_p , say $V = \{0, 1, \dots, p-1\}$ and $V' = \{0', 1', \dots, (p-1)'\}$. Let r be a positive integer dividing $p-1$ and $H(p, r)$ the unique subgroup of Z_p^* of order r . Define the graph $G(2p, r)$ to have vertex set $V \cup V'$ and edge set $\{xy' : x - y \in H(p, r)\}$.

Proposition 2.5. *Let p be a prime, and let X be a connected valency seven symmetric graph of order $2p$. Then one of the following situations occurs:*

- (1) $X \cong K_{7,7}$, the complete bipartite graph of order 14, and $\text{Aut}(K_{7,7}) = (S_7 \times S_7) \rtimes \mathbb{Z}_2$;
- (2) $X \cong G(2p, 7)$ with $p \equiv 1 \pmod{7}$, and $\text{Aut}(G(2p, 7)) = (\mathbb{Z}_p \rtimes \mathbb{Z}_7) \rtimes \mathbb{Z}_2$.

Finally, we introduce the so called Cayley graph. For a finite group G and a subset S of G such that $S = S^{-1}$ and $1 \notin S$, the *Cayley graph* $\text{Cay}(G, S)$ on G with respect to S is defined to have vertex set G and edge set $\{\{g, sg\} : g \in G, s \in S\}$. Given $g \in G$, right multiplication $x \mapsto xg$ (for $x \in G$) is a permutation $R(g)$ on G , and the homomorphism from G to $\text{Sym}(G)$ taking each g to $R(g)$ is called the *right regular representation* of G . The image $R(G) = \{R(g) : g \in G\}$ of G is a regular permutation group on G , and is isomorphic to G , which can therefore be regarded as a subgroup of the automorphism group $\text{Aut}(\text{Cay}(G, S))$. In particular, the Cayley graph $\text{Cay}(G, S)$ is vertex-transitive. Moreover, the group $\text{Aut}(G, S) = \{\alpha \in \text{Aut}(G) : S^\alpha = S\}$ is a subgroup of $\text{Aut}(\text{Cay}(G, S))$, indeed of the stabilizer $\text{Aut}(\text{Cay}(G, S))_1$ of the vertex 1. Also, a Cayley graph $\text{Cay}(G, S)$ is said to be *normal* if $R(G)$ is normal in $\text{Aut}(\text{Cay}(G, S))$. By [33], Propositions 1.3 and 1.5, a Cayley graph $\text{Cay}(G, S)$ is normal if and only if $\text{Aut}(\text{Cay}(G, S))_1 = \text{Aut}(G, S)$, or equivalently, if and only if $\text{Aut}(\text{Cay}(G, S))$ is isomorphic to the semidirect product $R(G) \rtimes \text{Aut}(G, S)$.

Now we introduce an infinite family of one-regular Cayley graphs on the dihedral group $D_{2n} = \langle a, b \mid a^n = b^2 = 1, b^{-1}ab = a^{-1} \rangle$. Let m and l be integers such that $l^6 + l^5 + l^4 + l^3 + l^2 + l + 1 \equiv 0 \pmod{m}$. Define

$$(2.2) \quad \mathcal{CD}_{2m}^l = \text{Cay}(D_{2m}, S),$$

where $S = \{b, ab, a^{l+1}b, a^{l^2+l+1}b, a^{l^3+l^2+l+1}b, a^{l^4+l^3+l^2+l+1}b, a^{l^5+l^4+l^3+l^2+l+1}b\}$.

By [10], we have the following propositions.

Proposition 2.6 ([10], Theorem 3.5). *Let n be a square-free integer and X a connected valency seven one-regular graph of order n . Then $n = 2 \cdot 7^t \cdot p_1 p_2 \dots p_s$, where $t \leq 1$, $s \geq 1$, and p_i 's are distinct primes such that $7 \mid (p_i - 1)$. Furthermore, X is isomorphic to one of \mathcal{CD}_n^l and there are exactly 6^{s-1} nonisomorphic such graphs of order n .*

Now we introduce the so called coset graph (see [22], [28]) constructed from a finite group G relative to a subgroup H of G and a union D of some double cosets of H in G such that $D^{-1} = D$. Denote by H_G the largest normal subgroup of G in H . The coset graph $\text{Cos}(G, H, D)$ of G with respect to H and D is defined to have vertex set $[G : H]$, the set of right cosets of H in G , and edge set $\{\{Hg, Hdg\} : g \in G, d \in D\}$. The action of G on $V(\text{Cos}(G, H, D))$ by right multiplication induces a vertex-transitive automorphism group, which is faithful if and only if $H_G = 1$. Furthermore, $\text{Aut}(G, H, D) = \{\alpha \in \text{Aut}(G) : H^\alpha = H, D^\alpha = D\}$ induces a group of automorphisms, which lies in the stabilizer of H in $\text{Aut}(\text{Cos}(G, H, D))$. Clearly, $\text{Cos}(G, H, D) \cong \text{Cos}(G, H^\alpha, D^\alpha)$ for every $\alpha \in \text{Aut}(G)$. Note that the concept of a coset graph is equivalent to the concept of an orbital graph (see [29]). Conversely, by [28] we have the following statement.

Proposition 2.7. *Let X be a graph and let G be a vertex-transitive subgroup of $\text{Aut}(X)$. Then X is isomorphic to a coset graph $\text{Cos}(G, H, D)$, where $H = G_u$ is the stabilizer of $u \in V(X)$ in G and D consists of all elements of G which map u to one of its neighbors. Further,*

- (1) X is connected if and only if D generates the group G ;
- (2) X is G -arc-transitive if and only if D is a single double coset. In particular, if $g \in G$ interchanges u and one of its neighbors, then $g^2 \in H$ and $D = HgH$;
- (3) the valency of X is equal to $|D|/|H| = |H : H \cap H^g|$.

3. CONSTRUCTIONS

In this section, we construct valency seven symmetric graphs of order $2pq$, where p and q are distinct primes.

Example 3.1. Let G be a subgroup of S_{14} such that $G \cong \text{PSL}(2, 13)$, and G contains the following elements:

$$\begin{aligned} a &= (1, 12)(2, 6)(3, 13)(4, 7)(8, 9)(10, 11), \\ b &= (1, 12, 2, 10, 14, 11, 6)(3, 9, 5, 8, 13, 4, 7), \\ g_2 &= (1, 6)(2, 4)(3, 8)(5, 7)(9, 10)(13, 14), \\ g_2 &= (1, 8)(3, 5)(4, 12)(6, 7)(9, 10)(11, 13). \end{aligned}$$

By Magma [1], $G = \langle a, b, g_i \rangle$ for each $1 \leq i \leq 2$ and $H = \langle a, b \rangle$. Define the following coset graphs:

$$\mathcal{C}_{78}^i = \text{Cos}(G, H, Hg_iH), \quad 1 \leq i \leq 2.$$

Again by Magma [1], the two coset graphs \mathcal{C}_{78}^i ($i = 1, 2$) are pairwise nonisomorphic connected valency seven 1-transitive graphs of order 78 with $\text{Aut}(\mathcal{C}_{78}^1) = \text{PSL}(2, 13)$ and $\text{Aut}(\mathcal{C}_{78}^2) = \text{PGL}(2, 13)$.

Lemma 3.2. *Each connected valency seven symmetric graph X of order 78 admitting $\text{PSL}(2, 13)$ as an arc-transitive automorphism group is isomorphic to \mathcal{C}_{78}^i ($i = 1, 2$). Furthermore, X is 1-transitive and $\text{Aut}(X) \cong \text{PSL}(2, 13)$ or $\text{PGL}(2, 13)$.*

Proof. Let $G = \text{PSL}(2, 13)$. As X is a G -arc-transitive graph of order 78, one has $|G_v| = 14$ for any vertex $v \in V(X)$, and by Proposition 2.4, we have $H = G_v \cong D_{14}$. The simplicity of G and the maximality of H imply that $H = N_G(H)$. Take an involution x in H , and set $\langle x \rangle = L$. Since G has one conjugacy class of involutions, by Proposition 2.4, $N_G(L) = D_{12}$. Clearly, $H \cap H^g = L$ and $N_H(L) \cong L$. Thus, there exists an involution g such that $g \in N_G(L)$ and $g \notin N_H(L)$. Furthermore, $g \notin H$, $|HgH|/|H| = 7$ and $\langle H, g \rangle = G$. This implies that $\text{Cos}(G, H, HgH)$ is a connected valency seven symmetric graph of order 78.

Let X be a connected valency seven symmetric graph of order 78 admitting $G = \text{PSL}(2, 13)$ as an arc-transitive automorphism group. Note that $G_v \cong D_{14}$ for any $v \in V(X)$. To complete the proof, it suffices to show that $X \cong \text{Cos}(G, H, HgH)$. By Magma [1], G has one conjugacy class of D_{14} and since G_v has seven subgroups isomorphic to \mathbb{Z}_2 , each of the subgroups fixes a vertex adjacent to v . By Proposition 2.7, one may assume that $X = \text{Cos}(G, H, HfH)$ such that $H \cap H^f = L$ and $f \in N_G(L)$. By [5], Theorem 2.1, f can be chosen to be a 2-element, and hence f is an involution in $N_G(L) \cong D_{12}$. By the connectivity of X , $f \notin N_H(L) \cong \mathbb{Z}_2$. Thus, f has six choices and by Magma [1], the six coset graphs $\text{Cos}(G, H, HfH)$ corresponding to the six involutions have two nonisomorphism classes. It follows that $X = \text{Cos}(G, H, HfH) \cong \mathcal{C}_{78}^1$ or \mathcal{C}_{78}^2 , as required. \square

Example 3.3. Let $G = S_8$. Then G has a subgroup $H \cong \mathbb{Z}_2^3 \times \text{SL}(3, 2)$ and an involution g such that $|HgH|/|H| = 7$ and $\langle H, g \rangle = G$. The coset graph $\text{Cos}(G, H, HgH)$ is denoted by \mathcal{C}_{30} .

Lemma 3.4. *Each connected valency seven symmetric graph X of order 30 admitting S_8 as an arc-transitive automorphism group is isomorphic to \mathcal{C}_{30} . Furthermore, X is 2-transitive and $\text{Aut}(X) \cong S_8$.*

Proof. Let $G = S_8$. Clearly, G has a maximal subgroup $T \cong A_8$ containing a maximal subgroup H such that $H \cong \mathbb{Z}_2^3 \times \text{SL}(3, 2)$. Let $L = \mathbb{Z}_2^3 \times S_4$ be a subgroup of H . By Magma [1], $N_G(L) = L \cdot \mathbb{Z}_2$ and $N_T(L) = L$, and by [19], one has $N_G(L) = S_2 \wr S_4$. Let $g \in N_G(L) \setminus L$ be an involution. Then $N_G(L) = L \cup Lg$, $L = H \cap H^g$, $|HgH|/|H| = 7$ and $\langle H, g \rangle = G$. It follows that the coset graph

$\text{Cos}(G, H, HgH)$ is a connected valency seven symmetric graph of order 30. (Note that H has yet another conjugacy class of order $|\mathbb{Z}_2^3 \rtimes S_4|$, which is not isomorphic to $\mathbb{Z}_2^3 \rtimes S_4$. By Magma [1], $N_G(L) = L$, no graph arises.)

Let X be a connected valency seven symmetric graph of order 30 admitting $G = S_8$ as an arc-transitive automorphism group. Then $G_v \cong \mathbb{Z}_2^3 \rtimes \text{SL}(3, 2)$ for any $v \in V(X)$. To complete the proof, it suffices to show that $X \cong \text{Cos}(G, H, HgH)$. Since G has one conjugacy class of $\mathbb{Z}_2^3 \rtimes \text{SL}(3, 2)$ and $\mathbb{Z}_2^3 \rtimes \text{SL}(3, 2)$ has seven subgroups isomorphic to $\mathbb{Z}_2^3 \rtimes S_4$, by Proposition 2.7, one may assume that $X = \text{Cos}(G, H, HfH)$ such that $H \cap H^f = L$ and $f \in N_G(L)$. Since $N_G(L) = L \cup Lg$, one has $f = lg$ for some $l \in L$. It follows that $HfH = HgH$, that is, $X = \text{Cos}(G, H, HfH) \cong \text{Cos}(G, H, HgH)$. By Magma [1], $\text{Aut}(X) = G$. \square

Example 3.5. Let $G = \text{Aut}(\text{PSL}(5, 2)) = \text{PSL}(5, 2).\mathbb{Z}_2$. Then G has a subgroup $H \cong \mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$ and an involution g such that $|HgH|/|H| = 7$ and $\langle H, g \rangle = G$. The coset graph $\text{Cos}(G, H, HgH)$ is denoted by \mathcal{C}_{310} .

Lemma 3.6. *Each connected valency seven symmetric graph X of order 310 admitting $\text{Aut}(\text{PSL}(5, 2))$ as an arc-transitive automorphism group is isomorphic to \mathcal{C}_{310} . Furthermore, X is 3-transitive and $\text{Aut}(X) \cong \text{Aut}(\text{PSL}(5, 2))$.*

Proof. By Atlas [3], $\text{Aut}(\text{PSL}(5, 2)) = \text{PSL}(5, 2).\mathbb{Z}_2$. Let $G = \text{Aut}(\text{PSL}(5, 2))$. Clearly, G has an index two maximal subgroup $T \cong \text{PSL}(5, 2)$ containing a maximal subgroup H such that $H \cong \mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$. Let $L = \mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times S_4)$ be a subgroup of H . By Magma [1], $N_G(L) = L.\mathbb{Z}_2$ and $N_T(L) = L$. Let $g \in N_G(L) \setminus L$ be an involution. Then $N_G(L) = L \cup Lg$, $L = H \cap H^g$, $|HgH|/|H| = 7$ and $\langle H, g \rangle = G$. It follows that the coset graph $\text{Cos}(G, H, HgH)$ is a connected valency seven symmetric graph of order 310. (Note that $\mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$ has yet another conjugacy class of order $|\mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times S_4)|$, which is not isomorphic to $\mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times S_4)$. By Magma [1], $N_G(L) = L$, no graph arises.)

Let X be a connected valency seven symmetric graph of order 310 admitting G as an arc-transitive automorphism group. Then $G_v \cong \mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$ for any $v \in V(X)$. To complete the proof, it suffices to show that $X \cong \text{Cos}(G, H, HgH)$. Since T has two conjugacy classes of maximal parabolic subgroups $\mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$, and has a graph automorphism g , which is of order 2, g fuses the two conjugacy classes of maximal parabolic subgroups. By Magma [1], $\mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$ has a conjugacy class of $\mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times S_4)$. By Proposition 2.7, one may assume that $X = \text{Cos}(G, H, HfH)$ so that $H \cap H^f = L$ and $f \in N_G(L)$. Since $N_G(L) = L \cup Lg$, one has $f = lg$ for some $l \in L$. It follows that $HfH = HgH$, that is, $X = \text{Cos}(G, H, HfH) \cong \text{Cos}(G, H, HgH)$. By Magma [1], $\text{Aut}(X) = G$. \square

4. MAIN RESULTS

In this section, we classify valency seven symmetric graphs of order $2pq$ for p and q primes. First, we consider valency seven symmetric graphs of order $4p$, where p is a prime.

Theorem 4.1. *Let p be a prime. Then X is a connected valency seven symmetric graph of order $4p$ if and only if $X \cong K_8$, a complete graph of order 8.*

Proof. For $p = 2$, K_8 is a unique symmetric graph of valency seven. For $p = 3$, by [21], there is no symmetric graph of valency seven. Thus, in what follows, we assume that $p \geq 5$. Let $A = \text{Aut}(X)$ and $v \in V(X)$. By Guo [15], $|A_v| \mid 2^{24} \cdot 3^4 \cdot 5^2 \cdot 7$, and hence $|A| \mid 2^s \cdot 3^t \cdot 5^r \cdot 7 \cdot p$ with $2 \leq s \leq 26$, $0 \leq t \leq 4$ and $0 \leq r \leq 2$. We divide our discussion into the following two cases. Let N be a minimal normal subgroup of A .

Assume that N is solvable. Then N is elementary abelian. By Proposition 2.1, N is semiregular on $V(X)$, and the quotient graph X_N of X relative to the orbits of N has valency seven. Since $|V(X)| = 4p$, A has no normal subgroup of order 4 or p .

It follows that $N \cong \mathbb{Z}_2$, forcing that $N \leq Z(A)$, the center of A . By Proposition 2.1, X_N is a connected valency seven symmetric graph of order $2p$ with A/N as an arc-transitive subgroup of $\text{Aut}(X_N)$. By Proposition 2.5, either $X_N \cong K_{7,7}$ or $X_N \cong G(2p, 7)$ with $7 \mid p - 1$. Take a minimal normal subgroup of A/N , say M/N . Let $X_N \cong K_{7,7}$. Clearly, $p = 7$. Suppose that M/N is solvable. Then $M/N \cong \mathbb{Z}_2, \mathbb{Z}_7$ or \mathbb{Z}_7^2 . If $M/N \cong \mathbb{Z}_2$ or \mathbb{Z}_7 , then A has a normal subgroup of order 4 or 7 because $N \cong \mathbb{Z}_2$, a contradiction. If $M/N \cong \mathbb{Z}_7^2$ then $M \cong \mathbb{Z}_2 \times \mathbb{Z}_7^2$. It is easy to see that M has two orbits on $V(X)$, and since M is abelian and $M_v \cong \mathbb{Z}_7$, one has $X \cong 2K_{7,7}$, a union of two copies of $K_{7,7}$, which contradicts the connectivity of X . Suppose that M/N is nonsolvable. Then $M/N \cong A_7$ or $A_7 \times A_7$. Obviously, M/N has two orbits on $V(X_N)$. Since $(M/N)_u \trianglelefteq (A/N)_u$ for any $u \in X_N$, by the primitivity of $(A/N)_u$ on the neighborhood of u one has $7 \mid |(M/N)_u|$, implying that $49 \mid |M/N|$. Thus, $M/N \cong A_7 \times A_7$. Let $B/N \cong A_7$ and $B/N \trianglelefteq M/N$. Similarly, B/N has two orbits on $V(X_N)$ and $7 \mid |(B/N)_w|$. Thus, $49 \mid |B/N|$, a contradiction. Let $X_N \cong G(2p, 7)$ with $7 \mid p - 1$. Then a normal Sylow p -subgroup of $\text{Aut}(X_N)$ must be PN/N because each Sylow p -subgroup of A/N is a Sylow p -subgroup of $\text{Aut}(X_N)$. It follows that $P \trianglelefteq A$ because P is characteristic in PN , which is impossible because A has no normal subgroup of order p .

If A has a solvable nontrivial normal subgroup, then A has a solvable minimal normal subgroup isomorphic to $\mathbb{Z}_2, \mathbb{Z}_2^2$ or \mathbb{Z}_p , which is impossible by the above

argument. Thus, in what follows we assume that A has no solvable nontrivial normal subgroups.

Now assume that N is nonsolvable. Then $N \cong T^m$, where T is a nonabelian simple group. By Proposition 2.1, N has at most two orbits on $V(X)$. Then $|N|$ is divisible by $2p \cdot 7$, and since $|N| \mid 2^s \cdot 3^t \cdot 5^r \cdot 7 \cdot p$ with $1 \leq s \leq 26$, $0 \leq t \leq 4$ and $0 \leq r \leq 2$. One has $N = T$ except $p = 7$.

If $p = 5$, then $|N|$ is a factor of $2^{26} \cdot 3^4 \cdot 5^3 \cdot 7$ and $|N|$ is divisible by $2 \cdot 5 \cdot 7$. By Table 1,

$$(4.1) \quad N \cong A_7, A_8, A_9, A_{10}, \text{PSL}(3, 4), \text{PSU}(3, 5), J_2, \text{PSp}(6, 2).$$

For $p = 7$, one has $7^2 \mid |N|$ and $|N| \mid 2^{26} \cdot 3^4 \cdot 5^2 \cdot 7^2$. If $N \cong T^2$, then $T \cong \text{PSL}(3, 4), \text{PSL}(2, 7), A_7, A_8, \text{PSL}(2, 8)$ by Table 1. Clearly, N has a normal subgroup isomorphic to T , say S . Since $S \trianglelefteq N$, one has $7 \mid |S_v|$ and S has an orbit of length 7, 14 or 28, implying that $49 \mid |S|$, a contradiction. Thus, $N = T$. In this case, $|N|$ has at most four primes, and $7^2 \mid |N|$. Again by Table 1, one has

$$(4.2) \quad N \cong \text{PSL}(2, 49).$$

Let $p > 7$. We first consider $N \cong \text{PSL}(2, p)$, the infinite family listed in Table 1. By the subgroup structure of $\text{PSL}(2, p)$, one has N_v is solvable and $|N_v| \mid 2^2 \cdot 3^2 \cdot 7$, and $5 \nmid |N_v|$. Then $|N|$ is a factor of $2^4 \cdot 3^2 \cdot 7 \cdot p$ and $|N|$ is divisible by $2 \cdot p \cdot 7$. Hence $|N| = |\text{PSL}(2, p)| = \frac{1}{2}p(p-1)(p+1)$ and $(\frac{1}{2}p(p+1), \frac{1}{2}(p-1)) = 1$. If $7 \mid p-1$, then $p+1 = 2^i \cdot 3^j$, where $1 \leq i \leq 4$, $0 \leq j \leq 2$. It follows that $p = 71$. Similarly, if $7 \mid p+1$, then $p = 13$. Combining with Table 1, N is one of the following:

$$(4.3) \quad \text{PSL}(2, 13), \text{PSL}(2, 71), \text{PSL}(2, 27), \text{PSU}(3, 8), Sz(8), A_{11}, M_{22}, \text{PSL}(2, 2^6),$$

$$(4.4) \quad \text{PSL}(4, 4), \text{PSL}(5, 2), {}^2D_4(2), G_2(4).$$

Since N is nonsolvable, N has at most two orbits. We may assume that N is a group listed in (4.1)–(4.4). Let N be transitive on $V(X)$. By Proposition 2.7, $X \cong \text{Cos}(N, H, HaH)$, where $H = N_v$, $a \in N \setminus H$ and $a^2 \in H$. By the Atlas [3], $N = A_7$ ($p = 5$) has no subgroup of order $|H| = |N|/|V(X)|$. Thus, $N \neq A_7$. Similarly, $N \neq \text{PSL}(2, 2^6)$ ($p = 13$). For $N = A_8$ ($p = 5$), $|N|/|V(X)|$ is not the order of the vertex stabilizer by Proposition 2.2, a contradiction. It follows that $N \neq A_8$. Similarly, $N \neq A_9$ ($p = 5$), A_{10} ($p = 5$), $\text{PSL}(3, 4)$ ($p = 5$), $\text{PSU}(3, 5)$ ($p = 5$), J_2 ($p = 5$), $\text{PSp}(6, 2)$ ($p = 5$), $\text{PSL}(2, 49)$, $\text{PSL}(2, 13)$ ($p = 13$), $\text{PSL}(2, 71)$ ($p = 71$), $\text{PSL}(2, 27)$ ($p = 13$), $\text{PSU}(3, 8)$ ($p = 19$), $Sz(8)$ ($p = 13$), A_{11} ($p = 11$), M_{22} ($p = 11$), $\text{PSL}(4, 4)$ ($p = 17$), $\text{PSL}(5, 2)$ ($p = 31$), ${}^2D_4(2)$ ($p = 17$), $G_2(4)$ ($p = 13$).

Let N have two orbits on $V(X)$. Then $|H| = |N|/\frac{1}{2}|V(X)|$. For $N = A_7$ ($p = 5$), by Proposition 2.2, $|N|/\frac{1}{2}|V(X)|$ is not the order of the vertex stabilizer, a contradiction. It follows that $N \neq A_7$. Similarly, $N \neq A_9$ ($p = 5$), A_{10} ($p = 5$), $\text{PSL}(3, 4)$ ($p = 5$), $\text{PSU}(3, 5)$ ($p = 5$), J_2 ($p = 5$), $\text{PSp}(6, 2)$ ($p = 5$), $\text{PSL}(2, 49)$, $\text{PSL}(2, 13)$ ($p = 13$), $\text{PSL}(2, 71)$ ($p = 71$), $\text{PSL}(2, 27)$ ($p = 13$), $\text{PSU}(3, 8)$ ($p = 19$), $Sz(8)$ ($p = 13$), M_{22} ($p = 11$), $\text{PSL}(4, 4)$ ($p = 17$), $\text{PSL}(5, 2)$ ($p = 31$), ${}^2D_4(2)$ ($p = 17$), $G_2(4)$ ($p = 13$). By the Atlas [3], $N = A_8$ has no subgroup of order $|H| = |N|/\frac{1}{2}|V(X)|$. Thus, $N \neq A_8$ ($p = 5$). Similarly, $N \neq \text{PSL}(2, 2^6)$ ($p = 13$). For $N = A_{11}$, one has $|H| = 2^6 \cdot 3^4 \cdot 5^2 \cdot 7$. By Proposition 2.2, $H \cong A_7 \times A_6$, and by [19], A_{11} has no subgroup which is isomorphic to $A_7 \times A_6$, a contradiction. This completes the proof. \square

Theorem 4.2. *Let X be a connected valency seven symmetric graph of order $2pq$, where $p > q$ are odd primes. Then X is 1-, 2- or 3-transitive. Furthermore, one of the following situations occurs:*

- (1) X is 1-transitive, and $X \cong C_{78}^i$ ($i = 1, 2$) with $\text{Aut}(C_{78}^1) \cong \text{PSL}(2, 13)$ and $\text{Aut}(C_{78}^2) \cong \text{PGL}(2, 13)$, or $X \cong \mathcal{CD}_{2pq}^l$ (defined in equation (2.2)) with $\text{Aut}(X) \cong D_{2pq} \rtimes \mathbb{Z}_7$ for some l satisfying $l^6 + l^5 + l^4 + l^3 + l^2 + l \equiv 0 \pmod{pq}$ —the number of pairwise nonisomorphic such graphs of order $2pq$ is

$$f(p, q) = \begin{cases} 1, & q = 7 \text{ and } 7 \mid p - 1; \\ 6, & 7 \mid q - 1 \text{ and } 7 \mid p - 1; \\ 0, & \text{otherwise.} \end{cases}$$

- (2) X is 2-transitive, and $X \cong C_{30}$ is a vertex bi-primitive graph with $\text{Aut}(X) \cong S_8$.
(3) X is 3-transitive, and $X \cong C_{310}$ is a vertex bi-primitive graph with $\text{Aut}(X) \cong \text{PSL}(5, 2) \cdot \mathbb{Z}_2$.

Proof. Let $A = \text{Aut}(X)$ and $v \in V(X)$. By Guo [15], $|A_v| \mid 2^{24} \cdot 3^2 \cdot 5^2 \cdot 7$, and hence $|A| = 2^s \cdot 3^t \cdot 5^r \cdot 7 \cdot q \cdot p$ with $1 \leq s \leq 25, 0 \leq t \leq 4$ and $0 \leq r \leq 2$. We first prove a claim.

Claim: If A has a normal subgroup of order q then $X \cong \mathcal{CD}_{2pq}^l$.

Let Q be a normal subgroup of A of order q . By Proposition 2.1, Q is semiregular on $V(X)$ and the quotient graph X_Q of X relative to Q is a symmetric graph of order $2p$ and valency seven with A/Q as an arc-transitive subgroup of $\text{Aut}(X_Q)$. By Proposition 2.5, one has $X_Q \cong K_{7,7}$ or $X_Q \cong G(2p, 7)$ with $7 \mid p - 1$.

Suppose that $X_Q \cong K_{7,7}$. Then $p = 7$ and $q = 3$ or 5 . Take a minimal normal subgroup of A/Q , say M/Q . Assume that M/Q is nonsolvable. Then $M/Q \cong A_7$ or $A_7 \times A_7$ because $A/Q \leq \text{Aut}(K_{7,7}) \cong (S_7 \times S_7) \rtimes \mathbb{Z}_2$. Obviously, M/Q has two orbits

on $V(X_Q)$ and $7 \mid |(M/Q)_w|$ for any $w \in V(X_Q)$, implying that $49 \mid |M/Q|$. Thus, $M/Q \cong A_7 \times A_7$. Let $B/Q \cong A_7$ and $B/Q \trianglelefteq M/Q$. Similarly, B/Q has two orbits on $V(X_Q)$ and $7 \mid |(B/Q)_w|$. Thus, $49 \mid |B/Q|$, a contradiction. Now assume that M/Q is solvable. Then $M/Q \cong \mathbb{Z}_2, \mathbb{Z}_7$ or \mathbb{Z}_7^2 . If $M/Q \cong \mathbb{Z}_2$ then X_M is a symmetric graph of order p and valency seven, a contradiction. If $M/Q \cong \mathbb{Z}_7$ then $M \cong \mathbb{Z}_{21}$ or \mathbb{Z}_{35} and M has two orbits on $V(X)$, implying that X is a bipartite graph. Let $R \leq M$ and $R \cong \mathbb{Z}_7$. Then $R \triangleleft A$, and since $R \leq M$, the quotient graph X_R is bipartite and of valency seven. However, $|X_R| = 6$ or 10 , a contradiction. If $M/Q \cong \mathbb{Z}_7^2$ then $M \cong Q \times \mathbb{Z}_7^2$ because $Q \cong \mathbb{Z}_3$ or \mathbb{Z}_5 . Since M is abelian and $M_v \cong \mathbb{Z}_7$, one has $X \cong 3K_{7,7}$ or $5K_{7,7}$, which contradicts the connectivity of X .

Thus, $X_Q \cong G(2p, 7)$ with $7 \mid p - 1$. By Proposition 2.5, X_Q is valency seven and 1-regular graph of order $2p$. Since A/Q is arc-transitive on X_Q , one has $A/Q = \text{Aut}(X_Q)$ and X is a valency seven and 1-regular graph of order $2pq$. By Proposition 2.6, $X \cong \mathcal{CD}_{2pq}^l$. This completes the proof of Claim.

If A has a normal subgroup of order 2, then the quotient graph has valency seven and odd order pq , a contradiction.

Let A have a normal subgroup P of order p . By Proposition 2.5, $X_P \cong K_{7,7}$ or $G(2q, 7)$ with $7 \mid q - 1$. Let $C := C_A(P)$. Clearly, $P \leq C$. If $P = C$ then $A/P \leq \text{Aut}(P) \cong \mathbb{Z}_{p-1}$, implying that A is abelian. It follows that A is regular on $V(X)$, which contradicts the fact that X is symmetric. Hence, $P < C$. Take a minimal normal subgroup of A/P , say M/P , in C/P . Suppose that M/P is solvable. By Proposition 2.1, M/P is semiregular on $V(X_P)$. Then $M/P \cong \mathbb{Z}_2$ or \mathbb{Z}_q , which implies that A has a normal subgroup of order 2 or q respectively; we have done two cases. Thus, M/P is nonsolvable, and hence $X_P \cong K_{7,7}$. Then $M/P \cong A_7$ or $A_7 \times A_7$. Obviously, M/P has two orbits on $V(X_P)$, and $7 \mid |(M/P)_u|$ for any $u \in V(X_P)$, implying that $49 \mid |M/P|$. Thus, $M/P \cong A_7 \times A_7$. Let $B/P \cong A_7$ and $B/P \trianglelefteq M/P$. Similarly, B/P has two orbits on $V(X_P)$ and $7 \mid |(B/P)_u|$. Thus, $49 \mid |B/P|$, a contradiction.

If A has a solvable nontrivial normal subgroup, then A has a solvable minimal normal subgroup isomorphic to $\mathbb{Z}_2, \mathbb{Z}_p$ or \mathbb{Z}_q , which was done by the above argument. Thus, in what follows we assume that A has no solvable nontrivial normal subgroups.

Let N be a minimal normal subgroup of A . Then $N \cong T^m$, where T is a nonabelian simple group. By Proposition 2.1, N has at most two orbits on $V(X)$. Since $pq \cdot 7 \mid |N|$ and $|N| \mid |A| = 2^s \cdot 3^t \cdot 5^r \cdot 7 \cdot q \cdot p$ with $1 \leq s \leq 25, 0 \leq t \leq 4$ and $0 \leq r \leq 2$, one has $N = T$ except for $p = 7$. Assuming that $p = 7$, one has $q = 3$ or 5 . Hence $7^2 \mid |N|$ and $|N| \mid 2^{25} \cdot 3^5 \cdot 5^3 \cdot 7^2$. If $N \cong T^2$, then $T \cong \text{PSL}(2, 7), \text{PSL}(2, 8), \text{PSL}(3, 4), A_7, A_8$ by Table 1. Clearly, N has a normal subgroup isomorphic to T , say S . Since $S \trianglelefteq N$, one has $7 \mid |S_v|$ and S has an orbit of length 7, $7q$ or $14q$, implying that $49 \mid |S|$, a contradiction. Thus, $N = T$. In this case, $|N|$ has at most four primes

$\{2, 3, 5, 7\}$, and $7^2 \mid |N|$. Again by Table 1, one has

$$(4.5) \quad N \cong \text{PSL}(2, 49).$$

Next, we assume that $p \neq 7$ and $N = T$. We first consider $N \cong \text{PSL}(2, p)$ ($p > 7$), the infinite family listed in Table 1. By the subgroup structure of $\text{PSL}(2, p)$, one has N_v is solvable, and by Proposition 2.2, $|N_v| \mid 2^2 \cdot 3^2 \cdot 7$ and $5 \nmid |N_v|$. Thus $|N| \mid 2^3 \cdot 3^2 \cdot 7 \cdot q \cdot p$, implying that N is at most five-prime factor simple group. Hence $|N| = |\text{PSL}(2, p)| = \frac{1}{2}p(p-1)(p+1)$ and $(\frac{1}{2}(p+1), \frac{1}{2}(p-1)) = 1$. For $q \leq 7$, if $7 \mid \frac{1}{2}(p+1)$, then $p-1 = 2^i \cdot 3^j \cdot q$, where $1 \leq i \leq 3, 0 \leq j \leq 2$. It follows that $p = 13, 41, 181$. Since $\text{PSL}(2, 181)$ is a six-prime factor simple group, one has $p = 13, 41$. Similarly, if $7 \mid \frac{1}{2}(p-1)$, then $p = 29, 71$. For $p > q > 7$, if $q \mid \frac{1}{2}(p+1)$, then $p-1 = 2^i \cdot 3^j \cdot 7$, where $1 \leq i \leq 3, 0 \leq j \leq 2$. It follows that $p = 43, 127$. Since $2^7 \mid |\text{PSL}(2, 127)|$, a contradiction. Thus $p = 43$. Similarly, if $q \mid \frac{1}{2}(p-1)$, then $p = 83, 97, 251, 503$. Hence $2^5 \mid |\text{PSL}(2, 97)|$ and $5^3 \mid |\text{PSL}(2, 251)|$, a contradiction. Thus $p = 83, 503$.

For $q = 3$, one has $3 \cdot 7 \cdot p \mid |N|$ and $|N| \mid 2^{25} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot p$. By Table 1, N is one of the following groups:

$$(4.6) \quad A_7, A_8, A_9, A_{10}, \text{PSL}(2, 27), \text{PSL}(3, 4), \text{PSU}(3, 5), \text{PSU}(3, 8), J_2,$$

$$(4.7) \quad D_4(2), \text{PSp}(6, 2), \text{PSp}(8, 2), A_{11}, A_{12}, M_{22}, \text{PSL}(2, 2^6), \text{PSL}(4, 4),$$

$$(4.8) \quad \text{PSL}(4, 4), \text{PSL}(5, 2), {}^2D_4(2), G_2(4), \text{PSL}(2, p) \quad (p = 13, 127).$$

For $q = 5$, one has $5 \cdot 7 \cdot p \mid |N|$ and $|N| \mid 2^{25} \cdot 3^4 \cdot 5^3 \cdot 7 \cdot p$. By Table 1, N is one of the following groups:

$$(4.9) \quad Sz(8), A_{11}, M_{22}, HS, \text{PSL}(2, 2^6), \text{PSL}(2, 5^3), \text{PSL}(5, 2), \text{PSL}(4, 4),$$

$$(4.10) \quad {}^2D_4(2), G_2(4), \text{PSL}(2, p) \quad (p = 29, 41, 71).$$

For $q \geq 7$, one has $7 \cdot q \cdot p \mid |N|$ and $|N| \mid 2^{25} \cdot 3^4 \cdot 5^2 \cdot 7 \cdot q \cdot p$. By Table 1, N is one of the following groups:

$$(4.11) \quad \text{PSL}(3, 8), {}^3D_4(2), \text{PSL}(2, 2^9), \text{PSp}(4, 8), M_{23}, M_{24}, J_1, \text{PSL}(3, 16),$$

$$(4.12) \quad \text{PSL}(2, p) \quad (p = 43, 83, 503).$$

We may assume that N is a group listed in (4.5)–(4.12). Let $G \leq A$ be a transitive subgroup of X . By Proposition 2.7, $X \cong \text{Cos}(G, H, HgH)$, where $H = G_v, g \in G \setminus H$, and $g^2 \in H$, implying that a normalizes $R = H \cap H^g$, that is, $g \in N_G(R) \setminus H$. Recall that N has at most two orbits on $V(X)$. First let N be transitive on $V(X)$. Take $G = N$.

If $N = A_7$ ($q = 3, p = 5$), then N_v has order $|N|/|V(X)| = 2^2 \cdot 3 \cdot 7$. However, A_7 has no subgroups of order $2^2 \cdot 3 \cdot 7$ by Atlas [3]. Thus, $N \neq A_7$. Similarly, $N \neq \text{PSL}(2, 127)$ ($q = 3, p = 127$), $\text{PSL}(2, 29)$ ($q = 5, p = 29$), $\text{PSL}(2, 41)$ ($q = 5, p = 41$), $\text{PSL}(2, 71)$ ($q = 5, p = 71$) and $\text{PSL}(2, 503)$ ($q = 251, p = 503$) by Proposition 2.4. If $N = A_9$ ($q = 3, p = 5$), A_{10} ($q = 3, p = 5$), $\text{PSU}(3, 8)$ ($q = 3, p = 19$), A_{11} ($q = 3, p = 11$) or ${}^2D_4(2)$ ($q = 3, p = 17$), then $3^3 \parallel |N_v|$ and $3^4 \nmid |N_v|$. By Proposition 2.2, it is not possible. If $N = A_8$ ($q = 3, p = 5$), then $|N_v| = 2^5 \cdot 3 \cdot 7$. By Proposition 2.2, there exists a vertex stabilizer whose order is $2^5 \cdot 3 \cdot 7$, a contradiction. Similarly, $N \neq \text{PSL}(2, 49)$ ($q = 3, p = 7$), $\text{PSL}(2, 49)$ ($q = 5, p = 7$), $\text{PSL}(2, 27)$ ($q = 3, p = 13$), $\text{PSL}(3, 4)$ ($q = 3, p = 5$), $\text{PSU}(3, 5)$ ($q = 3, p = 5$), J_2 ($q = 3, p = 5$), $\text{PSp}(6, 2)$ ($q = 3, p = 5$), $\text{PSp}(8, 2)$ ($q = 3, p = 5$), A_{11} ($p = 5, q = 11$), M_{22} ($q = 3, p = 11$), $\text{PSL}(2, 2^6)$ ($q = 3, p = 13$), $\text{PSL}(2, 2^6)$ ($q = 5, p = 13$), $\text{PSL}(4, 4)$ ($q = 3, p = 17$), $\text{PSL}(4, 4)$ ($q = 5, p = 17$), $\text{PSL}(5, 2)$ ($q = 3, p = 31$), $\text{PSL}(5, 2)$ ($q = 5, p = 31$), $D_4(2)$ ($q = 3, p = 5$), HS ($q = 5, p = 11$), $\text{PSL}(2, 5^3)$ ($q = 5, p = 31$), ${}^2D_4(2)$ ($q = 5, p = 17$), $G_2(4)$ ($q = 3, p = 13$), $G_2(4)$ ($q = 5, p = 13$), $Sz(8)$ ($q = 5, p = 13$), $\text{PSL}(2, 71)$ ($q = 3, p = 71$), $\text{PSL}(3, 8)$ ($q = 7, p = 73$), ${}^3D_4(2)$ ($q = 7, p = 17$), $\text{PSL}(2, 2^9)$ ($q = 19, p = 73$), $\text{PSp}(4, 8)$ ($q = 7, p = 13$), M_{23} ($q = 11, p = 23$), M_{24} ($q = 11, p = 23$), J_1 ($q = 11, p = 19$) and $N \neq \text{PSL}(3, 16)$ ($q = 13, p = 17$).

Suppose that $N = A_{12}$ ($q = 3, p = 11$). Then $|N_v| = |N|/|V(X)| = 2^8 \cdot 3^4 \cdot 5^2 \cdot 7$. By Proposition 2.2, $N_v \cong S_7 \times S_6$. By [19], one concludes that N has no subgroup which is isomorphic to $S_7 \times S_6$, a contradiction.

Suppose that $N = \text{PSL}(2, 13)$ ($q = 3, p = 13$). Then $|N_v| = 2 \cdot 7$. By Proposition 2.2, $N_v \cong D_{14}$, and by Proposition 2.4, N_v is a maximal subgroup. By Example 3.1, $X \cong \mathcal{C}_{78}^1$ or \mathcal{C}_{78}^2 .

Suppose that $N = \text{PSL}(2, 43)$ ($q = 11, p = 43$). Then $|N_v| = 2 \cdot 3 \cdot 7$. By Proposition 2.2, $N_v \cong F_{42}$, and by Proposition 2.4, one concludes that N has a unique conjugacy class D_{42} which has order 42. Clearly, it is isomorphic to F_{42} , a contradiction.

Suppose that $N = \text{PSL}(2, 83)$ ($q = 41, p = 83$). Then $|N_v| = 2 \cdot 3 \cdot 7$. By Proposition 2.2, $N_v \cong F_{42}$. By Proposition 2.4, one concludes that N has a unique maximal subgroup conjugacy class D_{84} which contains subgroups of order 42. Clearly, the subgroups of order 42 of D_{84} are isomorphic to D_{42} or \mathbb{Z}_{42} . They are not isomorphic to F_{42} , a contradiction.

Suppose that $N = M_{22}$ ($q = 5, p = 11$). Then $|N_v| = 2^6 \cdot 3^2 \cdot 7$. By Atlas [3], the unique maximal subgroup class of M_{22} which has order divided by $2^6 \cdot 3^2 \cdot 7$ is $L_3(4)$; again by Atlas [3], $L_3(4)$ has no subgroup of order $2^6 \cdot 3^2 \cdot 7$, a contradiction.

Now let N have two orbits on $V(X)$. If $N \neq A_7$, then N_v has order $|N|/\frac{1}{2}|V(X)| = 2^3 \cdot 3 \cdot 7$. However, A_7 has no subgroups of order $2^3 \cdot 3 \cdot 7$ by Atlas [3]. Thus, $N \neq A_7$. Similarly, $N \neq \text{PSL}(2, 13)$ ($q = 3, p = 13$), $\text{PSL}(2, 27)$ ($q = 3, p = 13$), $\text{PSL}(2, 127)$

$(q = 3, p = 127)$, $\text{PSL}(2, 29)$ ($q = 5, p = 29$), $\text{PSL}(2, 41)$ ($q = 5, p = 41$), $\text{PSL}(2, 71)$ ($q = 5, p = 71$) and $\text{PSL}(2, 43)$ ($q = 11, p = 43$) by Proposition 2.4. If $N = A_9$ ($q = 3, p = 5$), A_{10} ($q = 3, p = 5$), A_{11} ($q = 3, p = 11$), $\text{PSU}(3, 8)$ ($q = 3, p = 19$) or ${}^2D_4(2)$ ($q = 3, p = 17$), then $3^3 \parallel |N_v|$. By Proposition 2.2, this is not possible. If $N = A_{12}$ ($q = 3, p = 11$), then $|N_v| = 2^9 \cdot 3^4 \cdot 5^2 \cdot 7$. By Proposition 2.2, there exists no vertex stabilizer whose order is $2^9 \cdot 3^4 \cdot 5^2 \cdot 7$, a contradiction. Similarly, $N \neq \text{PSL}(2, 49)$ ($q = 3, p = 7$), $\text{PSL}(2, 49)$ ($q = 5, p = 7$), $\text{PSL}(2, 503)$ ($q = 251, p = 503$), $\text{PSL}(3, 4)$ ($q = 3, p = 5$), $\text{PSU}(3, 5)$ ($q = 3, p = 5$), J_2 ($q = 3, p = 5$), $\text{PSp}(6, 2)$ ($q = 3, p = 5$), $\text{PSp}(8, 2)$ ($q = 3, p = 5$), $Sz(8)$ ($q = 5, p = 13$), A_{11} ($p = 5, q = 11$), M_{22} ($q = 3, p = 11$), M_{22} ($q = 5, p = 11$), $\text{PSL}(2, 2^6)$ ($q = 3, p = 13$), $\text{PSL}(2, 2^6)$ ($q = 5, p = 13$), $\text{PSL}(4, 4)$ ($q = 3, p = 17$), $\text{PSL}(4, 4)$ ($q = 5, p = 17$), $\text{PSL}(5, 2)$ ($q = 3, p = 31$), HS ($q = 5, p = 11$), ${}^2D_4(2)$ ($q = 5, p = 17$), $D_4(2)$ ($q = 3, p = 5$), $G_2(4)$ ($q = 3, p = 13$), $G_2(4)$ ($q = 5, p = 13$), $\text{PSL}(2, 5^3)$ ($q = 5, p = 31$), $\text{PSL}(2, 71)$ ($q = 3, p = 71$), $\text{PSL}(2, 71)$ ($q = 5, p = 71$), $\text{PSL}(3, 8)$ ($q = 7, p = 73$), ${}^3D_4(2)$ ($q = 7, p = 17$), $\text{PSp}(4, 8)$ ($q = 7, p = 13$), $\text{PSL}(2, 2^9)$ ($q = 19, p = 73$), M_{23} ($q = 11, p = 23$), M_{24} ($q = 11, p = 23$), J_1 ($q = 11, p = 19$) and $N \neq \text{PSL}(3, 16)$ ($q = 13, p = 17$).

Suppose that $N = \text{PSL}(2, 83)$ ($q = 41, p = 83$). Then $|N_v| = 2^2 \cdot 3 \cdot 7$. By Proposition 2.2, $N_v \cong F_{42} \times \mathbb{Z}_2$. By Proposition 2.4, one concludes that N has a unique conjugacy class D_{84} which has order 84. Clearly, it is isomorphic to $F_{42} \times \mathbb{Z}_2$, a contradiction.

Suppose that $N = A_8$ ($q = 3, p = 5$). Then $|N_v| = |N|/\frac{1}{2}|V(X)| = 2^6 \cdot 3 \cdot 7$. By Proposition 2.2, $N_v \cong \mathbb{Z}_2^3 \times \text{SL}(3, 2)$. In this case, N has two orbits on $V(X)$, and $N_v \cap N_v^g \cong \mathbb{Z}_2^3 \rtimes S_4$. Let $C = C_A(N)$. Since N is simple, $C \cap N = 1$ and $CN = C \times N \trianglelefteq A$. Since $A/CN \lesssim \text{Out}(N)$, we have $A = (C \times N) \cdot O$ with $O \lesssim \text{Out}(N)$, where $\text{Out}(N)$ is the outer automorphism group of N . Hence $|A| \mid 2^{25} \cdot 3^5 \cdot 5^3 \cdot 7$ and $|N| = 2^6 \cdot 3^2 \cdot 5 \cdot 7$. Then $|C| \mid 2^{19} \cdot 3^3 \cdot 5^2$. If C is insoluble, by [3], pages 12–14, then C has a minimal normal insoluble subgroup $M \cong A_5, A_6$ or A_5^2 . Then $NM \trianglelefteq CN$ has at most two orbits on $V(X)$. Then $|(MN)_v| = |MN|/|V(X)| = 2^7 \cdot 3^2 \cdot 5 \cdot 7$ or $|MN|/\frac{1}{2}|V(X)| = 2^8 \cdot 3^2 \cdot 5 \cdot 7$ for $M \cong A_5$. By Proposition 2.2, there exists no vertex stabilizer whose order is $|(MN)_v|$, a contradiction. For $M \cong A_6$ or A_5^2 , one has $3^3 \mid |(MN)_v|$. By Proposition 2.2, this is a contradiction. Thus, C is solvable. Clearly, C is not semiregular on $V(X)$. If it were, X_C would be a connected valency seven graph of order $2pq/|C|$, yielding that $2 \nmid |C|$. Furthermore, $C \not\cong \mathbb{Z}_3$ or \mathbb{Z}_5 , because there is no connected valency seven symmetric graph of order 6 or 10. If C has at most two orbits on $V(X)$, then $|C| = 15$ or 30 . Let $R \cong \mathbb{Z}_5 \leq C$. Then $R \triangleleft A$, and then X_C is a connected valency seven graph of order 6, a contradiction. Thus, $C = 1$ and $A \leq \text{Aut}(A_8)$. Further, $A = S_8$. By Example 3.3, $X \cong \mathcal{C}_{30}$. Hence

N_v is a maximal subgroup of N , and N has two orbits on $V(X)$. Then X is a vertex bi-primitive 3-arc transitive graph.

Suppose that $N = \text{PSL}(5, 2)$ ($q = 5$, $p = 31$). Then $|N_v| = 2^{10} \cdot 3^2 \cdot 7$. By Proposition 2.2, $N_v \cong \mathbb{Z}_2^6 \rtimes (\text{SL}(2, 2) \times \text{SL}(3, 2))$. Let $C = C_A(N)$. Similarly to the above proof, one has $A = (C \times N).O$ with $O \lesssim \text{Out}(N)$, where $\text{Out}(N)$ is the outer automorphism group of N . Hence $|A| \mid 2^{25} \cdot 3^4 \cdot 5^3 \cdot 7 \cdot 31$ and $|N| = 2^{10} \cdot 3^2 \cdot 5 \cdot 7 \cdot 31$. Then $|C| \mid 2^{15} \cdot 3^2 \cdot 5^2$. If C is insolvable, by [3], pages 12–14, then C has a minimal normal insolvable subgroup $M \cong A_5, A_6$ or A_5^2 . Then $NM \trianglelefteq CN$ has at most two orbits on $V(X)$. For $M \cong A_5$, one has $3^3 \mid |(MN)_v|$. By Proposition 2.2, this is a contradiction. Thus, $M \not\cong A_5$. If $M \cong A_6$, then $|(MN)_v| = |MN|/|V(X)| = 2^{13} \cdot 3^4 \cdot 5 \cdot 7$ or $|MN|/\frac{1}{2}|V(X)| = 2^{12} \cdot 3^4 \cdot 5 \cdot 7$. By Proposition 2.2, there exists no vertex stabilizer whose order is $|(MN)_v|$, a contradiction. Similarly, $M \not\cong A_5^2$. Thus, C is solvable. Clearly, C is not semiregular on $V(X)$. If it were, X_C would be a connected valency seven graph of order $2pq/|C|$, yielding that $2 \nmid |C|$. Furthermore, $C \not\cong \mathbb{Z}_{31}$ because there is no connected valency seven symmetric graph of order 10. If $C \cong \mathbb{Z}_5$, by Proposition 2.5, there is no connected valency seven symmetric graph of order 62 because $7 \nmid p - 1$ with $p = 31$. Thus C has at most two orbits on $V(X)$, then $|C| = 5p$ or $10p$. Let $R \cong \mathbb{Z}_p < C$. Then $R \triangleleft A$, and then X_R is a connected valency seven graph of order 10, a contradiction. Thus, $C = 1$ and $A \leq \text{Aut}(N)$. Further, $A \cong \text{Aut}(\text{PSL}(5, 2)) \cdot \mathbb{Z}_2$ because $\text{Out}(N) = \mathbb{Z}_2$. By Example 3.5, $X \cong \mathcal{C}_{310}$. Hence N_v is a maximal subgroup of N , and N has two orbits on $V(X)$. Then X is a vertex bi-primitive 3-arc transitive graph. This completes the proof. \square

Acknowledgements. The authors are indebted to the anonymous referees for many valuable comments and constructive suggestions.

References

- [1] *W. Bosma, J. Cannon, C. Playoust*: The Magma algebra system. I. The user language. *J. Symb. Comput.* *24* (1997), 235–265. [zbl](#) [MR](#) [doi](#)
- [2] *Y. Cheng, J. Oxley*: On weakly symmetric graphs of order twice a prime. *J. Comb. Theory, Ser. B* *42* (1987), 196–211. [zbl](#) [MR](#) [doi](#)
- [3] *J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson*: Atlas of Finite Groups. Maximal Subgroups and Ordinary Characters for Simple Groups. Oxford University Press, Eynsham, 1985. [zbl](#) [MR](#)
- [4] *D. Ž. Djoković, G. L. Miller*: Regular groups of automorphisms of cubic graphs. *J. Comb. Theory, Ser. B* *29* (1980), 195–230. [zbl](#) [MR](#) [doi](#)
- [5] *X. G. Fang, C. E. Praeger*: Finite two-arc transitive graphs admitting a Suzuki simple group. *Commun. Algebra* *27* (1999), 3727–3754. [zbl](#) [MR](#) [doi](#)
- [6] *X. Fang, J. Wang, M. Y. Xu*: On 1-arc-regular graphs. *Eur. J. Comb.* *23* (2002), 785–791. [zbl](#) [MR](#) [doi](#)
- [7] *Y.-Q. Feng, M. Ghasemi, D.-W. Yang*: Cubic symmetric graphs of order $8p^3$. *Discrete Math.* *318* (2014), 62–70. [zbl](#) [MR](#) [doi](#)

- [8] *Y.-Q. Feng, J. H. Kwak*: Cubic symmetric graphs of order a small number times a prime or a prime square. *J. Comb. Theory, Ser. B* *97* (2007), 627–646. [zbl](#) [MR](#) [doi](#)
- [9] *Y.-Q. Feng, J. H. Kwak, M.-Y. Xu*: Cubic s -regular graphs of order $2p^3$. *J. Graph Theory* *52* (2006), 341–352. [zbl](#) [MR](#) [doi](#)
- [10] *Y.-Q. Feng, Y.-T. Li*: One-regular graphs of square-free order of prime valency. *Eur. J. Comb.* *32* (2011), 265–275. [zbl](#) [MR](#) [doi](#)
- [11] *A. Gardiner, C. E. Praeger*: On 4-valent symmetric graphs. *Eur. J. Comb.* *15* (1994), 375–381. [zbl](#) [MR](#) [doi](#)
- [12] *A. Gardiner, C. E. Praeger*: A characterization of certain families of 4-valent symmetric graphs. *Eur. J. Comb.* *15* (1994), 383–397. [zbl](#) [MR](#) [doi](#)
- [13] *D. Gorenstein*: Finite Simple Groups. An Introduction to Their Classification. The University Series in Mathematics, Plenum Press, New York, 1982. [zbl](#) [MR](#) [doi](#)
- [14] *S.-T. Guo, Y.-Q. Feng*: A note on pentavalent s -transitive graphs. *Discrete Math.* *312* (2012), 2214–2216. [zbl](#) [MR](#) [doi](#)
- [15] *S. Guo, Y. Li, X. Hua*: (G, s) -transitive graphs of valency 7. *Algebra Colloq.* *23* (2016), 493–500. [zbl](#) [MR](#) [doi](#)
- [16] *S.-T. Guo, J. Shi, Z.-J. Zhang*: Heptavalent symmetric graphs of order $4p$. *South Asian J. Math.* *1* (2011), 131–136. [zbl](#)
- [17] *X.-H. Hua, Y.-Q. Feng, J. Lee*: Pentavalent symmetric graphs of order $2pq$. *Discrete Math.* *311* (2011), 2259–2267. [zbl](#) [MR](#) [doi](#)
- [18] *Y. Li, Y.-Q. Feng*: Pentavalent one-regular graphs of square-free order. *Algebra Colloq.* *17* (2010), 515–524. [zbl](#) [MR](#) [doi](#)
- [19] *M. W. Liebeck, C. E. Praeger, J. Saxl*: A classification of the maximal subgroups of the finite alternating and symmetric groups. *J. Algebra* *111* (1987), 365–383. [zbl](#) [MR](#) [doi](#)
- [20] *P. Lorimer*: Vertex-transitive graphs: symmetric graphs of prime valency. *J. Graph Theory* *8* (1984), 55–68. [zbl](#) [MR](#) [doi](#)
- [21] *B. D. McKay*: Transitive graphs with fewer than twenty vertices. *Math. Comput.* *33* (1979), 1101–1121. [zbl](#) [MR](#) [doi](#)
- [22] *R. C. Miller*: The trivalent symmetric graphs of girth at most six. *J. Comb. Theory, Ser. B* *10* (1971), 163–182. [zbl](#) [MR](#) [doi](#)
- [23] *J.-M. Oh*: A classification of cubic s -regular graphs of order $14p$. *Discrete Math.* *309* (2009), 2721–2726. [zbl](#) [MR](#) [doi](#)
- [24] *J.-M. Oh*: A classification of cubic s -regular graphs of order $16p$. *Discrete Math.* *309* (2009), 3150–3155. [zbl](#) [MR](#) [doi](#)
- [25] *J. Pan, B. Ling, S. Ding*: One prime-valent symmetric graphs of square-free order. *Ars Math. Contemp.* *15* (2018), 53–65.
- [26] *J. Pan, B. Lou, C. Liu*: Arc-transitive pentavalent graphs of order $4pq$. *Electron. J. Comb.* *20* (2013), Research Paper P36, 9 pages. [zbl](#) [MR](#)
- [27] *P. Potočník*: A list of 4-valent 2-arc-transitive graphs and finite faithful amalgams of index $(4, 2)$. *Eur. J. Comb.* *30* (2009), 1323–1336. [zbl](#) [MR](#) [doi](#)
- [28] *G. Sabidussi*: Vertex-transitive graphs. *Monatsh. Math.* *68* (1964), 426–438. [zbl](#) [MR](#) [doi](#)
- [29] *C. C. Sims*: Graphs and finite permutation groups. *Math. Z.* *95* (1967), 76–86. [zbl](#) [MR](#) [doi](#)
- [30] *M. Suzuki*: Group Theory I. Grundlehren der Mathematischen Wissenschaften 247, Springer, Berlin, 1982. [zbl](#) [MR](#)
- [31] *R. A. Wilson*: The Finite Simple Groups. Graduate Texts in Mathematics 251, Springer, London, 2009. [zbl](#) [MR](#) [doi](#)
- [32] *J. Xu, M. Xu*: Arc-transitive Cayley graphs of valency at most four on abelian groups. *Southeast Asian Bull. Math.* *25* (2001), 355–363. [zbl](#) [MR](#) [doi](#)
- [33] *M.-Y. Xu*: Automorphism groups and isomorphisms of Cayley digraphs. *Discrete Math.* *182* (1998), 309–319. [zbl](#) [MR](#) [doi](#)

- [34] *J.-X. Zhou, Y.-Q. Feng*: Tetravalent s -transitive graphs of order twice a prime power. *J. Aust. Math. Soc.* 88 (2010), 277–288.



Authors' address: Xiao-Hui Hua, Li Chen, College of Mathematics and Information Science, Henan Normal University, Xinxiang 453007, P. R. China, e-mail: xhhua@htu.cn.