

Alireza K. Asboei; Seyed S. S. Amiri

The small Ree group  ${}^2G_2(3^{2n+1})$  and related graph

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 59 (2018), No. 3, 271–276

Persistent URL: <http://dml.cz/dmlcz/147395>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## The small Ree group ${}^2G_2(3^{2n+1})$ and related graph

ALIREZA K. ASBOEI, SEYED S. S. AMIRI

*Abstract.* Let  $G$  be a finite group. The main supergraph  $\mathcal{S}(G)$  is a graph with vertex set  $G$  in which two vertices  $x$  and  $y$  are adjacent if and only if  $o(x) \mid o(y)$  or  $o(y) \mid o(x)$ . In this paper, we will show that  $G \cong {}^2G_2(3^{2n+1})$  if and only if  $\mathcal{S}(G) \cong \mathcal{S}({}^2G_2(3^{2n+1}))$ . As a main consequence of our result we conclude that Thompson’s problem is true for the small Ree group  ${}^2G_2(3^{2n+1})$ .

*Keywords:* main supergraph; simple Ree group; Thompson’s problem

*Classification:* 20D08, 05C25

### 1. Introduction

Let  $G$  be a finite group and  $x \in G$ . The order of  $x$  is denoted by  $o(x)$ . The set of all element orders of  $G$  is denoted by  $\pi_e(G)$  and the set of all prime factors of  $|G|$  is denoted by  $\pi(G)$ . It is clear that the set  $\pi_e(G)$  is closed and partially ordered by divisibility, and hence it is uniquely determined by  $\mu(G)$ , the subset of its maximal elements. Let  $i \in \pi_e(G)$ . Set  $m_i = m_i(G) = |\{g \in G : o(g) = i\}|$ , and  $\text{nse}(G) = \{m_k(G) : k \in \pi_e(G)\}$  be the set of the numbers of elements with the same order.

We define the graph  $\mathcal{S}(G)$  with vertex set  $G$  such that two vertices  $x$  and  $y$  are adjacent if and only if  $o(x) \mid o(y)$  or  $o(y) \mid o(x)$ . This graph is called *main supergraph* of *power graph*  $G$  and was introduced in [8]. The power graph  $\mathcal{P}(G)$  of a group  $G$  is the graph with group elements as vertex set and two elements are adjacent if one is a power of the other. The main properties of this graph were investigated by P. J. Cameron in [3] and I. Chakrabarty et al. in [4]. The *proper main supergraph*  $\mathcal{S}^*(G)$  is the graph constructed from  $\mathcal{S}(G)$  by removing the identity element of  $G$ . We write  $x \sim y$  when two vertices  $x$  and  $y$  are adjacent.

We say that groups  $G_1$  and  $G_2$  are of the same *order type* if and only if  $m_t(G_1) = m_t(G_2)$  for all  $t$ . By the definition of the main supergraph, it is clear that if  $G_1$  and  $G_2$  are groups with the same order type, then  $\mathcal{S}(G_1) \cong \mathcal{S}(G_2)$ . The converse of this result is not generally correct. To prove this, we consider  $G_1 = C_4 \times C_4$  and  $G_2 = C_2 \times C_2 \times C_4$ . Since  $G_1$  and  $G_2$  are 2-groups, we have  $\mathcal{S}(G_1) \cong \mathcal{S}(G_2)$ . But  $m_4(G_1) = 12 > 8 = m_4(G_2)$  and  $m_2(G_1) = 3 < 7 = m_2(G_2)$ .

In 1987, J. G. Thompson, see [9, Problem 12.37], posed the following problem: **Thompson's problem.** Suppose that  $G_1$  and  $G_2$  are two groups of the same order type. If  $G_1$  is solvable, is it true that  $G_2$  is also necessarily solvable?

Obviously, if  $G_1$  and  $G_2$  are the same order type, then  $\text{nse}(G_1) = \text{nse}(G_2)$  and  $|G_1| = |G_2|$ . Therefore, if a group  $G$  has been uniquely determined by its order and  $\text{nse}(G)$ , then Thompson's problem is true for  $G$ . In [6], the authors proved that Thompson's problem is true for the small Ree group  ${}^2G_2(q)$ , where  $q \pm \sqrt{3q} + 1$  is a prime number ( $q = 3^{2n+1}$  and  $n$  is a natural number) by its  $\text{nse}$  and order.

Clearly, for two groups  $G_1$  and  $G_2$  that are the same order type, we have  $\mathcal{S}(G_1) \cong \mathcal{S}(G_2)$ . Therefore, if a group  $G$  has been uniquely determined by  $\mathcal{S}(G)$ , then Thompson's problem is true for  $G$ . If  $G$  is the alternating group of degrees  $p$ ,  $p + 1$  or  $p + 2$  or the symmetric group of degree  $p$ , where  $p$  is prime, then it is proved that these groups are uniquely determined by their main supergraph, see [1]. Also, in [2], it is proved that the groups  $\text{PSL}_2(p)$ ,  $\text{PGL}_2(p)$ , where  $p$  is prime, and all of the sporadic simple groups are uniquely determined by their main supergraph. In this paper, we remove the assumption  $q \pm \sqrt{3q} + 1$  is a prime number in [6] and as the main result, conclude that Thompson's problem is true for  ${}^2G_2(q)$ . In fact, we prove the following theorem.

**Theorem 1.1.** *Let  $G$  be a finite group. If  $\mathcal{S}(G) \cong \mathcal{S}({}^2G_2(3^{2n+1}))$ , where  $n$  is a natural number, then  $G \cong {}^2G_2(3^{2n+1})$ .*

As noted above, as an immediate consequence of Theorem 1.1, we have that

**Corollary 1.2.** *If  $G$  is a finite group with the same type as  ${}^2G_2(3^{2n+1})$ , then  $G$  is isomorphic to  ${}^2G_2(3^{2n+1})$ .*

We construct the *prime graph* of  $G$ , which is denoted by  $\Gamma(G)$ , as follows: the vertex set is  $\pi(G)$  and two distinct vertices  $p$  and  $q$  are joined by an edge if and only if  $G$  has an element of order  $pq$ ,  $p \neq q$ . Let  $t(G)$  be the number of connected components of  $\Gamma(G)$  and let  $\pi_1, \pi_2, \dots, \pi_{t(G)}$  be the connected components of  $\Gamma(G)$ . If  $2 \in \pi(G)$ , then we always suppose  $2 \in \pi_1$ .

Throughout this paper we denote by  $\varphi(n)$ , where  $n$  is a natural number, Euler's totient function. Let  $r$  be a prime number and  $\text{Syl}_r(G)$  be the set of Sylow  $r$ -subgroups of group  $G$ . We denote by  $P_r$  a Sylow  $r$ -subgroup of  $G$  and  $n_r(G)$  is the number of Sylow  $r$ -subgroups of  $G$ , that is,  $n_r(G) = |\text{Syl}_r(G)|$ . The other notations and terminologies in this paper are standard, and the reader is referred to [14] if necessary.

## 2. Preliminary results

We first quote some lemmas that are used in deducing the theorem of this paper.

**Lemma 2.1** ([7]). *Let  $G$  be a finite group and  $m$  be a positive integer dividing  $|G|$ . If  $L_m(G) = \{g \in G : g^m = 1\}$ , then  $m \mid |L_m(G)|$ .*

**Lemma 2.2** ([11]). *Let  $R$  be the small Ree group  ${}^2G_2(3^{2n+1})$ , where  $n$  is a natural number. Then  $\pi_e(G)$  exactly consists of divisors of  $6, 9, q - 1, (q + 1)/2$  and  $q \pm \sqrt{3q} + 1$ .*

**Lemma 2.3.** *Let  $H$  be a finite simple group. Then  $5 \nmid |H|$  holds if and only if  $H$  is isomorphic to one of the following simple groups:*

- (a)  $Z_p, p \neq 5$ ;
- (b)  $\text{PSL}_n(q), n = 2, 3$ , where  $q = p^f$  ( $f$  is odd),  $p \neq 5$  and  $p \neq 5k \pm 1$  for some  $k > 0$ ;
- (c)  $G_2(q)$ , where  $q = p^f$  ( $f$  is odd),  $p \neq 5$  and  $p \neq 5k \pm 1$  for some  $k > 0$ ;
- (d)  $\text{PSU}_3(q)$ , where  $q = p^f$  ( $f$  is odd),  $p \neq 5$  and  $p \neq 5k \pm 1$  for some  $k > 0$ ;
- (e)  ${}^3D_4(q)$ , where  $q = p^f$  ( $f$  is odd),  $p \neq 5$  and  $p \neq 5k \pm 1$  for some  $k > 0$ ;
- (f)  ${}^2G_2(3^{2n+1})$ , where  $n$  is a natural number.

PROOF: See [15, Lemma 2.5] or [10]. □

**Definition 2.1.** A finite group  $G$  is a *Frobenius group* if it has a proper nontrivial subgroup  $H$  such that  $H \cap H^g = 1$  for all  $g \in G - H$ . The subgroup  $H$  with these properties is called a *Frobenius complement* of  $G$ . The *Frobenius kernel* of  $G$ , with respect to  $H$ , is defined by  $K = (G - \bigcup_{g \in G} H^g) \cup \{1\}$ . A group  $G$  is a *2-Frobenius group* if there exists a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $K$  and  $G/H$  are Frobenius groups with kernels  $H$  and  $K/H$ , respectively.

We quote some known results about Frobenius group and 2-Frobenius group which are useful in the sequel.

**Lemma 2.4** ([5]). *Let  $G$  be a 2-Frobenius group of even order, i.e.,  $G$  is a finite group and has a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $K$  and  $G/H$  are Frobenius groups with kernels  $H$  and  $K/H$ , respectively. Then:*

- (a)  $t(G) = 2, \pi_1 = \pi(G/K) \cup \pi(H)$  and  $\pi_2 = \pi(K/H)$ ;
- (b)  $G/K$  and  $K/H$  are cyclic,  $|G/K| \mid (|K/H| - 1), (|G/K|, |K/H|) = 1$  and  $G/K \leq \text{Aut}(K/H)$ .

**Lemma 2.5** ([5]). *Suppose that  $G$  is a Frobenius group of even order and  $H, K$  are the Frobenius kernel and the Frobenius complement of  $G$ , respectively. Then  $t(G) = 2$ , and the prime graph components of  $G$  are  $\pi(H)$  and  $\pi(K)$ .*

**Lemma 2.6** ([13, Theorem A]). *If  $G$  is a finite group such that  $t(G) \geq 2$ , then  $G$  has one of the following structures:*

- (a)  $G$  is a Frobenius group or a 2-Frobenius group;
- (b)  $G$  has a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $\pi(H) \cup \pi(G/K) \subseteq \pi_1$  and  $K/H$  is a non-abelian simple group. In particular,  $H$  is nilpotent,  $G/K \leq \text{Out}(K/H)$  and the odd order components of  $G$  are the odd order components of  $K/H$ .

### 3. Proof of Theorem 1.1

In this section,  $q = 3^{2n+1}$ , where  $n$  is a natural number. Now, we prove the theorem stated in the introduction.

PROOF: By the definition of the main supergraph and our assumption, we have  $|G| = |{}^2G_2(q)|$  (note that  $|{}^2G_2(q)| = q^3(q^3 + 1)(q - 1)$ , by [14, page 137]). Also, by  $\mathcal{S}({}^2G_2(q)) \cong \mathcal{S}(G)$  and the definition of the proper main supergraph, we have  $\mathcal{S}^*({}^2G_2(q)) \cong \mathcal{S}^*(G)$ .

We will show that  $q - \sqrt{3q} + 1$ ,  $q + \sqrt{3q} + 1$ ,  $q - 1$  (or  $q + 1$ ) are all mutually coprime. Let  $r \mid (q - \sqrt{3q} + 1)$  and  $r \mid (q - 1)$ , where  $r$  is a prime number. Since  $r \mid (q - \sqrt{3q} + 1)$ , we have  $r \mid (q - \sqrt{3q} + 1)(q + \sqrt{3q} + 1) = q^2 - q + 1 = q^2 - (q - 1)$ . On the other hand,  $r \mid (q - 1)$ . It follows that  $r \mid q^2$ , which is a contradiction. Similarly, if  $r \mid (q + \sqrt{3q} + 1)$  and  $r \mid (q - 1)$ , then we get a contradiction.

Now, let  $r \mid (q - \sqrt{3q} + 1)$  and  $r \mid (q + 1)$ , where  $r$  is a prime number. Since  $r \mid (q + \sqrt{3q} + 1)$ , we have  $r \mid (q + \sqrt{3q} + 1)(q - \sqrt{3q} + 1) = q^2 - q + 1 = q^2 + 2 - (q + 1)$ . Therefore,  $r \mid (q^2 + 2)$ . On the other hand,  $r \mid (q + 1)$ . It follows that  $r \mid (q^2 + q)$ . Since  $r \mid (q^2 + 2)$  and  $r \mid (q^2 + q)$ , we have  $r \mid (q - 2)$ . Hence,  $r \mid 3$ . Because  $q = 3^{2n+1}$  and  $r \mid (q + 1)$ , we get a contradiction. Similarly, if  $r \mid (q + \sqrt{3q} + 1)$  and  $r \mid (q - 1)$ , then we get a contradiction.

By Lemma 2.2,  $\mu({}^2G_2(q)) = \{6, 9, q - 1, (q + 1)/2, q \pm \sqrt{3q} + 1\}$ . Thus  ${}^2G_2(q)$  has not any element of order  $rp$ , where  $r \in \pi(q^3(q^2 - 1)(q + \sqrt{3q} + 1))$  and  $p \in \pi(q - \sqrt{3q} + 1)$ . Also it has not any element of order  $rp$ , where  $r \in \pi(q^3(q^2 - 1)(q - \sqrt{3q} + 1))$  and  $p \in \pi(q + \sqrt{3q} + 1)$ . It follows that  $\mathcal{S}^*(G)$  is a disconnected graph with three connected components. We denote them by  $T_+$ ,  $T_-$  and  $T_0$  such that the vertices of  $T_+$  are elements  $x \in G$  with  $o(x) \mid (q + \sqrt{3q} + 1)$ , the vertices of  $T_-$  are elements  $x \in G$  with  $o(x) \mid (q - \sqrt{3q} + 1)$  and the vertices of  $T_0$  are elements  $x \in G$  with  $o(x) \mid q^3(q^2 - 1)$ .

Let  $x$  be an arbitrary vertex of  $T_+$  such that  $o(x) = r$ , where  $r$  is a prime and let  $y$  be an arbitrary vertex of  $T_-$  such that  $o(y) = s$ , where  $s$  is a prime. If  $rs \in \pi_e(G)$ , then there exists  $z \in G$  such that  $o(z) = rs$ . By definition of  $\mathcal{S}^*(G)$ , we have  $x \sim z$  and  $y \sim z$ . Thus  $T_+$  and  $T_-$  are connected in  $\mathcal{S}^*(G)$ , a contradiction. It follows that  $rs \notin \pi_e(G)$ . Therefore,  $r$  and  $s$  are not joined by an edge in prime graph  $G$ . Similarly, we can prove it for  $T_+$  and  $T_0$  and also for  $T_-$  and  $T_0$ . Thus  $\Gamma(G)$  has at least three connected components.

Since  $t(G) \geq 3$ , Lemmas 2.4 (a) and 2.5 show that  $G$  is neither a Frobenius group nor a 2-Frobenius group. By Lemma 2.6,  $G$  has a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $H$  and  $G/K$  are  $\pi_1$ -groups and  $K/H$  is a non-abelian simple group,  $|G/K|$  divides  $|\text{Out}(K/H)|$ .

By Lemma 2.3,  $|G| = |{}^2G_2(q)|$  is coprime to 5. Again by Lemma 2.3, since  $|K/H| \mid |G|$ ,  $K/H$  is isomorphic to one the following groups:  $\text{PSL}_2(f)$ ,  $\text{PSL}_3(f)$ ,  $\text{PSU}_3(f)$ ,  $G_2(f)$ ,  ${}^3D_4(f)$ , where  $f \equiv \pm 2 \pmod{5}$  ( $f$  is a power of prime  $p$ ) and  ${}^2G_2(f)$ , where  $f = 3^{2m+1} \geq 27$ .

We will show that the order of a Sylow 2-subgroup of  $G$  is 8. As noted at the beginning of the proof,  $|G| = |{}^2G_2(q)| = q^3(q^3 + 1)(q - 1) = q^3(q^2 - q + 1)(q^2 - 1)$ , where  $q = 3^{2n+1}$ . Clearly,  $2 \nmid q^3(q^2 - q + 1)$ . Since  $q^2 - 1 = (q - 1)(q + 1) = (3^{2n+1} - 1)(3^{2n+1} + 1) = 8(3^{2n} + 3^{2n-1} + \dots + 1)(3^{2n} - 3^{2n-1} + 3^{2n-2} - \dots + 1)$  and  $2 \nmid (3^{2n} + 3^{2n-1} + \dots + 1)(3^{2n} - 3^{2n-1} + 3^{2n-2} - \dots + 1)$ , we have  $|P_2| = 8$ .

By [14, Sections 4.3.3, 4.6.2],  $|G_2(f)| = f^6(f+1)^2(f-1)^2(f^2-f+1)(f^2+f+1)$  and  $|{}^3D_4(f)| = f^{12}(f^8 + f^4 + 1)(f^6 - 1)(f^2 - 1)$ . Clearly, the order of a Sylow 2-subgroup of  $G_2(f)$  or  ${}^3D_4(f)$  is greater than 8. Therefore, we can rule out the cases  $G_2(f)$  and  ${}^3D_4(f)$ .

If  $K/H$  is isomorphic to  $\text{PSL}_3(f)$  or  $\text{PSU}_3(f)$ , then the order of  $K/H$  is divisible by  $(f \pm 1)(f^2 - 1)$ . When  $f$  is odd, this is always divisible by 16 and so  $f$  must be even. Thus  $K/H$  is isomorphic to one of the groups:  $\text{PSL}_2(f)$  with  $f \equiv \pm 2 \pmod{5}$ ,  $\text{PSL}_3(2^u)$  with  $u \geq 2$ ,  $\text{PSU}_3(2^u)$  with  $u \geq 2$  and  ${}^2G_2(f)$ . Since 16 divides the order of  $\text{PSL}_3(2^u)$ ,  $\text{PSU}_3(2^u)$ ,  $K/H$  is isomorphic to  $\text{PSL}_2(f)$  or  ${}^2G_2(f)$ .

Let  $K/H$  be isomorphic to  $\text{PSL}_2(f)$  and let  $f = p^m$ , where  $p$  is a prime number and  $m$  a natural number. By the above discussion,  $q \pm \sqrt{3q} + 1$  are odd order components of  $K/H$ .

If  $p = 2$ , then  $f + 1$  and  $f - 1$  are the odd order components of  $\text{PSL}_2(f)$ , so  $q + \sqrt{3q} + 1 = f + 1$  and  $q - \sqrt{3q} + 1 = f - 1$ , which is impossible.

If  $p \neq 2$ , then the odd order components of  $\text{PSL}_2(f)$  are  $f$  and  $(f \pm 1)/2$ . Thus  $q + \sqrt{3q} + 1 = f$  and  $q - \sqrt{3q} + 1 = (f + 1)/2$ , or  $q + \sqrt{3q} + 1 = f$  and  $q - \sqrt{3q} + 1 = (f - 1)/2$ .

If the latter case holds,  $q - 3\sqrt{3q} + 2 = 0$ . This equation has no solutions in positive integer. Then the former case occur in which we have that  $q - 3\sqrt{3q} = 0$ . It follows that  $q = 27$  and  $f = 37$ . Therefore,  $K/H = \text{PSL}_2(37)$ . In this case  $\mathcal{S}^*({}^2G_2(27))$  has three components such that two components are complete graphs ( $T_+$  and  $T_-$ ). We show that the vertices of  $T_+$  or  $T_-$  are elements of order  $37 = 27 + \sqrt{3 \cdot 27} + 1$ . We know that order of  $T_+$  or  $T_-$  is  $m_{37} = 1633531536$ .

First, let  $x$  and  $y$  be two vertices of  $T_+$  or  $T_-$  such that  $o(x) = r$  and  $o(y) = s$ , where  $r \neq s$  and  $r, s \in \pi(G)$ . Since  $T_+$  and  $T_-$  are complete, we have  $x \sim y$ , a contradiction. Let  $r$  be a prime and the vertices of  $T_+$  or  $T_-$  be all of  $x \in G$  such that  $o(x) = r, r^2, \dots$ , or  $r^k$  (note that  $\exp(P_r) = r^k$ ). Then with considering  $m = |P_r|$  in Lemma 2.1,  $|P_r| \mid (1 + m_r + m_{r^2} + \dots + m_{r^k}) = 1 + m_{37} = 1633531537$ . It follows that  $r = 37$ . Hence, the vertices of  $T_+$  or  $T_-$  are  $x \in G$  such that  $o(x) = 37^k$ , where  $k \geq 1$  is an integer.

Since  $|G| = 2^3 \cdot 3^9 \cdot 7 \cdot 13 \cdot 19 \cdot 37$ , we have  $37^2 \notin \pi_e(G)$ . Therefore, the vertices of  $T_+$  or  $T_-$  are all of elements of order 37 in  $G$ . Therefore,  $G$  has not any element of order  $37r$ , where  $r \in \pi(G)$ .

By Lemma 2.6 (b),  $|G/K|$  divides  $|\text{Out}(K/H)| = |\text{Out}(\text{PSL}_2(37))| = 2$ . Since  $|K/H| = |\text{PSL}_2(37)| = 2^2 \cdot 3^2 \cdot 19 \cdot 37$ ,  $|G| = |{}^2G_2(27)| = 2^3 \cdot 3^9 \cdot 7 \cdot 13 \cdot 19 \cdot 37$  and  $|G| = |G/K| \cdot |K/H| \cdot |H|$ , we have  $|H| = 3^7 \cdot 7 \cdot 13$  or  $2 \cdot 3^7 \cdot 7 \cdot 13$ . Thus  $|H| \mid 2 \cdot 3^7 \cdot 7 \cdot 13$ . Since  $H \trianglelefteq G$ , we have  $n_{13}(H) = n_{13}(G) = m_{13}(G)/12$ . Since  $G$  has not any element of order  $37 \cdot 13$ ,  $P_{37}$  acts fixed point freely on the elements of order 13. Thus  $37 = |P_{37}| \mid m_{13}(G) = m_{13}(H)$ . By Sylow's theorem  $n_{13}(H) \mid |H|$ . This implies that  $2^2 \cdot 3^9 \cdot 7 \cdot 19 \cdot 37 \leq m_{13}(G) = m_{13}(H) < 2 \cdot 3^7 \cdot 7 \cdot 13$ , which is a contradiction.

By the above discussion,  $K/H$  is isomorphic to  ${}^2G_2(f)$ , where  $f = 3^{2m+1}$  and  $m$  is a natural number. Hence,  $t(K/H) = 3$  and  $f \pm \sqrt{3f} + 1$  are odd order components of  $K/H$  (see [12], Table Id). On the other hand,  $q \pm \sqrt{3q} + 1$  are also

the odd order components of  $K/H$ . This implies that  $q \pm \sqrt{3q} + 1 = f \pm \sqrt{3f} + 1$ . Consequently,  $f = q$ . Therefore,  $K/H \cong {}^2G_2(q)$ . Since  $|G| = |K/H| = |{}^2G_2(q)|$ , we deduce that  $G \cong {}^2G_2(q)$ .  $\square$

## REFERENCES

- [1] Asboei A. K., Amiri S. S. S., *Some alternating and symmetric groups and related graphs*, Beitr. Algebra Geom. **59** (2018), no. 1, 21–24.
- [2] Asboei A. K., Amiri S. S. S., *Some results on the main supergraph of finite groups*, accepted in Algebra Discrete Math.
- [3] Cameron P. J., *The power graph of a finite group II*, J. Group Theory **13** (2010), no. 6, 779–783.
- [4] Chakrabarty I., Ghosh S., Sen M. K., *Undirected power graphs of semigroups*, Semigroup Forum **78** (2009), no. 3, 410–426.
- [5] Chen G. Y., *On structure of Frobenius group and 2-Frobenius group*, J. Southwest China Normal Univ. **20** (1995), no. 5, 485–487 (Chinese).
- [6] Ebrahimzadeh B., Iranmanesh A., Parvizi Mosaed H., *A new characterization of Ree group  ${}^2G_2(q)$  by the order of group and the number of elements*, Int. J. Group Theory **6** (2017), no. 4, 1–6.
- [7] Frobenius G., *Verallgemeinerung des Sylow'schen Satzes*, Berl. Ber. (1895), 981–993 (German).
- [8] Hamzeh A., Ashrafi A. R., *Automorphism groups of supergraphs of the power graph of a finite group*, European J. Combin. **60** (2017), 82–88.
- [9] Mazurov V. D., Khukhro E. I., *Unsolved Problems in Group Theory*, Kourvka Notebook, Novosibirsk, Inst. Mat. Sibirsk. Otdel. Akad., 2006.
- [10] Shi W.-J., *A characterization of  $U_3(2^n)$  by their element orders*, Xinan Shifan Daxue Xuebao Ziran Kexue Ban **25** (2000), no. 4, 353–360.
- [11] Ward H. N., *On Ree's series of simple groups*, Trans. Amer. Math. Soc. **121** (1966), 62–89.
- [12] Weisner L., *On the number of elements of a group which have a power in a given conjugate set*, Bull. Amer. Math. Soc. **31** (1925), no. 9–10, 492–496.
- [13] Williams J. S., *Prime graph components of finite groups*, J. Algebra **69** (1981), no. 2, 487–513.
- [14] Wilson R. A., *The Finite Simple Groups*, Graduate Texts in Mathematics, 251, Springer, London, 2009.
- [15] Zhang Q., Shi W., Shen R., *Quasirecognition by prime graph of the simple groups  $G_2(q)$  and  ${}^2B_2(q)$* , J. Algebra Appl. **10** (2011), no. 2, 309–317.

A. K. Asboei:

DEPARTMENT OF MATHEMATICS, FARHANGIAN UNIVERSITY, TARBAT-E-MOALLEM ST, FARAHZADI BLV, 1998963341, TEHRAN, IRAN

*E-mail:* a.khalili@cfu.ac.ir

S. S. S. Amiri:

DEPARTMENT OF MATHEMATICS, BABOL BRANCH, ISLAMIC AZAD UNIVERSITY, 47471-37381, BABOL, IRAN

*E-mail:* salehiss@baboliau.ac.ir

(Received March 17, 2018, revised May 20, 2018)