

John Donnelly

The group ring $\mathbb{K}F$ of Richard Thompson's Group F has no minimal non-zero ideals

Archivum Mathematicum, Vol. 55 (2019), No. 1, 23–30

Persistent URL: <http://dml.cz/dmlcz/147647>

Terms of use:

© Masaryk University, 2019

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

**THE GROUP RING $\mathbb{K}F$ OF RICHARD THOMPSON'S
GROUP F HAS NO MINIMAL NON-ZERO IDEALS**

JOHN DONNELLY

ABSTRACT. We use a total order on Thompson's group F to show that the group ring $\mathbb{K}F$ has no minimal non-zero ideals.

1. INTRODUCTION

We define Richard Thompson's group F to be the group of right fractions of the monoid P which is given by the presentation

$$\langle x_0, x_1, x_2, \dots \mid x_n x_m = x_m x_{n+1} \text{ for } n > m \rangle.$$

Geoghegan has conjectured that the group F is an example of a finitely presented, nonamenable group which has no free subgroup on two generators [2]. In [1], Brin and Squier show that the group F has no free subgroup on two generators. However, the question of whether or not the group F is amenable has been open for over twenty years [2].

Let \mathbb{K} denote a field. It is shown in [1] that the group F is totally ordered. Using this fact we can show that the group ring $\mathbb{K}F$ is cancellative, and consequently does not have any zero-divisors. Thus, the set of all nonzero elements in $\mathbb{K}F$ forms a multiplicative monoid \mathcal{H} whose identity is the identity 1_F of the group F . We leave it to the reader to check that if \mathcal{H} is (left/right) amenable, then the group F is amenable.

Thus, one can ask whether or not the multiplicative monoid \mathcal{H} is right amenable. In [3], Frey gives necessary conditions that any minimal ideal of a semigroup S must satisfy for S to be right amenable. In particular, Frey shows that if S is a right amenable semigroup, \mathcal{L} is a minimal left ideal of S , and \mathcal{R} is a minimal right ideal of S , then

- (i) \mathcal{L} is a two-sided ideal of S .
- (ii) $\mathcal{R} \subseteq \mathcal{L}$.
- (iii) \mathcal{R} is a group.
- (iv) There exists a semigroup T such that \mathcal{L} is isomorphic to $\mathcal{R} \oplus T$, and such that for all $z_1, z_2 \in T$, $z_1 z_2 = z_1$.

2010 *Mathematics Subject Classification*: primary 20N99.

Key words and phrases: Thompson's Group F , amenability, minimal ideal.

Received December 4, 2016, revised November 2018. Editor J. Trlifaj.

DOI: 10.5817/AM2019-1-23

Frey also shows that if S is a semigroup containing a minimal left ideal \mathcal{L} and a minimal right ideal \mathcal{R} , then S is right amenable if and only if \mathcal{R} is an amenable group.

Thus, one can ask what the minimal ideals of \mathcal{H} are, and whether or not they satisfy the conditions stated above. In this paper, we use a total ordering on the group F to show that \mathcal{H} has no minimal left, right, or two-sided ideals.

2. A TOTAL ORDERING ON THE GROUP F

We denote the set of generators $\{x_0, x_1, x_2, \dots, x_n, \dots\}$ of P (and consequently, of F) by Σ , and we define the set $\Sigma_n = \{x_m \in \Sigma \mid m \geq n\}$. Given an element $q \in P$, we let $|q|$ denote the length of a word over Σ representing q . Every element of the group F can be represented uniquely by a normal form

$$x_{i_1}^{b_1} x_{i_2}^{b_2} x_{i_3}^{b_3} \dots x_{i_m}^{b_m} x_{j_k}^{-d_k} \dots x_{j_3}^{-d_3} x_{j_2}^{-d_2} x_{j_1}^{-d_1}$$

where

- (i) for each t , and for each r , we have that $b_t, d_r > 0$;
- (ii) $i_1 < i_2 < \dots < i_m$ and $j_1 < j_2 < \dots < j_k$;
- (iii) if there exists some i such that both x_i and x_i^{-1} are generators in the normal form, then x_{i+1} or x_{i+1}^{-1} is a generator in the normal form as well.

Given two generators x_i and x_j of P , then we define $x_i < x_j$ if and only if $i < j$. We can now use the shortlex ordering on the set of normal forms for the elements of the monoid P to get a total ordering $<_P$ on the monoid P . We use the ordering $<_P$ on P to define an ordering $<_F$ on all of the group F in the following way: Given $g \in F$ such that g has normal form xy^{-1} , with $x, y \in P$, then $g <_F 1_F$ if and only if $x <_P y$. We extend this to compare all elements of the group F by defining for each distinct pair $g, h \in F$ that $g <_F h$ if and only if $gh^{-1} <_F 1_F$. We will prove that $<_F$ is a well defined total ordering on the group F .

Let $g, h \in F$. Assume that gh^{-1} has normal form ab^{-1} , where $a, b \in P$. Since ab^{-1} is in normal form, then ba^{-1} is in normal form. Moreover, since $hg^{-1} = (gh^{-1})^{-1} = (ab^{-1})^{-1} = ba^{-1}$, then hg^{-1} has normal form ba^{-1} . Note that in case (i) below, since $a = b$ and ab^{-1} is in normal form, then a and b are empty words and consequently ab^{-1} is the identity element of F . Therefore, if gh^{-1} has normal form ab^{-1} , where $a, b \in P$, then

- (i) $g = h$ if and only if $a = b$;
- (ii) $gh^{-1} <_F 1_F$ if and only if $a <_P b$;
- (iii) $hg^{-1} <_F 1_F$ if and only if $b <_P a$.

Since for each pair of elements $a, b \in P$, exactly one of $a = b$, $a <_P b$, or $b <_P a$ must hold, then given two elements $g, h \in F$, exactly one of $g = h$, $gh^{-1} <_F 1_F$, or $hg^{-1} <_F 1_F$ must hold. Thus, given two distinct elements $g, h \in F$, then either $gh^{-1} <_F 1_F$, in which case $g <_F h$, or else $hg^{-1} <_F 1_F$, in which case $h <_F g$.

Thus, it follows that $<_F$ is well defined and linear.

Lemma 1. *Let $w_1, w_2 \in P$ be such that $w_1 <_P w_2$. If x_m is any generator of the monoid P , then $x_m w_1 <_P x_m w_2$.*

Proof. Let $|w_1| = h$ and $|w_2| = k$. If $|w_1| < |w_2|$, then we see that $|x_m w_1| = h + 1 < k + 1 = |x_m w_2|$, which implies that $x_m w_1 <_P x_m w_2$.

Assume that $|w_1| = k = |w_2|$. Let w_1 have normal form $v x_{a_1} x_{a_2} \dots x_{a_t}$, and let w_2 have normal form $v x_{b_1} x_{b_2} \dots x_{b_t}$, where v is a (possibly empty) word over Σ , and $a_1 < b_1$.

Assume that $k = 1$. In this case, v is empty, $w_1 = x_{a_1}$, and $w_2 = x_{b_1}$, with $a_1 < b_1$. First assume that $m \leq a_1 < b_1$. In this case, $x_m w_1$ has normal form $x_m x_{a_1}$ and $x_m w_2$ has normal form $x_m x_{b_1}$. Since $a_1 < b_1$, then $x_m w_1 <_P x_m w_2$. Next assume that $a_1 < m \leq b_1$. In this case $x_m w_1$ has normal form $x_{a_1} x_{m+1}$, and $x_m w_2$ has normal form $x_m x_{b_1}$. Since $a_1 < m$, then $x_m w_1 <_P x_m w_2$. Finally, assume that $a_1 < b_1 < m$. In this case $x_m w_1$ has normal form $x_{a_1} x_{m+1}$, and $x_m w_2$ has normal form $x_{b_1} x_{m+1}$. Since $a_1 < b_1$, then $x_m w_1 <_P x_m w_2$.

Now assume that $k \geq 2$, and that for each $j \in \{1, \dots, k-1\}$, if $u_1, u_2 \in P$ are such that $|u_1| = |u_2| = j$ and $u_1 <_P u_2$, then for each generator x_m of P , $x_m u_1 <_P x_m u_2$.

Assume that $|v| \geq 1$, and that $x_m v = v x_{m+|v|}$. Since $a_1 < b_1$, then it follows that $x_{a_1} x_{a_2} \dots x_{a_t} <_P x_{b_1} x_{b_2} \dots x_{b_t}$. Therefore, by our induction hypothesis we have that $x_{m+|v|} x_{a_1} x_{a_2} \dots x_{a_t} <_P x_{m+|v|} x_{b_1} x_{b_2} \dots x_{b_t}$. Thus, $x_{m+|v|} x_{a_1} x_{a_2} \dots x_{a_t}$ has normal form $\sigma x_{i_1} x_{i_2} \dots x_{i_q}$. Similarly, we see that $x_{m+|v|} x_{b_1} x_{b_2} \dots x_{b_t}$ has normal form $\sigma x_{j_1} x_{j_2} \dots x_{j_q}$, where σ is a (possibly empty) word over Σ , and $i_1 < j_1$. Therefore, $x_m w_1$ has normal form $v \sigma x_{i_1} x_{i_2} \dots x_{i_q}$, and $x_m w_2$ has normal form $v \sigma x_{j_1} x_{j_2} \dots x_{j_q}$. Since $i_1 < j_1$, then $x_m w_1 <_P x_m w_2$.

Now Assume that $|v| \geq 1$, and that $x_m v = u x_{m+|u|} z$, where z is some nonempty word over $\Sigma_{m+|u|}$, and where u is some (possibly empty) word over Σ . In this case, $x_m w_1$ has normal form $u x_{m+|u|} z x_{a_1} x_{a_2} \dots x_{a_t}$, and $x_m w_2$ has normal form $u x_{m+|u|} z x_{b_1} x_{b_2} \dots x_{b_t}$. Since $a_1 < b_1$, then it follows that $x_m w_1 <_P x_m w_2$.

Finally, assume that v is empty. In this case, w_1 has normal form $x_{a_1} x_{a_2} \dots x_{a_k}$, and w_2 has normal form $x_{b_1} x_{b_2} \dots x_{b_k}$, where $a_1 < b_1$. First assume that $m \leq a_1 < b_1$. In this case, $x_m w_1$ has normal form $x_m x_{a_1} x_{a_2} \dots x_{a_k}$, and $x_m w_2$ has normal form $x_m x_{b_1} x_{b_2} \dots x_{b_k}$. Since $a_1 < b_1$, then $x_m w_1 <_P x_m w_2$. Next assume that $a_1 < m \leq b_1$. In this case, $x_m w_1$ has normal form $x_{a_1} \beta$, where β is a word over Σ_{a_1} of length k , and $x_m w_2$ has normal form $x_m x_{b_1} x_{b_2} \dots x_{b_k}$. Since $a_1 < m$, then $x_m w_1 <_P x_m w_2$. Finally, assume that $a_1 < b_1 < m$. In this case, $x_m w_1$ has normal form $x_{a_1} \rho_1$, where ρ_1 is a word over Σ_{a_1} of length k , and $x_m w_2$ has normal form $x_{b_1} \rho_2$, where ρ_2 is a word over Σ_{b_1} of length k . Since $a_1 < b_1$, then $x_m w_1 <_P x_m w_2$. \square

Lemma 2. *Let $w_1, w_2 \in P$ be such that $w_1 <_P w_2$. If x_m is any generator of the monoid P , then $w_1 x_m <_P w_2 x_m$.*

Proof. Let $|w_1| = h$ and $|w_2| = k$. If $|w_1| < |w_2|$, then we see that $|w_1 x_m| = h + 1 < k + 1 = |w_2 x_m|$, which implies that $w_1 x_m <_P w_2 x_m$.

Assume that $|w_1| = k = |w_2|$. Let w_1 have normal form $x_{a_1} x_{a_2} \dots x_{a_k}$, and let w_2 have normal form $x_{b_1} x_{b_2} \dots x_{b_k}$. First assume that $k = 1$. In this case, $w_1 = x_{a_1}$ and $w_2 = x_{b_1}$, with $a_1 < b_1$. If $m < a_1 < b_1$, then $w_1 x_m$ has normal form $x_m x_{a_1+1}$, and $w_2 x_m$ has normal form $x_m x_{b_1+1}$, which implies that $w_1 x_m <_P w_2 x_m$. Assume

that $a_1 \leq m < b_1$. Thus, w_1x_m has normal form $x_{a_1}x_m$, and w_2x_m has normal form $x_mx_{b_1+1}$. If $a_1 < m$, then $w_1x_m = x_{a_1}x_m <_P x_mx_{b_1+1} = w_2x_m$. If $a_1 = m$, then $w_1x_m = x_mx_m <_P x_mx_{b_1+1} = w_2x_m$. Assume that $a_1 < b_1 \leq m$. In this case, we see that w_1x_m has normal form $x_{a_1}x_m$, and w_2x_m has normal form $x_{b_1}x_m$, which implies that $w_1x_m = x_{a_1}x_m <_P x_{b_1}x_m = w_2x_m$.

Now assume that $k \geq 2$, and that for each $j \in \{1, \dots, k-1\}$, if $u_1, u_2 \in P$ are such that $|u_1| = |u_2| = j$ and $u_1 <_P u_2$, then for each generator x_m of P , $u_1x_m <_P u_2x_m$.

Assume that $a_1 = b_1 \leq m$. In this case, w_1 has normal form $x_{a_1}\sigma_1$, and w_2 has normal form $x_{a_1}\sigma_2$, where σ_1 and σ_2 are words over Σ_{a_1} . Since $x_{a_1}\sigma_1 = w_1 <_P w_2 = x_{a_1}\sigma_2$, then it must be the case that $\sigma_1 <_P \sigma_2$. Therefore, it follows by our induction hypothesis that $\sigma_1x_m <_P \sigma_2x_m$. Thus, σ_1x_m has normal form $vx_{c_1}x_{c_2} \dots x_{c_t}$, and σ_2x_m has normal form $vx_{e_1}x_{e_2} \dots x_{e_t}$, where v is a word over Σ_{a_1} , and $c_1 < e_1$. Therefore, $x_{a_1}\sigma_1x_m$ has normal form $x_{a_1}vx_{c_1}x_{c_2} \dots x_{c_t}$, and $x_{a_1}\sigma_2x_m$ has normal form $x_{a_1}vx_{e_1}x_{e_2} \dots x_{e_t}$, where $c_1 < e_1$. Since $w_1x_m = x_{a_1}\sigma_1x_m$ and $w_2x_m = x_{a_1}\sigma_2x_m$, then it follows that $w_1x_m <_P w_2x_m$.

Now assume that $a_1 < b_1 \leq m$. In this case w_1x_m has normal form $x_{a_1}\beta_1$, where β_1 is a word over Σ_{a_1} , and w_2x_m has normal form $x_{b_1}\beta_2$, where β_2 is a word over Σ_{b_1} . Thus, $w_1x_m <_P w_2x_m$.

Assume that $a_1 = m < b_1$. In this case, w_1x_m has normal form $x_mx_m\rho$, where ρ is a word over Σ_m , and w_2x_m has normal form $x_mx_{b_1+1}x_{b_2+1} \dots x_{b_k+1}$. Since $m < b_1 < b_1 + 1$, then $w_1x_m <_P w_2x_m$.

Assume that $a_1 < m < b_1$. In this case, w_1x_m has normal form $x_{a_1}\alpha$, where α is a word over Σ_{a_1} , and w_2x_m has normal form $x_mx_{b_1+1}x_{b_2+1} \dots x_{b_k+1}$. Since $a_1 < m$, then $w_1x_m <_P w_2x_m$.

Finally, assume that $m < a_1 < b_1$. In this case, w_1x_m has normal form $x_mx_{a_1+1}x_{a_2+1} \dots x_{a_k+1}$, and w_2x_m has normal form $x_mx_{b_1+1}x_{b_2+1} \dots x_{b_k+1}$. This implies that $w_1x_m <_P w_2x_m$. \square

Given $a, b, c \in P$, with $a <_P b$, then by using Lemmas 1 and 2 as induction base steps, one can use induction on the length $|c|$ to show that $ca <_P cb$ and $ac <_P bc$. We now extend this property to the ordering $<_F$ by showing that for all $g, h, d \in F$, if $g <_F h$, then $dg <_F dh$ and $gd <_F hd$. Again, we note that in case (i) below, since $a = b$ and ab^{-1} is in normal form, then a and b are empty words and consequently ab^{-1} is the identity element 1_P of P .

Lemma 3. *Let $a, b, c, d \in P$ be such that $ab^{-1} = cd^{-1}$, and such that ab^{-1} is in normal form. Then*

- (i) $a = b$ if and only if $c = d$;
- (ii) $a <_P b$ if and only if $c <_P d$;
- (iii) $b <_P a$ if and only if $d <_P c$.

Proof. Let 1_P denote the identity element of the monoid P . Since $a = b$ if and only if $cd^{-1} = ab^{-1} = 1_P$, and since $cd^{-1} = 1_P$ if and only if $c = d$, then $a = b$ if and only if $c = d$.

Assume that $a \neq b$. When rewriting the normal form ab^{-1} to get the word cd^{-1} , we multiply a on the right by a (possibly empty) word u over Σ , and we multiply

b^{-1} on the left by the (possibly empty) word u^{-1} over Σ^{-1} . In particular, the word u consists of all the generators from c that cancel when we multiply c with d^{-1} , and then simplify to rewrite cd^{-1} in the normal form ab^{-1} . Thus, $c = au$ and $d = bu$. If $a <_P b$, then it follows by the comments above that $c = au <_P bu = d$. If $b <_P a$, then again it follows by the comments above that $d = bu <_P au = c$.

Now assume that $c <_P d$. Since $<_P$ is a linear ordering on P , then exactly one of the following is true: $a <_P b$, $a = b$, or $b <_P a$. If $a = b$, then $c = d$, a contradiction. If $b <_P a$, then it follows from the argument given above that $d <_P c$, a contradiction. Thus, $a <_P b$. A similar argument shows that if $d <_P c$, then $b <_P a$. \square

Lemma 4. *Let $c, d \in P$. Then $c <_P d$ if and only if $c <_F d$.*

Proof. Let cd^{-1} have normal form ab^{-1} , where $a, b \in P$. Assume that $c <_P d$. Since $c <_P d$ and cd^{-1} has normal form ab^{-1} , then it follows by Lemma 3 that $a <_P b$. Thus, by definition of $<_F$, we have that $ab^{-1} <_F 1_F$, which implies that $cd^{-1} <_F 1_F$, which in turn implies that $c <_F d$.

Conversely, assume that $c <_F d$. By definition of $<_F$, we have that $cd^{-1} <_F 1_F$. Since $ab^{-1} = cd^{-1}$, then $ab^{-1} <_F 1_F$, which implies that $a <_P b$. Therefore, by Lemma 3, we have that $c <_P d$. \square

Lemma 5. *Let $g, h \in F$ be such that $g <_F h$. Then for any $d \in F$, $gd <_F hd$.*

Proof. Since $g <_F h$, then it follows by definition of $<_F$ that $gh^{-1} <_F 1_F$. Thus, we see that $(gd)(d^{-1}h^{-1}) = gh^{-1} <_F 1_F$, which implies that $gd <_F hd$. \square

Lemma 6. *Let $g \in F$ and $c, d \in P$. If $g <_F c$ and $c <_F d$, then $g <_F d$.*

Proof. Let g have normal form ab^{-1} , where $a, b \in P$. Since $ab^{-1} <_F c$, then it follows by Lemma 5 that $a <_F cb$. Since $a <_F cb$, then it follows that $a <_P cb$. Similarly, since $c <_F d$, then $c <_P d$. Since $a <_P cb$ and $c <_P d$, then $a <_P cb <_P db$. Therefore, by definition of $<_F$, we have that $ab^{-1}d^{-1} <_F 1_F$, which implies that $ab^{-1} <_F d$. Hence, $g <_F d$. \square

Lemma 7. *Let $g, h \in F$. If $g <_F h$ and $h <_F 1_F$, then $g <_F 1_F$.*

Proof. Let g have normal form ab^{-1} , where $a, b \in P$, and let h have normal form cd^{-1} , where $c, d \in P$. Since $cd^{-1} <_F 1_F$, then $c <_F d$. Since $ab^{-1} <_F cd^{-1}$, then it follows by Lemma 5 that $ab^{-1}d <_F c$. Since $ab^{-1}d <_F c$ and $c <_F d$, then it follows by Lemma 6 that $ab^{-1}d <_F d$. Thus, by Lemma 5, we have that $ab^{-1} <_F 1_F$. Hence, $g <_F 1_F$. \square

Lemma 8. *Let $g, h, d \in F$. If $g <_F h$ and $h <_F d$, then $g <_F d$.*

Proof. Since $g <_F h$, then it follows by Lemma 5 that $gd^{-1} <_F hd^{-1}$. Since $h <_F d$, then it follows that $hd^{-1} <_F 1_F$. Therefore, it follows by Lemma 7 that $gd^{-1} <_F 1_F$. Thus, it follows by definition of $<_F$ that $g <_F d$. \square

Lemma 9. *Let $g, h \in F$, and let $b \in P$. If $g <_F h$, then $bg <_F bh$.*

Proof. Let gh^{-1} have normal form cd^{-1} , where $c, d \in P$. Since $g <_F h$, then it follows that $gh^{-1} <_F 1_F$. Since gh^{-1} has normal form cd^{-1} , then it follows that $cd^{-1} <_F 1_F$. Therefore, $c <_F d$, which implies that $c <_P d$. Thus, it follows that $bc <_P bd$, which implies that $bc <_F bd$, and therefore that $bcd^{-1}b^{-1} <_F 1_F$. Again, since gh^{-1} has normal form cd^{-1} , then it follows that $bgh^{-1}b^{-1} <_F 1_F$, and therefore that $bg <_F bh$. \square

Lemma 10. *Let $g, h \in F$, and let $b \in P$. If $g <_F h$, then $b^{-1}g <_F b^{-1}h$.*

Proof. If $b^{-1}g = b^{-1}h$, then $g = h$, a contradiction. Thus, $b^{-1}g \neq b^{-1}h$. Suppose that $b^{-1}h <_F b^{-1}g$. Thus, it follows by Lemma 9 that $b(b^{-1}h) <_F b(b^{-1}g)$, which implies that $h <_F g$, a contradiction. Hence, $b^{-1}g <_F b^{-1}h$. \square

Lemma 11. *Let $g, h, u \in F$. If $g <_F h$, then $ug <_F uh$.*

Proof. Let u have normal form ab^{-1} , where $a, b \in P$. Since $g <_F h$, then it follows by Lemma 10 that $b^{-1}g <_F b^{-1}h$. Therefore, since $b^{-1}g <_F b^{-1}h$, then it follows by Lemma 9 that $ab^{-1}g <_F ab^{-1}h$. Hence, $ug <_F uh$. \square

Lemma 12. *Let $g_1, g_2, h_1, h_2 \in F$ be such that $g_1 <_F g_2$ and $h_1 <_F h_2$. Then it follows that $g_1h_1 <_F g_2h_2$.*

Proof. Since $g_1 <_F g_2$, then it follows by Lemma 5 that $g_1h_1 <_F g_2h_1$. Similarly, since $h_1 <_F h_2$, then it follows by Lemma 11 that $g_2h_1 <_F g_2h_2$. Therefore, since $g_1h_1 <_F g_2h_1$ and $g_2h_1 <_F g_2h_2$, then it follows by Lemma 8 that $g_1h_1 <_F g_2h_2$. \square

3. THE MAIN RESULT

Lemma 13. *Let $g_1, g_2 \in F$. Assume that g_1 has normal form $a_1b_1^{-1}$, and that g_2 has normal form $a_2b_2^{-1}$, where $a_1, a_2, b_1, b_2 \in P$. If $|a_1| + |b_2| < |a_2| + |b_1|$, then $g_1 <_F g_2$.*

Proof. Let $b_1^{-1}b_2$ have normal form cd^{-1} , where $c, d \in P$. Each generator x_i in the normal form of b_2 which cancels when multiplying b_1^{-1} and b_2 to put $b_1^{-1}b_2$ in normal form will cancel with exactly one of the generators x_j^{-1} in the normal form of b_1^{-1} . That is, any generators from b_2 and b_1^{-1} which cancel when transforming $b_1^{-1}b_2$ into normal form will cancel in pairs. Thus, if k generators cancel from the normal form of b_2 , then k generators cancel from the normal form of b_1^{-1} . Therefore, we see that $|c| = |b_2| - k$, and that $|d| = |b_1| - k$. Since $a_1, c \in P$, then there is no cancellation of generators when multiplying a_1 and c . Thus, we see that $|a_1c| = |a_1| + |c|$. Similarly, we see that $|a_2d| = |a_2| + |d|$. Therefore, we have that $|a_1c| = |a_1| + |c| = |a_1| + |b_2| - k < |a_2| + |b_1| - k = |a_2| + |d| = |a_2d|$. Since $|a_1c| < |a_2d|$, then it follows that $a_1c <_F a_2d$. Thus, $a_1b_1^{-1}b_2 = a_1cd^{-1} <_F a_2$, which implies that $a_1b_1^{-1} <_F a_2b_2^{-1}$. Hence, $g_1 <_F g_2$. \square

Theorem 1. *Let \mathcal{H} denote the multiplicative monoid of nonzero elements in the group ring $\mathbb{K}F$. Then \mathcal{H} has no minimal ideals.*

Proof. Suppose, to the contrary, that \mathcal{I} is a minimal two-sided ideal of \mathcal{H} . Since \mathcal{H} is a cancellative monoid, then \mathcal{I} is a principal ideal. Let $\hat{g} = \sum_{i=1}^m r_i g_i \in \mathcal{H}$ be such that $\mathcal{I} = \mathcal{H}\hat{g}\mathcal{H}$. By renumbering if necessary, we may assume that $g_1 <_F g_2 <_F \cdots <_F g_m$. Since $\sum_{i=1}^m r_i g_i = g_1 \sum_{i=1}^m r_i (g_1^{-1} g_i)$, then $\mathcal{I} = \mathcal{H} \sum_{i=1}^m r_i (g_1^{-1} g_i) \mathcal{H}$. In particular, we may assume that $g_1 = 1_F$. Let g_m have normal form cd^{-1} , where $c, d \in P$. Let $\mathcal{J} = \mathcal{H}(\hat{g})(1_F + c)\mathcal{H} = \mathcal{H}(\sum_{i=1}^m r_i g_i + \sum_{i=1}^m r_i (g_i c))\mathcal{H}$. Since $1_F = g_1 <_F g_m = cd^{-1}$, then $1_F \leq_F d <_F c$. Thus, 1_F and $g_m c$ are the smallest and largest elements, respectively, of F used to write $(\hat{g})(1_F + c)$ as a sum in the group ring $\mathbb{K}F$. Since $(\hat{g})(1_F + c) \in \mathcal{I}$, then \mathcal{J} is a subideal of \mathcal{I} . Since \mathcal{I} is minimal, then it must be the case that $\hat{g} \in \mathcal{I} = \mathcal{J}$. Thus, there exist $\sum_{j=1}^l s_j h_j, \sum_{k=1}^e t_k q_k \in \mathcal{H}$

such that $\hat{g} = \left(\sum_{j=1}^l s_j h_j \right) (\hat{g})(1_F + c) \left(\sum_{k=1}^e t_k q_k \right)$. Again, by renumbering if necessary,

we may assume that $h_1 \leq_F h_2 \leq_F \cdots \leq_F h_l$ and $q_1 \leq_F q_2 \leq_F \cdots \leq_F q_e$. Since F is totally ordered, and since 1_F and g_m are the smallest and largest elements, respectively, of F used to write \hat{g} as a sum in the group ring $\mathbb{K}F$, then it follows that $h_1 q_1 = 1_F$, and $h_l g_m c q_e = g_m$. This implies that $h_1 = q_1^{-1}$.

First assume that $h_1 = q_1 = 1_F$. Since $1_F <_F c$, $1_F = q_1 \leq_F q_e$, and $1_F = h_1 \leq_F h_l$, then $g_m <_F h_l g_m c q_e$, a contradiction. Therefore, $q_1 \neq 1_F$. In this case we have that $1_F <_F q_1$ or $1_F <_F h_1$. We may assume that $1_F <_F q_1$ (the proof for the case that $1_F <_F h_1$ is similar). Since $1_F <_F q_1$, then $h_1 = q_1^{-1} <_F 1_F$. Let h_l have normal form ab^{-1} , and let q_e have normal form yx^{-1} , where $a, b, x, y \in P$. Since $1_F <_F q_1 \leq_F q_e$, then $|x| \leq |y|$. Similarly, since $1_F <_F c$, then $|c| \geq 1$. Since $h_l g_m c q_e = g_m$, then $h_l = g_m q_e^{-1} c^{-1} g_m^{-1} = (cd^{-1})(xy^{-1})(c^{-1})(dc^{-1})$. As in the proof of Lemma 13, when transforming h_l into normal form, any generators which cancel must cancel in pairs. Thus, if k generators cancel in c, x , or d , then k generators must cancel in d^{-1}, y^{-1} , or either of the copies of c^{-1} . Thus, $|a| = |c| + |x| + |d| - k$, and $|b| = |d| + |y| + 2|c| - k$. Therefore, we see that $|a| + |y| = (|c| + |x| + |d| - k) + |y| < |d| + |y| + 2|c| + |x| - k = |b| + |x|$. It follows by Lemma 13 that $h_l = ab^{-1} <_F xy^{-1} = q_e^{-1}$. Thus, it follows that $h_l <_F q_e^{-1} \leq_F q_1^{-1} = h_1$, which is a contradiction. Hence, \mathcal{H} does not have a minimal two-sided ideal.

A similar argument shows that \mathcal{H} has neither a minimal left ideal nor a minimal right ideal. \square

REFERENCES

- [1] Brin, M., Squier, C., *Groups of piecewise linear homeomorphisms of the real line*, Invent. Math. **79** (3) (1985), 485–498.
- [2] Cannon, J.W., Floyd, W.J., Parry, W.R., *Introductory notes on Richard Thompson's groups*, Enseign. Math. **42** 3–4) (1996), 215–256.
- [3] Frey, A.H., *Studies on Amenable Semigroups*, Ph.D. thesis, University of Washington, 1960.

DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF SOUTHERN INDIANA,
8600 UNIVERSITY BOULEVARD,
EVANSVILLE, INDIANA 47712
E-mail: jrdonnelly@usi.edu