

Bikram Banerjee

Chern rank of complex bundle

Commentationes Mathematicae Universitatis Carolinae, Vol. 60 (2019), No. 3, 401–413

Persistent URL: <http://dml.cz/dmlcz/147858>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2019

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Chern rank of complex bundle

BIKRAM BANERJEE

Abstract. Motivated by the work of A. C. Naolekar and A. S. Thakur (2014) we introduce notions of upper chern rank and even cup length of a finite connected CW-complex and prove that upper chern rank is a homotopy invariant. It turns out that determination of upper chern rank of a space X sometimes helps to detect whether a generator of the top cohomology group can be realized as Euler class for some real (orientable) vector bundle over X or not. For a closed connected d -dimensional complex manifold we obtain an upper bound of its even cup length. For a finite connected even dimensional CW-complex with its upper chern rank equal to its dimension, we provide a method of computing its even cup length. Finally, we compute upper chern rank of many interesting spaces.

Keywords: Chern class; characteristic rank; cup length; chern rank

Classification: 57R20

1. Introduction

In [3] J. Korbaš introduced the idea of *characteristic rank* of a smooth closed connected manifold X of dimension d . He defined *characteristic rank* of a d -dimensional smooth closed connected manifold X as the largest integer k such that every cohomology class of $H^i(X; \mathbb{Z}_2)$, $i \leq k$, can be expressed as a polynomial of the Stiefel–Whitney classes of the tangent bundle of X . In the same paper [3] J. Korbaš also used *characteristic rank* to get a bound for \mathbb{Z}_2 -cup length of a manifold X . The \mathbb{Z}_2 -cup length, denoted by $\text{Cup}(X)$ of a space X is defined to be the largest integer t such that there exist cohomology classes $x_i \in H^*(X; \mathbb{Z}_2)$, $\deg(x_i) \geq 1$, so that the cup product $x_1 x_2 \cdots x_t \neq 0$. Later in 2014, A. C. Naolekar and A. S. Thakur in [7] generalized the notion of *characteristic rank* to a real vector bundle ξ over a finite connected CW-complex X . If ξ is a real n -plane bundle over X then they defined *characteristic rank* (briefly $\text{char rank } \xi$) of ξ over X to be the largest integer k such that every cohomology class $x \in H^i(X; \mathbb{Z}_2)$, $i \leq k \leq \dim X$, can be expressed as a polynomial of Stiefel–Whitney classes of ξ . They also defined *upper characteristic rank* (X) of a finite connected CW-complex X as the maximum of $\text{char rank } \xi$ as ξ varies over all real vector bundles over X and thus by naturality of Stiefel–Whitney classes *upper characteristic rank* becomes a homotopy invariant. In [7] *characteristic rank* of real vector bundles over product of spheres $S^m \times S^n$, the real and complex projective spaces, the

spaces $S^1 \times \mathbb{C}P^n$, the Dold manifold $P(m, n)$, the Moore space $M(\mathbb{Z}_2, n)$ and the stunted projective space $\mathbb{R}P^n/\mathbb{R}P^m$ were computed. Moreover, some general facts about *characteristic rank* of real vector bundles were also proved.

This motivates us to define *chern rank* of a complex vector bundle over X . Throughout, by a (topological) space we mean a finite connected CW-complex and $H^*(X)$ ($\tilde{H}^*(X)$) denotes the graded (reduced) integral cohomology ring of X . We begin with the following definition.

Definition 1.1. Let ξ be a complex n -plane bundle over a finite connected CW-complex X . By *chern rank* ξ we mean the largest even integer $2k$, where $0 \leq 2k \leq \dim X$, such that every cohomology class $x \in H^{2i}(X)$, $i \leq k$, can be expressed as a polynomial of Chern classes of ξ . The upper chern rank (X) (in brief *uch rank* (X)) is defined to be the maximum chern rank ξ where ξ varies over all complex vector bundles over X , that is,

$$\text{uch rank}(X) = \max\{\text{chern rank } \xi : \xi \text{ is a complex vector bundle over } X\}.$$

From the naturality of Chern classes it follows that if X and Y are homotopy equivalent then $\text{uch rank}(X) = \text{uch rank}(Y)$. We note that determining *upper chern rank* of a topological space X sometimes helps to detect whether a generator of the top cohomology group can be realized as Euler class for some real (orientable) vector bundle over X or not. If for a $2n$ dimensional closed connected smooth manifold X the only nontrivial even dimensional reduced cohomology group is $\tilde{H}^{2n}(X)$ and $\text{uch rank}(X) = 2n$ then clearly a generator of $\tilde{H}^{2n}(X)$ can be realized as Euler class for some real (orientable) vector bundle over X . For example we will see that $\text{uch rank}(S^1 \times S^3) = 4$ (cf. Corollary 3.2) and consequently a generator of $\tilde{H}^4(S^1 \times S^3)$ can be realized as an Euler class of some real (orientable) vector bundle over $S^1 \times S^3$. Also for a finite connected CW-complex consisting of only even dimensional cells *upper chern rank* of X gives a lower bound for *upper characteristic rank* of X (cf. Lemma 2.2).

If X is a finite connected CW-complex, we denote by r_X the smallest even integer such that $\tilde{H}^{r_X}(X) \neq 0$. For X is a CW-complex with $\tilde{H}^{2i}(X) = 0 \ \forall i$, we define $r_X = \dim X + 2$ if X is even dimensional and $r_X = \dim X + 1$ otherwise. Clearly, for any complex vector bundle ξ over X ,

$$r_X - 2 \leq \text{chern rank } \xi \leq \text{uch rank}(X).$$

For a finite connected CW-complex X we define the even cup length (denoted by $\text{Cup}_E(X)$) of X to be the largest integer t such that the cup product $x_1 \cdot x_2 \cdots x_t \neq 0$ where each $x_i \in H^*(X)$ is of even degree and $\deg(x_i) \geq 2$. If X consists of only even dimensional cells then clearly $1 + \text{Cup}_E(X)$ is a suitable lower bound of $\text{Cat}(X)$ where $\text{Cat}(X)$ denotes the *Lyusternik-Shnirel'man* category. For a closed connected d -dimensional complex manifold we obtain a bound for $\text{Cup}_E(X)$ using chern rank. In particular we prove the following theorem.

Theorem 1.2. *Let X be a closed connected d -dimensional complex manifold such that $H^{2i}(X)$ is a free \mathbb{Z} -module for all i . If ξ is a complex vector bundle over X and there exists some nonzero even integer $2k \leq \text{chern rank } \xi$ such that every monomial $c_{i_1}(\xi) \cdots c_{i_r}(\xi)$, $1 \leq i_t \leq k$, of total degree $2d$ is zero then*

$$\text{Cup}_E(X) \leq 1 + \frac{2(d - k - 1)}{r_X}.$$

If X is a finite connected even dimensional CW-complex with

$$\text{uch rank}(X) = \dim X$$

then the following theorem tells us that $\text{Cup}_E(X)$ can be computed as the maximal length of nonzero product of Chern classes of a suitable complex vector bundle ξ over X .

Theorem 1.3. *Let X be an even dimensional finite connected CW-complex. If $\text{uch rank}(X) = \dim X$ then there exists a complex vector bundle ξ such that*

$$\text{Cup}_E(X) = \max\{k: \exists i_1, i_2, \dots, i_k \geq 1 \text{ with } c_{i_1}(\xi) \cdot c_{i_2}(\xi) \cdots c_{i_k}(\xi) \neq 0\}.$$

Finally, we compute uch rank of projective spaces $\mathbb{F}P^n$ (\mathbb{F} is real, complex or quaternionic). We give a full description of uch rank of product of spheres $S^m \times S^n$ where m, n are even integers and in the case where m is even and n is an odd integer. If m and n are both odd integers then we compute uch rank of $S^m \times S^n$ for some special cases. We also give computation of uch rank of X where X is wedge sum of spheres $S^m \vee S^n$, $\mathbb{R}P^n \times S^{2m}$, $\mathbb{C}P^n \times S^{2m}$, complex Stiefel manifolds $V_k(\mathbb{C}^n)$, $1 < k < n$, for $n - k$ is even or $n - k \neq 2^t - 1$, $t > 0$, and stunted complex projective spaces $\mathbb{C}P^n / \mathbb{C}P^m$.

2. Some general facts and proofs of Theorem 1.2 and Theorem 1.3

We recall that if X is a finite connected CW-complex then r_X denotes the smallest even integer such that $\tilde{H}^{r_X}(X) \neq 0$. For any X , $\tilde{H}^{2i}(X) = 0 \ \forall i$, we define $r_X = \dim X + 2$ if X is even dimensional and $r_X = \dim X + 1$ if X is odd dimensional CW-complex. We start with the following lemma.

Lemma 2.1. *Let ξ and η be two complex vector bundles over a finite connected CW-complex X .*

- (1) *If $\bar{\xi}$ is the conjugate bundle of ξ then*

$$\text{chern rank } \xi = \text{chern rank } \bar{\xi}.$$

- (2) *If $\omega = \text{Hom}(\xi, \mathbb{C})$, the dual bundle of ξ then*

$$\text{chern rank } \xi = \text{chern rank } \omega.$$

- (3) *If $c_{r_X}(\xi) = 0$ then $\text{chern rank } \xi = r_X - 2$.*

- (4) If $\widetilde{H}^{r_X}(X)$ is not cyclic then $\text{uch rank}(X) = r_X - 2$.
- (5) If $c(\xi) = 1$ then $\text{chern rank } \xi = r_X - 2$.
- (6) If $c(\eta) = 1$ then $\text{chern rank}(\xi \oplus \eta) = \text{chern rank } \xi$. Moreover,

$$\widetilde{K}(X) = 0 \text{ implies } \text{uch rank}(X) = r_X - 2.$$

- (7) If ξ and η are stably isomorphic then $\text{chern rank } \xi = \text{chern rank } \eta$.
- (8) There exists a complex vector bundle ξ' such that

$$\text{chern rank}(\xi \oplus \xi') = r_X - 2.$$

PROOF: (1) follows from the fact that the Chern class $c_k(\overline{\xi}) = (-1)^k c_k(\xi)$. As X is compact we may assume that ξ admits an Hermitian metric. Consequently $\omega = \text{Hom}(\xi, \mathbb{C})$ becomes canonically isomorphic to $\overline{\xi}$. Hence

$$\text{chern rank } \xi = \text{chern rank } \omega,$$

proving (2). Assertions (3) and (4) are obvious and (5) follows from (3). Assertion (6) follows from the fact that if $c(\eta) = 1$ then $c(\xi \oplus \eta) = c(\xi)$ and again $\widetilde{K}(X) = 0$ implies $c(\eta) = 1$ for any complex vector bundle η over X . To prove the statement (7), suppose ξ and η are stably isomorphic. Then $\xi \oplus \varepsilon^m \cong \eta \oplus \varepsilon^n$ for some m and n and hence $c(\xi) = c(\eta)$. Finally, as X is compact so for any bundle ξ over X there exists a bundle ξ' over X such that $\xi \oplus \xi' \cong \varepsilon^k$ for some k . Thus (8) follows from (5). □

Lemma 2.2. *If X is a finite connected CW-complex consisting of only even dimensional cells then upper characteristic rank $(X) \geq \text{uch rank}(X) + 1$.*

PROOF: It is clear that the coefficient homomorphism $H^*(X; \mathbb{Z}) \rightarrow H^*(X; \mathbb{Z}_2)$ becomes an epimorphism as X consists of only even dimensional cells. Now it is known that if ξ is a complex vector bundle over X then the coefficient homomorphism maps the total Chern class $c(\xi)$ onto the total Stiefel–Whitney class $w(\xi_R)$, see [5], Problem 14-B. Hence the proof follows. □

If $X = \Sigma Y$, where ΣY denotes the reduced suspension of Y then $\widetilde{H}^*(X) \cong \widetilde{H}^*(Y) \otimes \widetilde{H}^*(S^1)$ and consequently the cup product of two positive degree cohomology classes of $\widetilde{H}^*(X)$ becomes zero. Thus we have the following lemma.

Lemma 2.3. *Suppose $X = \Sigma Y$ and let $k_X = \max\{2k: H^{2j}(X) \text{ is cyclic, } 0 \leq j \leq k, 2k \leq \dim X\}$. Then $\text{uch rank}(X) \leq k_X$.*

In the above lemma trivial groups are considered to be cyclic. We note that if X is ordinary (nonreduced) suspension, then it is covered by two open contractible subsets, hence the cup product is trivial in this case as well and Lemma 2.3 applies.

Lemma 2.4. *Let $f: X \rightarrow Y$ be a map where X, Y are finite connected CW-complexes and let $f^*: H^*(Y) \rightarrow H^*(X)$ be a surjection. Then $\text{chern rank } f^*(\xi) \geq \min\{\text{chern rank } \xi, \dim X - 1\}$ for any complex vector bundle ξ over Y .*

PROOF: If $\dim X \geq \dim Y$ then from the naturality of Chern classes it follows that $\text{chern rank } f^*(\xi) \geq \text{chern rank } \xi$. Let $\dim X < \dim Y$. Now if $\dim X < \text{chern rank } \xi$ then clearly $\text{chern rank } f^*(\xi) = \dim X$ if $\dim X$ is even and $\text{chern rank } f^*(\xi) = \dim X - 1$ if $\dim X$ is odd. Again if $\dim X \geq \text{chern rank } \xi$ then $\text{chern rank } f^*(\xi) \geq \text{chern rank } \xi$. Combining all the above cases we get $\text{chern rank } f^*(\xi) \geq \min\{\text{chern rank } \xi, \dim X - 1\}$. \square

Let us consider the projective space $\mathbb{F}P^n$ where $\mathbb{F} = \mathbb{C}$ or \mathbb{H} , the complex or quaternionic numbers, respectively. If L and M denote the canonical (complex and quaternionic) line bundles over $\mathbb{C}P^n$ and $\mathbb{H}P^n$, respectively, then the Chern classes $c_1(L)$ and $c_2(M)$ are generators of $H^*(\mathbb{C}P^n)$ and $H^*(\mathbb{H}P^n)$, respectively. Hence we get the following theorem.

Theorem 2.5. *If $X = \mathbb{C}P^n$ or $\mathbb{H}P^n$ then $\text{uch rank}(X) = 2n$ or $4n$, respectively.*

Now we look at the chern rank of complex vector bundles over spheres. It follows from Theorem 2.5 that there exist complex vector bundles ξ_1 (line bundle) and ξ_2 (2-plane complex bundle) over $S^2 = \mathbb{C}P^1$ and $S^4 = \mathbb{H}P^1$, respectively, such that $c_1(\xi_1)$ and $c_2(\xi_2)$ are generators of $H^2(S^2)$ and $H^4(S^4)$, respectively. Thus $\text{chern rank } \xi_i = 2$ or 4 for $i = 1$ or 2 . Consequently $\text{uch rank}(S^{2n}) = 2n$ if $n = 1$ or 2 . In this context we want to state Bott integrality theorem which will be used in the sequel.

Theorem 2.6 (Bott integrality theorem [2], Chapter 20, Corollary 9.8). *Let $a \in H^{2n}(S^{2n})$ be a generator. Then for each complex vector bundle ξ over S^{2n} , the n th Chern class $c_n(\xi)$ is a multiple of $(n - 1)! a$, and for each m with $m \equiv 0 \pmod{(n - 1)!}$ there exists a unique $\xi \in \widetilde{K}(S^{2n})$ with $c_n(\xi) = ma$.*

Now it follows from Theorem 2.6 that if ξ is any complex vector bundle over S^{2n} where $n \neq 1$ or 2 then $c_n(\xi)$ cannot be a generator of $H^{2n}(S^{2n})$ and consequently for any complex vector bundle ξ over S^{2n} ($n \neq 1$ or 2) $\text{chern rank } \xi = 2n - 2$. We note that if n is odd then clearly $\text{uch rank}(S^n) = n - 1$. Combining these we get the following theorem.

Theorem 2.7. *If n is odd then $\text{uch rank}(S^n) = n - 1$, $\text{uch rank}(S^{2n}) = 2n$ if $n = 1$ or 2 and $\text{uch rank}(S^{2n}) = 2n - 2$ if $n \neq 1$ or 2 .*

If X and Y are two closed connected smooth orientable manifolds then the following theorem tells us that under suitable conditions *upper chern rank* of the product space $X \times Y$ is strictly less than $\dim(X \times Y)$.

Theorem 2.8. *Let X and Y be closed connected smooth orientable manifolds.*

- (1) *If $\widetilde{K}(X)$, $\widetilde{K}(Y)$ and $\widetilde{K}(X \wedge Y)$ are all trivial then*

$$\text{uch rank}(X \times Y) < \dim(X \times Y).$$

- (2) *If $\widetilde{KO}(X)$, $\widetilde{KO}(Y)$ and $\widetilde{KO}(X \wedge Y)$ are all trivial then*

$$\text{uch rank}(X \times Y) < \dim(X \times Y).$$

PROOF: (1) Let $\dim X = m$ and $\dim Y = n$. If $m + n$ is odd then it is trivial. So we assume $m + n$ is even. We note that as $X \times Y$ is orientable smooth manifold therefore $H_{m+n-1}(X \times Y)$ becomes torsion free and thus $H^{m+n}(X \times Y) \cong \mathbb{Z}$. We consider the inclusion followed by the quotient map $X \vee Y \hookrightarrow X \times Y \rightarrow X \wedge Y$. This yields the exact sequence $\tilde{K}(X \wedge Y) \rightarrow \tilde{K}(X \times Y) \rightarrow \tilde{K}(X \vee Y)$. Now $\tilde{K}(X \wedge Y) = 0$ and again $\tilde{K}(X) = 0 = \tilde{K}(Y)$ implies $\tilde{K}(X \vee Y) \cong \tilde{K}(X) \oplus \tilde{K}(Y) = 0$. Thus $\tilde{K}(X \times Y) = 0$ and consequently every complex vector bundle over $X \times Y$ becomes stably trivial. Thus for any complex vector bundle ξ over $X \times Y$ the total Chern class $c(\xi) = 1$ while $H^{m+n}(X \times Y) \neq 0$ and thus $\text{chern rank } \xi < \dim(X \times Y)$.

(2) Let $m + n$ be even. As before we get $\tilde{K}\widehat{O}(X \times Y) = 0$ and so for any real vector bundle η over $X \times Y$ the total Stiefel–Whitney class $w(\eta) = 1$. If possible $\text{uch rank}(X \times Y) = m + n$, there exists a complex vector bundle ξ over $X \times Y$ such that $\text{chern rank}(\xi) = m + n$. Let a be a generator of $H^{m+n}(X \times Y) \cong \mathbb{Z}$ and so a can be expressed as a polynomial of Chern classes $c_i(\xi)$. Let $a = P(c_1(\xi) \cdot c_2(\xi) \cdots c_t(\xi))$, $t \leq (m + n)/2$. Now if

$$f: H^*(X \times Y; \mathbb{Z}) \rightarrow H^*(X \times Y; \mathbb{Z}_2)$$

be the canonical coefficient homomorphism then $f(a)$ becomes the generator of $H^{m+n}(X \times Y; \mathbb{Z}_2) \cong \mathbb{Z}_2$ and again

$$\begin{aligned} f(a) &= f(P(c_1(\xi) \cdot c_2(\xi) \cdots c_t(\xi))) \\ &= P(f(c_1(\xi)) \cdot f(c_2(\xi)) \cdots f(c_t(\xi))) \\ &= P(\omega_2(\xi_R) \cdot \omega_4(\xi_R) \cdots \omega_{2t}(\xi_R)) = 0, \end{aligned}$$

a contradiction. Thus $\text{uch rank}(X \times Y) < \dim(X \times Y)$. □

If X is a closed connected complex manifold of complex dimension d and ξ is a complex vector bundle over X then the following theorem tells us that under certain given conditions $\text{chern rank } \xi$ can be predicted.

Theorem 2.9. *Let X be a closed connected complex manifold of complex dimension d . If $r_X \leq d$ and $H^{r_X}(X) \cong \mathbb{Z}$ then for any complex vector bundle ξ over X , $\text{chern rank } \xi$ is either less than $2d - r_X$ or $2d$.*

PROOF: Every complex manifold of complex dimension d is a $2d$ dimensional smooth orientable manifold. The triviality of $H^1(X), H^2(X), \dots, H^{r_X-1}(X)$ implies $H_1(X), H_2(X), \dots, H_{r_X-2}(X)$ are all trivial and hence by Poincaré duality the cohomology groups $H^{2d-1}(X), H^{2d-2}(X), \dots, H^{2d-r_X+2}(X)$ are trivial. Let ξ be a complex vector bundle over X such that $\text{chern rank } \xi \geq 2d - r_X$, $2d - r_X \geq r_X$. We only have to show that any cohomology class of $H^{2d}(X) \cong \mathbb{Z}$ can be expressed as a polynomial of Chern classes.

As $\text{chern rank}(\xi) \geq 2d - r_X \geq r_X$ therefore $H^{r_X}(X) = \langle c_{r_X/2}(\xi) \rangle$. Now as X is a closed connected \mathbb{Z} -orientable manifold so there exists some $\beta \in H^{2d-r_X}(X)$ such that $c_{r_X/2}(\xi) \cdot \beta$ is a generator of $H^{2d}(X)$, while β can be expressed as

a polynomial of Chern classes of ξ and consequently $c_{r_X/2}(\xi) \cdot \beta$ can be expressed as a polynomial of Chern classes of ξ . This completes the proof. \square

We recall that $\text{Cup}_E(X)$, the even cup length of X is the largest integer t such that the cup product $x_1 \cdot x_2 \cdots x_t \neq 0$ where each x_i is an even degree cohomology class with $\deg(x_i) \geq 2$. If X is a closed connected d -dimensional complex manifold then Theorem 1.2 gives a bound for $\text{Cup}_E(X)$. Proofs of Theorems 1.2 and 1.3 are similar to the proofs of Theorem 1.2 and 1.3 of [7], respectively.

PROOF OF THEOREM 1.2: Let $\text{Cup}_E(X) = t$ and $x_1 \cdot x_2 \cdots x_t \neq 0$ be a maximal string of nonzero cup product. We claim that $x_1 \cdot x_2 \cdots x_t \in H^{2d}(X)$. If not then $x_1 \cdot x_2 \cdots x_t \in H^{2d-2l}(X)$ for some $l > 0$. Now as $H^{2i}(X)$ is torsion free for all i , therefore the cup product pairing $H^{2d-2l}(X) \times H^{2l}(X) \rightarrow \mathbb{Z}$ is nonsingular and hence there exists $y \in H^{2l}(X)$ ($y \neq 0$) such that $x_1 \cdot x_2 \cdots x_t \cdot y \in H^{2d}(X)$ is a nonzero element. This contradicts the maximality of $x_1 \cdot x_2 \cdots x_t$.

Now we rearrange $x_1 \cdot x_2 \cdots x_t$ as $y_1 \cdot y_2 \cdots y_m \cdot z_1 \cdot z_2 \cdots z_n$ such that $\deg(y_i) = i$, $\deg(z_j) = j$ with $i \leq 2k$ and $j \geq 2k + 2$. If possible, suppose

$$x_1 \cdot x_2 \cdots x_t = y_1 \cdot y_2 \cdots y_m.$$

As $i \leq 2k \leq \text{chern rank}(\xi)$, therefore, $y_1 \cdot y_2 \cdots y_m$ is a polynomial in Chern classes $c_1(\xi), \dots, c_k(\xi)$ laying in $H^{2d}(X)$. Hence it is a sum of monomials in Chern classes each of which is zero and thus $y_1 \cdot y_2 \cdots y_m = 0$. Consequently, the string $z_1 \cdot z_2 \cdots z_n$ must exist.

Let $a = y_1 \cdot y_2 \cdots y_m$ and $b = z_1 \cdot z_2 \cdots z_n$. As $\deg(b) \geq 2k + 2$ therefore $\deg(a) \leq 2d - 2(k + 1)$ and

$$\begin{aligned} \text{Cup}_E(X) = m + n &\leq \frac{\deg(a)}{r_X} + \frac{\deg(b)}{2k + 2} = \frac{2(k + 1) \deg(a) + r_X \deg(b)}{2r_X(k + 1)} \\ &= \frac{2(k + 1) \deg(a) + r_X(2d - \deg(a))}{2r_X(k + 1)} \\ &= \frac{(2(k + 1) - r_X) \deg(a) + 2dr_X}{2r_X(k + 1)} \\ &\leq \frac{(2(k + 1) - r_X)(d - (k + 1)) + dr_X}{r_X(k + 1)} \\ &= \frac{r_X(k + 1) + 2(k + 1)(d - k - 1)}{r_X(k + 1)} = 1 + \frac{2(d - k - 1)}{r_X}. \end{aligned}$$

\square

PROOF OF THEOREM 1.3: As $\text{uch rank}(X) = \dim X$ therefore there exists a complex vector bundle ξ over X with $\text{chern rank} \xi = \dim X$. Let $\text{Cup}_E(X) = t$ and

$$x_1 \cdot x_2 \cdots x_i \cdots x_t \neq 0$$

be a maximal string of nonzero cup product. As chern rank $\xi = \dim X$ hence x_i can be expressed as a polynomial of Chern classes of ξ and consequently $x = x_1 \cdot x_2 \cdots x_t$ can be expressed as a sum of integral multiples of monomials of Chern classes $c_1(\xi), c_2(\xi), \dots, c_r(\xi)$, $2r \leq \max \deg(x_i)$, each of length at least t . But as monomials of Chern classes of length greater than t vanish therefore there must exist a monomial $c_{i_1}(\xi) \cdot c_{i_2}(\xi) \cdots c_{i_t}(\xi)$ of length t with $c_{i_1}(\xi) \cdot c_{i_2}(\xi) \cdots c_{i_t}(\xi) \neq 0$. \square

3. Some computations

In this final section we compute uch rank of some important spaces.

Theorem 3.1. *Let $X = S^m \times S^n$.*

(1) *If m, n are even integers and $m < n$ then*

$$\text{uch rank}(X) = \begin{cases} m - 2 & \text{if } m \neq 2, 4, \\ n - 2 & \text{if } m = 2, 4 \text{ and } n \neq 2, 4, \\ m + n & \text{if } m = 2, n = 4. \end{cases}$$

(2) *If m, n are even integers and $m = n$ then $\text{uch rank}(X) = m - 2$.*

(3) *If m is odd and n is even then*

$$\text{uch rank}(X) = \begin{cases} n - 2 & \text{if } n \neq 2, 4, \\ m + n - 1 & \text{if } n = 2, 4. \end{cases}$$

(4) *If m and n are odd integers and $m + n = 2$ or 4 then $\text{uch rank}(X) = m + n$.*

(5) *If $m, n \equiv 3 \pmod{8}$ then $\text{uch rank}(X) = m + n - 2$ and if $n \equiv 5 \pmod{8}$ then $\text{uch rank}(S^1 \times S^n) = n - 1$.*

PROOF: (1) We note that $\tilde{H}^i(S^m \times S^n)$ is nontrivial if $i = m, n$ or $m + n$. We observe that the inclusion map $i: S^m \hookrightarrow S^m \times S^n$ and projection $p: S^m \times S^n \rightarrow S^m$ induces isomorphisms on the m th cohomology groups, respectively. Thus if $m \neq 2, 4$ and ξ is a complex vector bundle over $S^m \times S^n$ with chern rank $\xi \geq m$ then $i^*(\xi)$ becomes a complex vector bundle over S^m and by naturality of Chern classes chern rank $i^*(\xi) \geq m$ which is a contradiction as $\text{uch rank}(S^m) = m - 2$ if $m \neq 2, 4$ (cf. Theorem 2.7). So it follows that $\text{uch rank}(S^m \times S^n) = m - 2$.

If $m = 2, 4$ and $n \neq 2, 4$ then by similar argument $\text{uch rank}(S^m \times S^n) \leq n - 2$. By Theorem 2.7, there exists a complex vector bundle γ over S^m with chern rank $\gamma = m$. Again as $p^*: H^m(S^m) \rightarrow H^m(S^m \times S^n)$ is an isomorphism, it follows that chern rank $p^*(\gamma) \geq m$. Thus $\text{uch rank}(S^m \times S^n) = n - 2$.

Finally, let $m = 2$ and $n = 4$. Note that there exist complex line bundle γ_1 and complex 2-plane bundle γ_2 over S^m and S^n , respectively, such that chern rank $\gamma_1 = 2$ and chern rank $\gamma_2 = 4$. Consider the projection maps $p_1: S^m \times S^n \rightarrow S^m$ and $p_2: S^m \times S^n \rightarrow S^n$. As $p_1^*: H^m(S^m) \rightarrow H^m(S^m \times S^n)$ and $p_2^*: H^n(S^n) \rightarrow H^n(S^m \times S^n)$ are isomorphisms so the total Chern class

$c(p_1^*(\gamma_1)) = 1 + a$ and $c(p_2^*(\gamma_2)) = 1 + b$ where a and b are generators of $H^m(S^m \times S^n)$ and $H^n(S^m \times S^n)$, respectively. Consider the Whitney sum $p_1^*(\gamma_1) \oplus p_2^*(\gamma_2)$ over $S^m \times S^n$ which is a 3-plane complex bundle over $S^m \times S^n$. Again $c(p_1^*(\gamma_1) \oplus p_2^*(\gamma_2)) = c(p_1^*(\gamma_1)) \cdot c(p_2^*(\gamma_2))$ and if a and b are generators of $H^m(S^m \times S^n)$ and $H^n(S^m \times S^n)$, respectively, then it follows from the cohomology ring structure of $H^*(S^m \times S^n)$ that $a \cdot b$ is a generator of $H^{m+n}(S^m \times S^n)$. Consequently it turns out that $\text{chern rank}(p_1^*(\gamma_1) \oplus p_2^*(\gamma_2)) = m + n$.

(2) The first nontrivial reduced integral cohomology group of $S^m \times S^m$ is $\tilde{H}^m(S^m \times S^m)$ which is free abelian of rank 2 and the proof follows from assertion (4) of Lemma 2.1.

(3) Here we notice that if m is odd and n is even then the only nontrivial even dimensional reduced integral cohomology group of $S^m \times S^n$ is $\tilde{H}^n(S^m \times S^n)$ and the proof is similar to the case of (1).

(4) As $S^m \times S^n$ is a closed connected $m + n$ dimensional smooth orientable manifold hence there exists a degree 1 map $f: S^m \times S^n \rightarrow S^{m+n}$. Thus $f_*: H_{m+n}(S^m \times S^n) \rightarrow H_{m+n}(S^{m+n})$ is an isomorphism and consequently $f^*: \text{Hom}(H_{m+n}(S^{m+n}); \mathbb{Z}) \rightarrow \text{Hom}(H_{m+n}(S^m \times S^n); \mathbb{Z})$ is an isomorphism. Again as $H_{m+n-1}(S^m \times S^n)$ is torsion free (as $S^m \times S^n$ is orientable) consequently $f^*: H^{m+n}(S^{m+n}) \rightarrow H^{m+n}(S^m \times S^n)$ becomes an isomorphism. Now the proof follows from the fact that $\text{uch rank}(S^{m+n}) = m + n$ if $m + n = 2$ or 4.

(5) If $m, n \equiv 3 \pmod{8}$ then $\widetilde{KO}(S^{m+n}) = 0 = \widetilde{KO}(S^n)$ and again as $m + n \equiv 6 \pmod{8}$ therefore $\widetilde{KO}(S^{m+n}) = \widetilde{KO}(S^m \wedge S^n) = 0$. By assertion (2) of Theorem 2.8 $\text{uch rank}(S^m \times S^n) < m + n$ and consequently $\text{uch rank}(S^m \times S^n) = m + n - 2$. If $n \equiv 5 \pmod{8}$ then every orientable real vector bundle over $S^1 \times S^n$ becomes stably trivial, see [6], Lemma 3.6, therefore there cannot exist any complex vector bundle ξ over $(S^1 \times S^n)$ such that $c_{(n+1)/2}(\xi)$ is a generator of $H^{n+1}(S^1 \times S^n)$ and thus $\text{uch rank}(S^1 \times S^n) = n - 1$. \square

We deduce the following corollary from part (4) of Theorem 3.1.

Corollary 3.2. *The upper chern rank of $S^1 \times S^1, S^1 \times S^3$ are 2 and 4, respectively.*

Remark. Note that $\text{uch rank}(S^1 \times S^1) = 2$ also follows from the fact that the first Chern class $c_1: \text{Vect}_{\mathbb{C}}^1(S^1 \times S^1) \rightarrow H^2(S^1 \times S^1) \cong \mathbb{Z}$ is an isomorphism ($\text{Vect}_{\mathbb{C}}^1(X)$ denotes the abelian group of isomorphism classes of complex line bundles over X with respect to tensor product operations).

Theorem 3.3. *Let $X = S^m \vee S^n$.*

(1) *If m, n are even integers and $m < n$ then*

$$\text{uch rank}(X) = \begin{cases} m - 2 & \text{if } m \neq 2, 4, \\ n - 2 & \text{if } m = 2 \text{ or } 4 \text{ and } n \neq 4, \\ n & \text{if } m = 2 \text{ and } n = 4. \end{cases}$$

(2) If m is odd and n is even integer then

$$\text{uch rank}(X) = \begin{cases} n - 2 & \text{if } m < n \text{ and } n \neq 2, 4, \\ n & \text{if } m < n \text{ and } n = 2 \text{ or } 4, \\ n - 2 & \text{if } m > n \text{ and } n \neq 2, 4, \\ m - 1 & \text{if } m > n \text{ and } n = 2 \text{ or } 4. \end{cases}$$

(3) If m, n are even integers and $m = n$ then $\text{uch rank}(X) = m - 2$.

PROOF: (1) Let $i_1: S^m \hookrightarrow S^m \vee S^n, i_2: S^n \hookrightarrow S^m \vee S^n$ be the inclusions and $r_1: S^m \vee S^n \rightarrow S^m, r_2: S^m \vee S^n \rightarrow S^n$ be the retraction maps. We consider the sequence of maps $S^m \hookrightarrow S^m \vee S^n \rightarrow S^m$ and $S^n \hookrightarrow S^m \vee S^n \rightarrow S^n$. Clearly $i_1^*: H^m(S^m \vee S^n) \rightarrow H^m(S^m), i_2^*: H^n(S^m \vee S^n) \rightarrow H^n(S^n)$ and $r_1^*: H^m(S^m) \rightarrow H^m(S^m \vee S^n), r_2^*: H^n(S^n) \rightarrow H^n(S^m \vee S^n)$ are isomorphisms. Now $\text{uch rank}(X) = m - 2$ if $m \neq 2, 4$, and it is equal to $n - 2$ if $m = 2, 4$ and $n \neq 4$, which follows by similar arguments as in part (1) of Theorem 3.1.

Let $m = 2, n = 4$ and $j: S^m \vee S^n \hookrightarrow S^m \times S^n$ is inclusion and p_1, p_2 are the projection maps: $p_1: S^m \times S^n \rightarrow S^m, p_2: S^m \times S^n \rightarrow S^n$. We consider the sequence of maps: $S^m \hookrightarrow S^m \vee S^n \hookrightarrow S^m \times S^n \rightarrow S^m$ and $S^n \hookrightarrow S^m \vee S^n \hookrightarrow S^m \times S^n \rightarrow S^n$. As $(j_k^* \circ j^*) \circ p_k^*$ is isomorphism, $k = 1$ or 2 , hence $i_1^* \circ j^*: H^m(S^m \times S^n) \rightarrow H^m(S^m)$ and $i_2^* \circ j^*: H^n(S^m \times S^n) \rightarrow H^n(S^n)$ are surjections and hence isomorphisms. Again as i_k^* is an isomorphism, $k = 1$ or 2 , so it follows that $j^*: H^m(S^m \times S^n) \rightarrow H^m(S^m \vee S^n)$ and $j^*: H^n(S^m \times S^n) \rightarrow H^n(S^m \vee S^n)$ are isomorphisms. Note that by part (1) of Theorem 3.1 $\text{uch rank}(S^m \times S^n) = m + n$ and therefore there exists a complex vector bundle ξ over $S^m \times S^n$ such that $\text{chern rank}(\xi) = m + n$. Clearly $\text{chern rank } j^*(\xi) = n$.

(2) We note that the only even dimensional nontrivial reduced cohomology group of $S^m \vee S^n$ is $\tilde{H}^n(S^m \vee S^n) \cong \tilde{H}^n(S^n)$ and the arguments are similar to the first case.

Proof of (3) follows from assertion (4) of Lemma 2.1 as the only even dimensional nontrivial reduced cohomology group $\tilde{H}^m(S^m \vee S^n)$ is free abelian of rank 2. □

Lemma 3.4. For any complex vector bundle ξ over $\mathbb{R}P^{2k}$ (or $\mathbb{R}P^{2k+1}$), $\text{chern rank } \xi$ is either 0 or $2k$ and

$$\text{uch rank}(\mathbb{R}P^{2k}) = 2k = \text{uch rank}(\mathbb{R}P^{2k+1}).$$

PROOF: The graded integral cohomology ring of $\mathbb{R}P^{2k}$ is given by

$$H^*(\mathbb{R}P^{2k}) \cong \mathbb{Z}[\alpha]/(2\alpha, \alpha^{k+1}), \quad \text{deg}(\alpha) = 2.$$

If ξ is a complex vector bundle over $\mathbb{R}P^{2k}$ with $c_1(\xi) = 0$ then $\text{chern rank } \xi = 0$ (for example we can take any trivial complex vector bundle) as $H^2(\mathbb{R}P^{2k}) \cong \mathbb{Z}_2$. On the contrary if $c_1(\xi) \neq 0$ then $H^{2i}(\mathbb{R}P^{2k}) = \langle (c_1(\xi))^i \rangle \cong \mathbb{Z}_2$ and consequently $\text{chern rank } \xi = 2k$. Now as $c_1: \text{Vect}_{\mathbb{C}}^1(\mathbb{R}P^{2k}) \rightarrow H^2(\mathbb{R}P^{2k})$ is an isomorphism

therefore there exists a complex line bundle ξ over $\mathbb{R}P^{2k}$ with $c_1(\xi) \neq 0$ and thus $\text{uch rank}(\mathbb{R}P^{2k}) = 2k$.

Again the graded integral cohomology ring of $\mathbb{R}P^{2k+1}$ is given by

$$H^*(\mathbb{R}P^{2k+1}) \cong \mathbb{Z}[\alpha, \beta]/(2\alpha, \alpha^{k+1}, \beta^2, \alpha\beta), \quad \deg(\alpha) = 2, \quad \deg(\beta) = 2k + 1$$

and the proof follows in similar fashion. □

Theorem 3.5. (1) *If $X = \mathbb{R}P^n \times S^{2m}$ then*

$$\text{uch rank}(X) = \begin{cases} 2(m+k) & \text{if } m = 2 \text{ and } n = 2k \text{ or } 2k + 1, \\ 2(m-1) & \text{if } m \neq 2. \end{cases}$$

(2) *If $X = \mathbb{C}P^n \times S^{2m}$ then*

$$\text{uch rank}(X) = \begin{cases} 2(m+n) & \text{if } m = 2, \\ 2(m-1) & \text{if } m \neq 2. \end{cases}$$

PROOF: (1) Let $n = 2k$. We consider the projection maps

$$p_1: \mathbb{R}P^{2k} \times S^{2m} \rightarrow \mathbb{R}P^{2k}$$

and $p_2: \mathbb{R}P^{2k} \times S^{2m} \rightarrow S^{2m}$. If a and b are generators of $H^2(\mathbb{R}P^{2k})$ and $H^{2m}(S^{2m})$, respectively, then the graded integral cohomology ring $H^*(\mathbb{R}P^{2k} \times S^{2m}) \cong \mathbb{Z}[\alpha, \beta]/(2\alpha, \alpha^{k+1}, \beta^2)$, $\deg(\alpha) = 2$, $\deg(\beta) = 2m$ where $\alpha = p_1^*(a)$ and $\beta = p_2^*(b)$.

If $m = 1$ then $\mathbb{R}P^{2k} \times S^2 \cong \mathbb{Z} \oplus \mathbb{Z}_2$ and by (4) of Lemma 2.1 $\text{uch rank}(X) = 0$. Let $m = 2$. Now it follows from Lemma 3.4 that there exists a complex line bundle ξ over $\mathbb{R}P^{2k}$ such that $\text{chern rank } \xi = 2k$ and there exists a complex vector bundle ξ' over S^4 such that $\text{chern rank } \xi' = 4$ (by Theorem 2.7). Let $a = c_1(\xi)$ and $b = c_2(\xi')$. Now we take the pull back bundles $v = p_1^*(\xi)$ and $\eta = p_2^*(\xi')$ over X and consider their Whitney sum $v \oplus \eta$. Clearly $c_1(v \oplus \eta) = c_1(v) = \alpha$ and $c_2(v \oplus \eta) = c_2(\eta) = \beta$ and consequently $\text{chern rank}(v \oplus \eta) = 2(m+k)$.

Finally let $m \neq 1, 2$. We note that $\text{chern rank } v \geq 2(m-1)$ and as β cannot be expressed as a product of cohomology classes of $H^*(X)$ with degree lower than $2m$ so $\text{chern rank } v = 2(m-1)$. Now if $\text{uch rank}(X) \geq 2m$, there exists a complex vector bundle γ over X such that $\text{chern rank } \gamma \geq 2m$. Let $i: S^{2m} \hookrightarrow \mathbb{R}P^{2k} \times S^{2m}$ be the inclusion map. As $i^* \circ p_2^* = \text{id}$ on $H^{2m}(S^{2m})$ thus it turns out that $i^*(\beta) = b$. Again as β cannot be expressed as a product of cohomology classes of $H^*(X)$ with degree lower than $2m$ therefore $c_m(\gamma)$ must be equal to β and so $c_m(i^*(\gamma)) = i^*c_m(\gamma) = i^*(\beta) = b$. Thus $\text{uch rank}(S^{2m}) = 2m$; which contradicts $\text{uch rank}(S^{2m}) = 2m - 2$ if $m \neq 1, 2$ (Theorem 2.7). This completes the proof for $m \neq 1, 2$.

If $n = 2k + 1$ then $H^*(\mathbb{R}P^{2k+1} \times S^{2m}) \cong \mathbb{Z}[\alpha, \beta, \lambda]/(2\alpha, \alpha^{k+1}, \lambda^2, \alpha \cdot \lambda, \beta^2)$ where $\deg(\alpha) = 2$, $\deg(\beta) = 2m$, $\deg(\lambda) = 2k + 1$ and the proof is similar to the case $n = 2k$.

(2) We note that the graded integral cohomology ring $H^*(\mathbb{C}P^n \times S^{2m}) \cong \mathbb{Z}[\alpha, \beta]/(\alpha^{n+1}, \beta^2)$ where $\deg(\alpha) = 2, \deg(\beta) = 2m$ and also if L is the canonical complex line bundle over $\mathbb{C}P^n$ then $\text{chern rank } L = 2n$. Now the proof follows by arguments as in (1). \square

Now we study complex vector bundles over complex Stiefel manifolds $V_k(\mathbb{C}^n)$ which consists of the orthonormal k -frames in \mathbb{C}^n .

Theorem 3.6. *Let $X = V_k(\mathbb{C}^n)$, where $1 < k < n$. Then $\text{uch rank}(X) = 4(n - k) + 2$ if $n - k$ is even or $n - k \neq 2^t - 1, t > 0$.*

PROOF: It is known that for any commutative ring with unit $R, H^*(V_k(\mathbb{C}^n); R) \cong \bigwedge(x_{2(n-k)+1}, x_{2(n-k)+3}, \dots, x_{2n-1})$, that is, the exterior algebra generated by $x_{2(n-k)+1}, x_{2(n-k)+3}, \dots, x_{2n-1}$ where $x_j \in H^j(V_k(\mathbb{C}^n); R)$, see [4], Proposition 5.11. We note that the first nontrivial dimensional reduced cohomology group of $V_k(\mathbb{C}^n)$ with integer coefficients is $\widetilde{H}^{4(n-k)+4}(V_k(\mathbb{C}^n); \mathbb{Z}) \cong \mathbb{Z}$. Also the integral cohomology structure of $V_k(\mathbb{C}^n)$ implies that the natural coefficient homomorphism $H^{4(n-k)+4}(V_k(\mathbb{C}^n); \mathbb{Z}) \rightarrow H^{4(n-k)+4}(V_k(\mathbb{C}^n); \mathbb{Z}_2)$ is an epimorphism where $H^{4(n-k)+4}(V_k(\mathbb{C}^n); \mathbb{Z}_2) \cong \mathbb{Z}_2$. Again it is well known that for any real vector bundle ξ over a space B , if $w_m(\xi), m > 0$ is the first nonzero Stiefel–Whitney class then m must be a power of 2, see [5], Problem 8-B. Now if $n - k (> 0)$ is even or $n - k \neq 2^t - 1, t > 0$, then $4(n - k) + 4$ cannot be a power of 2 and consequently for any vector bundle ξ over $V_k(\mathbb{C}^n), 1 < k < n, w_{4(n-k)+4}(\xi) = 0$. Thus for any complex vector bundle η over $V_k(\mathbb{C}^n), 1 < k < n; c_{2(n-k)+2}(\eta)$ cannot be a generator of $H^{4(n-k)+4}(V_k(\mathbb{C}^n); \mathbb{Z})$ as under the natural coefficient homomorphism $H^{4(n-k)+4}(V_k(\mathbb{C}^n); \mathbb{Z}) \rightarrow H^{4(n-k)+4}(V_k(\mathbb{C}^n); \mathbb{Z}_2)$, which is an epimorphism, the Chern class $c_{2(n-k)+2}(\eta)$ is mapped to the Stiefel–Whitney class $w_{4(n-k)+4}(\eta_R)$ and hence $\text{uch rank}(V_k(\mathbb{C}^n)) = 4(n - k) + 2$. \square

Theorem 3.7. *If $X = \mathbb{C}P^n/\mathbb{C}P^m$, where $m \geq 1, n \geq m + 2$ then*

$$\text{uch rank}(X) = \begin{cases} 2m & \text{if } m \neq 1, \\ 4 & \text{if } m = 1. \end{cases}$$

PROOF: First we observe that the first nontrivial cohomology group of X is $H^{2m+2}(X)$ and if $i: S^{2m+2} \hookrightarrow X$ is the inclusion map then $i^*: H^{2m+2}(X) \rightarrow H^{2m+2}(S^{2m+2})$ is an isomorphism. Now if $m \neq 1$ then $\text{uch rank}(S^{2m+2}) = 2m$ (cf. Theorem 2.7) and consequently $\text{uch rank}(X) = 2m$.

Next we consider the case when $m = 1$. Now $\mathbb{C}P^3/\mathbb{C}P^1 = S^4 \cup_{f_1} e^6$ where $f_1: S^5 \rightarrow S^4$ is the attaching map and e^6 denotes a 6-cell. It is well known that $\pi_5(S^4) \cong \mathbb{Z}_2$ and generated by $[\Sigma^2 f]$, where $\Sigma^2 f$ denotes the double suspension of the Hopf map $f: S^3 \rightarrow S^2$. Let α be a generator of $H^2(\mathbb{C}P^\infty; \mathbb{Z}_2)$ where $H^*(\mathbb{C}P^\infty; \mathbb{Z}_2) \cong \mathbb{Z}_2[\alpha]$. We note that the action of Steenrod square operation Sq^2 on α^2 is trivial. Let us consider the quotient map $q: \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty/\mathbb{C}P^1$. Now it follows from the naturality of Steenrod squaring operation that $Sq^2(x)$ is trivial where x is the generator of $H^4(\mathbb{C}P^\infty/\mathbb{C}P^1; \mathbb{Z}_2)$. Again applying naturality property of Steenrod squares with the inclusion map $i_1: \mathbb{C}P^3/\mathbb{C}P^1 \hookrightarrow \mathbb{C}P^\infty/\mathbb{C}P^1$ it

follows that the action of Sq^2 on the generator of $H^4(\mathbb{C}P^3/\mathbb{C}P^1; \mathbb{Z}_2) \cong \mathbb{Z}_2$ is trivial. Suppose now the attaching map $f_1: S^5 \rightarrow S^4$ was not null-homotopic. Then f_1 must be homotopic to the double suspension of the Hopf map $f: S^3 \rightarrow S^2$ as $\pi_5(S^4) \cong \mathbb{Z}_2$. Thus $\mathbb{C}P^3/\mathbb{C}P^1 = C_{\Sigma^2 f} = \Sigma^2 C_f$, where C_f is the associated mapping cone of $f: S^3 \rightarrow S^2$. Again as Steenrod square operations are invariant under suspension it follows that the action of Sq^2 on the generator of $H^4(\mathbb{C}P^3/\mathbb{C}P^1; \mathbb{Z}_2)$ is nontrivial, a contradiction. Consequently f_1 must be null-homotopic. Thus $\mathbb{C}P^3/\mathbb{C}P^1 \approx S^4 \vee S^6$.

Now by Theorem 3.3 (1), $\text{uchrank}(\mathbb{C}P^3/\mathbb{C}P^1) = \text{uchrank}(S^4 \vee S^6) = 4$. Again we consider the inclusion map $j: \mathbb{C}P^3/\mathbb{C}P^1 \hookrightarrow \mathbb{C}P^n/\mathbb{C}P^1$. As a map $j^*: H^k(\mathbb{C}P^n/\mathbb{C}P^1) \rightarrow H^k(\mathbb{C}P^3/\mathbb{C}P^1)$ induces isomorphisms for $k \leq 6$ so it follows that $\text{uchrank}(\mathbb{C}P^n/\mathbb{C}P^1) \leq 4$. Finally we note that a map $j^*: \widetilde{K}(\mathbb{C}P^n/\mathbb{C}P^1) \rightarrow \widetilde{K}(\mathbb{C}P^3/\mathbb{C}P^1)$ induces epimorphism in reduced K -groups, see [1], Theorem 7.2, and so $\text{uchrank}(\mathbb{C}P^n/\mathbb{C}P^1) = 4$. This completes the proof. \square

Acknowledgement. The author thanks Professor Goutam Mukherjee, Professor Gregory Arone and Dr. Prateep Chakraborty for their valuable suggestions and comments.

REFERENCES

- [1] Adams J. F., *Vector fields on spheres*, Ann. of Math. (2) **75** (1962), no. 3, 603–632.
- [2] Husemoller D., *Fibre Bundles*, McGraw-Hill Book Co., New York, 1966.
- [3] Korbaš J., *The cup length of oriented Grassmannians vs a new bound for zero-cobordant manifolds*, Bull. Belg. Math. Soc. Simon Stevin **17** (2010), no. 1, 69–81.
- [4] McCleary J., *A User's Guide to Spectral Sequences*, Cambridge Studies in Advanced Mathematics, 58, Cambridge University Press, Cambridge, 2001.
- [5] Milnor J., Stasheff J., *Characteristic Classes*, Annals of Mathematics Studies, 76, Princeton University Press, Princeton, University of Tokyo Press, Tokyo, 1974.
- [6] Naolekar A. C., *Realizing cohomology classes as Euler classes*, Math. Slovaca **62** (2012), no. 5, 949–966.
- [7] Naolekar A. C., Thakur A. S., *Note on the characteristic rank of vector bundles*, Math. Slovaca **64** (2014), no. 6, 1525–1540.

B. Banerjee:

DEPARTMENT OF MATHEMATICS, RANAGHAT COLLEGE, RANAGHAT, NADIA,
WEST BENGAL 741201, INDIA

E-mail: pbikraman@rediffmail.com

(Received May 9, 2018, revised October 10, 2018)