

Stephen J. Gardiner; Tomas Sjödin

On a conjecture of Král concerning the subharmonic extension of continuously differentiable functions

*Mathematica Bohemica*, Vol. 145 (2020), No. 1, 71–73

Persistent URL: <http://dml.cz/dmlcz/148065>

## Terms of use:

© Institute of Mathematics AS CR, 2020

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A CONJECTURE OF KRÁL CONCERNING  
THE SUBHARMONIC EXTENSION OF  
CONTINUOUSLY DIFFERENTIABLE FUNCTIONS

STEPHEN J. GARDINER, Dublin, TOMAS SJÖDIN, Linköping

Received August 24, 2018. Published online March 5, 2019.  
Communicated by Dagmar Medková

*Abstract.* This note verifies a conjecture of Král, that a continuously differentiable function, which is subharmonic outside its critical set, is subharmonic everywhere.

*Keywords:* subharmonic function; extension theorem

*MSC 2010:* 31B05

## 1. INTRODUCTION

A classical result of Radó (see Theorem 12.14 of [9]) says that if  $f$  is continuous on an open set  $\Omega \subset \mathbb{C}$  and holomorphic on  $\{z \in \Omega : f(z) \neq 0\}$ , then  $f$  is holomorphic on all of  $\Omega$ . An analogue for harmonic functions due to Král (see [6]) says that if  $u : \Omega \rightarrow \mathbb{R}$  is  $C^1$  on an open set  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 2$  and harmonic on  $\{x \in \Omega : \nabla u(x) \neq 0\}$ , then  $u$  is harmonic on all of  $\Omega$ . (A short proof of this result was recently given in [8].) Král conjectured in [7] that his result could be strengthened by substituting “subharmonic” for “harmonic” throughout. However, the methods of [6] and [8] are not applicable to subharmonic functions. The purpose of this note is to verify this conjecture.

## 2. MAIN RESULT

**Theorem 1.** *If  $u$  is  $C^1$  on an open set  $\Omega \subset \mathbb{R}^N$  and subharmonic on  $\{x \in \Omega : \nabla u(x) \neq 0\}$ , then  $u$  is subharmonic on all of  $\Omega$ .*

The idea of the proof below comes from the theory of viscosity solutions of partial differential equations, which is expounded in [2], [3]. In fact, Theorem 1 may readily be deduced from results in [5] concerning viscosity solutions of the  $p$ -Laplace equation (cf. [4] for a generalization of Král's original result to  $p$ -harmonic functions). However, we will instead give a self-contained argument, partially inspired by [5], that uses only some basic properties of subharmonic functions. A convenient background reference is [1].

*Proof.* Let  $\varepsilon > 0$  and  $B$  be an open ball  $\{x: \|x - x_1\| < r\}$  such that  $\bar{B} \subset \Omega$ . By taking the Poisson integral of  $u$  in  $B$  and adding the polynomial

$$x \mapsto \varepsilon \left( 1 + \frac{r^2 - \|x - x_1\|^2}{2N} \right),$$

we obtain a function  $h_\varepsilon \in C(\bar{B})$  satisfying

$$\begin{cases} \Delta h_\varepsilon = -\varepsilon & \text{in } B, \\ h_\varepsilon = u + \varepsilon & \text{on } \partial B. \end{cases}$$

It will be enough to show that  $h_\varepsilon \geq u$  in  $B$ , since we can then let  $\varepsilon$  tend to 0 to arrive at the required spherical mean value inequality for  $u$ .

The set

$$O = \{(x, y) \in \bar{B} \times \bar{B}: h_\varepsilon(x) - u(y) > \frac{1}{2}\varepsilon\}$$

is relatively open in  $\bar{B} \times \bar{B}$  and contains  $\{(x, x): x \in \partial B\}$ . Thus, the quantity  $\|x - y\|^4$  is bounded away from zero on  $\partial(B \times B) \setminus O$ , and we may choose  $c > 0$  large enough so that  $w > 0$  on  $\partial(B \times B)$ , where

$$w(x, y) = h_\varepsilon(x) - u(y) + c\|x - y\|^4, \quad x, y \in \bar{B}.$$

We suppose, for the sake of contradiction, that the minimum value of the continuous function  $w$  on  $\bar{B} \times \bar{B}$  is attained at some point  $(x_0, y_0) \in B \times B$ .

Setting  $y = y_0$  in the inequality

$$(1) \quad h_\varepsilon(x) - u(y) + c\|x - y\|^4 \geq h_\varepsilon(x_0) - u(y_0) + c\|x_0 - y_0\|^4, \quad x, y \in \bar{B},$$

we see that  $h_\varepsilon \geq \varphi$ , where

$$\varphi(x) = h_\varepsilon(x_0) + c(\|x_0 - y_0\|^4 - \|x - y_0\|^4), \quad x \in \bar{B}.$$

Further,  $h_\varepsilon - \varphi$  is smooth and attains its minimum value at  $x_0$ , so

$$\frac{\partial^2(h_\varepsilon - \varphi)}{\partial x_i^2}(x_0) \geq 0, \quad i = 1, \dots, N$$

and hence

$$\Delta\varphi(x_0) \leq \Delta h_\varepsilon(x_0) = -\varepsilon.$$

In particular,  $x_0 \neq y_0$  since  $\Delta\varphi(y_0) = 0$ .

Similarly, setting  $x = x_0$  in (1), we see that  $u \leq \psi$ , where

$$\psi(y) = u(y_0) + c(\|x_0 - y\|^4 - \|x_0 - y_0\|^4), \quad y \in \bar{B}.$$

Since  $u - \psi$  is  $C^1$  and attains its maximum value 0 at  $y_0$ , and also  $x_0 \neq y_0$ , we see that  $\nabla u(y_0) = \nabla\psi(y_0) \neq 0$ . By hypothesis, the formula

$$v(s) = w(x_0 + s, y_0 + s) = h_\varepsilon(x_0 + s) - u(y_0 + s) + c\|x_0 - y_0\|^4$$

defines a function which is superharmonic on some neighbourhood of 0 in  $\mathbb{R}^N$ . Since  $v$  attains a local minimum at 0, it must be constant near 0. However, this leads to the contradictory conclusion that  $\Delta u = -\varepsilon < 0$  near  $y_0$ .

The theorem now follows, because

$$\min_{\bar{B}}(h_\varepsilon - u) = \min_{x \in \bar{B}} w(x, x) \geq \min_{\bar{B} \times \bar{B}} w = \min_{\partial(B \times B)} w \geq 0.$$

□

### References

- [1] *D. H. Armitage, S. J. Gardiner*: Classical Potential Theory. Springer Monographs in Mathematics. Springer, London, 2001. [zbl](#) [MR](#) [doi](#)
- [2] *L. A. Caffarelli, X. Cabré*: Fully Nonlinear Elliptic Equations. Colloquium Publications 43. AMS, Providence, 1995. [zbl](#) [MR](#) [doi](#)
- [3] *M. G. Crandall, H. Ishii, P.-L. Lions*: User's guide to viscosity solutions of second order partial differential equations. Bull. Am. Math. Soc., New Ser. 27 (1992), 1–67. [zbl](#) [MR](#) [doi](#)
- [4] *P. Juutinen, P. Lindqvist*: A theorem of Radó's type for the solutions of a quasi-linear equation. Math. Res. Lett. 11 (2004), 31–34. [zbl](#) [MR](#) [doi](#)
- [5] *P. Juutinen, P. Lindqvist, J. J. Manfredi*: On the equivalence of viscosity solutions and weak solutions for a quasi-linear equation. SIAM J. Math. Anal. 33 (2001), 699–717. [zbl](#) [MR](#) [doi](#)
- [6] *J. Král*: Some extension results concerning harmonic functions. J. Lond. Math. Soc., II. Ser. 28 (1983), 62–70. [zbl](#) [MR](#) [doi](#)
- [7] *J. Král*: A conjecture concerning subharmonic functions. Čas. Pěst. Mat. 110 (1985), page 415. (In Czech.)
- [8] *A. V. Pokrovskii*: A simple proof of the Radó and Král theorems on removability of the zero locus for analytic and harmonic functions. Dopov. Nats. Akad. Nauk Ukr., Mat. Pryr. Tekh. Nauky 2015 (2015), 29–31. [zbl](#) [MR](#) [doi](#)
- [9] *W. Rudin*: Real and Complex Analysis. McGraw-Hill Book Co., New York, 1987. [zbl](#) [MR](#)

*Authors' addresses:* *Stephen J. Gardiner*, School of Mathematics and Statistics, University College Dublin, Science Centre-North Belfield, Dublin 4, Ireland, e-mail: [stephen.gardiner@ucd.ie](mailto:stephen.gardiner@ucd.ie); *Tomas Sjödin*, Department of Mathematics, Campus Valla, House B, Room 3A:681, Linköping University, 581 83, Linköping, Sweden, e-mail: [tomas.sjodin@liu.se](mailto:tomas.sjodin@liu.se).