

Applications of Mathematics

Duong Trong Luyen

Nontrivial solutions to boundary value problems for semilinear Δ_γ -differential equations

Applications of Mathematics, Vol. 66 (2021), No. 4, 461–478

Persistent URL: <http://dml.cz/dmlcz/148968>

Terms of use:

© Institute of Mathematics AS CR, 2021

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

NONTRIVIAL SOLUTIONS TO BOUNDARY VALUE PROBLEMS
FOR SEMILINEAR Δ_γ -DIFFERENTIAL EQUATIONS

DUONG TRONG LUYEN, Ho Chi Minh City

Received December 15, 2019. Published online February 8, 2021.

Abstract. In this article, we study the existence of nontrivial weak solutions for the following boundary value problem:

$$-\Delta_\gamma u = f(x, u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

where Ω is a bounded domain with smooth boundary in \mathbb{R}^N , $\Omega \cap \{x_j = 0\} \neq \emptyset$ for some j , Δ_γ is a subelliptic linear operator of the type

$$\Delta_\gamma := \sum_{j=1}^N \partial_{x_j} (\gamma_j^2 \partial_{x_j}), \quad \partial_{x_j} := \frac{\partial}{\partial x_j}, \quad N \geq 2,$$

where $\gamma(x) = (\gamma_1(x), \gamma_2(x), \dots, \gamma_N(x))$ satisfies certain homogeneity conditions and degenerates at the coordinate hyperplanes and the nonlinearity $f(x, \xi)$ is of subcritical growth and does not satisfy the Ambrosetti-Rabinowitz (AR) condition.

Keywords: Δ_γ -Laplace problem; Cerami condition; variational method; weak solution; Mountain Pass Theorem

MSC 2020: 35J70, 35J20, 35J25, 35D30

1. INTRODUCTION

In the last decades, the boundary value problem for semilinear elliptic equations

$$(1.1) \quad -\Delta u = f(x, u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

has been studied by many authors. The following AR condition, introduced in [2],

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.02–2020.13.

(AR) For some $\theta > 2$, and $R > 0$, we have

$$\theta F(x, \xi) \leq f(x, \xi)\xi \quad \forall |\xi| \geq R, \quad \forall x \in \Omega,$$

where $F(x, \xi) = \int_0^\xi f(x, \tau) d\tau$, plays an important role in their studies. With this (AR) condition, one can use the classical version of the Mountain Pass Theorem of Ambrosetti and Rabinowitz to study the existence of solutions (see, for example, [1], [2], [28] and references therein). Although the (AR) condition is quite natural and important, there are many problems where the nonlinear term $f(x, \xi)$ does not satisfy the (AR) condition, for example,

$$f(x, \xi) = \xi \ln(1 + |\xi|).$$

For this reason, in recent years, some authors have studied problem (1.1) by trying to drop the (AR) condition, for instance, Schechter and Zou [29], Miyagaki and Souto [27], Lam and Lu [14], [15], Liu [18], Liu and Wang [19].

Boundary value problems for nonlinear degenerate elliptic differential equations were treated in [10] and subsequently in [8], [11]. In [33], the critical exponent phenomenon was observed for a model of the Grushin-type operators. The results were then generalized in [30] for a large class of semilinear degenerate elliptic differential equations. Recently, in [31], Thuy and Tri have extended the research to a more complicated class of nonlinear degenerate elliptic differential operators. Very recently, in [12] the authors investigated the Δ_γ -Laplace operator under the additional assumption that the operator is homogeneous of degree two with respect to a semigroup of dilations in \mathbb{R}^N .

In this paper, we study the existence of nontrivial weak solutions to the following problem:

$$(1.2) \quad -\Delta_\gamma u = f(x, u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

where Ω is a bounded domain with smooth boundary in \mathbb{R}^N , $\Omega \cap \{x_j = 0\} \neq \emptyset$ for some j , and Δ_γ is a subelliptic operator of the form

$$\Delta_\gamma := \sum_{j=1}^N \partial_{x_j} (\gamma_j^2 \partial_{x_j}), \quad \gamma = (\gamma_1, \gamma_2, \dots, \gamma_N): \mathbb{R}^N \rightarrow \mathbb{R}^N.$$

Here, the functions $\gamma_j: \mathbb{R}^N \rightarrow \mathbb{R}$ are assumed to be continuous, different from zero and of class C^1 in $\mathbb{R}^N \setminus \Pi$, where

$$\Pi := \left\{ x = (x_1, x_2, \dots, x_N) \in \mathbb{R}^N : \prod_{j=1}^N x_j = 0 \right\}.$$

Moreover, we assume the following properties:

(i) There exists a semigroup of dilations $\{\delta_t\}_{t>0}$ such that

$$\begin{aligned} \delta_t: \mathbb{R}^N &\longrightarrow \mathbb{R}^N, \\ (x_1, \dots, x_N) &\longmapsto \delta_t(x_1, \dots, x_N) = (t^{\varepsilon_1} x_1, \dots, t^{\varepsilon_N} x_N), \end{aligned}$$

where $1 = \varepsilon_1 \leq \varepsilon_2 \leq \dots \leq \varepsilon_N$, such that γ_j is δ_t -homogeneous of degree $\varepsilon_j - 1$, i.e.,

$$\gamma_j(\delta_t(x)) = t^{\varepsilon_j - 1} \gamma_j(x) \quad \forall x \in \mathbb{R}^N, \quad \forall t > 0, \quad j = 1, \dots, N.$$

The number

$$\tilde{N} := \sum_{j=1}^N \varepsilon_j$$

is called the *homogeneous dimension* of \mathbb{R}^N with respect to $\{\delta_t\}_{t>0}$.

(ii)

$$\gamma_1 = 1, \quad \gamma_j(x) = \gamma_j(x_1, x_2, \dots, x_{j-1}), \quad j = 2, \dots, N.$$

(iii) There exists a constant $\varrho \geq 0$ such that

$$0 \leq x_k \partial_{x_k} \gamma_j(x) \leq \varrho \gamma_j(x) \quad \forall k \in \{1, 2, \dots, j-1\} \quad \forall j = 2, \dots, N,$$

and for every $x \in \overline{\mathbb{R}}_+^N := \{(x_1, \dots, x_N) \in \mathbb{R}^N : x_j \geq 0 \quad \forall j = 1, 2, \dots, N\}$.

(iv) Equalities $\gamma_j(x) = \gamma_j(x^*)$ ($j = 1, 2, \dots, N$) are satisfied for every $x \in \mathbb{R}^N$, where

$$x^* = (|x_1|, \dots, |x_N|) \quad \text{if} \quad x = (x_1, x_2, \dots, x_N).$$

The class of Δ_γ -operators includes many degenerate elliptic operators such as the Grushin-type operator

$$G_\alpha := \Delta_x + |x|^{2\alpha} \Delta_y, \quad \alpha \geq 0,$$

where (x, y) denotes a point of $\mathbb{R}^{N_1} \times \mathbb{R}^{N_2}$ (see [9]), and the operator of the form

$$P_{\alpha, \beta} := \Delta_x + \Delta_y + |x|^{2\alpha} |y|^{2\beta} \Delta_z, \quad (x, y, z) \in \mathbb{R}^{N_1} \times \mathbb{R}^{N_2} \times \mathbb{R}^{N_3},$$

where α, β are non-negative real numbers (see [31]). Many aspects of the theory of degenerate elliptic differential operators are presented in monographs [35], [36] (see also some recent results in [3], [7], [13], [20], [21], [22], [23], [24], [26], [32], [34]).

To study problem (1.2), we make the following assumptions:

- (A1) f is a real Carathéodory function on $\Omega \times \mathbb{R}$ such that there exist $q_1 \in (2, 2_\gamma^*)$ (where $2_\gamma^* := 2\tilde{N}/(\tilde{N} - 2)$), $\omega \in \mathcal{K}_{2, q_1}$ (see Definition 2.2) and some positive constant C_0 such that

$$|f(x, \xi)| \leq |\omega(x)\xi|^{q_1-1} + C_0 \quad \forall (x, \xi) \in \Omega \times \mathbb{R};$$

- (A2) there exist $C \in [0, \infty)$ and $h \in L^1(\Omega)$ such that $|f(x, \xi)| \leq h(x)$ for every x in Ω and $|\xi| \leq C$;
- (A3) there exists a non-positive function k in $L^{q_2}(\Omega)$, $q_2 > \tilde{N}/2$ such that $k(x) \leq f(x, \xi)/\xi$ for every $(x, \xi) \in \Omega \times \mathbb{R}$;
- (A4) $f(x, 0) = 0$ for every x in Ω and the following limit holds uniformly for a.e. $x \in \Omega$:

$$\lim_{\xi \rightarrow 0} \frac{f(x, \xi)}{\xi} = 0;$$

- (A5) $\lim_{\xi \rightarrow \infty} f(x, \xi)/\xi = \infty$ or $\lim_{\xi \rightarrow -\infty} f(x, \xi)/\xi = \infty$ a.e. in Ω ;
- (A6) $f(x, \xi)/\xi$ is increasing in $\xi \geq C$ and decreasing in $\xi \leq -C$ for every x in Ω .

Our main result is given by the following theorem.

Theorem 1.1. *Suppose that f satisfies (A1)–(A6). Then the boundary value problem (1.2) has a nontrivial weak solution.*

Remark 1.2. Suppose that f is continuous on $\overline{\Omega} \times \mathbb{R}$ and satisfies the following conditions:

- (A1') There exist $q_3 \in (2, 2_\gamma^*)$ and a positive real number C_1 such that

$$|f(x, \xi)| \leq C_1(1 + |\xi|^{q_3-1}) \quad \forall (x, \xi) \in \Omega \times \mathbb{R}.$$

- (A4') $f(x, 0) = 0$ for every x in Ω and $\lim_{\xi \rightarrow 0} f(x, \xi)/\xi = 0$ uniformly in Ω .

- (A5') $\lim_{\xi \rightarrow \infty} f(x, \xi)/\xi = \infty$ or $\lim_{\xi \rightarrow -\infty} f(x, \xi)/\xi = \infty$ uniformly in Ω .

Then f satisfies (A1)–(A5). Therefore, our theorem improves the corresponding results in [31], [35], [36].

This article is organized as follows. In Section 2 we present some definitions and preliminary results. In Section 3 we give the proof of our results.

2. PRELIMINARY RESULTS

2.1. Function spaces and embedding theorem. For a function u of class C^1 , we put

$$\nabla_\gamma u := (\gamma_1 \partial_{x_1} u, \gamma_2 \partial_{x_2} u, \dots, \gamma_N \partial_{x_N} u), \quad |\nabla_\gamma u| := \left(\sum_{j=1}^N |\gamma_j \partial_{x_j} u|^2 \right)^{1/2}.$$

If Ω is a bounded domain in \mathbb{R}^N , we denote by

$$S_{\gamma,0}^2(\Omega)$$

the closure of $C_0^1(\Omega)$ with respect to the norm

$$\|u\|_{S_{\gamma,0}^2(\Omega)} := \left(\int_\Omega |\nabla_\gamma u|^2 dx \right)^{1/2}.$$

From Proposition 3.2 and Theorem 3.3 in [12], we have the following embedding result.

Proposition 2.1. *Assume Ω is a bounded domain in \mathbb{R}^N , $\tilde{N} > 2$. Then the embedding $S_{\gamma,0}^2(\Omega) \hookrightarrow L^p(\Omega)$ is continuous for any $p \in [1, 2_\gamma^*]$, i.e., there exists $C_p > 0$ such that $\|u\|_{L^p(\Omega)} \leq C_p \|u\|_{S_{\gamma,0}^2(\Omega)}$ for all $u \in S_{\gamma,0}^2(\Omega)$. Moreover, $S_{\gamma,0}^2(\Omega)$ is compactly embedded into $L^p(\Omega)$ only for $p \in [1, 2_\gamma^*)$, where $L^p(\Omega)$ denotes the Lebesgue space with the standard norm $\|\cdot\|_{L^p(\Omega)}$.*

Definition 2.2. Let σ be a measurable function on Ω . We put

$$T_\sigma u = \sigma u \quad \forall u \in S_{\gamma,0}^2(\Omega).$$

We say that

- (i) σ is of class $\mathcal{C}_{2,p}$, if T_σ is a continuous mapping from $S_{\gamma,0}^2(\Omega)$ into $L^p(\Omega)$,
- (ii) σ is of class $\mathcal{K}_{2,p}$, if T_σ is a compact mapping from $S_{\gamma,0}^2(\Omega)$ into $L^p(\Omega)$.

We have the following results.

Theorem 2.3. *Let us assume that $\omega_1 \in \mathcal{C}_{2,p_1}$, $\omega_2 \in \mathcal{C}_{2,p_2}$ such that ω_1 and ω_2 are non-negative, where $p_1, p_2 \in [1, \infty)$, $p_1 < p_2$. Put*

$$\omega = \omega_1^{p_1(p_2-p)/(p(p_2-p_1))} \omega_2^{p_2(p-p_1)/(p(p_2-p_1))} \quad \text{for any } p \in (p_1, p_2).$$

Then $\omega \in \mathcal{C}_{2,p}$.

P r o o f. From $\omega_1 \in \mathcal{C}_{2,p_1}$, $\omega_2 \in \mathcal{C}_{2,p_2}$, hence there exists a positive real number C_2 such that

$$(2.1) \quad \left(\int_{\Omega} \omega_i^{p_i} |u|^{p_i} dx \right)^{1/p_i} \leq C_2 \|u\|_{S_{\gamma,0}^2(\Omega)} \quad \forall u \in S_{\gamma,0}^2(\Omega), \quad i = 1, 2.$$

By $p = p_1(p_2 - p)/(p_2 - p_1) + p_2(p - p_1)/(p_2 - p_1)$, applying Hölder's inequality and (2.1), we get

$$\begin{aligned} \left(\int_{\Omega} \omega^p |u|^p dx \right)^{1/p} &= \left(\int_{\Omega} \omega_1^{p_1(p_2-p)/(p_2-p_1)} |u|^{p_1(p_2-p)/(p_2-p_1)} \right. \\ &\quad \left. \times \omega_2^{p_2(p-p_1)/(p_2-p_1)} |u|^{p_2(p-p_1)/(p_2-p_1)p_2} dx \right)^{1/p} \\ &\leq \left\{ \left(\int_{\Omega} \omega_1^{p_1} |u|^{p_1} dx \right)^{(p_2-p)/(p_2-p_1)} \left(\int_{\Omega} \omega_2^{p_2} |u|^{p_2} dx \right)^{(p-p_1)/(p_2-p_1)} \right\}^{1/p} \\ &\leq C_2 \|u\|_{S_{\gamma,0}^2(\Omega)} \quad \forall u \in S_{\gamma,0}^2(\Omega). \end{aligned}$$

The proof of Theorem 2.3 is complete. \square

Theorem 2.4. *Let us assume that $\omega \in \mathcal{C}_{2,p}$ and θ are measurable functions on Ω such that $\omega \geq 0$ and $|\theta| \leq \omega^\alpha$, where $p \in [1, 2_\gamma^*)$, $\alpha \in (0, 1)$. Then θ is of class $\mathcal{K}_{2,p}$.*

P r o o f. Since ω is of class $\mathcal{C}_{2,p}$, T_ω is continuous from $S_{\gamma,0}^2(\Omega)$ into $L^p(\Omega)$ and by Proposition 2.1, we have

$$(2.2) \quad \left(\int_{\Omega} |u|^p \omega^p dx \right)^{1/p} \leq C_p \|u\|_{S_{\gamma,0}^2(\Omega)} \quad \forall u \in S_{\gamma,0}^2(\Omega).$$

By $\omega^\alpha(x) \leq 1 + \omega(x)$ for every x in Ω and $1, \omega \in \mathcal{C}_{2,p}$ hence ω^α belongs to $\mathcal{C}_{2,p}$. Thus θ is of class $\mathcal{C}_{2,p}$. Let M be a positive real number and $\{u_n\}_{n=1}^\infty$ be a sequence in $S_{\gamma,0}^2(\Omega)$, such that $\|u_n\|_{S_{\gamma,0}^2(\Omega)} \leq M$ for all n . By Proposition 2.1, we have (by passing to a subsequence if necessary)

$$(2.3) \quad \begin{aligned} u_n &\rightharpoonup u \text{ weakly in } S_{\gamma,0}^2(\Omega) \text{ as } n \rightarrow \infty, \\ u_n &\rightarrow u \text{ strongly in } L^p(\Omega) \text{ as } n \rightarrow \infty. \end{aligned}$$

Therefore

$$\|u\|_{S_{\gamma,0}^2(\Omega)} \leq \liminf_{n \rightarrow \infty} \|u_n\|_{S_{\gamma,0}^2(\Omega)} \leq M.$$

We shall prove that $\{T_\theta(u_n)\}_{n=1}^\infty$ converges to $T_\theta(u)$ in $L^p(\Omega)$. Let ε be a positive real number. Choose a positive real number δ such that

$$(2.4) \quad (2C_p M)^p \delta^{(\alpha-1)p} < \frac{\varepsilon^p}{2}.$$

Put $\Omega' = \{x \in \Omega : \omega(x) > \delta\}$. From (2.2) and (2.4), we get that

$$\begin{aligned}
 (2.5) \quad \int_{\Omega} |\theta(u_n - u)|^p dx &= \int_{\Omega} |u_n - u|^p |\theta|^p dx \\
 &\leq \int_{\Omega'} |u_n - u|^p \omega^{\alpha p} dx + \int_{\Omega \setminus \Omega'} |u_n - u|^p \omega^{\alpha p} dx \\
 &\leq \delta^{(\alpha-1)p} \int_{\Omega'} |u_n - u|^p \omega^p dx + \delta^{\alpha p} \int_{\Omega \setminus \Omega'} |u_n - u|^s dx \\
 &\leq \delta^{(\alpha-1)p} \int_{\Omega} |u_n - u|^p \omega^p dx + \delta^{\alpha p} \int_{\Omega} |u_n - u|^p dx \\
 &\leq \delta^{(\alpha-1)p} (C_p \|u_n - u\|_{S_{\gamma,0}^2(\Omega)})^p + \delta^{\alpha p} \int_{\Omega} |u_n - u|^p dx \\
 &\leq \delta^{(\alpha-1)p} (2C_p M)^p + \delta^{\alpha p} \int_{\Omega} |u_n - u|^p dx \\
 &\leq \frac{\varepsilon^p}{2} + \delta^{p\alpha} \int_{\Omega} |u_n - u|^p dx.
 \end{aligned}$$

By (2.3), there exists an integer n_0 such that

$$(2.6) \quad \int_{\Omega} |u_n - u|^p dx \leq \delta^{-\alpha p} \frac{\varepsilon^p}{2} \quad \forall n \geq n_0.$$

Combining (2.5) and (2.6), we complete the proof. \square

Corollary 2.5. *Let $p \in [1, 2_{\gamma}^*)$, $\eta \in (2p\tilde{N}/(2\tilde{N} - p(\tilde{N} - 2)), \infty)$ and $\theta \in L^{\eta}(\Omega)$. Then θ is of class $\mathcal{K}_{2,p}$.*

Proof. Let $\alpha \in (0, 1)$ such that $\alpha\eta = 2p\tilde{N}/(2\tilde{N} - p(\tilde{N} - 2))$ and $\omega = |\theta|^{1/\alpha}$. Then ω is in $L^{2p\tilde{N}/(2\tilde{N} - p(\tilde{N} - 2))}(\Omega)$. By

$$\frac{2\tilde{N} - p(\tilde{N} - 2)}{2\tilde{N}} + \frac{p(\tilde{N} - 2)}{2\tilde{N}} = 1,$$

applying Hölder's inequality, we obtain

$$\begin{aligned}
 \int_{\Omega} |\omega u|^p dx &\leq \left(\int_{\Omega} |\omega|^{2p\tilde{N}/(2\tilde{N} - p(\tilde{N} - 2))} dx \right)^{2\tilde{N} - p(\tilde{N} - 2)/2\tilde{N}} \\
 &\quad \times \left(\int_{\Omega} |u|^{2\tilde{N}/(\tilde{N} - 2)} dx \right)^{p(\tilde{N} - 2)/2\tilde{N}} \quad \forall u \in S_{\gamma,0}^2(\Omega),
 \end{aligned}$$

which implies that T_{ω} is continuous at 0 in $S_{\gamma,0}^2(\Omega)$. Thus T_{ω} is a linear continuous map from $S_{\gamma,0}^2(\Omega)$ into $L^p(\Omega)$. By Theorem 2.4, we have θ is of class $\mathcal{K}_{2,p}$. \square

Example 2.6. Let $\gamma = (1, 1, |x_1 x_2|)$, $\Omega = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3: |x| < 1\}$ and $p = 3$. Then $N = 3$, $\tilde{N} = 5$; $2p\tilde{N}/(2\tilde{N} - p(\tilde{N} - 2)) = 2 \cdot 3 \cdot 5/(5 \cdot 2 - 3(5 - 2)) = 30 < 40$. Put $\omega_0 = |x|^{-1/50} \sin(2017|x|)$, then ω_0 is in $L^{40}(\Omega)$. Thus by Corollary 2.5, ω_0 is of class $\mathcal{K}_{2,3}$.

Corollary 2.7. Let $p \in [1, 2_\gamma^*)$, α be in $(0, 1)$ and $\eta \in \mathcal{C}_{2,2}$. Then

$$\theta = \eta^{\alpha 2(2_\gamma^* - p)/(p(2_\gamma^* - 2))}$$

is of class $\mathcal{K}_{2,p}$.

Proof. Put $\omega_1 = \eta$, $\omega_2 = 1$, $p_1 = 2$, $p_2 = 2_\gamma^*$. By Proposition 2.1, $\omega_2 \in \mathcal{C}_{2,2_\gamma^*}$. By Theorem 2.3, we see that $\eta^{2(2_\gamma^* - p)/(p(2_\gamma^* - 2))} \in \mathcal{C}_{2,p}$. Thus by Theorem 2.4, $\eta^{\alpha 2(2_\gamma^* - p)/(p(2_\gamma^* - 2))}$ is of class $\mathcal{K}_{2,p}$. \square

Example 2.8. Let $\gamma = (1, 1, |x_1 x_2|)$, $\Omega = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3: |x| < 1\}$, $p = 3$, $\alpha = \frac{3}{4}$ and $\eta(x) = (1 - |x|^2)^{-1/2}$ for every x in Ω , hence $\eta \in \mathcal{C}_{2,2}$, $\tilde{N} = 5$. Note that $2_\gamma^* = 2\tilde{N}/(\tilde{N} - 2) = \frac{10}{3}$ and

$$\alpha \frac{2(2_\gamma^* - p)}{p(2_\gamma^* - 2)} = 2.$$

Put $\theta(x) = (1 - |x|^2)^{-1}$. Then $\theta \in \mathcal{K}_{2,3}$.

Example 2.9. Let $\gamma = (1, 1, |x_1 x_2|)$, $\Omega = \{x \in \mathbb{R}^3: |x| < 1\}$, $p = 3$, $\alpha = \frac{3}{4}$ and $\varrho(x) = (\frac{1}{2} - |x|^2)^2(1 - |x|^2)^{-1}$ for every x in Ω . By Example 2.8, $\varrho \in \mathcal{K}_{2,3}$. Put $a(x) = \varrho(x)^2 = (\frac{1}{2} - |x|^2)^4(1 - |x|^2)^{-2}$. Thus a is not integrable on Ω .

Theorem 2.10. Let $p \in [1, 2_\gamma^*)$, ω be in $\mathcal{K}_{2,p}$, a function $g \in L^{p/(p-1)}(\Omega)$ and f be a Carathéodory function from $\Omega \times \mathbb{R}$ into \mathbb{R} . Assume

$$|f(x, \xi)| \leq |\omega(x)|^{p-1} |\xi|^{p-1} + g(x) \quad \forall (x, \xi) \in \Omega \times \mathbb{R}.$$

Then $\Phi_1(u) \in C^1(S_{\gamma,0}^2(\Omega), \mathbb{R})$ and

$$\langle \Phi_1'(u), v \rangle = \int_{\Omega} f(x, u(x))v(x) dx$$

for all $v \in S_{\gamma,0}^2(\Omega)$, where

$$\Phi_1(u) = \int_{\Omega} F(x, u(x)) dx,$$

and $F(x, \xi) = \int_0^\xi f(x, \tau) d\tau$.

Proof. With a slight modification, the proof of this lemma is similar to the one of Lemma 2.3 in [25]. We omit the details. \square

Define the Euler-Lagrange functional associated with problem (1.2) as follows:

$$\Phi(u) = \frac{1}{2} \int_{\Omega} |\nabla_{\gamma} u|^2 dx - \int_{\Omega} F(x, u(x)) dx.$$

From Theorem 2.10 and the fact that f satisfies (A1), we have that Φ is well-defined on $S_{\gamma,0}^2(\Omega)$ and $\Phi \in C^1(S_{\gamma,0}^2(\Omega), \mathbb{R})$ with

$$\langle \Phi'(u), v \rangle = \int_{\Omega} \nabla_{\gamma} u \cdot \nabla_{\gamma} v dx - \int_{\Omega} f(x, u(x))v(x) dx$$

for all $v \in S_{\gamma,0}^2(\Omega)$.

Recall that a function $u \in S_{\gamma,0}^2(\Omega)$ is called a weak solution of problem (1.2) if

$$\int_{\Omega} \nabla_{\gamma} u \cdot \nabla_{\gamma} v dx = \int_{\Omega} f(x, u(x))v(x) dx \quad \forall v \in S_{\gamma,0}^2(\Omega).$$

Hence, the weak solutions of problem (1.2) are critical points of the functional Φ .

2.2. Mountain Pass Theorem.

Definition 2.11. Let \mathbf{X} be a real Banach space with its dual space \mathbf{X}^* and $\Phi \in C^1(\mathbf{X}, \mathbb{R})$. For $c \in \mathbb{R}$ we say that Φ satisfies the $(C)_c$ condition if for any sequence $\{u_n\}_{n=1}^{\infty} \subset \mathbf{X}$ with

$$\Phi(u_n) \rightarrow c \quad \text{and} \quad (1 + \|u_n\|_{\mathbf{X}})\|\Phi'(u_n)\|_{\mathbf{X}^*} \rightarrow 0,$$

there exists a subsequence $\{u_{n_k}\}_{k=1}^{\infty}$ that converges strongly in \mathbf{X} .

We will use the following version of the Mountain Pass Theorem:

Lemma 2.12 (see [5], [6]). *Let \mathbf{X} be a real Banach space and let $\Phi \in C^1(\mathbf{X}, \mathbb{R})$ satisfy the $(C)_c$ condition for any $c \in \mathbb{R}$, $\Phi(0) = 0$ and*

- (i) *there exist constants $\varrho, \alpha > 0$ such that $\Phi(u) \geq \alpha$ for all $u \in \mathbf{X}$, $\|u\|_{\mathbf{X}} = \varrho$;*
- (ii) *there exists a $u_1 \in \mathbf{X}$, $\|u_1\|_{\mathbf{X}} \geq \varrho$ such that $\Phi(u_1) \leq 0$.*

Then $\beta := \inf_{\lambda \in \Lambda} \max_{0 \leq t \leq 1} \Phi(\lambda(t)) \geq \alpha$ is a critical value of Φ , where

$$\Lambda := \{\lambda \in C([0; 1], \mathbf{X}) : \lambda(0) = 0, \lambda(1) = u_1\}.$$

3. PROOF OF THE MAIN THEOREM

We prove Theorem 1.1 by verifying that all conditions of Lemma 2.12 are satisfied. First, we check condition (i) in Lemma 2.12.

Lemma 3.1. *Assume that f satisfies conditions (A1) and (A4). Then there exist $\varrho, \alpha > 0$ such that*

$$\Phi(u) \geq \alpha \quad \forall u \in S_{\gamma,0}^2(\Omega), \quad \|u\|_{S_{\gamma,0}^2(\Omega)} = \varrho.$$

Proof. Suppose by contradiction that

$$\inf \left\{ \Phi(u) : u \in S_{\gamma,0}^2(\Omega), \|u\|_{S_{\gamma,0}^2(\Omega)} = \frac{1}{n} \right\} \leq 0 \quad \forall n \in \mathbb{N}.$$

Then, there exists a sequence $\{u_n\}_{n=1}^\infty$ in $S_{\gamma,0}^2(\Omega)$ such that $\|u_n\|_{S_{\gamma,0}^2(\Omega)} = 1/n$ and $\Phi(u_n) < 1/n^3$. Hence, we have

$$\frac{1}{n} > \frac{\Phi(u_n)}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}^2} = \frac{1}{2} - \int_{\Omega} \frac{F(x, u_n(x))}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}^2} dx,$$

and thus

$$(3.1) \quad \int_{\Omega} \frac{F(x, u_n(x))}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}^2} dx > \frac{1}{2} - \frac{1}{n}.$$

By (A4), for each $\varepsilon > 0$, we can find a number $\delta > 0$ such that

$$(3.2) \quad |F(x, \xi)| \leq \varepsilon \xi^2 \quad \text{for } |\xi| \leq \delta.$$

From (A1), we deduce that

$$(3.3) \quad |F(x, \xi)| \leq \frac{1}{q_1} |\omega(x)|^{q_1-1} |\xi|^{q_1} + C(\delta) |\xi|^{q_1} \quad \text{for } |\xi| \geq \delta.$$

It follows from (3.2), (3.3), Proposition 2.1 and Hölder's inequality that

$$\begin{aligned} \left| \int_{\Omega} F(x, u_n(x)) dx \right| &\leq \varepsilon \int_{\Omega} |u_n(x)|^2 dx + \frac{1}{q_1} \int_{\Omega} |\omega(x)|^{q_1-1} |u_n(x)|^{q_1} dx \\ &\quad + C(\delta) \int_{\Omega} |u_n(x)|^{q_1} dx \\ &\leq \frac{1}{q_1} \left(\int_{\Omega} |\omega(x) u_n(x)|^{q_1} dx \right)^{(q_1-1)/q_1} \left(\int_{\Omega} |u_n(x)|^{q_1} dx \right)^{1/q_1} \\ &\quad + \varepsilon \|u_n\|_{L^2(\Omega)}^2 + C(\delta) \|u_n\|_{L^{q_1}(\Omega)}^{q_1} \\ &\leq \varepsilon C_2^2 \|u_n\|_{S_{\gamma,0}^2(\Omega)}^2 + \frac{C_{q_1}^{q_1}}{q_1} \|u_n\|_{S_{\gamma,0}^2(\Omega)}^{q_1} + C(\delta) C_{q_1}^{q_1} \|u_n\|_{S_{\gamma,0}^2(\Omega)}^{q_1}, \end{aligned}$$

hence

$$\left| \int_{\Omega} \frac{F(x, u_n(x))}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}^2} dx \right| \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

which yields a contradiction to (3.1). Lemma 3.1 is proved. \square

Next, we check condition (ii) in Lemma 2.12.

Lemma 3.2. *Let ϱ be as in Lemma 3.1 and assume that f satisfies conditions (A3) and (A5). Then there exists u_1 in $S_{\gamma,0}^2(\Omega) \setminus B(0, \varrho)$ such that $\Phi(u_1) < 0$.*

Proof. We limit ourselves to consider the case

$$(3.4) \quad \lim_{\xi \rightarrow \infty} \frac{f(x, \xi)}{\xi} = \infty.$$

Take a point $u \in S_{\gamma,0}^2(\Omega)$ such that $\|u\|_{S_{\gamma,0}^2(\Omega)} = 1$, $u > 0$ and $\|u\|_{L^2(\Omega)} \neq 0$. Then, for any constant $R > 0$

$$\Phi(Ru) = \frac{R^2}{2} - \int_{\Omega} F(x, Ru(x)) dx.$$

Since we are assuming (3.4), there exists a number $M > 0$ such that $f(x, \xi) \geq 4\xi/\|u\|_{L^2(\Omega)}^2$ for $\xi \geq M$; moreover, from (A3) we get

$$\begin{aligned} F(x, \xi) &\geq \int_0^M k(x)\tau d\tau = k(x)\frac{M^2}{2} \quad \text{for } 0 \leq \xi \leq M, \\ F(x, \xi) &= \int_0^M f(x, \tau) d\tau + \int_M^\xi f(x, \tau) d\tau \\ &\geq k(x)\frac{M^2}{2} + \frac{2\xi^2}{\|u\|_{L^2(\Omega)}^2} - \frac{2M^2}{\|u\|_{L^2(\Omega)}^2} \quad \text{for } \xi \geq M. \end{aligned}$$

As $0 \leq Ru \leq M$ on $\Omega_{M/R}^u := \{x \in \Omega: |u(x)| \leq M/R\}$, we have

$$\begin{aligned} \int_{\Omega} F(x, Ru(x)) dx &= \int_{\Omega_{M/R}^u} F(x, Ru(x)) dx + \int_{\Omega \setminus \Omega_{M/R}^u} F(x, Ru(x)) dx \\ &\geq \int_{\Omega \setminus \Omega_{M/R}^u} \frac{2R^2 u^2(x)}{\|u\|_{L^2(\Omega)}^2} dx + M^2 \int_{\Omega} k(x) dx - \frac{2M^2}{\|u\|_{L^2(\Omega)}^2} \text{meas}(\Omega), \end{aligned}$$

where $\text{meas}(\cdot)$ denotes the Lebesgue measure of a set in \mathbb{R}^N . By Lebesgue's theorem, there exists a number R_0 such that

$$\int_{\Omega \setminus \Omega_{M/R_0}^u} u^2(x) dx \geq \frac{\|u\|_{L^2(\Omega)}^2}{2}.$$

Therefore, if $R \geq R_0$, then

$$\int_{\Omega} F(x, Ru(x)) \, dx \geq R^2 + M^2 \int_{\Omega} k(x) \, dx - \frac{2M^2}{\|u\|_{L^2(\Omega)}^2} \text{meas}(\Omega).$$

Consequently, if

$$R > \max \left\{ 2\sqrt{-M^2 \int_{\Omega} k(x) \, dx + \frac{2M^2}{\|u\|_{L^2(\Omega)}^2} \text{meas}(\Omega)}, R_0 \right\},$$

then

$$\Phi(Ru) \leq -M^2 \int_{\Omega} k(x) \, dx + \frac{2M^2}{\|u\|_{L^2(\Omega)}^2} \text{meas}(\Omega) - \frac{R^2}{2} < 0.$$

Thus, Φ satisfies the condition (ii) in Lemma 2.12. \square

Lemma 3.3. *Assume that f satisfies conditions (A2) and (A6). Then there exists a positive real number C_3 such that*

$$f(x, s)s - 2F(x, s) \leq f(x, t)t - 2F(x, t) + C_3h(x) \quad \forall x \in \Omega, |s| \leq |t|.$$

Proof. Since (A6) holds, by Lemma 2.3 in [18], we have that, for any $x \in \Omega$,

$$\xi \mapsto f(x, \xi)\xi - 2F(x, \xi)$$

is increasing in $\xi \geq C$ and decreasing in $\xi \leq -C$. Hence,

$$f(x, s)s - 2F(x, s) \leq f(x, t)t - 2F(x, t) \quad \forall x \in \Omega, C \leq s \leq t.$$

Let $x \in \Omega$ and $\xi \in [-C, C]$. By (A2), we get that

$$|f(x, \xi)| \leq h(x), \quad |F(x, \xi)| \leq \int_0^{\xi} h(x) \, d\tau \leq Ch(x).$$

Thus,

$$\begin{aligned} f(x, s)s - 2F(x, s) &\leq f(x, t)t - 2F(x, t) + 6Ch(x) && \forall x \in \Omega, 0 \leq s \leq t \leq C, \\ f(x, s)s - 2F(x, s) &\leq f(x, C)C - 2F(x, C) + 6Ch(x) \\ &\leq f(x, t)t - 2F(x, t) + 6Ch(x) && \forall x \in \Omega, 0 \leq s \leq C \leq t. \end{aligned}$$

Thus we get the lemma for $0 \leq s \leq t$. Similarly we obtain it for $t \leq s \leq 0$. The proof of Lemma 3.3 is complete. \square

We now show the main lemma of this paper.

Lemma 3.4. *Assume that f satisfies conditions (A1)–(A3), (A5) and (A6). Then Φ satisfies the $(C)_c$ condition for all $c \in \mathbb{R}$.*

Proof. Let $\{u_n\}_{n=1}^\infty \subset S_{\gamma,0}^2(\Omega)$ be a $(C)_c$ sequence, i.e.,

$$(3.5) \quad \Phi(u_n) \rightarrow c \quad \text{as } n \rightarrow \infty, \quad \lim_{n \rightarrow \infty} (1 + \|u_n\|_{S_{\gamma,0}^2(\Omega)}) \|\Phi'(u_n)\|_{(S_{\gamma,0}^2(\Omega))^*} = 0,$$

hence,

$$(3.6) \quad \begin{aligned} \lim_{n \rightarrow \infty} \int_{\Omega} \left(\frac{1}{2} f(x, u_n(x)) u_n(x) - F(x, u_n(x)) \right) dx \\ = \lim_{n \rightarrow \infty} \left(\Phi(u_n) - \frac{1}{2} \langle \Phi'(u_n), u_n \rangle \right) = c. \end{aligned}$$

We first show that $\{u_n\}_{n=1}^\infty$ is bounded in $S_{\gamma,0}^2(\Omega)$ by a contradiction argument. By passing to a subsequence if necessary, we can assume that $\|u_n\|_{S_{\gamma,0}^2(\Omega)} > 1$ and

$$(3.7) \quad \|u_n\|_{S_{\gamma,0}^2(\Omega)} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Setting

$$w_n = \frac{u_n}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}},$$

we get $\|w_n\|_{S_{\gamma,0}^2(\Omega)} = 1$, so we can extract a subsequence relabelled $\{w_n\}_{n=1}^\infty$ such that $\{w_n\}_{n=1}^\infty$ converges weakly to w in $S_{\gamma,0}^2(\Omega)$. Since Ω is bounded, Proposition 2.1 implies that

$$(3.8) \quad \begin{aligned} w_n &\rightarrow w \text{ strongly in } L^p(\Omega), \quad 1 \leq p < 2_\gamma^* \text{ as } n \rightarrow \infty, \\ w_n &\rightarrow w \text{ a.e. in } \Omega \text{ as } n \rightarrow \infty. \end{aligned}$$

Now, we consider two possible cases: $w = 0$ or $w \neq 0$.

Case 1: If $w = 0$, then for any $n \in \mathbb{N}$ there exists $t_n \in [0, 1]$ such that

$$(3.9) \quad \Phi(t_n u_n) = \max\{\Phi(s u_n) : s \in [0, 1]\}.$$

Fix a positive integer m and put $v_n = (4m)^{1/2} w_n$ for every positive integer n . Therefore, we have

$$(3.10) \quad \begin{aligned} v_n &\rightharpoonup 0 \text{ weakly in } S_{\gamma,0}^2(\Omega) \text{ as } n \rightarrow \infty, \\ v_n &\rightarrow 0 \text{ a.e. in } \Omega \text{ as } n \rightarrow \infty, \\ v_n &\rightarrow 0 \text{ strongly in } L^p(\Omega), \quad 1 \leq p < 2_\gamma^* \text{ as } n \rightarrow \infty, \\ \omega v_n &\rightarrow 0 \text{ strongly in } L^{q_1}(\Omega) \text{ as } n \rightarrow \infty. \end{aligned}$$

By (A1), applying Hölder's inequality, we get

$$\begin{aligned}
(3.11) \quad \left| \int_{\Omega} F(x, v_n(x)) \, dx \right| &\leq \int_{\Omega} |\omega(x)v_n(x)|^{q_1-1} |v_n(x)| \, dx + C_0 \int_{\Omega} |v_n(x)| \, dx \\
&\leq \left(\int_{\Omega} |\omega(x)v_n(x)|^{q_1} \, dx \right)^{(q_1-1)/q_1} \left(\int_{\Omega} |v_n(x)|^{q_1} \, dx \right)^{1/q_1} \\
&\quad + C_0 \int_{\Omega} |v_n(x)| \, dx.
\end{aligned}$$

From (3.10) and (3.11), we have

$$\left| \int_{\Omega} F(x, v_n(x)) \, dx \right| \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

hence,

$$\lim_{n \rightarrow \infty} \int_{\Omega} F(x, v_n(x)) \, dx = 0.$$

Since $\lim_{n \rightarrow \infty} (4m)^{1/2} \|u_n\|_{S_{\gamma,0}^2(\Omega)}^{-1} = 0$, there exists an integer N_m such that

$$\Phi(t_n u_n) \geq \Phi(v_n) = 2m - \int_{\Omega} F(x, v_n(x)) \, dx \geq m \quad \forall n \geq N_m,$$

that is, $\lim_{n \rightarrow \infty} \Phi(t_n u_n) = \infty$. Since $\Phi(0) = 0$ and $\lim_{n \rightarrow \infty} \Phi(u_n) = c$, it implies $t_n \in (0, 1)$ for any sufficiently large n and

$$\begin{aligned}
&\int_{\Omega} |\nabla_{\gamma}(t_n u_n)|^2 \, dx - \int_{\Omega} f(x, t_n u_n(x)) t_n u_n(x) \, dx \\
&= \langle \Phi'(t_n u_n), t_n u_n \rangle = t_n \frac{d}{dt} \Big|_{t=t_n} \Phi(t u_n) = 0.
\end{aligned}$$

Therefore, by Lemma 3.3, we get

$$\begin{aligned}
&\int_{\Omega} \left(\frac{1}{2} f(x, u_n(x)) u_n(x) - F(x, u_n(x)) \right) \, dx \\
&\geq \int_{\Omega} \left(\frac{1}{2} f(x, t_n u_n(x)) t_n u_n(x) - F(x, t_n u_n(x)) \right) \, dx - C_3 \|h\|_{L^1(\Omega)} \\
&= \int_{\Omega} \left(\frac{1}{2} |\nabla_{\gamma} t_n u_n|^2 - F(x, t_n u_n(x)) \right) \, dx - C_3 \|h\|_{L^1(\Omega)} \\
&= \Phi(t_n u_n) - C_3 \|h\|_{L^1(\Omega)} \rightarrow \infty,
\end{aligned}$$

which contradicts (3.6).

Case 2: If $w \neq 0$, the Lebesgue measure of the set $\Theta = \{x \in \Omega: w(x) \neq 0\}$ is positive. We have $\lim_{n \rightarrow \infty} |u_n(x)| = \infty$ for every x in Θ . Hence, by (A5) we deduce

$$(3.12) \quad \frac{f(x, u_n(x))}{u_n(x)} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

By (3.8) and Theorem 4.9 in [4], there exists a function $w_0 \in L^p(\Omega)$ such that

$$|w_n| \leq w_0 \quad \forall n, \text{ a.e. on } \Omega.$$

Now, using condition (A3) we have

$$\frac{f(x, \xi u_n(x))}{\xi u_n(x)} \cdot \frac{\xi |u_n(x)|^2}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}^2} \geq \xi k(x) w_0^2(x) \quad \forall x \in \Omega, \xi \in (0, 1);$$

thus, since $k w_0^2 \in L^1(\Omega)$ (as $k \in L^{q_2}(\Omega)$ for some $q_2 > \tilde{N}/2$), using (3.12), (A3) and the fact that $\Phi(u_n) \rightarrow c$, we deduce via the generalized Fatou lemma that

$$\begin{aligned} \frac{1}{2} &= \liminf_{n \rightarrow \infty} \left[\frac{1}{2} - \frac{\Phi(u_n)}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}^2} \right] = \liminf_{n \rightarrow \infty} \int_{\Omega} \frac{F(x, u_n(x))}{\|u_n\|_{S_{\gamma,0}^2(\Omega)}^2} dx \\ &= \liminf_{n \rightarrow \infty} \int_{\Omega} \int_0^1 \frac{f(x, \xi u_n(x))}{\xi u_n(x)} \cdot \xi |w_n(x)|^2 d\xi dx \quad (\text{since } w \equiv 0 \text{ on } \Omega \setminus \Theta) \\ &= \liminf_{n \rightarrow \infty} \int_{\Theta} \int_0^1 \frac{f(x, \xi u_n(x))}{\xi u_n(x)} \cdot \xi |w_n(x)|^2 d\xi dx \\ &\geq \int_{\Theta} \int_0^1 \liminf_{n \rightarrow \infty} \frac{f(x, \xi u_n(x))}{\xi u_n(x)} \cdot \xi |w_n(x)|^2 d\xi dx = \infty, \end{aligned}$$

which is impossible. In any case, we obtain a contradiction. Thus, the sequence $\{u_n\}_{n=1}^{\infty}$ is bounded in $S_{\gamma,0}^2(\Omega)$. Therefore, we can (by passing to a subsequence if necessary) suppose that

$$(3.13) \quad \begin{aligned} u_n &\rightharpoonup u \text{ weakly in } S_{\gamma,0}^2(\Omega) \text{ as } n \rightarrow \infty, \\ u_n &\rightarrow u \text{ a.e. in } \Omega \text{ as } n \rightarrow \infty, \\ u_n &\rightarrow u \text{ strongly in } L^p(\Omega), \quad 1 \leq p < 2_{\gamma}^* \text{ as } n \rightarrow \infty, \\ \omega u_n &\rightarrow \omega u \text{ strongly in } L^{q_1}(\Omega) \text{ as } n \rightarrow \infty. \end{aligned}$$

Thus by (A1), we have

$$(3.14) \quad \begin{aligned} \left| \int_{\Omega} f(x, u_n(x))(u_n(x) - u(x)) dx \right| &\leq C_0 \int_{\Omega} |u_n(x) - u(x)| dx \\ &+ \int_{\Omega} |u_n(x) - u(x)| |\omega(x) u_n(x)|^{p-1} dx \leq C_0 \|u_n - u\|_{L^1(\Omega)} \\ &+ \left(\int_{\Omega} |u_n(x) - u(x)|^{q_1} dx \right)^{1/q_1} \left(\int_{\Omega} |\omega(x) u_n(x)|^{q_1} dx \right)^{(q_1-1)/q_1}. \end{aligned}$$

In view of (3.13), we can conclude that

$$\int_{\Omega} f(x, u_n(x))(u_n(x) - u(x)) dx \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus,

$$(3.15) \quad \int_{\Omega} (f(x, u_n(x)) - f(x, u(x)))(u_n(x) - u(x)) dx \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

It follows from $\lim_{n \rightarrow \infty} \Phi'(u_n) = 0$ and (3.13) that

$$(3.16) \quad \langle \Phi'(u_n) - \Phi'(u), u_n - u \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

By (3.15) and (3.16), we obtain

$$\int_{\Omega} |\nabla_{\gamma} u_n - \nabla_{\gamma} u|^2 dx \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Therefore, we conclude that $u_n \rightarrow u$ strongly in $S_{\gamma,0}^2(\Omega)$. The proof of Lemma 3.4 is complete. \square

Proof of Theorem 1.1. By Lemmas 3.1, 3.2, and 3.4, all conditions of Lemma 2.12 are satisfied. Thus, problem (1.2) has a nontrivial weak solution \square

Example 3.5. Let $\gamma = (1, 1, |x_1 x_2|)$, $N = 3$, $\Omega = \{x \in \mathbb{R}^3: |x| < 1\}$, $\tilde{N} = 5$, $q_1 = 3$,

$$\begin{aligned} \omega_0(x) &= |x|^{-1/50} \sin(2017|x|) \quad \forall x \in \Omega, \\ \omega_1(x) &= \left(\frac{1}{2} - |x|^2\right)^2 (1 - |x|^2)^{-1} \quad \forall x \in \Omega, \\ \varphi_0(t) &= \begin{cases} t^2(1 - |t|) & \text{if } |t| \leq 1, \\ 0 & \text{if } |t| \in \mathbb{R} \setminus [-1, 1], \end{cases} \\ \varphi_1(t) &= \begin{cases} 0 & \text{if } |t| \leq 1, \\ t(|t| - 1) \ln(1 + |t|) & \text{if } |t| \in (1, 2], \\ t \ln(1 + |t|) & \text{if } |t| > 2, \end{cases} \\ f(x, t) &= \omega_0^2(x) \varphi_0(t) + \omega_1^2(x) \varphi_1(t) \quad \forall (x, t) \in \Omega \times \mathbb{R}. \end{aligned}$$

Let $\omega = |\omega_0| + \omega_1$, $C = 1$, $h(x) = |x|^{-1/50}$, $k(x) = -h(x)$ for every x in Ω . We see that $h \in L^1(\Omega)$ and $k \in L^{\tilde{N}/2}(\Omega)$. By Examples 2.6 and 2.8, ω is in $\mathcal{K}_{2,3}$. Thus f satisfies conditions (A1)–(A5).

We have $f(x, t)/t = \omega_1^2(x)(|t| - 1) \ln(1 + |t|)$ for every $t \in [-2, 2] \setminus [-1, 1]$ and $f(x, t)/t = \omega_1^2(x) \ln(1 + |t|)$ for every $t \in \mathbb{R} \setminus [-2, 2]$. Thus f satisfies (A6). Therefore, we can apply Theorem 1.1 to f with $C = 1$. We have that ω^2 is not integrable on Ω . Hence, the results in [2], [3], [12], [16], [17], [18], [22], [25] cannot be applied to solve of problem (1.2) in this case.

Acknowledgements. The author warmly thanks the anonymous referees for the careful reading of the manuscript and for their useful and nice comments.

References

- [1] *A. Ambrosetti, A. Malchiodi*: Nonlinear Analysis and Semilinear Elliptic Problems. Cambridge Studies in Advanced Mathematics 104. Cambridge University Press, Cambridge, 2007. [zbl](#) [MR](#) [doi](#)
- [2] *A. Ambrosetti, P. H. Rabinowitz*: Dual variational methods in critical point theory and applications. *J. Funct. Anal.* *14* (1973), 349–381. [zbl](#) [MR](#) [doi](#)
- [3] *C. T. Anh, B. K. My*: Existence of solutions to Δ_λ -Laplace equations without the Ambrosetti-Rabinowitz condition. *Complex Var. Elliptic Equ.* *61* (2016), 137–150. [zbl](#) [MR](#) [doi](#)
- [4] *H. Brezis*: Functional Analysis, Sobolev Spaces and Partial Differential Equations. Universitext. Springer, New York, 2011. [zbl](#) [MR](#) [doi](#)
- [5] *G. Cerami*: An existence criterion for the critical points on unbounded manifolds. *Rend., Sci. Mat. Fis. Chim. Geol.* *112* (1978), 332–336. (In Italian.) [zbl](#) [MR](#)
- [6] *G. Cerami*: Sull’esistenza di autovalori per un problema al contorno non lineare. *Ann. Mat. Pura Appl., IV. Ser.* *124* (1980), 161–179. (In Italian.) [zbl](#) [MR](#) [doi](#)
- [7] *B. Franchi, E. Lanconelli*: An embedding theorem for Sobolev spaces related to non-smooth vector fields and Harnack inequality. *Commun. Partial Differ. Equations* *9* (1984), 1237–1264. [zbl](#) [MR](#) [doi](#)
- [8] *N. Garofalo, E. Lanconelli*: Existence and nonexistence results for semilinear equations on the Heisenberg group. *Indiana Univ. Math. J.* *41* (1992), 71–98. [zbl](#) [MR](#) [doi](#)
- [9] *V. V. Grushin*: On a class of hypoelliptic operators. *Math. USSR, Sb.* *12* (1970), 458–476; translation from *Mat. Sb. (N.S.)* *83* (1970), 456–473. [zbl](#) [MR](#) [doi](#)
- [10] *D. S. Jerison*: The Dirichlet problem for the Kohn Laplacian on the Heisenberg group II. *J. Funct. Anal.* *43* (1981), 224–257. [zbl](#) [MR](#) [doi](#)
- [11] *D. S. Jerison, J. M. Lee*: The Yamabe problem on CR manifolds. *J. Diff. Geom.* *25* (1987), 167–197. [zbl](#) [MR](#) [doi](#)
- [12] *A. E. Kogoj, E. Lanconelli*: On semilinear Δ_λ -Laplace equation. *Nonlinear Anal., Theory Methods Appl., Ser. A* *75* (2012), 4637–4649. [zbl](#) [MR](#) [doi](#)
- [13] *A. E. Kogoj, E. Lanconelli*: Linear and semilinear problems involving Δ_λ -Laplacians. *Electron. J. Differ. Equ.* *2018* (2018), Article ID 25, 167–178. [zbl](#) [MR](#)
- [14] *N. Lam, G. Lu*: N -Laplacian equations in \mathbb{R}^N with subcritical and critical growth without the Ambrosetti-Rabinowitz condition. *Adv. Nonlinear Stud.* *13* (2013), 289–308. [zbl](#) [MR](#) [doi](#)
- [15] *N. Lam, G. Lu*: Elliptic equations and systems with subcritical and critical exponential growth without the Ambrosetti-Rabinowitz condition. *J. Geom. Anal.* *24* (2014), 118–143. [zbl](#) [MR](#) [doi](#)
- [16] *G. Li, C. Wang*: The existence of a nontrivial solution to a nonlinear elliptic problem of linking type without the Ambrosetti-Rabinowitz condition. *Ann. Acad. Sci. Fenn., Math.* *36* (2011), 461–480. [zbl](#) [MR](#) [doi](#)
- [17] *S. Li, S. Wu, H.-S. Zhou*: Solutions to semilinear elliptic problems with combined nonlinearities. *J. Differ. Equations* *185* (2002), 200–224. [zbl](#) [MR](#) [doi](#)
- [18] *S. Liu*: On superlinear problems without the Ambrosetti and Rabinowitz condition. *Nonlinear Anal., Theory Methods Appl., Ser. A* *73* (2010), 788–795. [zbl](#) [MR](#) [doi](#)
- [19] *Z. Liu, Z.-Q. Wang*: On the Ambrosetti-Rabinowitz superlinear condition. *Adv. Nonlinear Stud.* *4* (2004), 563–574. [zbl](#) [MR](#) [doi](#)
- [20] *D. T. Luyen*: Two nontrivial solutions of boundary-value problems for semilinear Δ_γ differential equations. *Math. Notes* *101* (2017), 815–823. [zbl](#) [MR](#) [doi](#)

- [21] *D. T. Luyen*: Existence of nontrivial solution for fourth-order semilinear Δ_γ -Laplace equation in \mathbb{R}^N . *Electron. J. Qual. Theory Differ. Equ.* *2019* (2019), Article ID 78, 12 pages. [zbl](#) [MR](#) [doi](#)
- [22] *D. T. Luyen, N. M. Tri*: Existence of solutions to boundary-value problems for semilinear Δ_γ differential equations. *Math. Notes* *97* (2015), 73–84. [zbl](#) [MR](#) [doi](#)
- [23] *D. T. Luyen, N. M. Tri*: Global attractor of the Cauchy problem for a semilinear degenerate damped hyperbolic equation involving the Grushin operator. *Ann. Pol. Math.* *117* (2016), 141–161. [zbl](#) [MR](#) [doi](#)
- [24] *D. T. Luyen, N. M. Tri*: Large-time behavior of solutions to degenerate damped hyperbolic equations. *Sib. Math. J.* *57* (2016), 632–649; translation from *Sib. Mat. Zh.* *57* (2016), 809–829. [zbl](#) [MR](#) [doi](#)
- [25] *D. T. Luyen, N. M. Tri*: Existence of infinitely many solutions for semilinear degenerate Schrödinger equations. *J. Math. Anal. Appl.* *461* (2018), 1271–1286. [zbl](#) [MR](#) [doi](#)
- [26] *D. T. Luyen, N. M. Tri*: Infinitely many solutions for a class of perturbed degenerate elliptic equations involving the Grushin operator. *Complex Var. Elliptic Equ.* *65* (2020), 2135–2150. [MR](#) [doi](#)
- [27] *O. H. Miyagaki, M. A. S. Souto*: Superlinear problems without Ambrosetti and Rabinowitz growth condition. *J. Differ. Equations* *245* (2008), 3628–3638. [zbl](#) [MR](#) [doi](#)
- [28] *P. H. Rabinowitz*: *Minimax Methods in Critical Point theory with Applications to Differential Equations*. Regional Conference Series in Mathematics 65. American Mathematical Society, Providence, 1986. [zbl](#) [MR](#) [doi](#)
- [29] *M. Schechter, W. Zou*: Superlinear problems. *Pac. J. Math.* *214* (2004), 145–160. [zbl](#) [MR](#) [doi](#)
- [30] *N. T. C. Thuy, N. M. Tri*: Some existence and nonexistence results for boundary value problems for semilinear elliptic degenerate operators. *Russ. J. Math. Phys.* *9* (2002), 365–370. [zbl](#) [MR](#)
- [31] *P. T. Thuy, N. M. Tri*: Nontrivial solutions to boundary value problems for semilinear strongly degenerate elliptic differential equations. *NoDEA, Nonlinear Differ. Equ. Appl.* *19* (2012), 279–298. [zbl](#) [MR](#) [doi](#)
- [32] *P. T. Thuy, N. M. Tri*: Long time behavior of solutions to semilinear parabolic equations involving strongly degenerate elliptic differential operators. *NoDEA, Nonlinear Differ. Equ. Appl.* *20* (2013), 1213–1224. [zbl](#) [MR](#) [doi](#)
- [33] *N. M. Tri*: Critical Sobolev exponent for degenerate elliptic operators. *Acta Math. Vietnam.* *23* (1998), 83–94. [zbl](#) [MR](#)
- [34] *N. M. Tri*: On Grushin’s equation. *Math. Notes* *63* (1998), 84–93; translation from *Mat. Zametki* *63* (1998), 95–105. [zbl](#) [MR](#) [doi](#)
- [35] *N. M. Tri*: *Semilinear Degenerate Elliptic Differential Equations: Local and Global Theories*. Lambert Academic Publishing, Saarbrücken, 2010.
- [36] *N. M. Tri*: *Recent Progress in the Theory of Semilinear Equations Involving Degenerate Elliptic Differential Operators*. Publ. House Sci. Technology, Hanoi, 2014.

Author’s address: Duong Trong Luyen, Division of Computational Mathematics and Engineering, Institute for Computational Science, Ton Duc Thang University, 19 Nguyen Huu Tho street, Tan Phong ward, District 7, Ho Chi Minh City, Vietnam; Faculty of Mathematics and Statistics, Ton Duc Thang University, 19 Nguyen Huu Tho street, Tan Phong ward, District 7, Ho Chi Minh City, Vietnam, e-mail: duongtrongluyen@tdtu.edu.vn.