

## Mařík, Jan: Scholarly works

---

Jan Mařík; Clifford E. Weil

Products of derivatives and approximate continuity

Proceedings of the Real Analysis Summer Symposium (Syracuse, N.Y., 1981), *Real Anal. Exchange* 1, 54–57

Persistent URL: <http://dml.cz/dmlcz/502129>

### Terms of use:

© Michigan State University Press, 1981

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*  
<http://dml.cz>

Jan Mařík and Clifford E. Weil, Department of Mathematics,  
Michigan State University, East Lansing, Michigan 48824

PRODUCTS OF DERIVATIVES AND APPROXIMATE CONTINUITY

This note contains the main points of a talk given by the first author at Real Analysis Symposium.

We introduce the following notation:  $\mathbb{R}$  is the real line;  $D$  is the system of all finite derivatives on  $\mathbb{R}$ ; if, moreover,  $n$  is a natural number, then  $D^n = \{f_1 \cdots f_n; f_1, \dots, f_n \in D\}$ ;  $C_{ap}$  is the system of all functions approximately continuous on  $\mathbb{R}$ .

It is well known that derivatives behave badly with respect to multiplication. Simple examples show that, for instance, the product of a derivative with a differentiable function need not be a derivative. We intend to show, roughly speaking, that operations performed on derivatives lead to derivatives only in exceptional cases. As an illustration we mention that if  $g$  is a function on  $\mathbb{R}$  such that the composite function  $g \circ f$  is in  $D$  for each  $f \in D$ , then  $g$  is linear. Another result pointing in the same direction is the following: Let  $f \in D$ , let  $g$  be a function strictly convex on an open interval containing  $f(\mathbb{R})$  and let  $g \circ f \in D$ . Then  $f \in C_{ap}$ . Taking  $g(y) = y^{-1}$  for each  $y \in (0, \infty)$  we see that if  $f \in D$ ,  $f > 0$  and if

also  $1/f \in D$ , then  $f \in C_{ap}$ . Using the Darboux property of derivatives we get: If  $f, g \in D$  and if  $fg$  is a nonzero constant, then  $f, g \in C_{ap}$ . This formulation suggests various generalizations. We succeeded to prove the following theorem:

Let  $f_1, \dots, f_n \in D$ ,  $f_j > 0$  and let  $f_1 \cdots f_n \in C_{ap}$ . Let  $a_1, \dots, a_n \in [0, 1]$ ,  $a_1 + \dots + a_n \leq 1$ . Then  $f_1^{a_1} \cdots f_n^{a_n} \in D \cap C_{ap}$ . In particular,  $f_j \in C_{ap}$  ( $j = 1, \dots, n$ ).

It is not surprising that products of derivatives behave badly with respect to addition. We see this well when we investigate sums of powers of derivatives. It is natural to ask, e.g., under what conditions the sum of the squares of two derivatives is in  $D^2$ . Because the product of a bounded derivative with a bounded approximately continuous functions is a derivative, the following holds: Let  $h \in D$ , let  $\varphi, \psi \in C_{ap}$  and let the functions  $h, \varphi$  and  $\psi$  be bounded. Let  $f = \varphi h$ ,  $g = \psi h$ . Then  $f^2 + g^2 \in D^2$ ; actually, it is the square of the derivative  $h(\varphi^2 + \psi^2)^{1/2}$ . It turns out that this example is not far from the general case. Namely: Let  $f, g \in D$ ,  $f^2 + g^2 \in D^2$ . Let  $h = (f^2 + g^2)^{1/2}$ . Then  $h \in D$ . If, moreover,  $h \geq 1$ , then  $f/h, g/h \in C_{ap}$ .

We get even somehow stronger results, when we add powers with different exponents. For instance: Let  $f, g, h \in D$ ,  $1 \leq f^2 + g^4 + h^4 \in D^2$  and  $g^4 + h^4 > 0$ . Then  $f, g, h \in C_{ap}$ .

It is probably not easy to give a simple characterization of a function in  $D^2$  or, in general, in  $D^n$ . However, we would like to mention some partial results connected with that problem.

Let  $f_j \in D$ ,  $f_j \geq 0$ , let  $A_j$  be a set closed in  $R$  and let  $g_j$  be its characteristic function ( $j = 1, \dots, n$ ). Then  $\sum_{j=1}^n f_j g_j \in D^2$ . In particular, the characteristic function  $g$  of a closed set  $A$  is always in  $D^2$ . A more detailed analysis shows that there are  $\varphi, \psi \in D$  such that  $0 \leq \varphi \leq 2$ ,  $0 \leq \psi \leq 2$  and  $\varphi\psi = g$ . If  $\emptyset \neq A \neq R$ , then 2 cannot be replaced by any smaller number.

With the help of A.M. Bruckner we have proven that every function of Baire class 1 that equals zero almost everywhere is in  $D^2$ .

Now let  $L = [0, \infty)$  and let  $D_L$  be the system of all derivatives on  $L$ . The meaning of  $D_L^n$  is now obvious. For each function  $F$  on  $L$  we define

$$T(F) = \sup \left\{ \limsup_{n \rightarrow \infty} \frac{F(y_n) - F(x_n)}{y_n - x_n} ; \right.$$

$$\left. 0 \leq x_n < y_n, y_n \rightarrow 0, \sup_n \frac{x_n}{y_n} < 1 \right\} .$$

We have always  $T(F) \geq \overline{F}^+(0)$ . Moreover it is easy to prove that  $T(F) = F'^+(0)$ , if  $F'^+(0)$  exists and is finite.

For each function  $f$  on  $L$  continuous on  $(0, \infty)$  such that the Lebesgue integral  $I = \int_0^1 f$  exists we

define  $S(f) = I$ , if  $I = \pm\infty$ , and  $S(f) = T(F)$ ,  
 where  $F(x) = \int_0^x f$ , if  $I \in \mathbb{R}$ . If  $f$  satisfies our  
 conditions and if  $f \in D_L$ , then, obviously,  $S(f) = f(0)$ .

Suppose that  $f$  is a function on  $L$  that is  
 positive and continuous on  $(0, \infty)$ . For each natural  
 number  $n$  define  $q_n = (S(f^{1/n}))^n$ . One of our results  
 says: Let  $n > 1$ . Then  $f \in D_L^n$  if and only if  
 $f(0) \geq q_n$ .

It is easy to see that  $q_2 \geq q_3 \geq \dots$ . If, for  
 instance,  $g$  is a nonconstant positive continuous  
 periodic function and if  $f(x) = g(1/x)$  for  $x \in (0, \infty)$ ,  
 then  $q_1 > q_2 > q_3 > \dots$ . This shows that  $D_L^n \neq D_L^{n+1}$   
 for each  $n$ .

Now let  $q = \lim q_n$ . If  $f(0) > q$ , then  $f \in D_L^n$   
 for some  $n$ ; if  $f(0) < q$ , then there is no such  $n$ .  
 If  $S(|\ln f|^a) < \infty$  for some  $a > 1$ , then  $q = \infty$  or  
 $q = \exp(S(\ln f))$ ; in general, if  $q < \infty$ , we have  
 $q = \lim(\exp(S(p \vee \ln f)))(p \rightarrow -\infty)$ .

The proofs of the assertions stated in this note  
 will be published later.

Finally, we would like to mention the following  
 problem: Is a bounded product of two (nonnegative)  
 derivatives the product of two (nonnegative) bounded  
 derivatives?