

Anna Helversen-Pasotto

Gamma-function and Gaussian-sum-function

In: Jarolím Bureš and Vladimír Souček (eds.): Proceedings of the Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1993. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 30. pp. [195]--200.

Persistent URL: <http://dml.cz/dmlcz/701518>

Terms of use:

© Circolo Matematico di Palermo, 1993

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Gamma-function and Gaussian-sum-function

A.Helversen-Pasotto

Let us recall that Euler gave the following integral presentation of the gamma-function

$$\Gamma(a) = \int_0^{\infty} e^{-t} t^a \frac{dt}{t}$$

and that the Gaussian sum of a multiplicative character A of a finite field \mathbf{F}_q is defined by

$$G(A) = \sum_{t \neq 0} E(t)A(t)$$

where E is the non-trivial additive character of \mathbf{F}_q defined by:

$$E(t) = e^{\frac{2\pi i \text{Tr}(t)}{p}}$$

for t element of \mathbf{F}_q , where $\text{Tr}(t) = t + t^p + \dots + t^{p^{n-1}}$ is the trace of t in the subfield \mathbf{F}_p of p elements, p prime and $q = p^n$.

Let us set

$$X = \{A : F_q^* \rightarrow C^* \mid A(t_1 t_2) = A(t_1)A(t_2) \text{ for } t_1, t_2 \in F_q^*\}$$

where F^* and C^* denote the multiplicative groups of F_q and of C , the field of complex numbers, respectively. The Gaussian-Sum-function G is defined on X and takes its values in C , or more precisely in the extension field of \mathbf{Q} , the field of rational numbers, obtained by adjunction of $e^{\frac{2\pi i}{p}}$ and $e^{\frac{2\pi i}{q-1}}$.

Lemma 1 *Let G be a finite abelian group, $|G|$ the number of its elements, and $A : G \rightarrow C^*$ a group morphism, then*

$$\sum_{g \in G} A(g) = \delta(A)|G|,$$

where $\delta(A) = 1$, if A is constant of value 1, and $\delta(A) = 0$, if not.

The proof is easy and well-known: If A is not constant of value 1, then there exists g_0 in G such that $A(g_0) \neq 1$; it follows that

$$\sum_g A(g) = \sum_g A(g_0 g) = A(g_0) \sum_g A(g)$$

so

$$(1 - A(g_0)) \sum_g A(g) = 0,$$

so

$$\sum_g A(g) = 0.$$

The following properties of Gaussian sums are consequences of Lemma 1 :

(P1) For A not constant of value 1, the absolute value of $G(A)$ equals the square-root of q ; for A constant of value 1 it is equal to 1.

(P2) For A in X not constant of value 1, we have

$$(G(A))^{-1} = q^{-1} A(-1) G(A^{-1}),$$

or more generally: for every A in X we have

$$G(A)G(A^{-1}) = qA(-1) - (q-1)\delta(A).$$

This can be generalized to

(P3) For A_1 and A_2 in X , we have

$$G(A_1)G(A_2) = b(A_1, A_2)G(A_1 A_2) + (q-1)A_1(-1)\delta(A_1 A_2)$$

where $b(A_1, A_2) = \sum_{t \neq 0,1} A_1(t)A_2(1-t)$ is so the called Jacobi sum.

The analogy between G and Γ appears by considering that

$$E(t_1 + t_2) = E(t_1)E(t_2)$$

for $t_1, t_2 \in F_q$ and

$$e^{-(t_1+t_2)} = e^{-t_1} e^{-t_2}$$

for t_1, t_2 in the real interval from 0 to ∞ , and that

$$A(t_1 t_2) = A(t_1)A(t_2)$$

for A in X and t_1, t_2 in F_q^* just as

$$(t_1 t_2)^a = t_1^a t_2^a$$

for t_1, t_2 reals, a in C .

The property (P3) of the Gaussian-sum-function is analogous to the well-known relation between gamma and beta function

$$\Gamma(a_1)\Gamma(a_2) = B(a_1, a_2)\Gamma(a_1 + a_2),$$

where a_1, a_2 are complex numbers of real part greater than zero and

$$B(a_1, a_2) = \int_0^1 t^{a_1-1}(1-t)^{a_2-1} dt$$

is the first Eulerian integral; the multiplication $A_1 A_2$ in the character-group X translates into the addition of the complex numbers a_1 and a_2 just as

$$t^{a_1} t^{a_2} = t^{a_1+a_2}$$

for t a real number and

$$A_1(t)A_2(t) = (A_1 A_2)(t)$$

for t in F_q .

The classical First Barnes' Lemma

$$\frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} \Gamma(a_1+s)\Gamma(a_2-s)\Gamma(a_3+s)\Gamma(a_4-s)ds = \frac{\Gamma(a_1 + a_2)\Gamma(a_2 + a_3)\Gamma(a_3 + a_4)\Gamma(a_4 + a_1)}{\Gamma(a_1 + a_2 + a_3 + a_4)}$$

(see for instance [6] for the hypothesis concerning the complex numbers $\{a_k, k = 1, 2, 3, 4\}$ and the path of integration) translates into the following identity for Gaussian sums, for which we will give an easy direct proof.

Proposition 1 For A_1, A_2, A_3, A_4 in X we have

$$\begin{aligned} & \frac{1}{q-1} \sum_{A \in X} G(A_1 A)G(A_2 A^{-1})G(A_3 A)G(A_4 A^{-1}) = \\ & = q(q-1)(A_1 A_3)(-1)\delta(A_1 A_2 A_3 A_4) + \frac{G(A_1 A_2)G(A_2 A_3)G(A_3 A_4)G(A_4 A_1)}{G(A_1 A_2 A_3 A_4)} \end{aligned}$$

Proof: Using the property (P3) we get

$$G(A_1A)G(A_2A^{-1}) = G(A_1A_2)b(A_1A, A_2A^{-1}) + (q - 1)(A_1A)(-1)\delta(A_1A_2)$$

and

$$G(A_3A)G(A_4A^{-1}) = G(A_3A_4)b(A_3A, A_4A^{-1}) + (q - 1)(A_3A)(-1)\delta(A_3A_4);$$

therefore the lefthand side of the identity becomes

$$\frac{1}{q - 1}(S_1 + S_2 + S_3 + S_4)$$

with

$$S_1 = G(A_1A_2)G(A_3A_4) \sum_{A \in X} b(A_1A, A_2A^{-1})b(A_3A, A_4A^{-1}),$$

$$S_2 = (q - 1)\delta(A_1A_2)A_1(-1)G(A_3A_4) \sum_{A \in X} A(-1)b(A_3A, A_4A^{-1}),$$

$$S_3 = (q - 1)\delta(A_3A_4)A_3(-1)G(A_1A_2) \sum_{A \in X} A(-1)b(A_1A, A_2A^{-1}),$$

$$S_4 = (q - 1)^3\delta(A_1A_2)\delta(A_3A_4)(A_1A_3)(-1);$$

now we compute the summation term in S_1 :

$$\sum_{A \in X} b(A_1A, A_2A^{-1})b(A_3A, A_4A^{-1}) = \sum_{s \neq 0, 1; t \neq 0, 1} A_1(s)A_2(1-s)A_3(t)A_4(1-t) \sum_A A\left(\frac{st}{(1-s)(1-t)}\right)$$

and using a dual form of Lemma 1 we get

$$\sum_A A\left(\frac{st}{(1-s)(1-t)}\right) = \left\{ \begin{array}{ll} q - 1 & \text{if } st = (1-s)(1-t) \\ 0 & \text{otherwise} \end{array} \right\} = \left\{ \begin{array}{ll} q - 1 & \text{if } t = 1 - s \\ 0 & \text{otherwise} \end{array} \right\}$$

therefore we get

$$\frac{1}{q - 1}(S_1 + S_2 + S_3 + S_4) = G(A_1A_2)G(A_3A_4) \sum_{s \neq 0, 1} (A_1A_4)(s)(A_2A_3)(1-s) + s_2 + s_3 + s_4,$$

where

$$s_2 = \delta(A_1A_2)A_1(-1)G(A_3A_4) \sum_{A \in X, s \neq 0, 1} A(-1)A_3(s)A_4(1-s)A\left(\frac{s}{1-s}\right)$$

$$s_3 = \delta(A_3A_4)A_3(-1)G(A_1A_2) \sum_{A \in X, s \neq 0,1} A(-1)A_1(s)A_2(1-s)A\left(\frac{s}{1-s}\right)$$

$$s_4 = (q-1)^2 \delta(A_1A_2)\delta(A_3A_4)(A_1A_3)(-1);$$

using the fact that $\frac{-s}{1-s} \neq 1$ we see that

$$\sum_{A \in X} A\left(\frac{-s}{1-s}\right) = 0$$

applying a dual version of Lemma 1; therefore $s_2 = s_3 = 0$; inverting the property (P3) we get

$$b(A_1A_4, A_2A_3) = \frac{G(A_1A_4)G(A_2A_3)}{G(A_1A_2A_3A_4)} + (q-1)\delta(A_1A_2A_3A_4)(A_1A_4)(-1)$$

and substituting the summation in the first term S_1 by this expression we obtain altogether

$$\begin{aligned} \frac{1}{q-1}(S_1 + S_2 + S_3 + S_4) &= \frac{G(A_1A_2)G(A_2A_3)G(A_3A_4)G(A_4A_1)}{G(A_1A_2A_3A_4)} \\ &+ (q-1)\delta(A_1A_2A_3A_4)(A_1A_4)(-1)G(A_1A_2)G(A_3A_4) \\ &+ (q-1)^2\delta(A_1A_2)\delta(A_3A_4)(A_1A_3)(-1); \end{aligned}$$

property (P2) allows easily to see that this expression is equal to the righthand side of the announced identity.

This simple and elementary proof seems to be new; at least no reference to it is known to the author who thanks Patrick Solé and Frédéric Testard for helpful comments during her seminar talks on the subject at Nice.

A different proof, using Mellin-transforms, has been found by Patrick Solé and the author, and has been adapted to the classical case by P.Solé; this seems to constitute a new proof of the classical Barnes' identity avoiding the use of the theorem of residues [2].

For other proofs, historical remarks, related questions and further references see [1, 2, 3, 4].

For elementary background concerning Gaussian sums see [5].

References

- [1] Greene J. : Hypergeometric functions over finite fields, *Trans.Amer.Math.Soc.* 301 (1987), 77-101
- [2] Helversen-Pasotto A., Sole P.: On Barnes' first lemma and its finite analogues, 1991, preprint
- [3] Helversen-Pasotto A.: L' identité de Barnes pour les corps finis, *C.R. Acad.Sci Paris*, t.286 (1978), Série A, 297-300
- [4] Helversen-Pasotto A.: On the structure constants of certain Hecke algebras, *Proc 10th Winter School "Geometry and Physics" Srni*, to appear in *Rendiconti del Circolo Mat. di Palermo*.
- [5] Ireland K., Rosen M.: *A classical introduction to modern number theory*, Springer-Verlag 1990, New York.
- [6] Slater L.J. : *Generalized hypergeometric functions*, Cambridge University Press 1966.

Anna Helversen-Pasotto
Laboratoire de Mathématiques
Université de Nice Sophia-Antipolis
C.N.R.S -U.A. Parc Valrose
F-06034-NICE-Cedex
France