

Zdeněk Frolík

A characterization of topologically complete spaces in the sense of E. Čech in terms of convergence of functions

Czechoslovak Mathematical Journal, Vol. 13 (1963), No. 1, 148–151

Persistent URL: <http://dml.cz/dmlcz/100557>

Terms of use:

© Institute of Mathematics AS CR, 1963

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A CHARACTERIZATION OF TOPOLOGICALLY COMPLETE SPACES
IN THE SENSE OF E. ČECH IN TERMS OF CONVERGENCE
OF FUNCTIONS

ZDENĚK FROLÍK, Praha

(Received February 2, 1961)

A characterization of topologically complete spaces (in the sense of E. Čech) analogous to the well known characterization of pseudocompact spaces in terms of convergence of continuous functions.

A space P is said to be topologically complete (in the sense of E. Čech) if P is completely regular and P is a G_δ in the Čech-Stone compactification $\beta(P)$ of P . In the present note, we shall give a characterization of topologically complete spaces analogous to the following characterization of pseudocompact spaces: If a decreasing sequence $\{f_n\}$ of continuous functions is pointwise convergent to zero, then $\{f_n\}$ is uniformly convergent.

All functions are supposed to be real-valued. If \mathfrak{F} is a family of functions on a set P , then the symbol $\mathfrak{F} \downarrow 0$ will be used to express that for every f_1 and f_2 there exists an f in \mathfrak{F} with $f \leq \min(f_1, f_2)$ and that for every x in P ,

$$\inf \{f(x); f \in \mathfrak{F}\} = 0.$$

All spaces under consideration are supposed to be completely regular. $B(P)$ denotes the family of all bounded continuous functions on a space P . The symbol $\alpha(P)$ will be used to denote the family of all subrings A of $B(P)$ satisfying the following two conditions

- (1) $f \in A \Rightarrow |f| \in A$.
- (2) For every x in P and every neighborhood U of x there exists an f in A such that $0 \leq f \leq 1$, $f(x) = 1$, $f[P - U] = 0$.

Definition. We shall say that a collection $\gamma \subset \alpha(P)$ has the property (V) if the following condition is fulfilled:

If $\mathfrak{F} \subset B(P)$, $\mathfrak{F} \downarrow 0$ and $\mathfrak{F} \cap C \downarrow 0$ for every C in γ , then for every $\varepsilon > 0$ there exists an f in \mathfrak{F} such that $\|f\| < \varepsilon$, i.e., there exists a sequence in \mathfrak{F} uniformly convergent to zero.

We shall say that a ring $A \in \alpha(P)$ has the property (V), if the collection $(A) \subset \gamma(P)$ has the property (V).

Example 1. A space P is compact if and only if $B(P)$ has the property (V).

Proof. Evidently the condition is necessary. To prove sufficiency suppose that there exists a maximal centered family \mathfrak{M} of closed subsets with $\bigcap \mathfrak{M} = \emptyset$. Consider the family \mathfrak{F} of all non-negative $f \in B(P)$ for which $f \geq 1$ on some $M \in \mathfrak{M}$. Clearly $\mathfrak{F} \downarrow 0$ and $\|f\| \geq 1$ for every f in \mathfrak{F} . Thus $B(P)$ does not have the property (V).

Theorem 1. Let m be a cardinal number. A space P is the intersection of m open sets in the Čech-Stone compactification $\beta(P)$ of P if and only if there exists a collection $\gamma \subset \alpha(P)$ with the property (V) such that the potency of γ is at most m .

Proof. First let us suppose that

$$P = \bigcap \mathfrak{M},$$

where \mathfrak{M} is a family of open subsets of $\beta(P)$ and the potency of \mathfrak{M} is at most m . For every M in \mathfrak{M} let $A(M)$ be the family consisting of restrictions to P of all $f \in B(\beta(P))$ with $f[\beta(P) - M] = (0)$. Clearly $A(M) \in \gamma(P)$ for all $M \in \mathfrak{M}$. It is easy to see that the collection $\{A(M); M \in \mathfrak{M}\}$ has the property (V). Indeed, if $\mathfrak{F} \subset \beta(P)$, $\mathfrak{F} \downarrow 0$ and $[\mathfrak{F} \cap A(M)] \downarrow 0$ for all $M \in \mathfrak{M}$, then $\mathfrak{F}^* \downarrow 0$, where \mathfrak{F}^* is the family of continuous extensions to $\beta(P)$ of all $f \in \mathfrak{F}$. Since $B(\beta(P))$ has the property (V), for every $\varepsilon > 0$ there exists a f^* in \mathfrak{F}^* with $\|f^*\| < \varepsilon$. If f is the restriction of f^* to P , then $f \in \mathfrak{F}$ and $\|f\| < \varepsilon$, which proves that the collection $\{A(M); M \in \mathfrak{M}\}$ has the property (V).

Conversely, let $\gamma \subset \alpha(P)$ be a collection with property (V) and let the potency of γ be at most m . For every C in γ let C^* be the family consisting of the continuous extensions to $\beta(P)$ of all $f \in C$. Put

$$K(C) = \{x; x \in \beta(P), f^* \in C^* \Rightarrow f^*(x) = 0\},$$

$$K = \bigcap \{K(C); C \in \gamma\},$$

$K(C)$ are compact subspaces of $\beta(P) - P$, and consequently, it is sufficient to prove

$$(3) \quad K = \beta(P) - P.$$

Clearly $K \subset \beta(P) - P$. Let us suppose that there exists a point x in $\beta(P) - (K \cup P)$. Let \mathfrak{F}^* be the family of all continuous non-negative functions f^* on $\beta(P)$ with $f^*(x) \geq 1$ and let \mathfrak{F} be the family consisting of the restrictions to P of all functions from \mathfrak{F}^* . Clearly $\mathfrak{F} \downarrow 0$ and $\|f\| \geq 1$ for every f in \mathfrak{F} . Let $C \in \gamma$. By our assumption there exists an f in C with $f^*(x) \neq 0$. Put

$$(4) \quad g = \max(0, f/f^*(x)).$$

Clearly $g \geq 0$ and $g^*(x) = 1$. If $y \in P$, then there exists a compact neighborhood F of y in $\beta(P)$ with $x \notin F$. According to condition (2) there exists a h in C with $h(y) = 1$, $h(P - F) = (0)$. Consider the function

$$(5) \quad k = \max(0, g - gh).$$

Clearly $k \in C$, $k(y) = 0$ and $k^*(x) = 1$. It follows that $(\mathfrak{F} \cap C) \downarrow 0$. But this is impossible, because γ has the property (V) and $\|f\| \geq 1$ for every f in \mathfrak{F} . This contradiction proves (3).

From the proof of the preceding Theorem 1 there follows at once theorem:

Theorem 2. *A space P is topologically complete in the sense of E. Čech if and only if there exists a decreasing sequence $\{A_n\}$ in $\alpha(P)$ with the property (V).*

Theorem 3. *A Lindelöf space P is topologically complete if and only if there exists a decreasing sequence $\{A_n\}$ in $\alpha(P)$ such that*

$$(6) \quad f_n \in A_n, \quad \{f_n\} \downarrow 0 \Rightarrow \lim_{n \rightarrow \infty} \|f_n\| = 0,$$

$$(7) \quad f \in A_n, \quad g \in A_{n+1}, \quad f \geq 0, \quad g \geq 0 \Rightarrow \min(f, g) \in A_{n+1}.$$

Proof. From the proof of Theorem 1 it follows at once that the condition is necessary. Conversely, let us suppose that there exists a sequence $\{A_n\}$ in $\alpha(P)$ satisfying (6). Let A_n^* be the family consisting of the continuous extensions of all $f \in A_n$ to $\beta(P)$. Put

$$(8) \quad K_n = \{x; x \in \beta(P), f^* \in A_n \Rightarrow f^*(x) = 0\},$$

$$(9) \quad K = \bigcup_{n=1}^{\infty} K_n.$$

The subspaces K_n of $\beta(P)$ being compact, it is sufficient to prove (3). Clearly $K \subset \beta(P) - P$. Suppose that there exists a point x in $\beta(P) - (P \cup K)$. First we shall construct sequences $\{f_k^n\}_{k=1}^{\infty}$ such that

$$(10) \quad f_k^n \in A_n, \quad \{f_k^n\}_{k=1}^{\infty} \downarrow 0 \quad (n = 1, 2, \dots).$$

Let n be a fixed positive integer. There exists an f in A_n such that $f^*(x) \neq 1$. Let g be the function defined by (4). For every y in P choose a compact neighborhood $F(y)$ of y in $\beta(P)$ with $x \notin F$. There exists a $h_y \in A_n$ such that $h_y(y) = 1$, $h_y[P - F(y)] = 0$. Put

$$r_y = \max(0, g - gh_y).$$

Clearly $r_y^*(x) = 1$, $r_y(y) = 0$ and $r_y \in A_n$. Since P is a Lindelöf space, there exists, for every $\varepsilon > 0$, a countable set $Y(\varepsilon) \subset P$ such that for any $y \in P$ there is a point $z \in Y(\varepsilon)$ with $r_z(y) < \varepsilon$. Let every $Y(1/j)$, $j = 1, 2, \dots$, be arranged in a sequence $\{z_i^j\}_{i=1}^{\infty}$; for $z = z_i^j$, denote r_z by r_i^j , and put

$$f_k^n = \min_{i, p \leq k} r_i^j \quad (k = 1, 2, \dots).$$

Clearly $f_k^n \in A_n$, $\{f_k^n\}_{k=1}^{\infty} \downarrow 0$.

We have proved that for every $n = 1, 2, \dots$ there exists a sequence $\{f_k^n\}_{k=1}^{\infty}$ in A_n with $\{f_k^n\}_{k=1}^{\infty} \downarrow 0$. Now put

$$f_n = \min_{i, j \leq n} f_i^j \quad (n = 1, 2, \dots).$$

According to (7), $f_n \in A_n$, and by construction $\{f_n\} \downarrow 0$ and $\|f_n\| = 1$, which contradicts (6). Thus (3) holds and P is topologically complete.

ХАРАКТЕРИЗАЦИЯ ТОПОЛОГИЧЕСКИ ПОЛНЫХ ПРОСТРАНСТВ
ПРИ ПОМОЩИ СХОДИМОСТИ ФУНКЦИЙ

ЗДЕНЕК ФРОЛИК (Zdeněk Frolik), Прага

Если \mathfrak{F} — множество непрерывных вещественных функций на пространстве P , то символ $\mathfrak{F} \downarrow 0$ обозначает, что

(1) Если $f_1, f_2 \in \mathfrak{F}$, то существует $f \in \mathfrak{F}$ так, что

$$f \leq \min(f_1, f_2).$$

(2) Для всякой точки $x \in P$

$$\inf \{f(x); f \in \mathfrak{F}\} = 0.$$

Через $B(P)$ обозначается множество всех ограниченных непрерывных вещественных функций на P ; $\alpha(P)$ обозначает множество всех подколец A кольца $B(P)$, имеющих следующие два свойства:

(a) $f \in A \Rightarrow |f| \in A$;

(б) Для всякой окрестности U всякой точки $x \in P$ существует $f \in A$ так, что $0 \leq f \leq 1, f(x) = 1, f[P - U] = (0)$.

Определение. Семейство $\gamma < \alpha(P)$ имеет свойство (V), если выполняется следующее условие:

Если $\mathfrak{F} \subset B(P), \mathfrak{F} \downarrow 0$ и также $(\mathfrak{F} \cap C) \downarrow 0$ для всякого $C \in \gamma$, то для всякого $\varepsilon > 0$ существует $f \in \mathfrak{F}$ так, что $\|f\| < \varepsilon$.

Доказываются следующие теоремы:

Теорема 1. *Вполне регулярное пространство P является пересечением m открытых множеств в чеховском компактном расширении тогда, и только тогда, если существует семейство $\gamma \subset \alpha(P)$ со свойством (V), имеющее мощность $\leq m$.*

Теорема 2. *Линделефовское пространство P является топологически полным в смысле Э. Чеха тогда, и только тогда, если существует невозрастающая последовательность $\{A_n\}$ в $\alpha(P)$ так, что*

(1) $f_n \in A_n, \{f_n\} \downarrow 0 \Rightarrow \lim \|f_n\| = 0$,

(2) $f \in A_n, g \in A_{n+1}, f \geq 0, g \geq 0 \Rightarrow \min(f, g) \in A_{n+1}$.