

V. R. Chandran

On a problem of B. Zelinka, II

Czechoslovak Mathematical Journal, Vol. 37 (1987), No. 1, 125–127

Persistent URL: <http://dml.cz/dmlcz/102141>

Terms of use:

© Institute of Mathematics AS CR, 1987

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A PROBLEM OF B. ZELINKA, II

V. R. CHANDRAN, Madras

(Received August 19, 1985)

In [2], B. Zelinka has posed the following problem viz whether there exists a commutative semi-group such that each tolerance relation is compatible with its element set? In [1] we have given an example of such a semi-group. The purpose of the present paper is to give a complete characterisation of B. Zelinka's problem. For definitions and notation refer [2]. Now we prove the following two theorems which completely characterize the problem of B. Zelinka.

Theorem 1. *Let $\langle S, * \rangle$ be a commutative semi-group with a multiplicatively zero element. Let $|S| \geq 3$. Then every tolerance relation in $\langle S, * \rangle$ is compatible with its element set if and only if $\langle S, * \rangle$ is a zero semi-group, i.e. the product of any two elements is zero.*

Proof. If part. Proof is exactly the same as in the example of [1]. Only if part. Let a, b be two distinct elements in S different from 0. Suppose $a * b \neq 0$, then $a * b = a$ or $a * b \neq a$. Case i. $a * b = a$. Define a tolerance relation ϱ in S as follows. $\varrho = \{(x, x) \mid x \in S\} \cup \{(a, b), (b, a), (0, b), (b, 0)\}$. We shall show that this tolerance relation ϱ is not compatible with its element set in S . For $b \varrho a$ and $0 \varrho b$ but $(b * 0, a * b) = (0, a) \notin \varrho$, a contradiction. Case ii. $a * b \neq a$. In this case define a tolerance relation T in S as follows. $T = \{(z, z) \mid z \in S\} \cup \{(a, b), (b, a), (0, a), (a, 0)\}$. Now $(a, b), (0, a) \in T$ but $(a * 0, b * a) = (0, a * b) \notin T$. Hence T is not compatible yielding a contradiction. Hence the product of any two distinct elements is zero. Now, it remains to prove that $a * a = 0$ for every $a \in S$. If $a * a \neq 0$, then $a * a = a$ or $a * a \neq a$. Since $|S| \geq 3$, S contains an element b different from 0 and a . Case iii. $a * a = a$. In this case define a tolerance relation ϱ' in $\langle S, * \rangle$ as follows. $\varrho' = \{(s, s) \mid s \in S\} \cup \{(a, b), (b, a), (0, b), (b, 0)\}$. Now $a \varrho' b$, $a \varrho' 0$. But $(a * a, b * 0) = (a, 0) \notin \varrho'$. Hence ϱ' is not compatible. Case iv. $a * a \neq a$. In this case, define a tolerance relation T' in $\langle S, * \rangle$ as follows. $T' = \{(s, s) \mid s \in S\} \cup \{(0, a), (a, 0)\}$. Now $(a, a) \in T'$ and $(0, a) \in T'$ but $(a * 0, a * a) = (0, a * a) \notin T'$, yielding a contradiction. Hence the theorem is proved.

Remark. There is meaning in taking $|S| \geq 3$. For now we give an example to show that the theorem is not true when $|S| = 2$. Let $S = \{a, b\}$. The multiplication

table of $\langle S, * \rangle$ is given below.

$*$	a	b
a	a	b
b	b	b

One can easily check that every tolerance relation in $\langle S, * \rangle$ is compatible and obviously $\langle S, * \rangle$ is not a zero semi-group.

Now, we are going to prove a theorem which is more powerful than theorem 1. We need the following definition.

Definition. A commutative semi-group $\langle S, * \rangle$ is called a *Zelinka semi-group* if and only if every tolerance relation in $\langle S, * \rangle$ is compatible with the elements of S .

Theorem 2. Let $\langle S, * \rangle$ be a commutative semi-group. Let $|S| \geq 3$. Then S is a Zelinka semi-group if and only if S contains a multiplicatively zero element and the product of any two elements in S is zero.

Proof. If part. Follows from theorem 1. Only if part. It suffices to prove that $\langle S, * \rangle$ contains a zero element, for then the result follows from theorem 1. Let a be any element in S . Now the set $a * S$ is either a single element set or contains more than one element.

Case i. $|a * S| = 1$. Let $a * S = \{x\}$ where $x \in S$. Now clearly x is a zero element in S for $x * S = (a * S) * S = a * (S * S) \subseteq a * S = \{x\}$. This implies x is a zero element in S and the result that the product of any two elements is zero follows from Theorem 1.

Case ii. $|a * S| > 1$. Then $a * S$ contains at least two distinct elements x, y . Clearly $x \neq y$.

Sub case a. $x \neq y, x, y \neq a$. Since $x, y \in a * S, a * b = x, a * c = y$ for some b, c in S . Now, let A be a tolerance relation defined as follows. $A = \{(s, s) \mid s \in S\} \cup \{(b, c), (c, b)\}$. Now, since A is compatible, $(a, a), (b, c) \in A$ implies $(a * b, a * c) = (x, y) \in A$. Hence the possibilities are $x = b, y = c$ or $x = c, y = b$.

Sub case a – i. $x = b, y = c$. Now we have $a * b = b, a * c = c$. Let B be a tolerance relation in $\langle S, * \rangle$ defined as follows: $B = \{(s, s) \mid s \in S\} \cup \{(a, c), (c, a), (a, b), (b, a)\}$. We shall show that B is not compatible. Now $(a, c) \in B$ and $(b, a) \in B$. But $(a * b, c * a) = (b, c) \notin B$ yielding a contradiction. The next possibility is $x = c$ and $y = b$.

Sub case a – ii. $x = c, y = b$. Now we have $a * c = b$ and $a * b = c$. Let C be a tolerance relation in $\langle S, * \rangle$ defined as follows. $C = \{(s, s) \mid s \in S\} \cup \{(a, b), (b, a)\} \cup \{(a, c), (c, a)\}$. Now $(a, b), (c, a) \in C$ but $(a * c, b * a) = (b, c) \notin C$, since $x = c, y = b$ and $x \neq y$ and $x, y \neq a$. Hence, C is not compatible, a contradiction.

Sub case b. $x \neq y$ and at least one of x, y equals a . W.l.o.g. assume that $x = a, y \neq a$. Now $a * b = a$ and $a * c = y$. Let D be a tolerance relation defined in $\langle S, * \rangle$ as follows, $D = \{(s, s) \mid s \in S\} \cup \{(a, b), (b, a), (a, c), (c, a)\}$. By assumption D is compatible. Hence, $(a, b), (c, a) \in D$ implies $(a * c, b * a) = (y, a) \in D$. Since $y \neq a$,

the possibilities are $y = b$ or $y = c$. We shall show that both these possibilities are impossible.

Sub case b – i. $y = b$. Then we have $a * y = a$ and $a * c = y$. Since $y \neq a$, clearly $y \neq c$ for $y = c$ implies $a * y = a * c$ which implies $a = y$ which is a contradiction. Let E be a tolerance relation defined in S as follows. $E = \{(s, s) \mid s \in S\} \cup \{(y, c), (c, y)\}$. Since E is compatible $(a, a) \in E$, $(y, c) \in E$ implies $(a * y, a * c) = (a, y) \in E$ which implies $y = a$ or $c = a$. Clearly $y \neq a$. Hence, other possibility is $c = a$. We shall show that this is also impossible. Let $c = a$. Now we have $a * y = a$ and $a * a = y$. Since $|S| \geq 3$, there exists an element d distinct from a and y . Now by assumption a, y, d are three distinct elements in S . Let F be a tolerance relation defined in $\langle S, * \rangle$ as follows. $F = \{(s, s) \mid s \in S\} \cup \{(y, d), (d, y)\}$. Now, $(a, a), (y, d) \in F$ and since F is compatible $(a * y, a * d) = (a, a * d) \in F$ which implies $a * d = a$. We shall show that $a * d = a$ is also not possible. For let G be a tolerance relation defined in $\langle S, * \rangle$ as follows. $G = \{(s, s) \mid s \in S\} \cup \{(a, d), (d, a)\}$. We shall show that G is not compatible. Now $(a, a), (a, d) \in G$. But $(a * a, a * d) = (y, a) \notin G$ yielding a contradiction.

Sub case b – ii. $y = c$. Then we have $a * b = a$ and $a * y = y$. Let H be a tolerance relation defined as follows. $H = \{(s, s) \mid s \in S\} \cup \{(b, y), (y, b)\}$. Since H is compatible, $(a, a), (y, b) \in H$ implies $(a * y, a * b) = (y, a) \in H$. The possibilities are $y = a$ or $y = b$ or $b = a$. Since $y \neq a$, $b = y$ or $b = a$. Since $x \neq y$, $x = b$ and $y = c$ we have $y \neq b$. Hence the remaining possibility is $b = a$. In this case, $a * a = a$ and $a * y = y$. Since $|S| \geq 3$, and $a \neq y$, there exist an element d different from a and y . Now let I be a tolerance relation defined in $\langle S, * \rangle$ as follows. $I = \{(s, s) \mid s \in S\} \cup \{(a, d), (d, a)\}$. Since I is compatible $(a, a), (a, d) \in I$ implies $(a * a, a * d) = (a, a * d) \in I$. Hence $a * d = a$ or d . We shall show that both the possibilities are impossible. Suppose $a * d = a$. Now define a tolerance relation J in $\langle S, * \rangle$ as follows. $J = \{(s, s) \mid s \in S\} \cup \{(d, y), (y, d)\}$. Now, $(a, a), (y, d) \in J$. But $(a * y, a * d) = (y, a) \notin J$ showing that J is not compatible, a contradiction. Next possibility is $a * d = d$. Let K be a tolerance relation defined in $\langle S, * \rangle$ as follows. $K = \{(s, s) \mid s \in S\} \cup \{(a, y), (y, a), (a, d), (d, a)\}$. Now $(a, y), (d, a) \in K$. But $(a * d, y * a) = (d, y) \notin K$, since a, y, d are distinct elements thus yielding a contradiction.

All these contradictions show that $a * S$ contains only one element say x . So by case (i), x is a zero element of $\langle S, * \rangle$. Now by theorem 1, $\langle S, * \rangle$ is a zero semi-group. (Q.E.D.).

Finally, I wish to express my thanks to the refereree.

References

- [1] V. R. Chandran: On a problem of B. Zelinka. Czechoslovak Math. J. 37(112), (1987), 124.
- [2] B. Zelinka: Tolerances in algebraic structures. Czechosl. Math. J. Vol. 25(100), (1975), 175–178.

Author's address: Ramanujan Institute for Advanced Study in Mathematics, Chepauk, Madras-5, India-(S).