

Christopher J. Duckenfield

\*-biregular rings

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 13 (1972), No. 3, 431--436

Persistent URL: <http://dml.cz/dmlcz/105431>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1972

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

\* -BIREGULAR RINGS

Christopher J. DUCKENFIELD, Gambier

Introduction. Regular rings were first defined by von Neumann [1] and used in connection with continuous geometries, there being an isomorphism between a continuous geometry and all principal left ideals of some regular ring. The theory was later expanded by introducing the notion of a \* -regular ring, and biregular rings were developed as a two-sided analogue to regularity. It is the purpose of this paper to develop a two-sided analogue to \* -regularity, and to produce an isomorphism theorem analogous to the above.

1. Regular, \* -regular and biregular rings.

1.1. Definition. An associative ring  $R$  with a unit is regular if  $axa = a$  is solvable in  $R$  for all  $a \in R$ .

1.2. Definition. A regular ring is \* -regular if there exists an involutory anti-automorphism  $a \rightarrow a^*$  of the ring onto itself, such that  $aa^* = 0$  if and only if  $a = 0$ .

If  $R$  is \* -regular an element  $a \in R$  for which  $a = a^*$  is called self-conjugate. Self-conjugate idempotents are called projections.

-----  
AMS, Primary: 16A30

Ref. Z. 2.723.2

We have the following properties (proved in [3]).

1.3. Theorem. If  $R$  is an associative ring with unit, then

i)  $R$  is regular if and only if every principal left ideal of  $R$  is generated by a unique idempotent.

ii)  $R$  is  $*$ -regular if and only if every principal left ideal of  $R$  is generated by a unique projection.

As a two-sided analogue to regularity we have the following.

1.4. Definition. A ring is said to be biregular if every principal ideal is generated by a central idempotent.

## 2. $*$ -Biregular rings.

In view of Theorem 1.3 we would expect that the defining criterion for a two-sided analogue to  $*$ -regularity would be that every principal two-sided ideal of such a ring be generated by a unique central projection. Our two-sided analogue to a  $*$ -regular ring will be defined as follows.

2.1. Definition. A ring is defined to be  $*$ -biregular if it is both biregular and  $*$ -regular.

2.2. Theorem. Every principal ideal in a  $*$ -biregular ring  $R$  is generated by a uniquely defined central projection.

Proof. Let  $I$  be a principal two-sided ideal in  $R$ . Then, since  $R$  is biregular,  $I$  is generated by a central idempotent  $e$ . We see immediately that  $(e^*)^2 = e^*$ , and that  $(ee^*)^2 = ee^*$ . Therefore  $(1-ee^*)ee^* = 0$ , and so  $(1-ee^*)ee^*(1-ee^*)^* = [(1-ee^*)e]$ .  $[(1-ee^*)e]^* = 0$ ,

which implies that  $e = ee^*e = ee^* = e^*e$ .  $e^*$  is central since, if  $x$  is an arbitrary member of  $R$ ,  $e^*x = (x^*e)^* = (ex^*)^* = xe^*$ . Obviously now,  $(e)R = (ee^*)R = I$ , and  $ee^*$  is a central projection.

If  $I = \rho R$ , where  $\rho$  is a central projection, then  $\rho = ee^*x$  and  $ee^* = \rho y$  for some  $x, y \in R$ . Then  $\rho = ee^*\rho = \rho ee^* = ee^*$ , and so we have uniqueness.

We can give a further description of the above projection  $ee^*$  by means of the following.

**2.3. Theorem.** If  $R$  is a  $*$ -biregular ring and  $I_a$  is a principal ideal of  $R$  generated by  $a$ , then the unique central projection which generates  $I_a$  is the least central element such that  $ad = a$ .

**Proof.**  $I_a$  is the set of all finite sums  $\sum_i x_i a y_i$ , where  $x_i, y_i \in R$ ,  $i = 1, 2, \dots$ . Also,  $I_a = eR$  where  $e$  is a central idempotent, and by the previous theorem,  $I_a = ee^*R$ , where  $ee^*$  is a central projection. Then  $a = ee^*z$  for some  $z \in R$  and therefore  $ae^* = ee^*ze^* = (ee^*)^2z = ee^*z = a$ . Thus  $a(ee^*) = a$  and  $ee^*$  is central.

Now let  $d$  be a central element such that  $ad = a$ . Then  $ee^* = \sum_i x_i a y_i = \sum_i x_i a d y_i = d \sum_i x_i a y_i = dee^*$ . Therefore we have  $ee^*R = dee^*R \subseteq dR$ , i.e.  $ee^* \in d$ .

The center of a biregular ring is biregular ([4], Theorem 4). We also prove the following result.

**2.4. Theorem.** The center of a  $*$ -regular ring is  $*$ -regular.

Proof. It is well known that the center of a regular ring is regular, and therefore we need only show that if  $a$  is in the center, then so is  $a^*$ . Let  $a \in Z$ , where  $Z$  is the center, and let  $x$  be an arbitrary element of  $R$ . Then  $a^*x = (x^*a)^* = (ax^*)^* = xa^*$ , i.e.  $a^*$  is central.

Therefore the center of a  $*$ -biregular ring is both biregular and  $*$ -regular, and we get

2.5. Theorem. The center of a  $*$ -biregular ring is  $*$ -biregular.

A  $*$ -regular ring is said to be complete if the lattice of its projections is complete, and Kaplansky [5] has shown that if a  $*$ -regular ring is complete then its projections form a continuous geometry. If the ring is commutative, then the principal one-sided ideals are in fact principal two-sided ideals. Therefore, if the center of a  $*$ -biregular ring is complete, the lattice of its principal ideals form a continuous geometry.

Morrison ([4], Theorem 5) has shown that there is an isomorphism between the principal ideals of the center of a biregular ring and the principal ideals of the ring itself. We therefore get the following.

2.6. Theorem. The lattice of the principal ideals of a  $*$ -biregular ring  $R$ , whose center is complete, is a continuous geometry, i.e. the central projections of a  $*$ -biregular ring form a continuous geometry.

This, of course, is the two-sided analogue to Kaplansky's result.

The following theorem is one of the main results of von Neumann [2].

2.7. Theorem. A complemented modular lattice admitting a homogeneous basis of rank  $\geq 4$  has orthocomplements if and only if it is isomorphic to the lattice of principal left ideals of some  $*$ -regular ring.

In a two-sided analogue to this theorem we would want to replace "the lattice of principal left ideals of some  $*$ -regular ring" by "the lattice of principal ideals of some  $*$ -biregular ring".

Now, a  $*$ -biregular ring is biregular, and the lattice of principal ideals of a biregular ring is a distributive, relatively complemented lattice (Anđrunakievich [6]). If the ring contains a unit (which is the case for a  $*$ -biregular ring, since a  $*$ -biregular ring is regular and a regular ring has a unit) then this lattice is a Boolean algebra. A Boolean algebra is certainly orthocomplemented and so we seek to prove the following

2.8. Theorem. A Boolean algebra  $B$  is isomorphic to the lattice of principal ideals of some  $*$ -biregular ring.

Proof. Every Boolean algebra  $B$  is isomorphic to the lattice of principal ideals of some Boolean ring  $R$  (Birkhoff, [7], p.155). Trivially, a Boolean ring is commutative, regular and biregular. The commutativity gives us that the identity mapping is an anti-automorphism  $a \rightarrow a^*$  of  $R$  onto itself. Also  $aa^* = 0$  implies  $a = a^2 = aa^* = 0$ , since every element of a Boolean ring is an idempotent. Therefore  $R$  is  $*$ -regular and biregular, and hence is  $*$ -bire-

gular.

R e f e r e n c e s

- [1] J.von NEUMANN: On regular rings, Proc.Nat.Acad.Sci.  
U.S.A.22(1936),707-713.
- [2] ----- : Continuous Geometry, Princeton,1960.
- [3] L. SKORNYAKOV: Complemente $\bar{d}$  Modular Lattices and Regular Rings, Oliver and Boyd,1964.
- [4] D. MORRISON: Biregular rings and the ideal lattice isomorphisms, Proc.Amer.Math.Soc.6(1955),46-49.
- [5] I. KAPLANSKY: Any orthocomplemente $\bar{d}$  complete modular lattice is a continuous geometry, Ann.Math.61 (1955),524-541.
- [6] V. ANDRUNAKIEVICH: Biregular rings, Matem.Sb.39(1956), 447-464.
- [7] G. BIRKHOFF: Lattice Theory, Amer.Math.Soc.,Coll.Series 1948,25.

Kenyon College  
Gambier  
Ohio 43022  
U.S.A.

(Obstatum 15.3. 1972)