

Urban Cegrell

On the space and dual space of functions representable by differences of subharmonic functions

Commentationes Mathematicae Universitatis Carolinae, Vol. 18 (1977), No. 4, 685--695

Persistent URL: <http://dml.cz/dmlcz/105812>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1977

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

18,4 (1977)

ON THE SPACE AND DUAL SPACE OF FUNCTIONS REPRESENTABLE BY
DIFFERENCES OF SUBHARMONIC FUNCTIONSUrban CEGRELL^{x)}, Uppsala

Abstract: The linear space of differences of subharmonic functions is given a Fréchet space topology. This space together with its dual space is studied. A decomposition theorem for functionals vanishing on the harmonic functions is given and the functionals which are carried by one point is determined. It follows that, in the subharmonic case, the stable polar set always is countable.

Key words: Subharmonic function, Fréchet space, positive functional, polar set of a function.

AMS: 31B05

Ref. Ž.: 7.58

1. Introduction. Let U be an open subset of \mathbb{R}^n , $n \geq 2$, and denote by $SH(U)$ the subharmonic functions on U . In this paper we study the linear space $\mathcal{D}SH(U)$, of functions which can be written as a difference of subharmonic functions. This subject has been treated by Arsove [1] and Kiselman [3]. We shall also study its dual space $\mathcal{D}SH'(U)$ here.

The corresponding function spaces made up by differences of convex or plurisubharmonic functions have been studied by Kiselman [3] and Cegrell [2].

x) Supported by the Swedish Natural Science Research Council Contract No. F 3435-007.

2. $\mathcal{D}'SH(U)$ and $\mathcal{D}'SH'(U)$. $\mathcal{D}'SH(U)$ is a Fréchet space with topology given by the seminorms $\|\varphi\|_K = \inf (\int_K |\varphi|_1 + |\varphi|_2; \varphi = \varphi_1 - \varphi_2, \varphi_1, \varphi_2 \in SH(U)), K \subset \subset U$. For a proof of this, see Schaefer [4], p. 221.

An equivalent topology on $\mathcal{D}'SH(U)$ is given by the seminorms

$$\|\varphi\|_{U'} = \inf (\int_{U'} |\varphi_1| + |\varphi_2|; \varphi = \varphi_1 - \varphi_2, \varphi_1, \varphi_2 \in SH(U'))$$

where U' is open and relatively compact in U .

That $\mathcal{D}'SH(U)$ is complete under this topology is a consequence of Theorem 2.1. Moreover, $\|\cdot\|_{U'}$ gives a weaker topology than $\|\cdot\|_K$ and since both turn $\mathcal{D}'SH(U)$ into a Fréchet space, they are equivalent.

Theorem 2.1. If $\varphi \in L^1_{loc}(U)$ and if $\varphi|_{U'} \in \mathcal{D}'SH(U')$ for every U' open and relatively compact in U then $\varphi \in \mathcal{D}'SH(U)$.

Proof. Arsove [1] Theorem 10.

Definition. A compact subset K of U is said to be a carrier for $\mu \in \mathcal{D}'SH'(U)$ if to every open U' containing K there is a constant c such that

$$|\mu(\varphi)| \leq c \|\varphi\|_{U'}, \quad \forall \varphi \in \mathcal{D}'SH(U).$$

Definition. A subset K of U is said to be a support for $\mu \in \mathcal{D}'SH'(U)$ if, for any open O with $K \subset \subset O$, μ vanishes on those functions in $\mathcal{D}'SH(U)$ which vanish on $U \cap O$.

Definition. E is a notation for the fundamental solution to the equation $\Delta f = \sigma_0$ in \mathbb{R}^n when σ_0 is the Dirac measure at zero

$$E(z) = \begin{cases} \frac{1}{2\pi} \log |z|, & n = 2 \\ c_n \frac{-1}{|z|^{n-2}} & n > 2. \end{cases}$$

Corollary 2.2. Every $\mu \in \mathcal{O}SH(U)$ has a compact support.

Proof. Let B be a carrier for μ and choose U' open with $B \subset U' \subset U$. Then there is a constant $c > 0$ so that $|\mu(\varphi)| \leq c \|\varphi\|_{U'}$, $\forall \varphi \in \mathcal{O}SH(U)$. Since an equivalent topology on $\mathcal{O}SH(U)$ is defined by the seminorms $\|\varphi\|_{U''}$ there is a constant d and an open U'' relatively compact in U so that $\|\varphi\|_{U'} \leq d \|\varphi\|_{U''}$, $\forall \varphi \in \mathcal{O}SH(U)$. Hence $|\mu(\varphi)| \leq c \cdot d \|\varphi\|_{U''}$, $\forall \varphi \in \mathcal{O}SH(U)$ so if $\varphi \in \mathcal{O}SH(U)$ with $\varphi|_{U''} = 0$ then $\mu(\varphi) = 0$, which means that U'' is a support for μ .

Corollary 2.3. $\mathcal{O}SH(\mathbb{R}^n)|_U$ is dense in $\mathcal{O}SH(U)$.

Proof. By the Hahn-Banach theorem it is enough to prove that if $\mu \in \mathcal{O}SH'(U)$ vanishes on $\mathcal{O}SH(\mathbb{R}^n)|_U$ then $\mu \equiv 0$. So let $\varphi \in \mathcal{O}SH(U)$ and $\mu \in \mathcal{O}SH'(U)$ vanishing on $\mathcal{O}SH(\mathbb{R}^n)|_U$ be given. Choose $\theta \in \mathcal{D}(U)$, $0 \leq \theta \leq 1$ with $\theta = 1$ near a compact support A , for μ . Then $\varphi - E * \theta \Delta \varphi$ is subharmonic on U and harmonic near A . So there is a $\psi \in \mathcal{O}CVX(\mathbb{R}^n)$ with $\psi = \varphi - E * \theta \Delta \varphi$ near A . (See Kiselman [3].) Hence

$$0 = \mu(\psi) = \mu(\varphi - E * \theta \Delta \varphi) = \mu(\varphi)$$

since $E * \theta \Delta \varphi \in SH(\mathbb{R}^n)$.

Theorem 2.4. Assume that $\mu \in \mathcal{O}SH'(U)$ and that A and B are compact supports for μ . Then $A \cap B$ is a support.

Proof. Given U_3 open and $\varphi \in \mathcal{O}SH(U)$ vanishing near \bar{U}_3

where $A \cap B \subset \subset U_3 \subset \subset U$. We have to prove that $\mu(\varphi) = 0$.
 Choose U_2 open so that $B \subset \subset U_2$; $\bar{U}_2 \cap A \cap \mathcal{U}U_3 = \emptyset$. Choose U_1
 open so that $B \subset \subset U_1 \subset \subset U_2$ and $\theta_1 \in \mathcal{D}(U_2)$ with $\theta_1 = 1$
 near \bar{U}_1 . Then $\varphi = E * \theta_1 \Delta \varphi + h$ near \bar{U}_1 where h is harmo-
 nic near \bar{U}_1 and where $E * \theta_1 \Delta \varphi$ is harmonic on an open set
 U_4 such that $A \subset \subset U_4 \subset \subset U$; $U_4 \cap \bar{U}_2 \cap \mathcal{U}U_3 = \emptyset$.

Choose now $\theta_2 \in \mathcal{D}(U_2)$; $\theta_2 = 1$ near \bar{U}_1 so that
 $\theta_2 \cdot h \in \mathcal{S}CVX(\mathbb{R}^n)$ and $\theta_3 \in \mathcal{D}(U_4)$, $\theta_3 = 1$ near A . Then
 $\theta_3 \cdot E * \theta_1 \Delta \varphi \in \mathcal{S}CVX(\mathbb{R}^n)$ and we define $f, g \in \mathcal{S}CVX(\mathbb{R}^n)$
 by

$$f = \theta_2 \cdot \theta_3 \cdot h; \quad g = \theta_3 \cdot E * \theta_1 \Delta \varphi \cdot$$

On $U_1 \cap U_3$ we have $g + f = \theta_3 E * \theta_1 \Delta \varphi + \theta_3 \cdot h = \theta_3 \varphi = 0$
 since φ vanishes on U_3 and since $\theta_3 = 0$ on $U_1 \cap \mathcal{U}U_3$, $g +$
 $+ f = 0$ on U_1 which contains B . Hence

$$0 = \mu(f + g) = \mu(E * \theta_1 \Delta \varphi + \theta_2 \cdot h) = \mu(\varphi)$$

since $\theta_2 = 1$ near \bar{U}_1 and the proof is complete.

Remark. Theorem 2.4 and Corollary 2.2 prove that every
 $\mu \in \mathcal{S}SH'(U)$ has a smallest compact support.

Definition. Let K be a compact subset of U . Then \hat{K} is
 defined by

$$\hat{K} = \{z \in U; \varphi(z) \leq \sup_{\xi \in K} \varphi(\xi) \quad \forall \varphi \in SH(U)\}$$

Lemma 2.5. Let K be compact in U . Then \hat{K} is compact in
 U . Given $\varepsilon > 0$, and U_1 an open neighbourhood of \hat{K} . Then the-
re is a continuous and subharmonic function φ on U such that
 $\varphi = 0$ on \hat{K} and $\varphi \geq \varepsilon$ on $\mathcal{U}U_1$.

Proof. Consider $\hat{K}_\varepsilon = \{z \in U; \varphi(z) \leq \sup_K \varphi \quad \forall \varphi \in SH(U) \cap$
 $\cap C(U)\}$. It is clear that $K \subset \hat{K}_\varepsilon \subset \hat{K}$, and if $z_0 \in \partial U$

$$-E(z - z_0) > \sup_{\xi \in K} [-E(\xi - z_0)] \text{ for } |z - z_0| < d(K, CU)$$

so it follows that \hat{K}_c is compact in U .

We claim that $\hat{K} = \hat{K}_c$. If $z_1 \notin \hat{K}$ then there is a $\varphi \in SH(U) \cap C(U)$ with $\varphi(z_1) > \sup_K \varphi$. Choose $\varphi_2 \in SH(U) \cap C(U)$ so that $\varphi_\varepsilon \searrow \varphi$, $\varepsilon \searrow 0$ on a compact set containing z_1 and K in its interior. Then there is an ε_0 so that $\sup_K \varphi_{\varepsilon_0} \leq \frac{1}{2} (\varphi(z_1) + \sup_K \varphi)$. But $\varphi_{\varepsilon_0}(z_1) \geq \varphi(z_1)$ so $z_1 \notin \hat{K}_c$.

To a given open set U_1 with $\hat{K} \subset \subset U_1 \subset \subset U$ it is easy to see that there are finitely many functions $\varphi_i \in SH(U) \cap C(U)$, $1 \leq i \leq m$ with $\sup_{1 \leq i \leq m} \varphi_i = 0$ on \hat{K} and $\sup_{1 \leq i \leq m} \varphi_i \geq 1$ on ∂U_1 .

Proposition 2.6. Let K be a carrier for $\mu \in \mathcal{D}'SH'(U)$. Then \hat{K} is a support for μ .

Proof. Let K be a carrier for $\mu \in \mathcal{D}'SH'(U)$ with $K = \hat{K}$. Choose an open set U_1 so that $\hat{K} \subset \subset U_1 \subset \subset U$ and let $\varphi \in \mathcal{D}'SH(U)$ with $\varphi|_{U_1} = 0$ be given. We have to prove that $\mu(\varphi) = 0$.

Since

$$\begin{aligned} \varphi &= \varphi_1 - \varphi_2 = \varphi_1 - E * \chi_{U_1} \Delta \varphi_1 - \\ &\quad - (\varphi_2 - E * \chi_{U_1} \Delta \varphi_1) \end{aligned}$$

we have a representation ψ_1 and ψ_2 of φ where ψ_1 and ψ_2 are continuous near K . Using Lemma 2.5 we can find an open set U_2 , $K \subset \subset U_2 \subset \subset U_1$ and a continuous subharmonic function ψ so that

$$\begin{aligned} \inf_{\partial U_2} \psi &> \sup_{U_2} - \psi_2 \\ \sup_{U_3} \psi &\leq \inf_{U_3} - \psi_2 \end{aligned}$$

where $K \subset \subset U_3 \subset \subset U_2$ so it follows that

$$g(z) = \begin{cases} \psi(z), & z \notin U_2 \\ \sup(\psi_1 - \psi_2), & z \in U_2 \end{cases}$$
 is continuous on U and subharmonic near $\mathcal{C}U_2$.

Now $\theta_1 = \psi_1 + g$, $\theta_2 = \psi_2 + g$ are subharmonic on U and $\theta_1 - \theta_2 = \varphi$ so since K is a carrier for μ we have

$$|\mu(\varphi)| \leq c_{U_3} \int_{U_3} |\theta_1| + |\theta_2| = 0$$

and the proof is complete.

Corollary 2.7. Let μ be a non-vanishing element in $\mathcal{C}SH(U)$. If A and B are carriers for μ then $\hat{A} \cap \hat{B} \neq \emptyset$.

3. Positive functionals on $\mathcal{C}SH(U)$

Definition. Denote by $\mathcal{C}SH'_+(U)$ the set of elements in $\mathcal{C}SH'(U)$ which only takes non-negative values on $SH(U)$.

Remark. Any real-valued linear map which is defined on $\mathcal{C}SH(U)$ and which is non-negative on $SH(U)$ is continuous (see Proposition 1.1 in Cegrell [2]). In particular, we have the following

Lemma 3.8. Let K be a compact subset of U . Then

$$\mathcal{C}SH(U) \ni \varphi \longmapsto \Delta \varphi \{K\} \quad (= \int_K \Delta \varphi)$$

is an element in $\mathcal{C}SH'(U)$.

Theorem 3.9. $\mu \in \mathcal{C}SH'(U)$. Then the following conditions are equivalent.

- 1) $\mu = \mu_1 - \mu_2$ where $\mu_1, \mu_2 \in \mathcal{C}SH'_+(U)$;
- 2) there is a compact subset, K , of U such that $\mu(\varphi)$ vanishes for all $\varphi \in \mathcal{C}SH(U)$ which are harmonic near K ;

- 3) μ vanishes on the harmonic functions;
 4) there is a compact subset, K, of U and a constant c so
 that $|\mu(\varphi)| \leq c \int_K \Delta \varphi \quad \forall \varphi \in SH(U)$.

Proof. 1) \implies 2). If $\mu \in \mathcal{D}'SH(U)$, let K be a carrier for μ . Then \hat{K} is a support for μ by Proposition 2.6. Given $\varphi \in \mathcal{D}'SH(U)$ which is harmonic near \hat{K} we can construct $\varphi_1, \varphi_2 \in SH(U)$ so that $\varphi_1 = -\varphi_2 = \varphi$ near \hat{K} . Hence $0 \leq \mu(\varphi_1) = \mu(\varphi) = \mu(-\varphi_2) \leq 0$ so $\mu(\varphi) = 0$.

2) \implies 3) is trivial.

3) \implies 4). Denote with $M(U)$ the Fréchet space of measures on U with topology defined by seminorms $\|t\|_K =$ total mass of f on K, $K \subset U$.

Let $\mathcal{H}a(U)$ be a notation for the harmonic functions on U, which form a closed subspace of $\mathcal{D}'SH(U)$. Let now j be a notation for the mapping

$$\mathcal{D}'SH(U)/\mathcal{H}a(U) \ni \varphi \xrightarrow{j} \Delta \varphi \in M(U).$$

That j is continuous follows from Lemma 3.8. Furthermore, j is a bijection so j^{-1} is continuous since both $\mathcal{D}'SH(U)/\mathcal{H}a(U)$ and $M(U)$ are Fréchet spaces.

Now since $\mu = 0$ on $\mathcal{H}a(U)$ we have

$$|\mu(\varphi)| \leq c \inf_{h \in \mathcal{H}a(U)} \|\varphi + h\|_K \quad \forall \varphi \in \mathcal{D}'SH(U)$$

for a fixed constant c and compact set K. But j^{-1} is continuous so there is another constant d and another compact set L in U so that

$$\inf_{h \in \mathcal{H}a(U)} \|\varphi + h\|_K \leq d \inf_{\substack{\varphi_1, \varphi_2 \in SH(U) \\ \varphi = \varphi_1 - \varphi_2}} \int_L \Delta \varphi_1 + \Delta \varphi_2 \quad \forall \varphi \in \mathcal{D}'SH(U).$$

In particular, $|\mu(\varphi)| \leq c \cdot d \int_L \Delta \varphi \quad \forall \varphi \in SH(U)$.

4) \implies 1). $\mu(\varphi) = c \int_K \Delta \varphi + \mu(\varphi) - c \int_K \Delta \varphi$ is the desired representation.

4. Functionals on SH(U) carried by one point. We shall now determine all functionals $\mu \in \mathcal{D}'\text{SH}'(U)$ which are carried by one point $z_0 \in U$. We can of course restrict ourselves to the case $z_0 = 0$. Then, if μ is carried by the origin, μ is also supported by the origin. Let B denote $\{z; |z| < 1\}$.

Lemma 4.10. Let $\varphi \in \text{SH}(B)$ and assume that φ is bounded below in a neighbourhood of zero. Then $\mu(\varphi) = 0 \quad \forall \mu \in \mathcal{D}'\text{SH}'(B)$ which are carried by zero.

Proof. Given $\varphi \in \text{SH}(B)$ bounded below near zero. Assume first that $\varphi \leq 0$ on $\{z; |z| \leq r\}$ where $0 < r < 1$. If we put

$$\psi_n = \begin{cases} \sup \left(\frac{1}{n^2} \log \left| \frac{z}{r} \right|, \varphi \right), & |z| \leq r \\ \frac{1}{n^2} \log \left| \frac{z}{r} \right|, & |z| > r \end{cases}$$

it follows that $\psi_n \in \text{SH}(B)$ and $\psi_n = \varphi$ near zero.

Put $\Theta_N = \sum_{\nu=1}^N \psi_\nu$. Then $(M > N)$

$$\begin{aligned} \|\Theta_M - \Theta_N\|_{B'} &= \int_{B'} \left| \sum_{\nu=N+1}^M \psi_\nu \right| \leq \\ &\leq \left(\int_B \left| \log \left| \frac{z}{r} \right| \right| \right) \sum_{\nu=N+1}^M \frac{1}{\nu^2} \rightarrow 0, \quad \min(M, N) \rightarrow +\infty \end{aligned}$$

for every B' relatively compact in B.

So it follows that Θ_N converges to a limit $\Theta \in \mathcal{D}'\text{SH}(U)$. Now $\mu(\Theta) = \lim_{N \rightarrow +\infty} \mu(\Theta_N) = \lim_{N \rightarrow +\infty} \sum_{\nu=1}^N \mu(\varphi)$ which gives $\mu(\varphi) = 0$. If we apply this to $\varphi - \sup_{|z| < \frac{1}{2}} \varphi$ the lemma follows.

Definition. Let $s_z(U)$ be the functions in $\mathcal{S}SH(U)$ which have a representation $\varphi = \varphi_1 - \varphi_2$ where $\varphi_1 + \varphi_2(z) > -\infty$ and let $\mathcal{F}_z(U)$ denote the closure of $s_z(U)$ in $\mathcal{S}SH(U)$.

Lemma 4.11. Let $\mu \in \mathcal{S}SH'(B)$ be carried by zero. Then $\mu(\varphi) = 0 \quad \forall \varphi \in \mathcal{F}_0$.

Proof. It is enough to prove that if $\varphi \in SH(B)$ with $\varphi(0) > -\infty$ then $\mu(\varphi) = 0$. Choose $\theta_n \in \mathcal{D}(B)$, $0 \leq \theta_n \leq 1$, $\theta_n \geq \theta_{n+1}$, $\theta_n = 1$ near zero, $\lim_{n \rightarrow \infty} \theta_n = 0$ outside zero. By Theorem 3.9 and Lemma 4.10 there is a constant c so that

$$|\mu(\varphi)| = |\mu(E * \theta_n \Delta \varphi)| \leq c \int_K \theta_n \Delta \varphi \rightarrow 0, \quad n \rightarrow \infty$$

since $\varphi(0) > -\infty$.

Definition. Denote by $T_z(\varphi)$ the functional

$$\mathcal{S}SH(B) \ni \varphi \mapsto \Delta \varphi \{z\} \quad (= \int \Delta \varphi \{z\}).$$

Lemma 4.12. $T_0(\varphi) = 0 \iff \varphi \in \mathcal{F}_0$.

Proof. \Leftarrow) Clear by Lemma 4.11.

\Rightarrow) Choose θ_n as in the proof of Lemma 4.11 and assume that $\varphi = \varphi_1 - \varphi_2 \in \mathcal{S}SH(B)$ with $T_0(\varphi_1) = T_0(\varphi_2)$. Then

$$E * \theta_n \Delta \varphi_1 \nearrow E \cdot T_0(\varphi_1), \quad n \rightarrow +\infty$$

$$E * \theta_n \Delta \varphi_2 \nearrow E \cdot T_0(\varphi_2), \quad n \rightarrow +\infty.$$

Put

$$\psi_1 = \varphi_1 - E \cdot T_0(\varphi_1), \quad \text{then } \psi_1, \psi_2 \in SH(B),$$

$$\psi_2 = \varphi_2 - E \cdot T_0(\varphi_2)$$

$$\varphi = \psi_1 - \psi_2 \quad \text{and}$$

$$\psi_1^n = \psi_1 - E * \theta_n \Delta \psi_1 \in s_0, \quad n \in \mathbb{N}$$

$$\psi_2^n = \psi_2 - E * \Theta_n \Delta \psi_2 \in \mathfrak{s}_0, \quad n \in \mathbb{N}.$$

For every compact subset K of U we have

$$\begin{aligned} & \| \psi_1 - \psi_2 - (\psi_1^n - \psi_2^n) \|_K \leq \| E * \Theta_n \Delta \psi_1 \|_K + \\ & + \| E * \Theta_n \Delta \psi_2 \|_K \leq \int_K (E \cdot T(\varphi_1) - E * \Theta_n \Delta \varphi_1) dz + \\ & + \int_K (E \cdot T(\varphi_1) - E * \Theta_n \Delta \varphi_2) dz \rightarrow 0 \quad n \rightarrow +\infty, \text{ which} \\ & \text{means that } \varphi \in \mathfrak{F}_0. \end{aligned}$$

Theorem 4.13. Let $\mu \in \sigma\text{SH}'(B)$ be carried by zero.

Then

$$\mu(\varphi) = \mu(E) \cdot T_0(\varphi) \quad \forall \varphi \in \sigma\text{SH}(B).$$

Proof. If $T_0(\varphi) = 0$ then $\varphi \in \mathfrak{F}_0$ by Lemma 4.12 and we have $\mu(\varphi) = 0$ by Lemma 4.11. Thus $\mu = \alpha \cdot T_0$ for some constant α and since $T_0(E) = 1$ the theorem follows.

Remark. The notation of polar and stable polar set for plurisubharmonic functions were introduced in Kiselman [3]. The polar set of a function $f \in \sigma\text{SH}(U)$ is

$$P(f) = \bigcap_{f_1, f_2} \{z \in U; (f_1 + f_2)(z) = -\infty\}; \quad f = f_1 - f_2, \\ f_1, f_2 \in \text{SH}(U)$$

and the stable polar set of f is

$$P_*(f) = \bigcup_{\omega} \bigcap_{g \in \omega} P(g) \quad (\omega \text{ varies over the neighbourhood of } f).$$

Now, $P_*(f) = \{z \in Z; T_z(f) \neq 0\}$ but $\{z \in U; T_z(f) \neq 0\}$ is a countable set so it follows that the stable polar set of any function $f \in \sigma\text{SH}(U)$ is countable.

References

- [1] ARSOVE, M.G.: Functions representable as differences of subharmonic functions, Trans. Amer. Math. Soc. 75

(1953), 327-365.

- [2] CEGRELL, U.: On the spaces of delta-convex and delta-plurisubharmonic functions and their duals. Manuscript.
- [3] KISELMAN, C.O.: Fonctions delta-convexes, delta-sousharmoniques et delta-plurisousharmoniques, Séminaire d'analyse (P. Lelong), 1975-76. To appear in Springer Lecture Notes in Mathematics.
- [4] SCHAEFER, H.H.: Topological vector spaces, Springer-Verlag 1971.

Uppsala Universitet
Matematiska Institutionen
Syslomanagatan 8, 75223 Uppsala
Sverige

(Oblatum 6.7. 1977)