

Aleksander Błaszczyk

A construction of the Gleason space

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 24 (1983), No. 2, 233--236

Persistent URL: <http://dml.cz/dmlcz/106222>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1983

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A CONSTRUCTION OF THE GLEASON SPACE

A. BŁASZCZYK

Abstract: A construction of the Gleason space.

Key-words: Extremally disconnected space, irreducible map, Gleason space.

Classification: 54G05

---

The Gleason space of a compact (Hausdorff) space  $X$  is the (unique up to homeomorphism) extremally disconnected compact space  $G(X)$  which has an irreducible mapping onto  $X$ ; see e.g. Comfort and Negreponitis [1] page 57 and for the original construction see Gleason [3]. The aim of this note is to show an easy and short construction of the Gleason space. This construction was inspired by some ideas of Mioduszewski [4].

Recall, a space  $X$  is extremally disconnected if for every open set  $U \subset X$ , the closure  $\text{cl}U$  is open. A continuous mapping  $f: X \xrightarrow{\text{onto}} Y$  will be called irreducible if for every closed set  $F \subset X$ ,  $\text{cl}f(F) = Y$  whenever  $F \neq \emptyset$ . If  $X$  is compact, then our definition coincides with the usual one which asserts that  $f$  is irreducible if there does not exist any proper closed subset  $F$  of  $X$  such that  $f$  carries  $F$  onto  $Y$ ; see [1] page 55.

Lemma 1. If  $T$  is a regular topology on a set  $X$ , then there exists a completely regular extremally disconnected

topology  $T^*$  on  $X$  such that  $T \subset T^*$  and the identity  $i: (X, T^*) \rightarrow (X, T)$  is irreducible, where  $(X, T)$  and  $(X, T^*)$  denote  $X$  endowed with the topology  $T$  and  $T^*$  respectively.

**Proof.** Let  $S$  be the set of all regular topologies on  $X$  containing  $T$  and such that

- (1) if  $T' \in S$ , then the identity  $i: (X, T') \rightarrow (X, T)$  is irreducible.

The set  $S$  is ordered by inclusion. Note that if  $L \subset S$  is a chain, then the topology  $T'$  generated by  $\cup L$  belongs to  $S$ . Indeed,  $T'$  is regular because  $\cup L$  is a base of  $T'$  and all topologies in  $L$  are regular. To show that  $i: (X, T') \rightarrow (X, T)$  is irreducible, suppose that  $X - U$  is dense in  $(X, T)$  for some  $U \in \cup L$ . Since  $U \in T''$  for some  $T'' \in L \subset S$ , we get a contradiction with condition (1). So, every chain in  $S$  is bounded. Hence, by the Kuratowski-Zorn Lemma, there exists in  $S$  a maximal element  $T^*$ . It remains to show that  $(X, T^*)$  is extremally disconnected; note that extremally disconnected regular spaces have bases consisting of closed-open sets, hence they are completely regular. Suppose,  $U \in T^*$  and  $\text{cl}U \notin T^*$ , where  $\text{cl}$  denotes the closure in  $T^*$ . Let  $T'$  be the topology generated by  $T^* \cup \{\text{cl}U\}$ . Clearly,  $T'$  is regular and  $T^* \cup \{\bigvee \text{cl}U : U \in T^*\}$  is a base of  $T'$ . If  $X - (\bigvee \text{cl}U)$  is dense in  $(X, T)$ , then  $\bigvee \text{cl}U = \emptyset$ . Indeed,  $X - (\bigvee U)$  is dense in  $(X, T)$  and  $U \cap V \in T^* \in S$ , hence  $U \cap V = \emptyset$ . Therefore  $T' \in S$ , and we get a contradiction with the maximality of  $T$ .

**Lemma 2.** If  $X$  is regular,  $f: X \xrightarrow{\text{onto}} Y$  is continuous and  $f|G$  is irreducible for some dense  $G \subset X$  such that  $f(G) = Y$ , then  $f$  is irreducible.

**Proof.** Suppose the contrary. Then  $f(X - U)$  is dense in

$Y$  for some open  $U \subset X$ ,  $U \neq \emptyset$ . Since  $X$  is regular, we can assume that  $U$  is regularly open, i.e.  $\text{Int } \text{cl}U = U$ . We claim that

(2)  $G - U$  is a dense subset of  $X - U$ .

Indeed, if  $V \subset X$  is regularly open and  $V \cap (X - U) \cap (G - U) = \emptyset$ , then  $G \cap V \subset U$ . Thus  $\text{cl}V \subset \text{cl}U$ , because  $G$  is dense in  $X$ . Since  $V$  and  $U$  are regularly open,  $V \cap (X - U) = \emptyset$ . Hence, the condition (2) is proved. Now, since  $f$  is continuous and  $f(X - U)$  is dense in  $Y$ ,  $f(G - U)$  is also dense in  $Y$ . But  $f|G$  is irreducible, so  $G \cap U = \emptyset$ ; a contradiction.

Construction of the Gleason space: Let  $X$  be a compact space. By Lemma 1, there exist a completely regular extremally disconnected space  $Y$  and an irreducible mapping  $f$  from  $Y$  onto  $X$ . Clearly,  $f$  has a continuous extension  $\bar{f}$  over the Čech-Stone compactification  $\beta Y$ . By Lemma 2,  $\bar{f}$  is irreducible. It is easy to check that  $\beta Y$  is extremally disconnected; see e.g. Engelking [2], page 453. Therefore  $\beta Y$  is the Gleason space of  $X$ .

Remark. By slight modifications, our construction can be extended to some wider classes of spaces. For such generalizations of the Gleason spaces (so called absolutes) see e.g. expository papers by Ponomarev and Šapiro [5] and Woods [6].

#### R e f e r e n c e s

- [1] W.W. COMFORT and S. NEGREPONTIS: The theory of ultrafilters, Springer-Verlag 1974.
- [2] R. ENGELKING: General Topology, Warszawa 1977.
- [3] A.M. GLEASON: Projective topological spaces, Illinois J. Math. 2(1958), 482-489.
- [4] J. MIODUSZEWSKI: On a method which leads to extremally

disconnected covers, Proceedings of the Third Prague Topological Symposium 1971, Academia Prague 1972, 309-311.

- [5] V.I. PONOMAREV and L.B. ŠAPIRO: Absolutes of topological spaces and their continuous maps, Uspekhi Math. 31 (1976), 121-136.
- [6] R.G. WOODS: A survey of absolutes of topological spaces, Topological Structures II, Part 2, Proceedings of the Symposium in Amsterdam 1978, 323-362.

Institute of Mathematics, Silesian University, ul. Bankowa 14,  
Katowice, Poland

(Oblatum 23.2. 1983)