

Tomáš Kepka; Petr Nĕmec
Torsion quasimodules

Commentationes Mathematicae Universitatis Carolinae, Vol. 25 (1984), No. 4, 699--717

Persistent URL: <http://dml.cz/dmlcz/106336>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1984

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

TORSION QUASIMODULES
T. KEPKA and P. NĚMEC

Abstract: Using the preradical approach, torsion and co-cyclic quasimodules are investigated. It is also shown how varieties of quasimodules are constructed from varieties of modules and 3-elementary commutative Moufang loops.

Key words: Commutative Moufang loop, quasimodule, preradical, variety of quasimodules.

Classification: 20N05

1. Introduction

A loop $Q(+)$ satisfying the identity $(x+x)(y+z) = (x+y)(x+z)$ is commutative and it is called a commutative Moufang loop. We denote by $\underline{C}(Q(+))$ the centre of $Q(+)$, i.e. $a \in \underline{C}(Q(+))$ iff $(a+x)y = a+(x+y)$ for all $x, y \in Q$. Then $\underline{C}(Q(+))$ is a normal subloop of $Q(+)$, $\exists x \in \underline{C}(Q(+))$ for every $x \in Q$ and we have the upper central series $0 = \underline{C}_0(Q(+)) \subseteq \underline{C}_1(Q(+)) \subseteq \underline{C}_2(Q(+)) \subseteq \dots \subseteq \underline{C}_n(Q(+)) \subseteq \dots$ of $Q(+)$, where $\underline{C}_{n+1}(Q(+))/\underline{C}_n(Q(+)) = \underline{C}(Q(+)/\underline{C}_n(Q(+)))$ for every $n = 0, 1, 2, \dots$. The loop $Q(+)$ is said to be nilpotent of class at most n if $\underline{C}_n(Q(+)) = Q$. Further, for all $x, y, z \in Q$, the associator $[x, y, z]$ is defined by $[x, y, z] = ((x+y)+z) - (x+(y+z))$ and we

denote by $\underline{A}(Q(+))$ the subloop generated by all associators. Then $\underline{A}(Q(+))$ is a normal subloop of $Q(+)$ and $3x = 0$ for every $x \in \underline{A}(Q(+))$. Moreover, we have the lower central series $Q = \underline{A}_0(Q(+)) \supseteq \underline{A}_1(Q(+)) \supseteq \underline{A}_2(Q(+)) \supseteq \dots \supseteq \underline{A}_n(Q(+)) \supseteq \dots$ of $Q(+)$, where $\underline{A}_{n+1}(Q(+))$ is the subloop generated by all associators $[x, y, z]$, $x \in \underline{A}_n(Q(+))$, $y, z \in Q$, for every $n = 0, 1, 2, \dots$. The loop $Q(+)$ is nilpotent of class at most n iff $\underline{A}(Q(+)) \subseteq \subseteq \underline{C}_{n-1}(Q(+))$ iff $\underline{A}_{n-1}(Q(+)) \subseteq \underline{C}(Q(+))$ and iff $\underline{A}_n(Q(+)) = 0$. As for details and further information concerning commutative Moufang loops, the reader is referred to [5].

Let $Q(+)$ be a commutative Moufang loop. A mapping f of Q into Q is said to be n -central, n being an integer, if $nx + f(x) \in \underline{C}(Q(+))$ for every $x \in Q$. Clearly, f is n -central iff it is m -central, where $m \in \{0, 1, 2\}$ and $n = 3k+m$. The zero endomorphism $x \rightarrow 0$ is 0-central, the automorphism $x \rightarrow -x$ is 1-central and the identical automorphism $x \rightarrow x$ is 2-central. As proved in [9], the set of all (0,1,2-)central endomorphisms of $Q(+)$ is an associative ring with unit.

Throughout the paper, let \underline{R} be an associative ring with unit, $\underline{\Phi}$ a ring homomorphism of \underline{R} onto the three-element field $\underline{Z}_3 = \{0, 1, 2\}$ and $\underline{I} = \text{Ker } \underline{\Phi}$. By a ($\underline{\Phi}$ -special unitary left \underline{R} -) quasimodule Q we mean a commutative Moufang loop $Q(+)$ equipped with scalar multiplication by elements of \underline{R} such that the usual module identities are satisfied, i.e. $r(xy) = rx+ry$, $(r+s)x = rx+sx$, $r(sx) = (rs)x$, $1x = x$ for all $r, s \in \underline{R}$, $x, y \in Q$ and, moreover, $tx \in \underline{C}(Q(+))$ for all $x \in Q$ and $t \in \underline{I}$. The last condition says that the endomorphism $x \rightarrow rx$ of $Q(+)$

is $(-\overline{\Phi}(r))$ -central for all $r \in \underline{R}$. Some information concerning quasimodules and constructions of non-associative quasimodules can be found in [9], [10] and [11].

Let Q be a quasimodule. A subquasimodule P of Q is normal in Q (i.e. P is a block of a congruence of Q) iff $P(+)$ is a normal subloop of $Q(+)$. Now it is easy to see that all the members of the upper central series as well as of the lower central series of $Q(+)$ are normal subquasimodules of Q . Hence Q is said to be nilpotent of class at most n iff the loop $Q(+)$ is so. Further, we shall say that Q is a primitive quasimodule if $rx = 0$ for all $r \in \underline{I}$ and $x \in Q$.

1.1 Example. Every commutative Moufang loop (abelian groups included) is a \underline{Z} -quasimodule, \underline{Z} being the ring of integers and $\overline{\Phi}$ the natural homomorphism of \underline{Z} onto \underline{Z}_3 .

1.2 Example. Let $Q(+)$ be a 3-elementary commutative Moufang loop. Put $rx = \overline{\Phi}(r)x$ for all $r \in \underline{R}$ and $x \in Q$. Then $Q = Q(+, rx)$ is a primitive quasimodule and we see that the classes of primitive quasimodules, \underline{Z}_3 -quasimodules and 3-elementary commutative Moufang loops are equivalent.

1.3 Example. Let $Q(+)$ be a non-associative commutative Moufang loop. Denote by R the ring of central endomorphisms of $Q(+)$. For every $f \in R$ there is a unique $n(f) \in \{0, 1, 2\}$ such that f is $n(f)$ -central and the mapping $f \rightarrow -n(f)$ is a ring homomorphism of R onto \underline{Z}_3 . Now, Q has an R -quasimodule structure.

1.4 Example. A quasigroup G is said to be trimedial if every subquasigroup of G generated by at most 3 elements is medial,

i.e. satisfies the identity $xy.uv = xu.yv$. Trimedial and medial quasigroups appear in many geometrical situations (see e.g. [1], [4], [15], [16]) and important classes of trimedial quasigroups are idempotent trimedial quasigroups (called also distributive quasigroups and determined by the identities $x.yz = xy.zx$, $yz.x = yx.zx$), symmetric trimedial quasigroups (better known as CH-quasigroups or Manin quasigroups and determined by the identities $xy = yx$, $x.xy = y$ and $xx.yz = xy.zx$) and idempotent symmetric trimedial quasigroups (distributive Steiner quasigroups known in an equivalent form as Hall triple systems). Now, let $R = \underline{\mathbb{Z}}[x, y, x^{-1}, y^{-1}]$. As proved in [12], the classes of pointed trimedial quasigroups and centrally pointed quasimodules are equivalent.

1.5 Proposition. Let n be a positive integer.

- (i) Every quasimodule which can be generated by at most n elements is nilpotent of class at most $m = \max(1, n-1)$.
- (ii) The free primitive quasimodule of rank n (and hence the free quasimodule of rank n) is nilpotent of class precisely m .

Proof. (i) See [9, Proposition 4.3]; the assertion is a consequence of the same result for commutative Moufang loops which is known as the Bruck-Slaby's theorem ([5, Theorem VIII, 10.1]).

(ii) See [2, Corollary IV.3.2].

1.6 Proposition. Let Q be a quasimodule. Then both $\underline{\mathbb{A}}(Q)$ and $Q/\underline{\mathbb{C}}(Q)$ are primitive.

Proof. Let $r \in \underline{\mathbb{I}}$. We have $rx \in \underline{\mathbb{C}}(Q)$ for all $x \in Q$ and it is clear that $Q/\underline{\mathbb{C}}(Q)$ is primitive. On the other hand, the mapping

$f: x \rightarrow rx$ is an endomorphism of $Q(+)$ and $\text{Im } f \subseteq \underline{G}(Q(+))$. Consequently, $\text{Im } f$ is associative, hence $\underline{A}(Q) \subseteq \text{Ker } f$ and $ry = 0$ for all $y \in \underline{A}(Q)$.

1.7 Proposition. (i) Every simple quasimodule is a module.

(ii) Every maximal subquasimodule of a nilpotent quasimodule is normal.

(iii) If the ring \underline{R} is left noetherian then every subquasimodule of a finitely generated quasimodule is finitely generated.

Proof. See [9, Lemma 4.8, Corollary 4.11, Proposition 4.6].

Let Q be a quasimodule. For all $a, b \in Q$, define a mapping $i_{a,b}$ by $i_{a,b}(x) = ((x+a)+b) - (a+b)$. Then $i_{a,b}$ is an automorphism of the loop $Q(+)$ and $i_{a,b}(x) = x + [x, a, b]$.

1.8 Lemma. Let P be a subquasimodule of a quasimodule Q . The following conditions are equivalent:

- (i) P is a normal subquasimodule of Q .
- (ii) $i_{a,b}(P) \subseteq P$ for all $a, b \in Q$.
- (iii) $[x, a, b] \in P$ for all $x \in P, a, b \in Q$.

Proof. Easy.

1.9 Lemma. Let Q be a quasimodule and $a, b \in Q$. Then $i_{a,b}$ is an automorphism of the quasimodule Q .

Proof. Let $r \in \underline{R}$ be arbitrary and $s = -\overline{\Phi}(r).1$. We have $(r+s)x \in \underline{G}(Q)$ for every $x \in Q$. Denote $c = (r+s)a$, $d = (r+s)b$. Then $si_{a,b}(x) + i_{a,b}(rx) = i_{a,b}((r+s)x) = (r+s)x$ and $(r+s)i_{a,b}(x) = i_{c,d}((r+s)x) = (r+s)x$. Consequently, $i_{a,b}(rx) = ri_{a,b}(x)$.

2. Preradicals

By a preradical p (for quasimodules) we mean any subfunctor of the identity functor, i.e. p assigns to each quasimodule Q a subquasimodule $p(Q)$ in such a way that $f(p(Q)) \subseteq p(P)$ whenever f is a homomorphism of Q into a quasimodule P . The basic properties of preradicals for quasimodules are the same as in the module case and the reader is referred to [3] and [9] for details. We shall also use the terminology introduced in [3]. Recall that a preradical p is said to be hereditary if $p(P) = P \cap p(Q)$ whenever P is a subquasimodule of a quasimodule Q . A preradical p is said to be cohereditary if $f(p(Q)) = p(P)$ whenever f is a surjective homomorphism of a quasimodule Q onto a quasimodule P . If p is a preradical then by 1.9 $p(Q)$ is a normal subquasimodule of Q for every quasimodule Q . Further, p is said to be a radical if $p(Q/p(Q)) = 0$ for every quasimodule Q .

2.1 Example. It is easy to see that for every integer $n \geq 0$, \underline{A}_n is a cohereditary radical. On the other hand, \underline{C} is not a preradical, since the centre is in general preserved only by surjective homomorphisms.

2.2 Example. For every quasimodule Q , let $\underline{B}(Q)$ denote the least normal subquasimodule of Q such that the corresponding factor is primitive. Then \underline{B} is a cohereditary radical. By 1.6, $\underline{B}(Q) \subseteq \underline{C}(Q)$.

2.3 Lemma. Let Q be a quasimodule generated by a set M . Then

$\underline{B}(Q)$ is just the subloop of $Q(+)$ generated by all rx , $r \in \underline{I}$, $x \in M$.

Proof. Denote by P that subloop. Since \underline{I} is an ideal, it is easy to verify that P is a subquasimodule and $ry \in P$ for all $r \in \underline{I}$, $y \in Q$. Further, P is normal and hence $P = \underline{B}(Q)$.

2.4 Example. For every quasimodule Q , let $\underline{D}(Q)$ denote the least normal subquasimodule such that the corresponding factor is a \underline{Z}_3 -module, i.e. a vector space over \underline{Z}_3 . Then \underline{D} is a cohereditary radical and, moreover, $\underline{D} = \underline{A} + \underline{B}$, i.e. $\underline{D}(Q) = \{x+y; x \in \underline{A}(Q), y \in \underline{B}(Q)\}$ for every quasimodule Q .

2.5 Example. For every quasimodule Q , let $\underline{J}(Q)$ denote the intersection of all maximal normal subquasimodules of Q ; $\underline{J}(Q) = Q$ if there are no such subquasimodules. Clearly, $\underline{J}(Q)$ is just the intersection of all $\text{Ker } f$, f ranging over all homomorphisms of Q into simple (quasi)modules. Thus \underline{J} is a radical and $\underline{A} \subseteq \underline{J} \subseteq \underline{D}$ (use 1.7).

2.6 Proposition. Let Q be a quasimodule.

(i) $\underline{J}(Q)$ is the intersection of all normal maximal subquasimodules of Q .

(ii) If Q is nilpotent then $\underline{J}(Q)$ is the intersection of all maximal subquasimodules of Q .

(iii) Let Q be finitely generated, $P \subseteq \underline{J}(Q)$ be a normal subquasimodule of Q and let f denote the natural homomorphism of Q onto Q/P . If M is a subset of Q such that $f(M)$ generates Q/P then Q is generated by M .

Proof. (i) and (ii) follow from 1.7(i),(ii), respectively.

(iii) Assume, on the contrary, that Q is not generated by M and let N be a finite set generating Q . Further, let K be a subset of N maximal with respect to the property that $M \cup K$ do not generate Q and take $a \in N \setminus K$. There is a subquasimodule G of Q maximal with respect to $M \cup K \subseteq G$ and $a \notin G$. It is easy to see that G is a maximal subquasimodule of Q and hence $P \subseteq G$, a contradiction.

Let \mathcal{F} be a filter of left ideals of the ring \underline{R} . For every quasimodule Q , let $p(Q)$ denote the set of all $x \in Q$ such that $(0:x) = \{r \in \underline{R}; rx = 0\} \in \mathcal{F}$. Then $p = p_{\mathcal{F}}$ is a hereditary preradical.

2.7 Proposition. There is a one-to-one correspondence between hereditary preradicals and filters of left ideals given by

$$\begin{aligned} \mathcal{F} &\longrightarrow p(Q) = \{x \in Q; (0:x) \in \mathcal{F}\}, \\ p &\longrightarrow \mathcal{F}_p = \{I \subseteq \underline{R}; p(\underline{R}/I) = \underline{R}/I\}. \end{aligned}$$

This correspondence induces a one-to-one correspondence between hereditary radicals and radical filters.

Proof. See [9, Proposition 3.2, Lemma 3.3, Lemma 3.4].

Let p be a preradical. Define a preradical \hat{p} by $\hat{p}(Q) = \bigcap \text{Ker } f$, $f: Q \rightarrow P$, $p(P) = 0$. Clearly, \hat{p} is a radical and it is just the least radical containing p .

2.8 Lemma. Let p be a preradical. Then a quasimodule Q is \hat{p} -torsion iff there are an ordinal number α and a chain Q_β , $0 \leq \beta \leq \alpha$, of normal subquasimodules of Q such that $Q_0 = 0$, $Q_\alpha = Q$ and $Q_{\beta+1}/Q_\beta = p(Q/Q_\beta)$ for every $0 \leq \beta < \alpha$, $Q_\beta = \bigcup_{\gamma < \beta} Q_\gamma$ for β limit.

Proof. Obvious.

2.9 Lemma. Let p be a hereditary preradical. Then \hat{p} is a hereditary radical.

Proof. See [9, Proposition 3.7].

Let A be a simple module. Then A is isomorphic to \underline{R}/I for a maximal left ideal I ; we denote by $\mathcal{F} = \mathcal{F}_I$ ($\mathcal{R} = \mathcal{R}_I$) the filter (radical filter) generated by I and we put $\underline{S}_A = p_{\mathcal{F}}$. By 2.7 and 2.9, $\hat{\underline{S}}_A = p_{\mathcal{R}}$.

The field \underline{Z}_3 considered as a module is simple and isomorphic to \underline{R}/I . We shall also use the notation $\underline{L} = \underline{S}_{\underline{Z}_3}$ and $\underline{K} = \underline{I}$. Finally, denote by \mathcal{F} (resp. \mathcal{R}) the filter (radical filter) generated by all maximal left ideals and put $\underline{S} = p_{\mathcal{F}}$, so that $\hat{\underline{S}} = p_{\mathcal{R}}$.

3. \underline{S} - and $\hat{\underline{S}}$ -torsion quasimodules

3.1 Proposition. A quasimodule Q is \underline{L} -torsion iff it is primitive.

Proof. Obvious.

3.2 Proposition. Let Q be a finitely generated primitive quasimodule. Then Q is finite and $|Q| = 3^n$ for some $n \geq 0$.

Proof. The field \underline{Z}_3 is clearly a noetherian ring and the result follows from 1.7(iii) by induction on the nilpotence class of Q .

3.3 Proposition. For every quasimodule Q , $\underline{A}(Q) \subseteq \underline{L}(Q) \subseteq \underline{K}(Q) \subseteq \hat{\underline{S}}(Q)$ and $\underline{A}(Q) \subseteq \underline{L}(Q) \subseteq \underline{S}(Q)$. Consequently, every \underline{K} -torsion-free quasimodule (and also every $\hat{\underline{S}}$ -torsionfree quasimodule) is a module.

Proof. This follows from 1.6.

3.4 Corollary. Let A be a simple module not isomorphic to \underline{Z}_3 . Then every \hat{S}_A -torsion quasimodule is a module.

Now, denote by \mathcal{Y} a representative set of simple modules such that $\underline{Z}_3 \in \mathcal{Y}$.

3.5 Proposition. Let Q be an \underline{S} -torsion quasimodule. Then Q is a direct sum of subquasimodules $\underline{S}_A(Q)$, $A \in \mathcal{Y}$. If $A \neq \underline{Z}_3$ then $\underline{S}_A(Q)$ is a module isomorphic to a direct sum of copies of A . If $A = \underline{Z}_3$ then $\underline{S}_A(Q)$ is a primitive quasimodule.

Proof. First, let $B \in \mathcal{Y}$ be arbitrary and let P be the subquasimodule generated by $\bigcup \underline{S}_A(Q)$, $A \in \mathcal{Y}$, $A \neq B$. Let \mathcal{F} be the filter generated by all maximal left ideals I such that \underline{R}/I is not isomorphic to B and let $a \in \underline{S}_B(Q) \cap P$. Then the cyclic module $\underline{R}a$ is both \underline{S}_B -torsion and p_f -torsion (both \underline{S}_B and p_f are hereditary and P is p_f -torsion), so that $a = 0$. Now, suppose that $B = \underline{Z}_3$. Then $(P + \underline{C}(Q)) / \underline{C}(Q)$ is both \underline{L} -torsion and p_f -torsion, hence it is a zero module and $P \subseteq \underline{C}(Q)$. In particular, P is a module and the sum $\underline{L}(Q) + P$ is direct. Finally, $\underline{A}(Q) \subseteq \underline{L}(Q)$ and $Q / \underline{A}(Q) = (\underline{L}(Q) + P) / \underline{A}(Q)$. From this, $Q = \underline{L}(Q) + P$ and the rest is clear.

3.6 Theorem. Suppose that the ring \underline{R} has primary decompositions. Let Q be an \hat{S} -torsion quasimodule. Then Q is a direct sum of subquasimodules $\hat{S}_A(Q)$, $A \in \mathcal{Y}$. If $A \neq \underline{Z}_3$ then $\hat{S}_A(Q)$ is a module.

Proof. We have $\underline{A}(Q) \subseteq \underline{L}(Q)$ and $Q / \underline{A}(Q)$ is generated by the image of $\bigcup \hat{S}_A(Q)$, $A \in \mathcal{Y}$. Hence Q is generated by this set and we can proceed in the same way as in the proof of 3.5.

3.7 Proposition. Let Q be a finite \underline{K} -torsion module. Then $|Q| =$

$= 3^n$ for some $n \geq 0$.

Proof. The assertion is an easy consequence of 3.2.

3.8 Lemma. Let I be an ideal of \underline{R} and let \mathcal{Q} be the radical filter generated by I . Then:

- (i) A left ideal K belongs to \mathcal{Q} iff for every sequence a_1, a_2, \dots of elements of I there is $n \geq 1$ with $a_n \dots a_1 \in K$.
- (ii) If I is finitely generated as a left ideal then a left ideal K belongs to \mathcal{Q} iff $I^n \subseteq K$ for some $n \geq 1$.

Proof. See e.g. [3, Corollary III.4.6, Proposition III.4.4].

3.9 Corollary. Let Q be a quasimodule. Then $x \in \underline{K}(Q)$ iff for every sequence a_1, a_2, \dots of elements of \underline{I} there is $n \geq 1$ with $a_n \dots a_1 x = 0$. Moreover, if \underline{I} is finitely generated as a left ideal then $x \in \underline{K}(Q)$ iff $\underline{I}^n x = 0$ for some $n \geq 1$.

3.10 Lemma. Let I be a finitely generated maximal left ideal of the ring \underline{R} such that I is an ideal and $A = \underline{R}/I$ is finite. Then every finitely generated \hat{S}_A -torsion module is finite.

Proof. Clearly, I^n/I^{n+1} is finitely generated and \underline{R}/I^n is finite for every $n \geq 1$. By 3.8(ii), every cyclic \hat{S}_A -torsion module is finite and the rest is clear.

3.11 Proposition. Suppose that \underline{I} is finitely generated as a left ideal. Then every finitely generated \underline{K} -torsion quasimodule Q is finite.

Proof. We shall proceed by induction on the nilpotence class n of Q . If $n \leq 1$ then Q is a module and the result follows from 3.10. Now, let $n \geq 2$. We have $\underline{A}_n(Q) = 0$, $\underline{A}_{n-1}(Q) \subseteq \underline{C}(Q)$ and $G = Q/\underline{A}_{n-1}(Q)$ is finite by the induction hypothesis. There are two finite subsets N and M of $\underline{A}_{n-2}(Q)$ and Q , respec-

tively, such that $(N + \underline{A}_{n-1}(Q)) / \underline{A}_{n-1}(Q) = \underline{A}_{n-2}(Q) / \underline{A}_{n-1}(Q)$ and $(M + \underline{A}_{n-1}(Q)) / \underline{A}_{n-1}(Q) = G$. Denote by P the subquasimodule generated by all the associators $[x, y, z]$, $x \in N$, $y, z \in M$. Then P is a finitely generated subquasimodule of $\underline{A}_{n-1}(Q)$ and hence P is a normal finitely generated submodule of Q . In particular, P is finite. On the other hand, if $u \in \underline{A}_{n-2}(Q)$ and $v, w \in Q$ are arbitrary, then $u = x + a$, $v = y + b$, $w = z + c$ for some $x \in N$, $y, z \in M$ and $a, b, c \in \underline{C}(Q)$. We have $[u, v, w] = [x, y, z] \in P$ and we see that $P = \underline{A}_{n-1}(Q)$. Thus both $\underline{A}_{n-1}(Q)$ and G are finite, so that Q is finite, too.

3.12 Proposition. Let I be a finitely generated maximal left ideal of \underline{R} such that I is an ideal and $A = \underline{R}/I$ is finite. Then every finitely generated \underline{S}_A -torsion quasimodule is finite.

Proof. By 3.4, 3.10 and 3.11.

3.13 Theorem. Suppose that every maximal left ideal of \underline{R} is an ideal, finitely generated as a left ideal, maximal ideals commute and every simple module is finite. Let Q be a finitely generated \underline{S} -torsion quasimodule. Then Q is finite and there are $A_1, \dots, A_n \in \mathcal{Y}$ such that Q is isomorphic to the product $\underline{S}_{A_1}(Q) \times \dots \times \underline{S}_{A_n}(Q)$.

Proof. The ring \underline{R} has primary decompositions and the result now follows from 3.6 and 3.12.

3.14 Remark. The assumptions of the preceding theorem are satisfied e.g. if \underline{R} is a finitely generated commutative ring.

3.15 Proposition. Suppose that \underline{R} is left noetherian and every simple module is finite. Then every finitely generated \underline{S} -torsion quasimodule is finite.

Proof. In the situation of Lemma 2.8, ω is finite by 1.7(iii)

and we can proceed by induction, using 3.5 and 3.2.

4. Cocyclic quasimodules

A quasimodule Q is said to be cocyclic if it contains a (non-zero) normal simple submodule A such that A is contained in every non-zero normal subquasimodule of Q .

4.1 Lemma. Let Q be a quasimodule and A be a normal simple subquasimodule of Q . Then $A \subseteq \underline{C}(Q)$.

Proof. Let $a \in A$ and $x, y \in Q$ be arbitrary. Denote by P the subquasimodule generated by a, x, y . Then P is a nilpotent quasimodule and $A \subseteq \underline{C}(P)$ by [9, Lemma 4.7]. Consequently, $(a+x)+y = a+(x+y)$ and we have proved that $A \subseteq \underline{C}(Q)$.

4.2 Proposition. Let Q be a cocyclic quasimodule and A be the normal simple submodule of Q . Then:

- (i) $A \subseteq \underline{C}(Q)$ and $\underline{S}(Q) = \underline{S}_A(Q)$.
- (ii) Q is subdirectly irreducible.
- (iii) Either $A \subseteq \underline{A}(Q)$ and A is isomorphic to \underline{Z}_3 or Q is a module.
- (iv) $\underline{C}(Q)$ is a cocyclic module.

Proof. Easy (use 4.1).

4.3 Corollary. A quasimodule Q is cocyclic iff $\underline{C}(Q) \neq 0$ and Q is subdirectly irreducible. In particular, a nilpotent (resp. finitely generated) quasimodule is cocyclic iff it is subdirectly irreducible.

4.4 Proposition. Suppose that \underline{R} is commutative and noetherian. Let Q be a cocyclic quasimodule and A the normal simple submodule of Q . Then:

- (i) Q is $\underline{\mathbb{S}}_A$ -torsion.
- (ii) If Q is finitely generated and A is finite then Q is finite.
- (iii) If Q is non-associative then A is isomorphic to $\underline{\mathbb{Z}}_3$ and Q is $\underline{\mathbb{K}}$ -torsion.
- (iv) If Q is finitely generated and non-associative then Q is finite.

Proof. First, let Q be a module. By [3, Proposition VI.3.4], $\underline{\mathbb{R}}$ is a stable ring and so the injective hull E of Q is $\underline{\mathbb{S}}_A$ -torsion. Now, suppose that A is isomorphic to $\underline{\mathbb{Z}}_3$. We have $A \subseteq \underline{\mathbb{C}}(Q)$ and $\underline{\mathbb{C}}(Q)$ is $\underline{\mathbb{K}}$ -torsion, since it is a cocyclic module. On the other hand, $Q/\underline{\mathbb{C}}(Q)$ is a primitive quasimodule and thus Q is $\underline{\mathbb{K}}$ -torsion. The rest is clear.

4.5 Example. Let α be an infinite limit ordinal number and $\mathbb{N} = \{a_0, a_1, \dots\}$ be the canonical basis of the vector space $Q = \underline{\mathbb{Z}}_3^{(\alpha)}$. Define a mapping t of \mathbb{N}^3 into Q by $t(a_\beta, a_{\beta+1}, a_{\beta+2}) = a_0$, $t(a_{\beta+1}, a_\beta, a_{\beta+2}) = -a_0$ for $1 \leq \beta \leq \alpha$ and $t(a_\beta, a_\gamma, a_\delta) = 0$ in all remaining cases. It is clear that t can be extended uniquely to a trilinear mapping T of Q^3 into Q such that $T(x, x, y) = T(T(x, y, z), u, v) = T(u, v, T(x, y, z)) = T(u, T(x, y, z), v) = 0$ for all $x, y, z, u, v \in Q$. Put $x * y = x + y + T(x, y, x-y)$ for all $x, y \in Q$. Then $Q' = Q(*)$ is a primitive quasimodule nilpotent of class 2 (see [14]). Moreover, $a \in \underline{\mathbb{C}}(Q')$ iff $T(a, x, y) + T(x, y, a) + T(y, a, x) = 0$ for all $x, y \in Q$. Now it is easy to check that we have $\underline{\mathbb{C}}(Q') = \underline{\mathbb{A}}(Q') = \{0, a_0, -a_0\}$. In particular, Q' is a cocyclic quasimodule. Thus for every infinite cardinal \aleph there is a cocyclic primitive quasimodule (nilpotent of class 2) of cardinality \aleph .

4.6 Example. Let $n \geq 4$, $Q = \underline{\mathbb{Z}}_3^{(n)}$, $a_1 = (1, 0, \dots, 0)$, \dots , $a_n =$

$= (0, \dots, 0, 1)$, $N = \{ a_1, \dots, a_n \}$. Define a mapping t of \mathbb{H}^3 into Q by $t(a_i, a_{i+1}, a_{i+2}) = a_n$, $t(a_{i+1}, a_i, a_{i+2}) = -a_n$ for every $1 \leq i \leq n-3$, $t(a_{n-2}, a_{n-1}, 1) = a_n$, $t(a_{n-1}, a_{n-2}, 1) = -a_n$, $t(a_{n-1}, 1, 2) = a_n$, $t(1, a_{n-1}, 2) = -a_n$. Then t can be extended uniquely to a trilinear mapping T of Q^3 into Q and we put $x * y = x + y + T(x, y, x - y)$. Then $Q' = Q(*)$ is a primitive quasimodule nilpotent of class 2 , $|Q'| = 3^n$ and it is not difficult to check that Q' is cocyclic, provided $n \neq 5$ and $n \neq 6k+1$. By [14] , for every $m \geq 1$, $m \neq 2, 3, 5$, there is a cocyclic primitive quasimodule of order 3^m , nilpotent of class 2. On the other hand, it is clear that there are no cocyclic primitive quasimodules of order 3^2 , 3^3 and it is proved in [8] that there is no cocyclic primitive quasimodule of order 3^5 .

5. Cohereditary radicals and varieties of quasimodules

By a variety of quasimodules we mean a non-empty class of quasimodules closed under cartesian products, subquasimodules and homomorphic images.

5.1 Proposition. (i) If q is a cohereditary radical then the class \mathcal{V}_q of all torsionfree quasimodules is a variety.

(ii) Let \mathcal{V} be a variety of quasimodules. For every quasimodule Q , let $q_{\mathcal{V}}(Q) = \bigcap \text{Ker } f$, $f: Q \rightarrow P$, $P \in \mathcal{V}$. Then $q_{\mathcal{V}}$ is a cohereditary radical.

(iii) The correspondence $q \rightarrow \mathcal{V}_q$ and $\mathcal{V} \rightarrow q_{\mathcal{V}}$ between cohereditary radicals and varieties of quasimodules is bijective.

Proof. Easy.

Let \mathcal{V} be a variety of quasimodules. Denote by \mathcal{V}_m (resp. \mathcal{V}_p) the class of all modules (resp. primitive quasimodules) contained in \mathcal{V} and put $L_{\mathcal{V}} = \alpha_{-}(\mathbb{R})$. Then both \mathcal{V}_m and \mathcal{V}_p

are varieties, $L_{\mathcal{V}}$ is an ideal of \underline{R} , $L_{\mathcal{V}}Q = 0$ for every quasimodule $Q \in \mathcal{V}$ and a module M belongs to \mathcal{V}_m iff $L_{\mathcal{V}}M = 0$.

5.2 Proposition. Let \mathcal{V} be a variety of quasimodules such that $L_{\mathcal{V}} \not\subseteq \underline{I}$. Then $\mathcal{V} = \mathcal{V}_m$ and $\mathcal{V}_p = 0$.

Proof. We have $\underline{R} = L_{\mathcal{V}} + \underline{I}$, so that $Q = \underline{R}Q = 0$ for every $Q \in \mathcal{V}_p$.

5.3 Proposition. Let \mathcal{V} be a variety of quasimodules and let $F \in \mathcal{V}$ be a quasimodule free in \mathcal{V} . Then $\underline{B}(F) \cap \underline{A}(F) = 0$.

Proof. Let X be a free basis of F and let f denote the natural homomorphism of F onto $G = F/\underline{A}(F)$. Then G is a free $\underline{R}/L_{\mathcal{V}}$ -module, $f|X$ is injective and $f(X)$ is a free basis of G . Now, let $a \in \underline{A}(F) \cap \underline{B}(F)$. By 2.3 there are $n \geq 0$, pairwise different $x_1, \dots, x_n \in X$ and elements $r_1, \dots, r_n \in \underline{I}$ with $a = r_1x_1 + \dots + r_nx_n$ (we have $r_ix_i \in \underline{G}(F)$). Consequently, $0 = r_1f(x_1) + \dots + r_nf(x_n)$, $r_1, \dots, r_n \in L$ and $a = 0$.

5.4 Proposition. Let \mathcal{V} be a variety of quasimodules. Then \mathcal{V} is just the variety generated by $\mathcal{V}_m \cup \mathcal{V}_p$.

Proof. This is an easy consequence of 5.3.

5.5 Proposition. Let \mathcal{U} and \mathcal{W} be varieties of modules and primitive quasimodules, respectively. Denote by \mathcal{V} the variety of quasimodules generated by $\mathcal{U} \cup \mathcal{W}$. Then $\mathcal{V}_m = \mathcal{U}$ and $\mathcal{V}_p = \mathcal{W}$.

Proof. Let $F \in \mathcal{V}$ be a free quasimodule of infinite countable rank. Since \mathcal{V} is generated by $\mathcal{U} \cup \mathcal{W}$, F is isomorphic to a subquasimodule of the product $G \times P$, $G \in \mathcal{U}$ and $P \in \mathcal{W}$ being free of infinite countable rank; we shall assume that F is a subquasimodule of $G \times P$. Consequently, $L_{\mathcal{U}}F = 0$ and we

see that $\mathcal{U} = \mathcal{V}_m$. On the other hand, $\underline{B}(F) \subseteq H = G \times 0$, $\underline{B}(F)$ is a normal subquasimodule of $G \times P$ and $F/\underline{B}(F)$ is isomorphic to a subquasimodule of $(H/\underline{B}(F)) \times P \in \mathcal{W}$. However, \mathcal{V}_p is generated by $F/\underline{B}(F)$ and therefore $\mathcal{W} = \mathcal{V}_p$.

Now, denote by \mathcal{Y} the dual lattice of the lattice of ideals of the ring \underline{R} and by \mathcal{P} the lattice of varieties of primitive quasimodules (i.e. the lattice of varieties of 3-elementary commutative Moufang loops). Let \mathcal{X} be the subset of $\mathcal{Y} \times \mathcal{P}$ formed by all couples (I, \mathcal{U}) , where either $\mathcal{U} = 0$, or $0 \neq \mathcal{U} \neq \mathcal{U}_m$ and $I \subseteq \underline{I}$.

5.6 Theorem. The lattice of varieties of quasimodules is isomorphic to the lattice \mathcal{X} .

Proof. Apply 5.2, 5.4 and 5.5.

5.7 Proposition. Let \underline{R} be left noetherian, $n \geq 0$ and \mathcal{V} be a variety of quasimodules nilpotent of class at most n . Then \mathcal{V} is finitely based (i.e. \mathcal{V} can be determined by a finite number of identities).

Proof. Using 1.7(iii), we can proceed in the same way as in the proof of [6, Theorem III].

5.8 Corollary. Let \underline{R} be left noetherian, $n \geq 0$ and \mathcal{V} be a variety of quasimodules nilpotent of class at most n . Then \mathcal{V} contains only countably many subvarieties.

By [13, §10], the lattice of varieties of primitive quasimodules nilpotent of class at most 2 is a three-element chain. Having some information on the lattice of ideals of \underline{R} (e.g. if \underline{R} is a commutative principal ideal ring, etc.) and using 5.6, we can describe the lattice of varieties of quasimodules nilpo-

tent of class at most 2 . Moreover, applying the methods developed in [7] for medial quasigroups, the results are transferable to various classes of trimedial quasigroups (cf. 1.4).

References

- [1] L.Bénéteau, Etude algébrique des espaces barycentres et des espaces planairement affines, Doctoral Thesis, Université Paul Sabatier, Toulouse 1974
- [2] L.Bénéteau, Free commutative Moufang loops and anticommutative graded rings, *J. Algebra* 67 (1980), 1-35
- [3] L.Bican, T.Kepka, P.Němec, Rings, modules, and preradicals, *Lecture Notes in Pure and Appl. Math.* 75, Marcel Dekker, Inc New York 1982
- [4] G.Bol, Gewebe und Gruppen, *Math. Ann.* 114 (1937), 414-431
- [5] R.H.Bruck, A Survey of Binary Systems, "Ergebnisse der Mathematik und ihrer Grenzgebiete", Band 20, Springer Verlag, Berlin - Heidelberg - New York 1958 (1966, 1971)
- [6] T.Evans, Identities and relations in commutative Moufang loops, *J. Algebra* 31 (1974), 508-513
- [7] J.Ježek, T.Kepka, Varieties of abelian quasigroups, *Czech. Math. J.* 27 (1977), 473-503
- [8] T.Kepka, Distributive Steiner quasigroups of order 3^5 , *Comment. Math. Univ. Carolinae* 19 (1978), 389-401
- [9] T.Kepka, Notes on quasimodules, *Comment. Math. Univ. Carolinae* 20 (1979), 229-247
- [10] T.Kepka, P.Němec, Quasimodules generated by three elements, *Comment. Math. Univ. Carolinae* 20 (1979), 249-266
- [11] T.Kepka, P.Němec, Trilinear constructions of quasimodules,

Comment. Math. Univ. Carolinae 21 (1980), 341-354

- [12] T.Kepka, P.Němec, Distributive groupoids and the finite basis property, *J. Algebra* 70 (1981), 229-237
- [13] T.Kepka, P.Němec, Commutative Moufang loops and distributive groupoids of small orders, *Czech. Math. J.* 31 (1981), 633-669
- [14] S.Klossek, Kommutative Spiegelungsräume, *Mitt. Math. Sem. Giessen*, Heft 117, Giessen 1975
- [15] J.I.Manin, *Cubic Forms: Algebra, Geometry, Arithmetic*, North-Holland Publ. Comp., Amsterdam - London - New York, 1974
- [16] J.-P.Soublin, Etude algébrique de la notion de moyenne, *J. Math. Pures et Appl.* 50 (1971), 53-264

Matematicko-fyzikální fakulta, Karlova universita, Sokolovská 83,
186 00 Praha 8, Czechoslovakia

(Oblatum 30.5. 1984)