

Andrzej Walendziak

Semimodularity in lower continuous strongly dually atomic lattices

Archivum Mathematicum, Vol. 32 (1996), No. 3, 163--165

Persistent URL: <http://dml.cz/dmlcz/107571>

Terms of use:

© Masaryk University, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

SEMIMODULARITY IN LOWER CONTINUOUS STRONGLY DUALY ATOMIC LATTICES

ANDRZEJ WALENDZIAK

ABSTRACT. For lattices of finite length there are many characterizations of semimodularity (see, for instance, Grätzer [3] and Stern [6]–[8]). The present paper deals with some conditions characterizing semimodularity in lower continuous strongly dually atomic lattices. We give here a generalization of results of paper [7].

1. PRELIMINARIES

Let L be a lattice. By $[a, b]$ ($a \leq b, a, b \in L$) we denote an interval, that is the set of all $c \in L$ for which $a \leq c \leq b$. For $a, b \in L$ we say that a is a lower cover of b and we write $a \prec b$ if and only if $a < b$ and $[a, b] = \{a, b\}$.

A lattice L is called strongly dually atomic (see [4]), if for any $a, b \in L$ with $a < b$ there is $p \in [a, b]$ such that $p \prec b$. A complete lattice L is lower continuous, if $a \vee \bigwedge C = \bigwedge \{a \vee c : c \in C\}$ for all $a \in L$ and for all chains C in L .

Semimodularity is usually defined as follows:

Definition. A lattice L is called (upper) semimodular, if for all $a, b \in L$, $a \wedge b \prec a$ implies $b \prec a \vee b$.

It is immediate that modular lattices and geometric lattices are semimodular. There are many semimodular lattices being neither modular nor geometric (see Birkhoff [1], Crawley-Dilworth [2], Grätzer [3] and Stern [8]).

2. RESULTS

First we put $J(L) := \{u \in L : u = a \vee b \text{ implies } u = a \text{ or } u = b\}$. The elements of $J(L)$ are called the join-irreducibles of L . In a strongly dually atomic lattice L the unique lower cover of a join-irreducible ($0 \neq u \in J(L)$) is denoted by u' . As a preparation we need.

1991 *Mathematics Subject Classification*: 06C10.

Key words and phrases: lower continuous lattices, strongly dually atomic lattices, semimodular and atomic lattices.

Received April 11, 1995.

Lemma. *Let L be a lower continuous strongly dually atomic lattice. If $p \prec q$ ($p, q \in L$), then there exists a join-irreducible $u \in J(L)$ such that $p \vee u = q$ and $p \wedge u = u'$.*

Proof. Consider the set $T := \{t \in L : p \vee t = q\}$. T is nonempty, since $q \in T$. Let C be a chain in T . The lower continuity yields

$$p \vee \bigwedge C = \bigwedge \{p \vee c : c \in C\} = q.$$

Thus $\bigwedge C \in T$, and T contains a minimal element u , by Zorn's lemma. Clearly, $u \in J(L)$, $p \vee u = q$ and from $u \not\leq p$ it follows that $p \wedge u \leq u'$.

Observe that $u' \leq p$. Indeed, if $u' \not\leq p$, then $p \vee u' = q$, that is $u' \in T$ and $u' < u$, contradicting the minimality of u . Thus we have $u' \leq p \wedge u$. Hence we obtain $p \wedge u = u'$ which completes the proof. \square

Remark 1. For lattices of finite length this lemma was proved in Stern [5] (see also [8], p. 25).

Our main result is the following

Theorem 1. *Let L be a lower continuous strongly dually atomic lattice. Then the following conditions are equivalent:*

- (i) L is semimodular,
- (ii) L satisfies the exchange property for join-irreducibles, i.e.,
for all $u, v \in J(L)$ and arbitrary $b \in L$, $v \leq b \vee u$ and $v \not\leq b \vee u'$
imply $u \leq b \vee v \vee u'$,
- (iii) $b \wedge u \prec u$ implies $b \prec b \vee u$ for all $u \in J(L)$ and $b \in L$.

Proof. Implication (i) \Rightarrow (ii) follows from the proof of Theorem of [7].

(ii) \Rightarrow (iii). Suppose that (iii) does not hold. Let $u \in J(L)$, $b, q \in L$ be elements such that $u \wedge b = u' \prec u$ and $b < q < b \vee u$. Since L is strongly dually atomic there is an element $p \in L$ such that $b \leq p \prec q < b \vee u$.

By Lemma we get the existence of a join-irreducible element $v \in J(L)$ with $p \vee v = q$. It follows that $v \leq b \vee u$ and $v \not\leq b = b \vee u'$. Applying (ii) we obtain that $u \leq b \vee v \vee u' = b \vee v$. Then $b \vee u \leq b \vee v \leq q$. This contradiction shows that (iii) holds.

(iii) \Rightarrow (i). Let $a, b \in L$ be elements for which $a \wedge b \prec a$. Without loss of generality we may assume that a, b are incomparable elements. By Lemma, there exists a join-irreducible element $u \in J(L)$ such that $(a \wedge b) \vee u = a$ and $b \wedge u = u'$. Applying (iii) we get that $b \prec b \vee u$. Since $a \vee b = (a \wedge b) \vee u \vee b = b \vee u$, we obtain $b \prec a \vee b$, which shows that L is semimodular. \square

Remark 2. Since every lattice of finite length is lower continuous and strongly dually atomic, from this theorem it follows Theorem of [7].

We recall that a lattice L is atomistic if every non-zero element of L is a join of atoms. In an atomistic lattice each join-irreducible element is an atom. Then, as the consequence of Theorem 1 we get the following result which is a generalization of Corollary of [7].

Theorem 2. *Let L be an atomistic lower continuous strongly dually atomic lattice. Then the following conditions are equivalent:*

- (i) L is semimodular,
- (ii) L satisfies the Steinitz-MacLane exchange property, that is, for all atoms $p, q \in L$ and for arbitrary $b \in L$, the relations $p \leq b \vee q$ and $p \not\leq b$ imply $q \leq b \vee p$,
- (iii) L has the covering property, i.e., $b \wedge p = 0$ implies $b \prec b \vee p$ for any atom $p \in L$ and for arbitrary $b \in L$.

REFERENCES

- [1] Birkhoff, G., *Lattice Theory*, 3rd edition, American Mathematical Society, Providence, RI, 1967.
- [2] Crawley, P., Dilworth, R. P., *Algebraic Theory of Lattices*, Prentice-Hall, Englewood Cliffs (N.J.), 1973.
- [3] Grätzer, G., *General Lattice Theory*, Birkhäuser Basel, 1978.
- [4] Richter, G., *The Kuroš-Ore Theorem, finite and infinite decompositions*, *Studia Sci. Math. Hungar.*, **17**(1982), 243-250.
- [5] Stern, M., *Exchange properties in lattices of finite length*, *Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe* 31 (1982), 15-26.
- [6] Stern, M., *Semimodularity in lattices of finite length*, *Discrete Math.* **41** (1982), 287-293.
- [7] Stern, M., *Characterizations of semimodularity*, *Studia Sci. Math. Hungar.* **25** (1990), 93-96.
- [8] Stern, M., *Semimodular Lattices*, B. G. Teubner Verlagsgesellschaft, Stuttgart-Leipzig, 1991.

ANDRZEJ WALENDZIAK
 DEPARTMENT OF MATHEMATICS
 AGRICULTURAL AND PEDAGOGICAL UNIVERSITY
 PL-08110 SIEDLCE, POLAND