

Eva Špáníková

On oscillation of differential systems of neutral type

Archivum Mathematicum, Vol. 40 (2004), No. 3, 263--271

Persistent URL: <http://dml.cz/dmlcz/107909>

Terms of use:

© Masaryk University, 2004

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**ON OSCILLATION OF DIFFERENTIAL
SYSTEMS OF NEUTRAL TYPE**

EVA ŠPÁNIKOVÁ

ABSTRACT. We study oscillatory properties of solutions of the systems of differential equations of neutral type.

1. INTRODUCTION

In this paper we consider the neutral differential systems of the form

$$(S) \quad \begin{aligned} [y_1(t) - a(t)y_1(g(t))] &= p_1(t)y_2(t) \\ y_2'(t) &= p_2(t)f(y_1(h(t))), \quad t \in R_+ = [0, \infty). \end{aligned}$$

The following conditions are assumed to hold throughout this paper:

- (a) $a : R_+ \rightarrow (0, \infty)$ is a continuous function;
- (b) $g : R_+ \rightarrow R_+$ is a continuous and increasing function and $\lim_{t \rightarrow \infty} g(t) = \infty$;
- (c) $p_i : R_+ \rightarrow R_+, i = 1, 2$ are continuous functions not identically equal to zero in every neighbourhood of infinity,

$$\int^{\infty} p_1(t) dt = \infty;$$

- (d) $h : R_+ \rightarrow R_+$ is continuous and increasing function and $\lim_{t \rightarrow \infty} h(t) = \infty$;

- (e) $f : R \rightarrow R$ is a continuous function, $uf(u) > 0$ for $u \neq 0$,

and $|f(u)| \geq K|u|$, where $0 < K = \text{const}$.

Let $p_1(t) \equiv 1$ on R_+ and $f(u) = u, u \in R$. Then the system (S) is equivalent to the equation

$$\frac{d^2}{dt^2}[y_1(t) - a(t)y_1(g(t))] - p_2(t)y_1(h(t)) = 0, \quad t \in R_+.$$

2000 *Mathematics Subject Classification*: 34K15, 34K40.

Key words and phrases: neutral differential system, oscillatory (nonoscillatory) solution.

Received July 19, 2002.

The oscillatory properties of the solutions of the equation

$$\frac{d^2}{dt^2}[y_1(t) - a(t)y_1(g(t))] + p_2(t)y_1(h(t)) = 0, \quad t \in R_+.$$

are studied in the paper [8].

The oscillatory theory of neutral differential systems have been studied for example in the papers [1-5], [7], [10,11] and in the references given therein. The more detailed list of publication of the presented topic is given in the monography [6], where the problem of existence of the solutions of neutral differential systems is also studied. The purpose of this paper is to establish some new criteria for the oscillation of the systems (S). Our results are new and extend and improve the know criteria for the oscillation of the differential systems of neutral type.

Let $t_0 \geq 0$. Denote

$$\tilde{t}_0 = \min \{t_0, g(t_0), h(t_0)\}.$$

A function $y = (y_1, y_2)$ is a solution of the system (S) if there exists a $t_0 \geq 0$ such that y is continuous on $[\tilde{t}_0, \infty)$, $y_1(t) - a(t)y_1(g(t))$, $y_2(t)$, are continuously differentiable on $[t_0, \infty)$ and y satisfies (S) on $[t_0, \infty)$.

Denote by W the set of all solutions $y = (y_1, y_2)$ of the system (S) which exist on some ray $[T_y, \infty) \subset R_+$ and satisfy

$$\sup\{|y_1(t)| + |y_2(t)| : t \geq T\} > 0 \quad \text{for any } T \geq T_y.$$

A solution $y \in W$ is nonoscillatory if there exists a $T_y \geq 0$ such that its every component is different from zero for all $t \geq T_y$. Otherwise a solution $y \in W$ is said to be oscillatory.

Denote

$$P_1(t) = \int_0^t p_1(x) dx, \quad t \geq 0.$$

For any $y_1(t)$ we define $z_1(t)$ by

$$(1) \quad z_1(t) = y_1(t) - a(t)y_1(g(t)).$$

2. SOME BASIC LEMMAS

The next Lemma 1 can be derived on the base of Lemma 1 in [5].

Lemma 1. *Let $y \in W$ be a solution of the system (S) with $y_1(t) \neq 0$ on $[t_0, \infty)$, $t_0 \geq 0$. Then y is nonoscillatory, $z_1(t)$, $y_2(t)$ are monotone on some ray $[T, \infty)$, $T \geq t_0$ and $z_1(t) \neq 0$ on $[T, \infty)$.*

Lemma 2 [9, Lemma 2]. *In addition to the conditions (a) and (b) suppose that*

$$1 \leq a(t) \quad \text{for } t \geq 0.$$

Let $y_1(t)$ be a continuous nonoscillatory solution of the functional inequality

$$y_1(t)[y_1(t) - a(t)y_1(g(t))] > 0$$

defined in a neighbourhood of infinity. Suppose that $g(t) > t$ for $t \geq 0$. Then $y_1(t)$ is bounded.

Lemma 3 [9, Lemma 3]. *Assume that*

$$q : R_+ \rightarrow R_+, \quad \delta : R_+ \rightarrow R \quad \text{are continuous functions,} \quad \lim_{t \rightarrow \infty} \delta(t) = \infty$$

and

$$\delta(t) < t \quad \text{for } t \geq 0, \quad \liminf_{t \rightarrow \infty} \int_{\delta(t)}^t q(s) ds > \frac{1}{e}.$$

Then the functional inequality

$$x'(t) + q(t)x(\delta(t)) \leq 0, \quad t \geq 0$$

cannot have an eventually positive solution and

$$x'(t) + q(t)x(\delta(t)) \geq 0, \quad t \geq 0$$

cannot have an eventually negative solution.

3. OSCILLATION THEOREMS

In this section we shall study the oscillation of the solutions of the system (S). In the next theorems $g^{-1}(t)$ and $h^{-1}(t)$ will denote the inverse functions of $g(t)$, $h(t)$ and $\alpha : R_+ \rightarrow R$ is a continuous function.

Theorem 1. *Suppose that*

$$h(t) \leq g(t), \quad t < \alpha(t), \quad h(\alpha(t)) < t \quad \text{for } t \geq 0$$

and

$$(2) \quad \liminf_{t \rightarrow \infty} \int_{h(\alpha(t))}^t K p_1(s) \int_s^{\alpha(s)} p_2(v) dv ds > \frac{1}{e},$$

$$(3) \quad \int_0^\infty \frac{p_2(s) ds}{a(g^{-1}(h(s)))} < \infty, \quad \limsup_{t \rightarrow \infty} \left\{ K P_1(t) \int_{h^{-1}(g(t))}^\infty \frac{p_2(s) ds}{a(g^{-1}(h(s)))} \right\} > 1.$$

Then every solution $y \in W$ of (S) with $y_1(t)$ bounded is oscillatory.

Proof. Let $y = (y_1, y_2) \in W$ be a nonoscillatory solution of (S) with $y_1(t)$ bounded. Without loss of generality we may suppose that $y_1(t)$ is positive and bounded for $t \geq t_0$. From the second equation of (S), (c), (d), (e) we get

$$y_2'(t) \geq 0 \quad \text{for sufficiently large } t \geq t_0.$$

In view of Lemma 1 we have two cases for sufficiently large $t_2 \geq t_1$:

- 1) $y_2(t) > 0, t \geq t_2$;
- 2) $y_2(t) < 0, t \geq t_2$.

Case 1. Because $y_2(t)$ is positive and nondecreasing we have

$$(4) \quad y_2(t) \geq L, \quad t \geq t_2, \quad 0 < L - \text{const.}$$

Integrating the first equation of (S) from t_2 to t and using (1) and (4) we get

$$(5) \quad z_1(t) - z_1(t_2) \geq L \int_{t_2}^t p_1(s) ds, \quad t \geq t_2.$$

From (5) and (c) we have $\lim_{t \rightarrow \infty} z_1(t) = \infty$. From (1) we have

$$z_1(t) < y_1(t), \quad t \geq t_2$$

and this contradicts the fact that $y_1(t)$ is bounded. The Case 1 cannot occur.

Case 2. We can consider two possibilities.

(A) Let $z_1(t) > 0$ for $t \geq t_3$, where $t_3 \geq t_2$ is sufficiently large. We have $z_1(t) < y_1(t)$ and using (e) we get

$$p_2(t)z_1(h(t)) \leq \frac{p_2(t)f(y_1(h(t)))}{K}, \quad t \geq t_4,$$

where $t_4 \geq t_3$ is sufficiently large.

Integrating the second equation of (S) from t to $\alpha(t)$ and then using the last inequality and $y_2(\alpha(t)) < 0$ we obtain

$$-y_2(t) \geq K \int_t^{\alpha(t)} p_2(s)z_1(h(s)) ds, \quad t \geq t_4.$$

Multiplying the last inequality by $p_1(t)$ and then using the monotonicity of $z_1(t)$ we have

$$(6) \quad z_1'(t) + \left(K p_1(t) \int_t^{\alpha(t)} p_2(s) ds \right) z_1(h(\alpha(t))) \leq 0, \quad t \geq t_4.$$

By condition (2) and Lemma 3 the inequality (6) cannot have an eventually positive solution. This is a contradiction.

(B) Let $z_1(t) < 0$ for $t \geq t_3$. From (1) and (e) we have

$$z_1(t) > -a(t)y_1(g(t)), \quad t \geq t_3$$

and

$$(7) \quad -\frac{Kp_2(t)z_1(g^{-1}(h(t)))}{a(g^{-1}(h(t)))} \leq Kp_2(t)y_1(h(t)) \leq p_2(t)f(y_1(h(t))), \quad t \geq t_4,$$

where $t_4 \geq t_3$ is sufficiently large.

In view of the second equation of (S) inequality (7) implies

$$(8) \quad y_2'(t) + \frac{Kp_2(t)z_1(g^{-1}(h(t)))}{a(g^{-1}(h(t)))} \geq 0, \quad t \geq t_4.$$

Integrating (8) from t to t^* and then letting $t^* \rightarrow \infty$ we get

$$(9) \quad y_2(t) \leq \int_t^\infty \frac{Kp_2(s)z_1(g^{-1}(h(s))) ds}{a(g^{-1}(h(s)))}, \quad t \geq t_4.$$

With regard to (3) we get

$$(10) \quad \frac{1}{K} < \limsup_{t \rightarrow \infty} \left\{ P_1(t) \int_{h^{-1}(g(t))}^\infty \frac{p_2(s) ds}{a(g^{-1}(h(s)))} \right\} \leq \limsup_{t \rightarrow \infty} \int_t^\infty \frac{P_1(s)p_2(s) ds}{a(g^{-1}(h(s)))}.$$

We claim that the condition (3) implies

$$(11) \quad \int_T^\infty \frac{P_1(s)p_2(s) ds}{a(g^{-1}(h(s)))} = \infty, \quad T \geq 0.$$

Otherwise if

$$\int_T^\infty \frac{P_1(s)p_2(s) ds}{a(g^{-1}(h(s)))} < \infty,$$

we can choose $T_1 \geq T$ such large that

$$\int_{T_1}^\infty \frac{P_1(s)p_2(s) ds}{a(g^{-1}(h(s)))} < \frac{1}{K},$$

which is a contradiction with (10).

Integrating $\int_T^t P_1(s)y_2'(s) ds$ by parts we have

$$(12) \quad \int_T^t P_1(s)y_2'(s) ds = P_1(t)y_2(t) - P_1(T)y_2(T) - z_1(t) + z_1(T).$$

In this case

$$(13) \quad z_1(t) \leq -M, \quad 0 < M - \text{const.}$$

Using the second equation of (S), (7) and (13) from (12) we get

$$\begin{aligned} \int_T^t P_1(s)y_2'(s) ds &= \int_T^t P_1(s)p_2(s)f(y_1(h(s))) ds \\ &\geq KM \int_T^t \frac{P_1(s)p_2(s) ds}{a(g^{-1}(h(s)))}, \quad t \geq T \geq t_4. \end{aligned}$$

The last inequality together with (12) implies

$$(14) \quad MK \int_T^t \frac{P_1(s)p_2(s) ds}{a(g^{-1}(h(s)))} \leq P_1(t)y_2(t) - P_1(T)y_2(T) - z_1(t) + z_1(T),$$

$$t \geq T \geq t_4.$$

Combining (11) with (14) we get $\lim_{t \rightarrow \infty} (P_1(t)y_2(t) - z_1(t)) = \infty$ and

$$-z_1(t) \geq -P_1(t)y_2(t), \quad t \geq t_5, \quad \text{where } t_5 \geq t_4 \text{ is sufficiently large.}$$

The last inequality together with (9) and the monotonicity of $z_1(t)$ implies

$$\begin{aligned} -z_1(t) &\geq -KP_1(t) \int_t^\infty \frac{p_2(s)z_1(g^{-1}(h(s))) ds}{a(g^{-1}(h(s)))} \\ &\geq -KP_1(t)z_1(t) \int_{h^{-1}(g(t))}^\infty \frac{p_2(s) ds}{a(g^{-1}(h(s)))}, \quad t \geq T \geq t_5 \end{aligned}$$

and

$$1 \geq KP_1(t) \int_{h^{-1}(g(t))}^\infty \frac{p_2(s) ds}{a(g^{-1}(h(s)))}, \quad t \geq t_5,$$

which contradicts (3). This case cannot occur. The proof is complete. \square

Theorem 2. *Suppose that*

$$1 \leq a(t), \quad t < g(t), \quad t < \alpha(t), \quad h(\alpha(t)) < t \quad \text{for } t \geq 0$$

and the conditions (2), (3) are satisfied. Then all solutions of (S) are oscillatory.

Proof. Let $y = (y_1, y_2) \in W$ be a nonoscillatory solution of (S). Without loss of generality we may suppose that $y_1(t)$ is positive for $t \geq t_0$. As in the proof of Theorem 1 we get two cases — Case 1 and Case 2.

Case 1. Analogously as in the Case 1 of the proof of Theorem 1 we can show that $\lim_{t \rightarrow \infty} z_1(t) = \infty$. By Lemma 2 $y_1(t)$ is bounded and from (1) $z_1(t) < y_1(t)$ for sufficiently large t . Then $z_1(t)$ is bounded, which is a contradiction. The Case 1 cannot occur.

Case 2. We can treat this case in the same way as in the proof of Theorem 1 we only remind that $h(t) < g(t)$ follows from the above conditions. The proof is complete. □

Theorem 3. *Suppose that*

$$t < g(t), \quad t < \alpha(t), \quad h(\alpha(t)) < t, \quad t < g(h(t)) \quad \text{for } t \geq 0,$$

$$(15) \quad \limsup_{t \rightarrow \infty} \int_{h^{-1}(g^{-1}(t))}^t K(P_1(t) - P_1(s))p_2(s)a(h(s)) ds > 1,$$

and conditions (2) and (3) hold. Then all solutions of (S) are oscillatory.

Proof. Let $y = (y_1, y_2) \in W$ be a nonoscillatory solution of (S). Without loss of generality we may suppose that $y_1(t)$ is positive for $t \geq t_0$. As in the proof of Theorem 1 we get two cases — Case 1 and Case 2.

Case 1. In this case

$$\begin{aligned} y_1(t) &> a(t)y_1(g(t)), & y_1(t) &> z_1(t), \\ y_1(h(t)) &> a(h(t))y_1(g(h(t))) &> a(h(t))z_1(g(h(t))) \end{aligned}$$

and

$$(16) \quad p_2(t)f(y_1(h(t))) \geq Kp_2(t)y_1(h(t)) > Kp_2(t)a(h(t))z_1(g(h(t))),$$

for $t \geq t_3$, where $t_3 \geq t_2$ is sufficiently large.

Combining the integral identity

$$z_1(t) = z_1(\xi) + (P_1(t) - P_1(\xi))y_2(\xi) + \int_{\xi}^t (P_1(t) - P_1(s))y_2'(s) ds$$

with (16) we get

$$z_1(t) \geq \int_{\xi}^t K(P_1(t) - P_1(s))p_2(s)a(h(s))z_1(g(h(s))) ds, \quad t > \xi \geq t_3.$$

Putting $\xi = h^{-1}(g^{-1}(t))$ and using the monotonicity of $z_1(t)$ from the last inequality we get

$$1 \geq \int_{h^{-1}(g^{-1}(t))}^t K(P_1(t) - P_1(s))p_2(s)a(h(s)) ds,$$

which contradicts the condition (15).

Case 2. We can treat this case in the same way as in the proof of Theorem 1. The proof is complete. \square

Remark 1. Theorems 1-3 remain true if we change the condition (3) by the condition

$$(3') \quad \int \frac{p_2(s) ds}{a(g^{-1}(h(s)))} = \infty$$

because the conditions (3') implies (11).

Example 1. We consider the system

$$(17) \quad \begin{aligned} \left[y_1(t) - \frac{1}{4} y_1(8t) \right]' &= t y_2(t) \\ y_2'(t) &= \frac{c}{t^3} y_1 \left(\frac{t}{4} \right), \quad t > 0, \end{aligned}$$

where c is a positive constant. In this example $a(t) = \frac{1}{4}$, $g(t) = 8t$, $p_1(t) = t$, $P_1(t) = \frac{t^2}{2}$, $p_2(t) = \frac{c}{t^3}$, $h(t) = \frac{t}{4}$, $f(t) = t$ and $K = 1$. We choose $\alpha(t) = 2t$ and calculate the conditions (2), (3) and (15) as follows

$$\begin{aligned} \liminf_{t \rightarrow \infty} \int_{\frac{t}{2}}^t s \int_s^{2s} \frac{c}{v^3} dv ds &= \frac{3c \ln 2}{8}, \\ \limsup_{t \rightarrow \infty} \left\{ \frac{t^2}{2} \int_{32t}^{\infty} \frac{4c ds}{s^3} \right\} &= \frac{c}{1024}, \\ \limsup_{t \rightarrow \infty} \int_{\frac{t}{2}}^t \left(\frac{t^2}{2} - \frac{s^2}{2} \right) \frac{c ds}{4s^3} &= \frac{c}{8} \left(\frac{3}{2} - \ln 2 \right). \end{aligned}$$

For $c > 1024$ all conditions of Theorem 3 are satisfied and so all solutions of (17) are oscillatory.

Acknowledgements. This research was supported by the grant No. 2/3205/23 of Scientific Grant Agency of Ministry of Education of Slovak Republic and Slovak Academy of Sciences.

REFERENCES

- [1] Foltynska, I. and Werbowski, J., *On the oscillatory behaviour of solutions of system of differential equations with deviating arguments*, Colloquia Math. Soc. J. B. **30**, Qualitative theory of Diff. Eq. Szegéd, (1979), 243–256.
- [2] Ivanov, A. F. and Marušiak, P., *Oscillatory properties of systems of neutral differential equations*, Hiroshima Math. J. **24** (1994), 423–434.
- [3] Kitamura, Y. and Kusano, T., *On the oscillation of a class of nonlinear differential systems with deviating argument*, J. Math. Anal. Appl. **66** (1978), 20–36.
- [4] Marušiak, P., *Oscillation criteria for nonlinear differential systems with general deviating arguments of mixed type*, Hiroshima Math. J. **20** (1990), 197–208.
- [5] Marušiak, P., *Oscillatory properties of functional differential systems of neutral type*, Czechoslovak Math. J. **43** (118) (1993), 649–662.
- [6] Marušiak, P. and Olach, R., *Functional differential equations*, Edis, Žilina, (2000) (In Slovak).
- [7] Mihalíková, B., *A note on the asymptotic properties of systems of neutral differential equations*, Proceedings of the International Scientific Conference of Mathematics, University of Žilina (2000), 133–139.
- [8] Mohamad, H. and Olach, R., *Oscillation of second order linear neutral differential equations*, Proceedings of the International Scientific Conference of Mathematics, University of Žilina (1998), 195–201.
- [9] Oláh, R., *Oscillation of differential equation of neutral type*, Hiroshima Math. J. **25** (1995), 1–10.
- [10] Špániková, E., *Oscillatory properties of solutions of three-dimensional differential systems of neutral type*, Czechoslovak Math. J. **50** (125), (2000), 879–887.
- [11] Špániková, E., *Oscillatory properties of solutions of neutral differential systems*, Fasc. Math. **31** (2001), 91–103.

DEPARTMENT OF APPLIED MATHEMATICS, THE UNIVERSITY OF ŽILINA
 J. M. HURBANA 15, 010 26 ŽILINA, SLOVAKIA
 E-mail: spanik@kam.utc.sk