

Alois Klíč

Value distribution of meromorphic functions and their derivatives

Časopis pro pěstování matematiky, Vol. 100 (1975), No. 4, 325--333

Persistent URL: <http://dml.cz/dmlcz/117882>

Terms of use:

© Institute of Mathematics AS CR, 1975

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

VALUE DISTRIBUTION OF MEROMORPHIC FUNCTIONS
AND THEIR DERIVATIVES

ALOIS KLÍČ, Praha

(Received January 23, 1974)

1. Definitions, basic theorems. In the whole paper, meromorphic functions are understood to be meromorphic in \mathbf{C} .

Let f be a meromorphic function, $n(r, f)$ let denote the number of poles of the function f that lie in the disc $|z| \leq r$ and $n(r, a)$ let denote the number of roots of the equation $f(z) = a$ in the disc $|z| \leq r$, each point counted with regard to its multiplicity. Usually it has been put $n(r, f) = n(r, \infty)$.

Let us set

$$N(r, f) = \int_0^r \frac{n(t, f) - n(0, f)}{t} dt + n(0, f) \ln r,$$

$$N(r, a) = \int_0^r \frac{n(t, a) - n(0, a)}{t} dt + n(0, a) \ln r,$$

$$m(r, f) = \frac{1}{2\pi} \int_0^{2\pi} \ln^+ |f(re^{i\varphi})| d\varphi,$$

$$m(r, a) = \frac{1}{2\pi} \int_0^{2\pi} \ln^+ \frac{1}{|f(re^{i\varphi}) - a|} d\varphi.$$

The function $T(r, f) = m(r, f) + N(r, f)$ is called *Nevanlinna characteristic function of f* . Further, let us denote by $\bar{n}(r, f)$ ($\bar{n}(r, a)$, $a \in \mathbf{C}$) the number of different poles (different roots of the equation $f(z) = a$, respectively) that lie in the disc $|z| \leq r$.

Analogously we define

$$\bar{N}(r, f) = \int_0^r \frac{\bar{n}(t, f) - \bar{n}(0, f)}{t} dt + \bar{n}(0, f) \ln r,$$

$$\bar{N}(r, a) = \int_0^r \frac{\bar{n}(t, a) - \bar{n}(0, a)}{t} dt + \bar{n}(0, a) \ln r.$$

1.1 Theorem. (First Main Theorem of the value distribution theory.) For any meromorphic function f , the equation

$$m(r, a) + N(r, a) = T(r, f) + \varepsilon(r, a)$$

holds for each $a \in \mathbf{C}$, where $\varepsilon(r, a) = O(1)$ for $r \rightarrow \infty$.

1.2 Theorem. (Second Main Theorem of the value distribution theory.) Let f be a nonconstant meromorphic function. If $a_1, a_2, \dots, a_q, q \geq 1$, are mutually distinct finite or infinite complex numbers, then

$$\sum_{v=1}^q m(r, a_v) \leq 2 T(r, f) - N_1(r) + S(r, f),$$

where $N_1(r) = N(r, 1/f') + 2 N(r, f) - N(r, f')$ and the remainder $S(r, f)$ satisfies the following conditions: $S(r, f) = o\{T(r, f)\}$ with at most the exception of a set \mathbf{E} of values (r) of finite Lebesgue measure. If f is of finite order, then $S(r, f) = o\{T(r, f)\}$ without exceptional intervals.

1.3 Definition. Let f be a meromorphic function, $a \in \mathbf{C} \cup \{\infty\}$. Let us set

$$\delta(a) = \delta(a, f) = \liminf_{r \rightarrow \infty} \frac{m(r, a)}{T(r, f)} = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{N(r, a)}{T(r, f)},$$

$$\Theta(a) = \Theta(a, f) = 1 - \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}(r, a)}{T(r, f)},$$

$$\vartheta(a) = \vartheta(a, f) = \liminf_{r \rightarrow \infty} \frac{N(r, a) - \bar{N}(r, a)}{T(r, f)}.$$

Recall that $\Theta(a) \geq \delta(a) + \vartheta(a)$. The quantity $\delta(a)$ is called the *deficiency* of the value a , $\vartheta(a)$ is called the *ramification index* of the point a . The value a is called *deficient value* (or *Nevanlinna exceptional value*) if $\delta(a) > 0$.

If the equation $f(z) = a, a \in \mathbf{C}$, has only a finite number of roots, then the value a is called *Picard exceptional value*. The function f must be transcendental. It is clear that every *Picard exceptional value* is *Nevanlinna exceptional value*, but the contrary is not true.

In the following we shall need the following theorems ($S(r, f)$ has the same meaning as in Theorem 1.2):

1.4 Theorem. (Milloux, see [5] or [2].) Let f be a meromorphic function, $k \in \mathbf{N}$ arbitrary. Then

$$(1) \quad T(r, f^{(k)}) \leq (k + 1) T(r, f) + S(r, f).$$

1.5 Theorem. (See [5].) Let f be an entire function, $k \in \mathbf{N}$ arbitrary. Then

$$(2) \quad T(r, f^{(k)}) \leq T(r, f) + S(r, f).$$

1.6 Theorem. (See [8] or [4].) Let f be a meromorphic function for which $S(r, f) = o\{T(r, f)\}$. Then

$$(3) \quad \frac{1}{n+1} \sum_{a \neq \infty} \delta(a, f) \leq \delta(0, f^{(n)})$$

for arbitrary $n \in \mathbf{N}$.

1.7 Note. The relation $S(r, f) = o\{T(r, f)\}$ is valid for every function of finite order, but it need not be fulfilled for functions of infinite order.

1.8 Theorem. (Hayman, see [2].) Let f be a transcendental meromorphic function, $k \in \mathbf{N}$ arbitrary. Then

$$(4) \quad T(r, f) \leq \left(2 + \frac{1}{k}\right) N\left(r, \frac{1}{f}\right) + \left(2 + \frac{2}{k}\right) \bar{N}\left(r, \frac{1}{f^{(k)} - 1}\right) + S(r, f).$$

Corollary. Either the function f assumes every finite value infinitely many times, or $f^{(k)}$ ($k \in \mathbf{N}$) assumes every nonzero finite value infinitely many times.

1.9 Theorem. (Milloux, see [3], p. 132.) Let f be a transcendental meromorphic function, $k \in \mathbf{N}$, $a \in \mathbf{C}$, $b \neq 0$. Then

$$(5) \quad T(r, f) \leq \bar{N}(r, f) + N\left(r, \frac{1}{f-a}\right) + \bar{N}\left(r, \frac{1}{f^{(k)}-b}\right) - N\left(r, \frac{f^{(k)}-b}{f^{(k+1)}}\right) + S(r, f).$$

1.10 Theorem. (See [1], [2].) Let f be a meromorphic function. Then the quantity $\Theta(a)$ vanishes for all except at most a countable set of values a . Furthermore,

$$(6) \quad \sum_a \{\delta(a) + \vartheta(a)\} \leq \sum_a \Theta(a) \leq 2.$$

1.11 Note. In the proof of Theorem 2.2 we shall need the following inequality which was obtained when proving Theorem 1.2 (see [1], [4], [8]):

$$(7) \quad N\left(r, \frac{1}{f'}\right) + \sum_{v=1}^q m(r, a_v) + S(r, f) \leq T(r, f') \leq N(r, f') + m(r, f) + S(r, f).$$

Here a_1, a_2, \dots, a_q are arbitrary finite complex numbers.

2. Some generalizations of Polya-Saxer theorem. The following theorem was proved by Polya and Saxer (1923).

2.1 Theorem. (See [6].) If an entire transcendental function has finite Picard exceptional value, then every its derivative assumes all finite nonzero values infinitely many times.

The Nevanlinna theory is a tool for finer investigation in this direction, and with its help it is possible to prove theorems that generalize, in different ways, the Polya-Saxer theorem.

The following generalization of the Polya-Saxer theorem is a consequence of the inequality (3)."

2.2 Theorem. *Let f be a meromorphic function for which $S(r, f) = o\{T(r, f)\}$ and $\delta(\infty, f) = 1$. If the function f has finite Nevanlinna exceptional value a (that means $\delta(a) > 0$), then every derivative of f assumes all finite nonzero values infinitely many times.*

Proof. First we shall prove (under our suppositions) that the relation $\delta(\infty, f) = 1$ implies the relation $\delta(\infty, f^{(k)}) = 1$, for arbitrary $k \in \mathbf{N}$. From the evident relation

$$\frac{N(r, f')}{T(r, f')} = \frac{N(r, f) + \bar{N}(r, f)}{T(r, f)} \cdot \frac{T(r, f)}{T(r, f')}$$

we get

$$(8) \quad 1 - \delta(\infty, f') = \overline{\lim}_{r \rightarrow \infty} \frac{N(r, f')}{T(r, f')} \leq [1 - \delta(\infty, f) + 1 - \Theta(\infty, f)] \overline{\lim}_{r \rightarrow \infty} \frac{T(r, f)}{T(r, f')}.$$

The inequality (7) yields a lower estimate for $\underline{\lim}_{r \rightarrow \infty} T(r, f')/T(r, f)$ (and thereby also an upper estimate for $\overline{\lim}_{r \rightarrow \infty} T(r, f)/T(r, f')$). From the inequality (7) we obtain easily

$$\begin{aligned} \underline{\lim}_{r \rightarrow \infty} \frac{T(r, f')}{T(r, f)} &\geq \underline{\lim}_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f'}\right) + \sum_1^q m(r, a_\nu)}{T(r, f)} + \\ &+ \underline{\lim}_{r \rightarrow \infty} \frac{S(r, f)}{T(r, f)} \geq \underline{\lim}_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f'}\right)}{T(r, f)} + \sum_{\nu=1}^q \delta(a_\nu, f). \end{aligned}$$

According to the suppositions of our theorem it is $\delta(a) > 0$. If we choose $a_\nu = a$ for any ν , then $\underline{\lim}_{r \rightarrow \infty} T(r, f')/T(r, f) \geq \delta(a) > 0$. Hence also

$$\overline{\lim}_{r \rightarrow \infty} \frac{T(r, f)}{T(r, f')} = \frac{1}{\underline{\lim}_{r \rightarrow \infty} \frac{T(r, f')}{T(r, f)}} < +\infty.$$

If $\delta(\infty, f) = 1$, then $\Theta(\infty, f) = 1$, and from the inequality (8) we get $\delta(\infty, f') = 1$. The validity of the relation $\delta(\infty, f^{(k)}) = 1$, for arbitrary $k \in \mathbf{N}$ is obtained by simple induction.

The relation (6) from Theorem 1.10 applied to $f^{(k)}$ gives, with respect to $\delta(\infty, f^{(k)}) = 1$,

$$\sum_{b \neq 0, \infty} \delta(b, f^{(k)}) + \delta(0, f^{(k)}) \leq 1.$$

From (3) we get

$$\sum_{b \neq 0, \infty} \delta(b, f^{(k)}) \leq 1 - \frac{1}{k+1} \sum_{a \neq \infty} \delta(a, f).$$

According to the suppositions, there exist $a \in \mathbf{C}$ that $\delta(a, f) > 0$. Then $\sum_{b \neq 0, \infty} \delta(b, f^{(k)}) < 1$. Thus $\delta(b, f^{(k)}) < 1$ for every finite nonzero complex value b .

This implies that the function f assumes the value b infinitely many times, for the function f is transcendental.

The corollary of Theorem 1.8 yields a further generalization of the Polya-Saxer theorem.

2.3 Theorem. *Let f be a transcendental meromorphic function. If the function f has finite Picard exceptional value, then every derivative of f assumes all finite nonzero values infinitely many times.*

3. In this section some further generalizations of the Polya-Saxer theorem will be proved.

3.1 Theorem. *Let f be a transcendental meromorphic function for which $\bar{N}(r, f) = o\{T(r, f)\}$ (that means $\Theta(\infty, f) = 1$). If the function f has finite Nevanlinna exceptional value, then every derivative of f assumes all finite nonzero value infinitely many times.*

3.2 Theorem. *Let f be a transcendental meromorphic function for which $S(r, f) = o\{T(r, f)\}$ and $\Theta(\infty, f) = 1$. If $\delta(a, f) > 0$, $a \in \mathbf{C}$, then for every finite nonzero complex number b and arbitrary $k \in \mathbf{N}$, the inequality*

$$(9) \quad \delta(b, f^{(k)}) \leq \Theta(b, f^{(k)}) \leq 1 - \frac{\delta(a, f)}{k+1}$$

holds.

3.3 Theorem. *Let f be an entire transcendental function for which $S(r, f) = o\{T(r, f)\}$. If $\delta(a, f) > 0$, $a \in \mathbf{C}$, then for every finite nonzero complex number b and arbitrary $k \in \mathbf{N}$, the inequality*

$$(10) \quad \delta(b, f^{(k)}) \leq \Theta(b, f^{(k)}) \leq 1 - \delta(a, f)$$

holds.

Proof of Theorem 3.1. We use the inequality (5), choosing the notation so that $\delta(a, f) > 0$. Let us suppose that the equation $f^{(k)}(z) = b$ has only a finite number of roots.

Then

$$\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right) = o\{T(r, f)\}.$$

Let us divide the inequality (5) by $T(r, f)$. We obtain the inequality

$$(11) \quad 1 \leq \frac{\bar{N}(r, f)}{T(r, f)} + \frac{N\left(r, \frac{1}{f - a}\right)}{T(r, f)} + \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f)} - \frac{\bar{N}\left(r, \frac{f^{(k)} - b}{f^{(k+1)}}\right)}{T(r, f)} + \frac{S(r, f)}{T(r, f)}.$$

In (11) we let $r \rightarrow \infty$, $r \notin E$, where the set E has finite Lebesgue measure. The set E is "the exceptional set" from the Nevanlinna Second Main Theorem. Recall that $S(r, f) = o\{T(r, f)\}$ for $r \rightarrow \infty$, $r \notin E$.

Since

$$\delta(a, f) > 0,$$

it is

$$(12) \quad \overline{\lim}_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f - a}\right)}{T(r, f)} = A < 1.$$

According to our notation $\delta(a, f) = 1 - A$. From (11), (12) we obtain the inequality

$$1 \leq \overline{\lim}_{\substack{r \rightarrow \infty \\ r \notin E}} \frac{N\left(r, \frac{1}{f - a}\right)}{T(r, f)} \leq \overline{\lim}_{r \rightarrow \infty} \frac{N\left(r, \frac{1}{f - a}\right)}{T(r, f)} = A.$$

This contradicts the inequality $A < 1$. Therefore the supposition that there is only a finite number of roots of the equation $f^{(k)}(z) = b$ is not correct, hence the function $f^{(k)}$ assumes the value b infinitely many times, QED.

Proof of Theorem 3.2. Again we use the inequality (5). Now we consider functions for which $S(r, f) = o\{T(r, f)\}$. From (11) we obtain the inequality

$$(13) \quad \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f)} \geq 1 - A.$$

The definition of the deficiency and the inequality (1) imply the following inequalities:

$$1 - A \leq \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f)} = (k + 1) \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{(k + 1)T(r, f) + S(r, f)} \leq \\ \leq (k + 1) \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f^{(k)})},$$

$$(k + 1) - (k + 1) \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f^{(k)})} \leq A + k = 1 - \delta(a, f) + k,$$

$$1 - \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f^{(k)})} \leq \frac{k + 1 - \delta(a, f)}{k + 1} = 1 - \frac{\delta(a, f)}{k + 1}.$$

The last inequality may be rewritten in the form

$$\delta(b, f^{(k)}) \leq \Theta(b, f^{(k)}) \leq 1 - \frac{\delta(a, f)}{k + 1} \quad \text{QED.}$$

Proof of Theorem 3.3. The proof of the inequality (10) is analogous to that of the inequality (9), we only use the stricter inequality (2) instead of (1).

From the inequalities (13) and (2) we get

$$1 - A \leq \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f)} = \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f) + S(r, f)} \leq \overline{\lim}_{r \rightarrow \infty} \frac{\bar{N}\left(r, \frac{1}{f^{(k)} - b}\right)}{T(r, f^{(k)})}.$$

Then

$$\Theta(b, f^{(k)}) \leq A = 1 - \delta(a, f),$$

which together with the well-known inequality $\delta(a) \leq \Theta(a)$ yields (10), QED.

4. Remarks.

4.1. The supposition in Theorem 3.1 about the existence of finite Nevanlinna exceptional value is essential. It will be seen in the following

Assertion. Let $f(z) = e^z + z$. Then $\delta(a, f) = 0$ is valid for every $a \in \mathbf{C}$, that means, f has not finite Nevanlinna exceptional value.

The function f is entire, hence the second supposition of Theorem 3.1 is evidently fulfilled. Its derivative $f'(z) = e^z + 1$ never assumes the nonzero value 1.

Proof of the assertion. We use the theorem which supplies in the terms of covering sufficient conditions for the validity of $\delta(a, f) = 0$. First let us recall some concepts. Let $w = f(z)$ be an entire function. Let us denote by \mathcal{F} the Riemann surface of the analytic function f^{-1} . Its ramification points lie just over the points $w_k \in \mathbf{C}$, $w_k = f(z_k)$, for which $f'(z_k) = 0$. Further let us denote by π the natural projection \mathcal{F} on \mathbf{C} ($\pi(\mathfrak{Z}) = w$, where \mathfrak{Z} is the algebraic element of the analytic function f^{-1} , with the centre at w).

Now, the following theorem (see [3], p. 431) is valid:

Theorem. *Let $w = f(z)$ be an entire function, \mathcal{F} the Riemann surface of f^{-1} , $a \in \mathbf{C}$ arbitrary. Let $\Lambda > 0$ and an η -neighbourhood $U(a, \eta)$ exist with the following properties: If $\mathcal{F}_v \subset \mathcal{F}$ is an arbitrary domain over $U(a, \eta)$ ($\pi(\mathcal{F}_v) = U(a, \eta)$), then over every point $w \in U(a, \eta)$ there lie just λ_v points and $1 \leq \lambda_v \leq \Lambda$ (every ramification point of the order m is counted $(m - 1)$ -times). Then $\delta(a, f) = 0$.*

Now let us construct the Riemann surface \mathcal{F} of the analytic function f^{-1} , where $f(z) = e^z + z$. It is $f'(z) = e^z + 1 = 0$ for $z_k = (2k + 1)\pi i$, $k = 0, \pm 1, \pm 2, \dots$. The ramification points lie over $w_k = e^{z_k} + z_k = -1 + (2k + 1)\pi i$. These ramification points are of the first order, for $f''(z) = e^z \neq 0$. The function $e^z + z$ maps conformally the strip (see [7], p. 481, example 7)

$$\Omega_k = \{z \in \mathbf{C}, (2k - 1)\pi < \text{Im } z < (2k + 1)\pi\}$$

onto $\mathbf{C} \setminus (p_1^{(k)} \cup p_2^{(k)})$, where

$$p_1^{(k)} = \{z \in \mathbf{C}, z = x + (2k + 1)\pi i, x \in (-\infty, -1)\},$$

$$p_2^{(k)} = \{z \in \mathbf{C}, z = x + (2k - 1)\pi i, x \in (-\infty, -1)\}.$$

Let \mathcal{F} be constructed so that the k -st sheet \mathcal{P}_k of the plane \mathbf{C} , which is cut along the rays $p_1^{(k)}$ and $p_2^{(k)}$, is connected in the usual way with the $(k + 1)$ -st sheet \mathcal{P}_{k+1} along $p_2^{(k)}$ and with the $(k - 1)$ -st sheet \mathcal{P}_{k-1} along $p_1^{(k)}$, $k = 0, \pm 1, \pm 2, \dots$. We obtain infinitely-many-sheeted surface. All domains lying over an arbitrary disc D with the centre at a point $w \neq w_k$, which contains none of the points w_k , are discs. All domains over an arbitrary disc D_k with the centre at w_k are discs except a single one which is a two-sheeted disc. In this domain, exactly two points lie over each point $w \in D_k$. We can choose $\Lambda = 2$ or $\Lambda = 1$, hence the function $w = e^z + z$ has no Nevanlinna exceptional value.

4.2. Let us compare the suppositions of Theorems 2.2 and 3.1.

a. Theorem 2.2 applies only to functions with $S(r, f) = o\{T(r, f)\}$, while in Theorem 3.1 this condition does not appear.

b. The supposition $\Theta(\infty, f) = 1$ in Theorem 3.1 is weaker than the supposition $\delta(\infty, f) = 1$ in Theorem 2.2, for there exist functions (see [3], p. 517, Theorem 6.4) for which $\Theta(\infty, f) = 1$ and $\delta(\infty, f) < 1$.

4.3. Theorem 3.1 is a consequence of the inequalities (9), (10) for functions with $S(r, f) = o\{T(r, f)\}$. The inequalities (9), (10) say: The number of points at which the function f assumes the value $a \in \mathbf{C}$, is inversely proportional to the multiplicity with which an arbitrary derivative of f assumes every nonzero finite value b .

4.4 Corollary of Theorem 3.3. Let f be a transcendental entire function for which $S(r, f) = o\{T(r, f)\}$. If there exists $a \in \mathbf{C}$ so that $\delta(a, f) = 1$, then

$$(14) \quad \sum_{b \neq 0, \infty} \Theta(b, f^{(k)}) = 0$$

for arbitrary $k \in \mathbf{N}$.

For a comparison we introduce a result due to Ullrich (see [4] or [8]). It is known that for entire functions the inequality

$$(3') \quad \sum_{a \neq \infty} \delta(a, f) \leq \delta(0, f^{(k)}), \quad k \in \mathbf{N} \text{ arbitrary}$$

holds, which is sharper than the inequality (3).

From (3'), analogously to the proof of Theorem 2.2 we obtain

$$\sum_{b \neq 0, \infty} \delta(b, f^{(k)}) \leq 1 - \sum_{a \neq \infty} \delta(a, f).$$

Now, if $\sum_{a \neq 0} \delta(a, f) = 1$, then

$$\sum_{b \neq 0, \infty} \delta(b, f^{(k)}) = 0$$

for arbitrary $k \in \mathbf{N}$.

References

- [1] *R. Nevanlinna*: Eindeutige analytische Funktionen, Berlin, 1953.
- [2] *W. K. Hayman*: Meromorphic functions, Oxford, 1964.
- [3] *A. A. Goldberg, I. V. Ostrovskij*: Распределение значений мероморфных функций, „Наука“, Moscow 1970.
- [4] *H. Wittich*: Neuere Untersuchungen über eindeutige analytische Funktionen, Springer-Verlag, 1955.
- [5] *H. Milloux*: Les fonctions meromorphes et leurs dérivées, Paris, 1940.
- [6] *W. Saxer*: Über die Picardschen Ausnahmewerte sukzessiver Derivierten, Math. Z., 17, 1923.
- [7] *I. Černý*: Základy analýzy v komplexním oboru, „Academia“, Praha 1967.
- [8] *E. Ullrich*: Über die Ableitung einer meromorphen Funktion, Sitzungsber. Preuss. Akad. Wiss., Math.-Phys. Kl. 1929.

Author's address: 166 28 Praha 6, Suchbátarova 1905 (katedra matematiky VŠCHT Praha).