

Cong Xin Wu; Qing-Ying Bu

Köthe dual of Banach sequence spaces $\ell_p[X]$ ($1 \leq p < \infty$) and Grothendieck space

Commentationes Mathematicae Universitatis Carolinae, Vol. 34 (1993), No. 2, 265–273

Persistent URL: <http://dml.cz/dmlcz/118580>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1993

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Köthe dual of Banach sequence spaces $\ell_p[X]$ ($1 \leq p < \infty$) and Grothendieck space

WU CONGXIN, BU QINGYING

Abstract. In this paper, we show the representation of Köthe dual of Banach sequence spaces $\ell_p[X]$ ($1 \leq p < \infty$) and give a characterization of that the spaces $\ell_p[X]$ ($1 < p < \infty$) are Grothendieck spaces.

Keywords: vector-valued sequence space; Köthe dual; GAK-space; Grothendieck space

Classification: 46B16

Let X be a Banach space and X^* its topological dual, and let B_X denote the closed unit ball of X . For $1 \leq p < \infty$, let

$$\ell_p(X) = \left\{ \bar{x} = (x_j) \in X^{\mathbb{N}} : \|\bar{x}\|_{\ell_p} = \left(\sum_{i=1}^{\infty} \|x_i\|^p \right)^{1/p} < \infty \right\},$$

$$\ell_p[X] = \left\{ \bar{x} = (x_j) \in X^{\mathbb{N}} : \text{for each } f \in X^*, \sum_{i \geq 1} |f(x_i)|^p < \infty \right\}.$$

And for each $\bar{x} \in \ell_p[X]$, let

$$\|\bar{x}\|_{(\ell_p)} = \sup \left\{ \left(\sum_{i \geq 1} |f(x_i)|^p \right)^{1/p} : f \in B_{X^*} \right\}.$$

Then $(\ell_p(X), \|\cdot\|_{\ell_p})$ and $(\ell_p[X], \|\cdot\|_{(\ell_p)})$ are Banach spaces (see [1], [2], [3]). For $\bar{x} \in X^{\mathbb{N}}$, let

$$\bar{x} (i \leq n) = (x_1, \dots, x_n, 0, 0, \dots),$$

$$\bar{x} (i > n) = (0, \dots, 0, x_{n+1}, x_{n+2}, \dots).$$

And let

$$\ell_p[X]_r = \{ \bar{x} \in \ell_p[X] : \lim_n \|\bar{x} (i > n)\|_{(\ell_p)} \} = 0.$$

If $\ell_p[X]_r = \ell_p[X]$, then $\ell_p[X]$ is said to be a GAK-space [4].

For a vector-valued sequence space $S(X)$ from X , define its Köthe dual with respect to the dual pair (X, X^*) (see [4]) as follows:

$$S(X)^\times |_{(X, X^*)} = \left\{ \bar{f} = (f_j) \in X^{*\mathbb{N}} : \text{for each } \bar{x} = (x_j) \in S(X), \sum_{i \geq 1} |f_i(x_i)| < \infty \right\}.$$

We denote $S(X)^\times |_{(X, X^*)}$ by $S(X)^\times$ simply if the meaning is clear from the context.

*The authors are supported by The Scientific Fund of China for Ph.D. instructors in Universities.

Lemma 1. For $1 \leq p < \infty$, $(\ell_p[X]_r)^\times = \ell_p[X]^\times$.

PROOF: It is easy to see that $\ell_p[X]^\times \subseteq (\ell_p[X]_r)^\times$. So we only need to prove that $(\ell_p[X]_r)^\times \subseteq \ell_p[X]^\times$.

For $\bar{x} = (x_j) \in \ell_p[X]$ and $t = (t_j) \in c_0$, let $t\bar{x} = (t_j x_j)$. Then $\|t\bar{x} (i > n)\|_{(\ell_p)} \leq \|\bar{x}\|_{(\ell_p)} \sup_{i>n} |t_i|$ implies that $t\bar{x} \in \ell_p[X]_r$. So for $\bar{f} = (f_j) \in (\ell_p[X]_r)^\times$, we have

$$\sum_{i \geq 1} |f_i(t_i x_i)| < \infty.$$

It follows from the fact that $t \in c_0$ was taken arbitrary that

$$\sum_{i \geq 1} |f_i(x_i)| < \infty.$$

Thus, $\bar{f} \in \ell_p[X]^\times$ and the proof is completed. □

Lemma 2. (1) For $1 \leq p < \infty$, $\ell_p[X]^\times \subseteq (\ell_p[X], \|\cdot\|_{(\ell_p)})^*$ and $(\ell_p[X]_r)^\times = (\ell_p[X]_r, \|\cdot\|_{(\ell_p)})^*$.

(2) Let $\|\cdot\|_{(\ell_p)}^*$ denote the dual norm of $\|\cdot\|_{(\ell_p)}$ on the dual space $(\ell_p[X], \|\cdot\|_{(\ell_p)})^*$. Then for each $\bar{x} \in \ell_p[X]$, we have

$$\|\bar{x}\|_{(\ell_p)} = \sup\{|\langle \bar{x}, \bar{f} \rangle| : \bar{f} \in \ell_p[X]^\times, \|\bar{f}\|_{(\ell_p)}^* \leq 1\},$$

where $\langle \bar{x}, \bar{f} \rangle = \sum_{i \geq 1} f_i(x_i)$.

PROOF: See Theorem 2.3 in [3]. □

Lemma 3. Every weak* unconditionally Cauchy series in X^* is weak unconditionally Cauchy series.

PROOF: See the proof of p. 49, Corollary 11 in [5]. □

Lemma 4. For $1 \leq p < \infty$,

$$\ell_p[X^*] = \left\{ \bar{f} = (f_j) \in X^{*\mathbb{N}} : \text{for each } x \in X, \sum_{i \geq 1} |f_i(x)|^p < \infty \right\}.$$

PROOF: Let

$$\Delta = \left\{ \bar{f} = (f_j) \in X^{*\mathbb{N}} : \text{for each } x \in X, \sum_{i \geq 1} |f_i(x)|^p < \infty \right\}.$$

By definition, we only need to prove that $\Delta \subseteq \ell_p[X^*]$.

Let $\bar{f} \in \Delta$ and $t_j \in \ell_q(1/p + 1/q = 1)$. Then $\sum_{i \geq 1} |f_i(t_i x)| < \infty$ for each $x \in X$. So the series $\sum_j t_j f_j$ is weak* unconditionally Cauchy in X^* and hence, it is weak unconditionally Cauchy by Lemma 3. That is, $\sum_{i \geq 1} |F(t_i f_i)| < \infty$ for each $F \in X^{**}$. Since (t_j) is arbitrary in ℓ_q , $\sum_{i \geq 1} |F(f_i)|^p < \infty$ and $\bar{f} = (f_j) \in \ell_p[X^*]$. The proof is completed. □

Lemma 5 (the principle of local reflexivity, [6]). *Let X be a normed space and Z^{**} a finite dimensional subspace of X^{**} . For $\{F_i\}_1^n \subseteq Z^{**}$, $\{f_i\}_1^n \subseteq X^*$ and $\varepsilon > 0$, there exists a linear map $T : Z^{**} \rightarrow X$ such that $\|T\| \leq 1$ and*

$$|f_i(TF_i) - F_i(f_i)| < \varepsilon, \quad i = 1, 2, \dots, n.$$

Proposition 6. $\ell_p[X^{**}]^\times |_{(X^{**}, X^*)} = \ell_p[X]^\times |_{(X, X^*)}$ ($1 \leq p < \infty$).

PROOF: It is easy to see that $\ell_p[X] \subseteq \ell_p[X^{**}]$ implies that

$$\ell_p[X^{**}]^\times |_{(X^{**}, X^*)} \subseteq \ell_p[X]^\times |_{(X, X^*)} .$$

So we only need to prove that

$$\ell_p[X]^\times |_{(X, X^*)} \subseteq \ell_p[X^{**}]^\times |_{(X^{**}, X^*)} .$$

Let $\bar{f} = (f_j) \in \ell_p[X]^\times |_{(X, X^*)}$ and $\bar{F} = (F_j) \in \ell_p[X^{**}]$. For a fixed $n \in \mathbb{N}$, by Lemma 5, there exists a linear map $T_n : \text{span}\{F_i\}_1^n \rightarrow X$ such that $\|T_n\| \leq 1$ and

$$|F_i(f_i)| \leq |f_i(T_n F_i)| + 1/n, \quad i = 1, 2, \dots, n.$$

Now we prove that $\{(T_n F_1, \dots, T_n F_n, 0, 0, \dots)\}_{n=1}^\infty$ is a bounded subset of $\ell_p[X]$. By Theorem 1.5 in [2], we have

$$\begin{aligned} & \| (T_n F_1, \dots, T_n F_n, 0, 0, \dots) \|_{(\ell_p)} \\ &= \sup \left\{ \left\| \sum_{i=1}^n s_i T_n F_i \right\| : s = (s_j) \in B_{\ell_q} \right\} (1/p + 1/q) = 1 \\ &\leq \sup \left\{ \|T_n\| \left\| \sum_{i=1}^n s_i F_i \right\| : s \in B_{\ell_q} \right\} \\ &\leq \sup \left\{ \left\| \sum_{i=1}^\infty s_i F_i \right\| : s \in B_{\ell_q} \right\} = \|\bar{F}\|_{(\ell_p)} . \end{aligned}$$

So $\{(T_n F_1, \dots, T_n F_n, 0, 0, \dots)\}_{n=1}^\infty$ is a bounded subset of $\ell_p[X]$ and hence, $\sigma(\ell_p[X], \ell_p[X]^\times |_{(X, X^*)})$ -bounded. Thus, we have

$$\sum_{i=1}^n |F_i(f_i)| \leq \sum_{i=1}^n |f_i(T_n F_i)| + 1 \leq \sup_{n \geq 1} \left\{ \sum_{i=1}^n |f_i(T_n F_i)| \right\} + 1.$$

Because $n \in \mathbb{N}$ is arbitrary, it follows that

$$\sum_{i=1}^\infty |F_i(f_i)| < \infty .$$

So we prove that $\bar{f} = (f_j) \in \ell_p[X^{**}]^\times |_{(X^{**}, X^*)}$ and this completes the proof. \square

Proposition 7. $(\ell_p[X]^\times |_{(X, X^*)}^\times |_{(X^*, X^{**})} = \ell_p[X^{**}] \ (1 \leq p < \infty).$

PROOF: By Proposition 6, it is easy to see that

$$\begin{aligned} \ell_p[X^{**}] &\subseteq (\ell_p[X^{**}]^\times |_{(X^{**}, X^*)}^\times |_{(X^*, X^{**})} \\ &= (\ell_p[X]^\times |_{(X, X^*)}^\times |_{(X^*, X^{**})} . \end{aligned}$$

So we only need to prove that

$$(\ell_p[X^{**}]^\times |_{(X^{**}, X^*)}^\times |_{(X^*, X^{**})} \subseteq \ell_p[X^{**}].$$

Let $\overline{F} = (F_j) \in (\ell_p[X^{**}]^\times |_{(X^{**}, X^*)}^\times |_{(X^*, X^{**})}$. Since $f \in X^*$ and $t = (t_j) \in \ell_q \ (1/p + 1/q = 1)$ implies that $(t_j f) \in \ell_p[X^{**}]^\times |_{(X^{**}, X^*)}$, $\sum_{i \geq 1} |F_i(t_i f)| < \infty$. Thus, $\sum_{i \geq 1} |F_i(f)|^p < \infty$ and hence, $\overline{F} \in \ell_p[X^{**}]$ by Lemma 4. The proof is completed. \square

Theorem 8. For $1 \leq p < \infty$, $\ell_p \overset{\vee}{\otimes} X$, the injective tensor product of ℓ_p and X , is isometrically isomorphic to the space $(\ell_p[X]_r, \|\cdot\|_{(\ell_p)})$.

PROOF: For each $u = \sum_{i=1}^n t^{(i)} \otimes x_i \in \ell_p \otimes X \ (t^{(i)} \in \ell_p, x_i \in X)$, define $\overline{x}_u = (\sum_{i=1}^n t_1^{(i)} x_i, \sum_{i=1}^n t_2^{(i)} x_i, \dots)$. Then

$$\begin{aligned} \|\overline{x}_u\|_{(\ell_p)} &= \sup \left\{ \left| \sum_{k \geq 1} s_k f \left(\sum_{i=1}^n t_k^{(i)} x_i \right) \right| : f \in B_{X^*}, s \in B_{\ell_q} \right\} \\ &= \sup \left\{ \left| \sum_{i=1}^n f(x_i) \langle t^{(i)}, s \rangle \right| : f \in B_{X^*}, s \in B_{\ell_q} \right\} \\ &= \lambda(u) \quad (\text{see [7, p. 223]}) \quad (1/p + 1/q = 1). \end{aligned}$$

Let $M = \sup_{1 \leq i \leq n} \|x_i\|$. It follows from the above equality that

$$\begin{aligned} \|\overline{x}_u \ (j > k)\|_{(\ell_p)} &= \sup \left\{ \left| \sum_{i=1}^n f(x_i) \langle t^{(i)}, s \ (j > k) \rangle \right| : f \in B_{X^*}, s \in B_{\ell_q} \right\} \\ &\leq M \sup \left\{ \sum_{i=1}^n |\langle t^{(i)}, s \ (j > k) \rangle| : s \in B_{\ell_q} \right\}. \end{aligned}$$

Since B_{ℓ_q} is weak* compact, Theorem 6.11 in [8] implies that

$$\lim_k \|\overline{x}_u \ (j > k)\|_{(\ell_p)} = 0.$$

So, $\overline{x}_u \in \ell_p[X]_r$ and we can define a map $\varphi : \ell_p \otimes X \rightarrow \ell_p[X]_r$ by $\varphi(u) = \overline{x}_u$. It is easy to see that φ is a linear isometrically isomorphic map from $\ell_p \otimes X$ to $\ell_p[X]_r$. Next, we only need to prove that φ is surjective.

For $\bar{x} = (x_1, \dots, x_n, 0, 0, \dots)$, if we let $u = \sum_{i=1}^n e_i \otimes x_i$ (where $e_i = (0, \dots, 0, 1^{(i)}, 0, 0, \dots)$), then $\bar{x} = \varphi(u)$. Notice that $\lim_n \bar{x}$ ($j \leq n$) = \bar{x} for each $\bar{x} \in \ell_p[X]_r$. So φ is surjective and the proof is completed. \square

For two Banach spaces X and Y , let $\mathcal{B}^\wedge(X, Y)$, $I(X, Y)$ and $N(X, Y)$ denote the class of integral bilinear functionals on $X \times Y$, the class of integral operators from X to Y and the class of nuclear operators from X to Y respectively (see p. 232 and p. 170 in [7]).

Theorem 9. *Let $1 \leq p < \infty$ and $1/p + 1/q = 1$. Then $\bar{f} = (f_j) \in \ell_p[X]^\times |_{(X, X^*)}$ if and only if there exist an $r = (r_j) \in \ell_1$ a bounded sequence $\{s^{(n)}\}_{n=1}^\infty$ of ℓ_q and a bounded sequence $\{h_n\}_{n=1}^\infty$ of X^* such that*

$$f_i = \sum_{n \geq 1} r_n s_i^{(n)} h_n, \quad i = 1, 2, \dots$$

PROOF: Necessity. Let $\bar{f} = (f_j) \in \ell_p[X]^\times$. By Lemma 1 and Lemma 2, $\bar{f} \in (\ell_p[X]_r, \|\cdot\|_{(\ell_p)})^*$. So Theorem 8 implies that there is an $\psi^* \in (\ell_p \overset{\vee}{\otimes} X)^*$ corresponding to \bar{f} . By Definition 6 in [7, p. 232], there is an $\psi \in \mathcal{B}^\wedge(\ell_p, X)$ corresponding to ψ^* . Furthermore, by Corollary 12 in [7, p. 237], there exists a $T_\psi \in I(\ell_p, X^*)$ corresponding to ψ . Since Corollary 10 in [7, p. 235] and Theorem 6 in [7, p.248] guarantee that $I(\ell_p, X^*) = N(\ell_p, X^*)$, there are an $r = (r_j) \in \ell_1$, a bounded sequence $\{s^{(n)}\}_{n=1}^\infty$ of ℓ_q and a bounded sequence $\{h_n\}_{n=1}^\infty$ of X^* such that

$$T_\psi(t) = \sum_{n \geq 1} r_n \langle t, s^{(n)} \rangle h_n, \quad \text{for } t \in \ell_p.$$

Now for each $i \geq 1$ and each $x \in X$, by the above corresponding relations, we have

$$T_\psi(e_i)(x) = \psi(e_i, x) = \psi^*(e_i \otimes x) = \langle \varphi(e_i \otimes x), \bar{f} \rangle = f_i(x).$$

Thus

$$f_i = T_\psi(e_i) = \sum_{n \geq 1} r_n s_i^{(n)} h_n, \quad i = 1, 2, \dots$$

Sufficiency. Let $M = \sup_{n \geq 1} \|s^{(n)}\|_q$ and $N = \sup_{n \geq 1} \|h_n\|$. Then, for each $\bar{x} = (x_j) \in \ell_p[X]$, we have

$$\sum_{i \geq 1} |s_i^{(n)} h_n(x_i)| \leq MN \|\bar{x}\|_{(\ell_p)}, \quad \text{for } n \geq 1.$$

And so

$$\sum_{i \geq 1} |f_i(x_i)| \leq \sum_{n \geq 1} |r_n| \sum_{i \geq 1} |s_i^{(n)} h_n(x_i)| < \infty.$$

Therefore, $\bar{f} \in \ell_p[X]^\times$ and the proof is completed. \square

Theorem 10. For $1 < p < \infty$, $(\ell_p[X]^\times, \|\cdot\|_{(\ell_p)}^*)$ is a GAK-space.

PROOF: Let $\bar{f} = (f_j) \in \ell_p[X]^\times$. Then by Theorem 9, there exist an $r = (r_j) \in \ell_1$, a bounded sequence $\{s^{(n)}\}_1^\infty$ of ℓ_q and a bounded sequence $\{h_n\}_1^\infty$ of X^* such that

$$f_i = \sum_{n \geq 1} r_n s_i^{(n)} h_n, \quad i = 1, 2, \dots$$

Without loss of generality, we can assume that $\|s^{(n)}\|_q \leq 1$ and $\|h_n\| \leq 1$ for $n \geq 1$. Thus, for $\bar{x} \in \ell_p[X]$ with $\|\bar{x}\|_{(\ell_p)} \leq 1$, we have

$$\sum_{i \geq 1} |s_i^{(n)} h_n(x_i)| \leq \|\bar{x}\|_{(\ell_p)} \leq 1 \quad \text{for } n \geq 1.$$

So

$$\left\{ \left(\sum_{i \geq 1} |s_i^{(n)} h_n(x_i)| \right)_{n \geq 1} : \|\bar{x}\|_{(\ell_p)} \leq 1 \right\} \subseteq B_{\ell_\infty}.$$

Let $\varepsilon > 0$. Then B_{ℓ_∞} is weak* compact implies that there exists an $n_0 \in \mathbb{N}$ such that

$$\sum_{n > n_0} |r_n| \sum_{i \geq 1} |s_i^{(n)} h_n(x_i)| < \varepsilon/2, \quad \bar{x} \in \ell_p[X], \quad \|\bar{x}\|_{(\ell_p)} \leq 1.$$

Since B_{ℓ_p} is weakly compact set and

$$\left\{ (h_n(x_i))_{i \geq 1} : \bar{x} \in \ell_p[X], \|\bar{x}\|_{(\ell_p)} \leq 1, n \geq 1 \right\} \subseteq B_{\ell_p},$$

there is a $k_0 \in \mathbb{N}$ such that for each $k > k_0$,

$$\sum_{i > k} |s_i^{(n)} h_n(x_i)| < \varepsilon/2 \|r\|_1$$

for $\bar{x} \in \ell_p[X]$ with $\|\bar{x}\|_{(\ell_p)} \leq 1$ and $n = 1, 2, \dots, n_0$. Thus, for each $\bar{x} \in \ell_p[X]$ with $\|\bar{x}\|_{(\ell_p)} \leq 1$ and each $k > k_0$, we have

$$\begin{aligned} \sum_{i > k} |f_i(x_i)| &\leq \sum_{n=1}^{n_0} |r_n| \sum_{i > k} |s_i^{(n)} h_n(x_i)| + \sum_{n > n_0} |r_n| \sum_{i > k} |s_i^{(n)} h_n(x_i)| \\ &\leq \left(\sum_{n=1}^{n_0} |r_n| \right) \varepsilon/2 \|r\|_1 + \sum_{n > n_0} |r_n| \sum_{i \geq 1} |s_i^{(n)} h_n(x_i)| < \varepsilon. \end{aligned}$$

So for $k > k_0$,

$$\begin{aligned} \|\bar{f}(j > k)\|_{(\ell_p)}^* &= \sup \left\{ |\langle \bar{x}, \bar{f}(j > k) \rangle| : \bar{x} \in \ell_p[X], \|\bar{x}\|_{(\ell_p)} \leq 1 \right\} \\ &= \sup \left\{ \left| \sum_{i > k} f_i(x_i) \right| : \|\bar{x}\|_{(\ell_p)} \leq 1 \right\} < \varepsilon. \end{aligned}$$

Therefore, $\lim_k \|\bar{f} (j > k)\|_{(\ell_p)^*}^* = 0$ and $\bar{f} \in (\ell_p[X], \|\cdot\|_{(\ell_p)^*})_r$. □

For $1 < p < \infty$, by Theorem 10 and [4, Proposition 4.9], we have

$$(*) \quad (\ell_p[X]^\times |_{(X, X^*)})^\times |_{(X^*, X^{**})} = (\ell_p[X]^\times |_{(X, X^*)}, \|\cdot\|_{(\ell_p)^*})^*.$$

Now, if we let $\|\cdot\|_{(\ell_p)^{**}}$ denote the dual norm of $\|\cdot\|_{(\ell_p)^*}$ on the dual space $(\ell_p[X]^\times |_{(X, X^*)}, \|\cdot\|_{(\ell_p)^*})^*$, then by Proposition 7 and Lemma 2, the norm $\|\cdot\|_{(\ell_p)}$ on the space $\ell_p[X^{**}]$ is equal to the norm $\|\cdot\|_{(\ell_p)^{**}}$.

Similarly as the proof of Theorem 3.6 in [3], we have the following two propositions.

Proposition 11. *Let $\bar{f}^{(n)} \in \ell_p[X]^\times$ ($1 \leq p < \infty$). Then that*

$$\sigma(\ell_p[X]^\times |_{(X, X^*)}, (\ell_p[X]^\times |_{(X, X^*)})^\times |_{(X^*, X^{**})}) - \lim_n \bar{f}^{(n)} = 0$$

is equivalent to

- (a) $\sigma(X^*, X^{**}) - \lim_n f_i^{(n)} = 0$ for $i \geq 1$; and
- (b) $\sup_{n \geq 1} \|\bar{f}^{(n)}\|_{(\ell_p)^*} < \infty$

if and only if $(\ell_p[X]^\times |_{(X, X^*)})^\times |_{(X^*, X^{**})}, \|\cdot\|_{(\ell_p)^{**}}$ is a GAK-space.

Proposition 12. *Let $\bar{f}^{(n)} \in (\ell_p[X]_r)^*$ ($1 \leq p < \infty$). Then*

$$\sigma((\ell_p[X]_r)^*, \ell_p[X]_r) - \lim \bar{f}^{(n)} = 0$$

if and only if $\sigma(X^*, X) - \lim_n f_i^{(n)} = 0$ for $i \geq 1$ and $\sup_{n \geq 1} \|\bar{f}^{(n)}\|_{(\ell_p)^*} < \infty$.

We say a Banach space X to be a Grothendieck space if every weak* null sequence on X^* is weak null sequence (see [7, p. 179]). Leonard [1] has proved that $\ell_p(X)$ ($1 < p < \infty$) is a Grothendieck space if and only if X is a Grothendieck space. Now we have

Theorem 13. *For $1 < p < \infty$. The Banach space $(\ell_p[X]_r, \|\cdot\|_{(\ell_p)})$ is a Grothendieck space if and only if*

- (i) X is a Grothendieck space; and
- (ii) $(\ell_p[X^{**}], \|\cdot\|_{(\ell_p)})$ is a GAK-space.

PROOF: Sufficiency. By (ii), $(\ell_p[X], \|\cdot\|_{(\ell_p)})$ is a GAK-space, i.e. $\ell_p[X]_r = \ell_p[X]$.

Let $\bar{f}^{(n)} \in (\ell_p[X], \|\cdot\|_{(\ell_p)})^*$ such that

$$\sigma(\ell_p[X]^*, \ell_p[X]) - \lim_n \bar{f}^{(n)} = 0.$$

By Proposition 12, we have

$$\sigma(X^*, X) - \lim_n f_i^{(n)} = 0, \quad i = 1, 2, \dots$$

and

$$\sup_{n \geq 1} \|\bar{f}^{(n)}\|_{(\ell_p)}^* < \infty.$$

By (i), we have

$$\sigma(X^*, X^{**}) - \lim_n f_i^{(n)} = 0, \quad i = 1, 2, \dots$$

By (ii) and Propositions 2, 6, 7, the space $((\ell_p[X]^*)^\times |_{(X^*, X^{**})}, \|\cdot\|_{(\ell_p)})$ is a GAK-space. So Proposition 11 guarantees that

$$\sigma(\ell_p[X]^*, (\ell_p[X]^*)^\times) - \lim_n \bar{f}^{(n)} = 0.$$

It follows from (*) that

$$\sigma(\ell_p[X]^*, \ell_p[X]^{**}) - \lim_n \bar{f}^{(n)} = 0.$$

and completes the sufficiency.

Necessity. To prove (i), let $f_n \in X^*$ ($n \geq 1$) such that

$$\sigma(X^*, X) - \lim_n f_n = 0.$$

Let $\bar{f}^{(n)} = (f_n, 0, 0, \dots)$ for $n \geq 1$. Then $\bar{f}^{(n)} \in (\ell_p[X]_r)^*$ and

$$\sigma((\ell_p[X]_r)^*, \ell_p[X]_r) - \lim_n \bar{f}^{(n)} = 0.$$

So

$$\sigma((\ell_p[X]_r)^*, (\ell_p[X]_r)^{**}) - \lim_n \bar{f}^{(n)} = 0$$

and hence, $\sigma(X^*, X^{**}) - \lim_n f_n = 0$. (i) follows.

For (ii), let $\bar{f}^{(n)} \in \ell_p[X]^\times |_{(X, X^*)}$ such that

$$\sigma(X^*, X^{**}) - \lim_n f_i^{(n)} = 0, \quad i = 1, 2, \dots$$

and

$$\sup_{n \geq 1} \|\bar{f}^{(n)}\|_{(\ell_p)}^* < \infty.$$

By Lemmas 1, 2 and Proposition 12, we have

$$\sigma((\ell_p[X]_r)^*, \ell_p[X]_r) - \lim_n \bar{f}^{(n)} = 0.$$

And hence,

$$\sigma((\ell_p[X]_r)^*, (\ell_p[X]_r)^{**}) - \lim_n \bar{f}^{(n)} = 0.$$

It follows from (*) that

$$\sigma(\ell_p[X]^\times |_{(X, X^*)}, (\ell_p[X]^\times |_{(X, X^*)})^\times |_{(X^*, X^{**})}) - \lim_n \bar{f}^{(n)} = 0.$$

So Propositions 6, 7, 11 imply that $(\ell_p[X]^{**}, \|\cdot\|_{(\ell_p)})$ is a GAK-space and (ii) follows. The proof is completed. □

Corollary 14. *If $\ell_p[X]_r$ ($1 < p < \infty$) is a Grothendieck space, then $\ell_p[X]$ is a GAK-space.*

REFERENCES

- [1] Leonard I.E., *Banach sequence spaces*, J. Math. Anal. Appl. **54** (1976), 245–265.
- [2] Wu Congxin, Bu Qingying, *The vector-valued sequence spaces $\ell_p(X)$ ($1 \leq p < \infty$) and Banach spaces not containing a copy of c_0* , A Friendly Collection of Mathematical Papers I, Jilin Univ. Press, Changchun, China, 1990, 9–16.
- [3] ———, *Banach sequence spaces $\ell_p[X]$ ($1 \leq p < \infty$) and their properties*, to appear.
- [4] Gupta M., Kamthan P.K., Patterson J., *Duals of generalized sequence spaces*, J. Math. Anal. Appl. **82** (1981), 152–168.
- [5] Diestel J., *Sequences and Series in Banach Spaces*, Graduate Texts in Math. 92, Springer-Verlag, 1984.
- [6] Simons S., *Locally reflexivity and (p, q) -summing maps*, Math. Ann. **198** (1972), 335–344.
- [7] Diestel J. Uhl J.J., *Vector Measures*, Amer. Math. Soc. Surveys 15, Providence, 1977.
- [8] Kamthan P.K., Gupta M., *Sequence Spaces and Series*, Lecture Notes 65, Dekker, New York, 1981.

DEPARTMENT OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN, 150006 CHINA

(Received September 23, 1992)