

Aleš Drápal; Tomáš Kepka; Michal Musílek
Group conjugation has non-trivial LD-identities

Commentationes Mathematicae Universitatis Carolinae, Vol. 35 (1994), No. 2, 219--222

Persistent URL: <http://dml.cz/dmlcz/118659>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Group conjugation has non-trivial LD-identities

ALEŠ DRÁPAL, TOMÁŠ KEPKA, MICHAL MUSÍLEK

Abstract. We show that group conjugation generates a proper subvariety of left distributive idempotent groupoids. This subvariety coincides with the variety generated by all cancellative left distributive groupoids.

Keywords: left distributivity, free group

Classification: 20N02, 20A99

Given a group G , define an operation $*$ on G by $x * y = xyx^{-1}$. $G(*)$ is an idempotent left distributive groupoid (i.e. $x * x = x$ and $x * (y * z) = (x * y) * (x * z)$ for any $x, y, z \in G$). It has been an open question, whether a non-trivial free idempotent groupoid occurs as a subgroupoid of $G(*)$ for some group G , particularly for G free. One easily verifies that

$$(\dagger) \quad ((x * y) * y) * x = (x * y) * ((y * x) * x)$$

holds in any group G .

We present here two left distributive idempotent groupoids on four elements which do not satisfy (\dagger) :

$*$	1	2	3	4		$*$	1	2	3	4
1	1	2	4	3		1	1	2	3	4
2	1	2	4	3		2	2	2	4	4
3	4	4	3	1		3	1	1	3	3
4	3	3	1	4		4	1	2	3	4

To see that the identity (\dagger) does not hold, put $x = 2$ and $y = 3$, in both cases.

Note that our result contrasts with an older result of Pierce [4], by which free left distributive idempotent groupoids satisfying $x * (x * y) = y$ can be obtained as subgroupoids of $G(*)$, if the group G is freely generated by involutions.

The presented groupoids have been found by a brute-force computer search. Each of these groupoids has just one non-trivial automorphism, and hence on the set $\{1, 2, 3, 4\}$ there exist 24 groupoids isomorphic to one of them. There are 2183 different idempotent left distributive groupoids on a four-element set, and these groupoids fall into 141 isomorphism classes. The above examples represent the only classes, in which the identity (\dagger) does not hold.

A groupoid $A(*)$ is said to be left (right) cancellative, if $a*b = a*c$ ($b*a = c*a$) implies $b = c$ for any $a, b, c \in A$. If a groupoid is both left and right cancellative, it is called cancellative.

In every idempotent left distributive groupoid $A(*)$ the elastic law $x*(y*x) = (x*y)*x$ holds and we have

$$\begin{aligned} &(((x*y)*y)*(x*y))*(((x*y)*y)*x) = \\ &((x*y)*y)*((x*y)*x) = (x*y)*(y*x) = \\ &(x*y)*((y*x)*(y*x)) = (x*y)*(((y*x)*y)*((y*x)*x)) = \\ &((x*y)*(y*(x*y)))*((x*y)*((y*x)*x)) = \\ &(((x*y)*y)*(x*y))*((x*y)*((y*x)*x)) \end{aligned}$$

for any $x, y \in A$.

Denote by $W(X)$ the absolutely free groupoid of terms with a base X and by \sim the congruence of $W(X)$ induced by the left distributive and idempotent laws. Then $W(X)/\sim$ is a free left distributive idempotent groupoid. We have proved:

Proposition 1. *If $\text{card}(X) \geq 2$, then $W(X)/\sim$ is not left cancellative.*

For terms $t_1, t_2, \dots, t_k \in W(X)$ write $t_1 t_2 \dots t_k$ in place of $t_1(t_2(\dots t_k))$ and define a relation \approx on $W(X)$ by

$$s \approx t \iff a_1 a_2 \dots a_k s \sim a_1 a_2 \dots a_k t$$

for some $a_1, a_2, \dots, a_k \in W(X)$.

Proposition 2. *The relation \approx is a congruence of $W(X)$ and $W(X)/\approx$ is left cancellative. Moreover, $W(X)/\approx$ is free in the variety generated by all left cancellative left distributive idempotent groupoids.*

PROOF: Note that we have $b_1 \dots b_r a_1 \dots a_k t \sim (b_1 \dots b_r a_1) \dots (b_1 \dots b_r a_k) (b_1 \dots b_k t)$ for any $b_1, \dots, b_r, a_1, \dots, a_k, t \in W(X)$. Therefore if $a_1 \dots a_k r \sim a_1 \dots a_k s$ and $b_1 \dots b_m s \sim b_1 \dots b_m t$ hold, then $b_1 \dots b_m a_1 \dots a_k r \sim b_1 \dots b_m a_1 \dots a_k s \sim (b_1 \dots b_m a_1) \dots (b_1 \dots b_m a_k) (b_1 \dots b_m s) \sim (b_1 \dots b_m a_1) \dots (b_1 \dots b_m a_k) (b_1 \dots b_m t) \sim b_1 \dots b_m a_1 \dots a_k t$ holds as well. This proves that \approx is an equivalence. To prove it is a congruence, one proceeds in a similar manner.

$W(X)/\approx$ is thus idempotent, left distributive and left cancellative. If $A = A(\cdot)$ is another left cancellative idempotent left distributive groupoid and $\varphi : W(X) \rightarrow A$ is a homomorphism, then $a_1 \dots a_k s \sim a_1 \dots a_k t$ implies $\varphi(a_1) \dots \varphi(a_k) \varphi(s) = \varphi(a_1) \dots \varphi(a_k) \varphi(t)$. As A is left cancellative, we obtain $\varphi(s) = \varphi(t)$, and we see that $\ker \varphi$ contains \approx . \square

Let V_g be the variety generated by all groupoids $G(*)$ for G a group and $*$ the conjugation. Further, let V_c denote the variety generated by all cancellative left distributive groupoids. From $x*(x*x) = (x*x)*(x*x)$ it follows that every

right cancellative groupoid is idempotent, and hence V_c contains only idempotent groupoids.

It is not complicated to prove that V_g and V_c coincide. However, first we shall describe a generator of V_g .

For a free group $F = F(X)$ with a base X , denote $F_0 = F_0(X)$ the subgroupoid of $F(*)$ generated by X .

Proposition 3. *Let $F = F(x, y)$ be the free group with two generators. Then the groupoid $F_0(*)$ generates the variety V_g .*

PROOF: The set of all $y^i x y^{-i}$, $i \geq 0$, is a free base of a subgroup it generates, and belongs to F_0 . Hence it suffices to prove that $F_0(X)$ generates V_g for a countable infinite set $X = \{x_1, x_2, \dots\}$. We shall show that if a $*$ -identity $t(y_1, \dots, y_k) = s(y_1, \dots, y_k)$ is not satisfied in a group G , it does not hold also in $F_0(X)$. Let $g_1, \dots, g_k \in G$ be such that $t(g_1, \dots, g_k) \neq s(g_1, \dots, g_k)$ and consider a group homomorphism $\varphi : F(X) \rightarrow G$ with $\varphi(x_i) = g_i$ for $1 \leq i \leq k$. Then $\psi = \varphi \upharpoonright F_0(X)$ is a homomorphism of $F_0(X)$ into $G(*)$ and hence $t(x_1, \dots, x_k) \neq s(x_1, \dots, x_k)$. \square

From the proof of the above proposition we also obtain:

Corollary. *$F_0(X)$ is free in V_g for any nonempty base X .*

Proposition 4. *$F_0(X)$ is a right cancellative groupoid for any nonempty set X .*

PROOF: Every $a \in F_0(X)$ is a conjugate of some $x \in X$. Hence a cannot be a non-trivial positive power of any $u \in F(X)$. If $a, b, c \in F_0(X)$ are such that $b * a = c * a$, then $c^{-1}b$ and a commute in $F(X)$. Hence $c^{-1}b$ and a are powers of some element $v \in F(X)$. But then $a = v^{\pm 1}$, and we can assume $a = v$. As the sum of all exponents is zero in $c^{-1}b$ and it is i in a^i , we see that $c = b$. \square

Proposition 5. *The varieties V_g and V_c coincide.*

PROOF: $V_g \subseteq V_c$ by Propositions 3 and 4. Let $A(*) \in V_c$ and suppose first that the left translations of A (i.e. the mappings $L_a : b \rightarrow a * b$) are permutations — such groupoids are often called left quasigroups. By a well known and easy construction, the mapping $a \rightarrow L_a$ is a homomorphism of $A(*)$ into $G(*)$, where G is the group generated by $\{L_a; a \in A\}$. Now, every left cancellative idempotent LD-groupoid can be embedded into a left distributive idempotent left quasigroup [2, Proposition 2.10]. If A is cancellative, then for $a \in A$ the mapping $a \rightarrow L_a$ is faithful, and hence A is in V_g . \square

Denote by V the variety of all idempotent LD-groupoids and by V_l (V_r) its subvariety generated by all left (right) cancellative groupoids. We have $V_g \subseteq V_l \subsetneq V$ and $V_g \subseteq V_r \subseteq V$. It seems to be an open problem which of the indicated inclusions are proper ones. We conjecture that V_g equals V_l and V_r coincides with V .

The free non-idempotent groupoids have received a lot of interest recently (for example, see [1] and [3]). We hope that this short contribution will help to focus interest also to the idempotent case.

The authors were recently informed that the identity (\dagger), the respective four-element groupoids and the Propositions 1 and 2 were found out independently by Larue [5], [6]. Note also that the congruence \approx appears already in [2].

REFERENCES

- [1] Dehornoy P., *Braid groups and left distributive structures*, Transactions AMS, to appear.
- [2] Kepka T., *Notes on left distributive groupoids*, Acta Univ. Carolinae - Math. et Ph. **22** (1981), 23–37.
- [3] Laver R., *The left distributive law and the freeness of an algebra of elementary embeddings*, Advances in Mathematics **91** (1992), 209–231.
- [4] Pierce R.S., *Symmetric groupoids*, Osaka J. Math. **15** (1978), 51–76.
- [5] Larue D.M., Unpublished notes, 1992.
- [6] ———, Ph. D. Thesis, University of Colorado, 1994.

Aleš Drápal and Tomáš Kepka
FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83,
186 00 PRAHA 8, CZECH REPUBLIC

Michal Musílek
U PLOVÁRNÝ 1415, 504 01 NOVÝ BYDŽOV, CZECH REPUBLIC

(Received November 15, 1993)