

Václav Slavík

A note on convex sublattices of lattices

Commentationes Mathematicae Universitatis Carolinae, Vol. 36 (1995), No. 1, 7--9

Persistent URL: <http://dml.cz/dmlcz/118725>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1995

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A note on convex sublattices of lattices

VÁCLAV SLAVÍK

Abstract. Let $CSub(\mathbf{K})$ denote the variety of lattices generated by convex sublattices of lattices in \mathbf{K} . For any proper variety \mathbf{V} , the variety $CSub(\mathbf{V})$ is proper. There are uncountably many varieties \mathbf{V} with $CSub(\mathbf{V}) = \mathbf{V}$.

Keywords: lattice, convex sublattice, variety

Classification: 06B25

Let A be a lattice. Denote by $Int(A)$ the lattice of all intervals of A and by $CSub(A)$ the lattice of all convex sublattices of A . The empty set is considered to be in both $Int(A)$ and $CSub(A)$. For a variety \mathbf{K} of lattices, let $CSub(\mathbf{K})$ denote the variety of lattices generated by $\{CSub(A); A \in \mathbf{K}\}$.

The aim of the paper is to show that there exist uncountably many varieties of lattices \mathbf{K} with $CSub(\mathbf{K}) = \mathbf{K}$ and that for any proper subvariety \mathbf{K} of lattices $CSub(\mathbf{K})$ is also proper. Thus we give a partial answer to the problem I. 10 posed in G. Grätzer [1].

Lemma 1. *If p is a lattice term in k variables and A_1, \dots, A_k convex sublattices of a lattice A , then*

$$p(A_1, \dots, A_k) = \bigcup \{p(I_1, \dots, I_k); I_j \subseteq A_j \text{ and } I_j \in Int(A)\}.$$

PROOF: (By induction on the length of p) Evidently, $p(I_1, \dots, I_k) \subseteq p(A_1, \dots, A_k)$ for any intervals $I_j \subseteq A_j$. We must show that every element of $p(A_1, \dots, A_k)$ belongs to $p(I_1, \dots, I_k)$ for some intervals I_j of A_j . If p is a variable, it is clear. Let x be an element of $t_1(A_1, \dots, A_k) \vee t_2(A_1, \dots, A_k)$ for some terms t_1, t_2 . We shall show that $x \in t_1(I_1, \dots, I_k) \vee t_2(I_1, \dots, I_k)$ for some intervals $I_j \subseteq A_j$. If either $t_1(A_1, \dots, A_k) = \emptyset$ or $t_2(A_1, \dots, A_k) = \emptyset$, then we get it by induction. In the opposite case there exist elements $a_1, b_1 \in t_1(A_1, \dots, A_k)$ and $a_2, b_2 \in t_2(A_1, \dots, A_k)$ such that $a_1 \wedge a_2 \leq x \leq b_1 \vee b_2$. By assumption there exist intervals J_j, K_j, L_j, M_j of A_j such that $a_1 \in t_1(J_1, \dots, J_k)$, $b_1 \in t_1(K_1, \dots, K_k)$, $a_2 \in t_2(L_1, \dots, L_k)$, $b_2 \in t_2(M_1, \dots, M_k)$. It is evident that $a_1, a_2, b_1, b_2, x \in t_1(I_1, \dots, I_k) \vee t_2(I_1, \dots, I_k)$, where $I_j = J_j \vee K_j \vee L_j \vee M_j$. The rest is easy. \square

For any lattice A , $Int(A)$ is a sublattice of $CSub(A)$. Combining this fact with Lemma 1 we obtain the following proposition.

Proposition 1. *Let A be a lattice and \mathbf{V} a variety of lattices. Then $\text{Int}(A) \in \mathbf{V}$ if and only if $\text{CSub}(A) \in \mathbf{V}$.*

For a bounded lattice B (with the least element u and the greatest element v), denote by $\mathcal{L}(B)$ the lattice pictured in Fig. 1. Denote by $\mathcal{L}_0(B)$ the lattices obtained from $\mathcal{L}(B)$ by excluding its least element.

Let \mathbf{V} be a variety of lattices. Denote by $\mathcal{L}(\mathbf{V})$ the class of all lattices L such that whenever $\mathcal{L}(B)$ is a sublattice of L , then $B \in \mathbf{V}$.




Figure 1: $\mathcal{L}(B)$

It is shown in [2] that $\mathcal{L}(\mathbf{V})$ is a variety of lattices. Moreover, $\mathcal{L}(\mathbf{V})$ is proper if \mathbf{V} is and $\mathcal{L}(\mathbf{V}) \neq \mathcal{L}(\mathbf{W})$ for any pair of varieties $\mathbf{V} \neq \mathbf{W}$. If variety \mathbf{V} is self-dual, then $\mathcal{L}(\mathbf{V})$ is self-dual, too.

Proposition 2. *Let A be a lattice and \mathbf{V} a self-dual variety of lattices. Then $A \in \mathcal{L}(\mathbf{V})$ if and only if $\text{Int}(A) \in \mathcal{L}(\mathbf{V})$.*

PROOF: Without loss of generality we may assume that A is a bounded lattice. The mapping h of A into $\text{Int}(A)$ defined by $h(a) = [0, a]$ (0 is the least element of A) is an embedding of A into $\text{Int}(A)$. So any variety containing $\text{Int}(A)$ must contain also the lattice A . Now suppose that $A \in \mathcal{L}(\mathbf{V})$. Let $\mathcal{L}(B)$ be a sublattice of $\text{Int}(A)$ for some bounded lattice B . For any element $b = [b_1, b_2] \in \mathcal{L}(B) \subseteq \text{Int}(A)$ different from the least element of $\mathcal{L}(B)$, denote $h(b) = (b_1, b_2)$. Clearly, the mapping h is an embedding of the partial lattice $\mathcal{L}_0(B)$ into $A^* \times A$, where A^* denotes the lattice dual to A . One can easily show that the sublattice of $A^* \times A$ generated by $h(\mathcal{L}_0(B))$ is isomorphic to $\mathcal{L}(B)$ (see [2]). Since $A^* \times A \in \mathcal{L}(\mathbf{V})$, we get $B \in \mathbf{V}$ and thus $\text{Int}(A) \in \mathcal{L}(\mathbf{V})$.

Since any proper variety \mathbf{V} of lattices is a subvariety of a proper self-dual variety \mathbf{W} and \mathbf{W} is a subvariety of a proper variety $\mathcal{L}(\mathbf{W})$ which is, by Propositions 1 and 2, closed under the formation of lattice of all convex sublattices, we get the following result.

Theorem 1. *For any proper variety \mathbf{V} of lattices, the variety $CSub(\mathbf{V})$ is proper.*

Since there exist uncountably many proper self-dual varieties of lattices (see [3], [4]) and $\mathcal{L}(\mathbf{V}) \neq \mathcal{L}(\mathbf{W})$ if $\mathbf{V} \neq \mathbf{W}$, we have, by Propositions 1 and 2, the following theorem.

Theorem 2. *There exist uncountably many self-dual varieties \mathbf{V} of lattices such that $CSub(\mathbf{V}) = \mathbf{V}$.*

REFERENCES

- [1] Grätzer G., *General Lattice Theory*, Birkhäuser Verlag, 1978.
- [2] Adams M.E., Sichler J., *Lattices with unique complementation*, *Pacif. J. Math.* **92** (1981), 1–13.
- [3] McKenzie R., *Equational bases for lattice theories*, *Math. Scand.* **27** (1970), 24–38.
- [4] Wille R., *Primitive subsets of lattices*, *Alg. Universalis* **2** (1972), 95–98.

AGRICULTURE UNIVERSITY, DEPARTMENT OF MATHEMATICS, 165 21 PRAHA 6 – SUCHBOL,
CZECH REPUBLIC

(Received August 2, 1994)