

Venu G. Menon

A note on topology of  $\mathcal{Z}$ -continuous posets

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 37 (1996), No. 4, 821--824

Persistent URL: <http://dml.cz/dmlcz/118890>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## A note on topology of $Z$ -continuous posets

VENU G. MENON

*Abstract.*  $Z$ -continuous posets are common generalizations of continuous posets, completely distributive lattices, and unique factorization posets. Though the algebraic properties of  $Z$ -continuous posets had been studied by several authors, the topological properties are rather unknown. In this short note an intrinsic topology on a  $Z$ -continuous poset is defined and its properties are explored.

*Keywords:*  $Z$ -continuous posets, intrinsic topology

*Classification:* 06B30, 06B35, 54F05

### Introduction

$Z$ -continuous posets were introduced by Wright, Wagner, and Thatcher [WWT] as a generalization of continuous lattices. The family of  $Z$ -continuous posets in fact includes completely distributive lattices ([R]), and unique factorization posets ([M]). The algebraic properties of  $Z$ -continuous posets had been studied by several authors eg. [BE], [N], [V1], [V2]. Though topological methods play an important role in the theory of continuous lattices from its inception, the topological properties of  $Z$ -continuous posets have never been studied. In this short note, we define an intrinsic topology on a  $Z$ -continuous poset, and point out some pleasant properties of this topology. Of course a lot more need to be done in this direction.

A *subset system*  $Z$  is a function which assigns to each poset  $P$  a set  $Z(P)$  of subsets of  $P$  such that (i) for all  $P$ , all singletons of  $P$  are in  $Z(P)$ , and (ii) if  $f : P \rightarrow Q$  is a monotone function between posets, and  $S$  is  $Z(P)$ , then  $f(S)$  is in  $Z(Q)$  ([WWT]). Some examples of the subset systems are all subsets, directed subsets, and finite subsets; see [V1] and [V2] for more examples. For  $S \in Z(P)$ ,  $\downarrow S$  is called a  $Z$ -ideal. The poset (ordered by inclusion) of all  $Z$ -ideals of a poset  $P$  is denoted by  $I_Z(P)$ . Let  $P$  be a poset. For  $x, y \in P$ ,  $x$  is said to be  $Z$ -waybelow  $y$  (written  $x \ll y$ ) if whenever  $y \leq \sup S$  for some  $S \in Z(P)$ , there exists an  $s \in S$  such that  $x \leq s$ . A poset is called  $Z$ -continuous if (i) it is  $Z$ -complete (meaning: for every  $S \in Z(P)$ ,  $\sup S$  exists), (ii) for every  $x \in P$ , the set  $\downarrow x = \{y : y \ll x\} \in I_Z(P)$ , and for every  $x \in P$ ,  $x = \sup \downarrow x$ . A  $Z$ -continuous poset is called *strongly  $Z$ -continuous* if the waybelow relation has the interpolation property; that is,  $x \ll y$  implies that there exists a  $z \in P$  such that  $x \ll z \ll y$ . If the subset system is union-complete, then any  $Z$ -continuous

poset is strongly  $Z$ -continuous ([V1]). The following table shows the most well known examples of  $Z$ -continuous posets. See [V2] for more examples.

<i>Subset system <math>Z</math></i>	<i><math>Z</math>-continuous poset</i>
All subsets	Completely distributive lattices [R]
Directed subsets	Continuous posets [COMP]
Finite subsets	Unique factoring posets [M]

### 1. Topology

**Definition 1.1.** For a poset  $P$ , let  $\sigma_Z(P)$  denote the set of all subsets  $V$  of  $P$  satisfying the following conditions: (i)  $V = \uparrow V$ , and (ii) whenever  $\sup S$  is in  $V$  for some  $S \in Z(P)$ , then there exists  $s \in S$  such that  $s \in V$ . Let  $\omega_Z(P) = \{P \setminus \uparrow x : x \in P\}$ . Let  $\lambda(P)$  denote the topology on  $P$  generated by  $\omega_Z(P) \cup \sigma_Z(P)$  as subbasic open sets.

If  $Z$  is the subset system of all subsets, this topology is the same as the interval topology, and if  $Z$  is the subset system of all directed subsets, then this topology is the same as the Lawson topology ([COMP]).

**Proposition 1.2.** If  $P$  is a strongly  $Z$ -continuous poset, then  $\lambda_Z(P)$  is a  $T_3$  topology.

PROOF: Since  $P \setminus \downarrow x \in \sigma_Z(P)$ ,  $\downarrow x$  is a closed set, and since  $P \setminus \uparrow x \in \omega_Z(P)$ ,  $\uparrow x$  is a closed set. Therefore  $\{x\} = \uparrow x \cap \downarrow x$  is closed, and hence  $\lambda_Z(P)$  is a  $T_1$  topology. Now we shall show that  $\lambda_Z(P)$  is regular. It is sufficient if we show that for each  $y \in P$ , and a subbasic open set  $U$  containing  $y$ , there exists an open set  $V$  such that  $y \in V$ , and the closure of  $V$  is contained in  $U$ .

Let  $y \in V$  where  $V \in \sigma_Z(P)$ . Since  $y = \sup \downarrow y$  and  $\downarrow y$  is a  $Z$ -ideal, there exists  $x \ll y$  such that  $x \in V$ . Therefore  $y \in \uparrow x \subseteq Cl(\uparrow x) \subseteq \uparrow x \subseteq V$ . Now we shall show that  $\uparrow x$  is an open set. Let  $\sup S \in \uparrow x$  for some  $Z$ -set  $S$  of  $P$ . By the interpolation property, there exists a  $z \in P$  such that  $x \ll z \ll \sup S$ . Then there exists  $s \in S$  such that  $x \ll z \leq s$ . This proves that  $\uparrow x$  is open.

Now let  $y \in P \setminus \uparrow x$ . Then  $x \not\leq y$ , and therefore there exists  $u \ll x$  such that  $u \not\leq y$ . By the interpolation property, there exists  $z$  such that  $u \ll z \ll x$ . Therefore  $y \in P \setminus \uparrow u \subseteq Cl(P \setminus \uparrow u) \subseteq P \setminus \uparrow z \subseteq P \setminus \uparrow x$ . This completes the proof of the proposition. □

For the remaining of this note, we assume the topology on a  $Z$ -continuous  $P$  poset is the  $\Lambda(P)$  topology. A function between two  $Z$ -continuous posets is called a *homomorphism* if it preserves the sups of  $Z$ -sets and is an upper adjoint. See [BE] and [V1].

**Proposition 1.3.** Let  $P, Q$  be  $Z$ -continuous posets. If  $f : P \rightarrow Q$  is a homomorphism, then  $f$  is continuous.

PROOF: Since  $f$  is an upper adjoint  $\inf f^{-1}(\uparrow t)$  exists for all  $t \in Q$ . Let  $s = \inf f^{-1}(\uparrow t)$ . Then, since upper adjoints preserves infs,  $f(s) = f(\inf f^{-1}(\uparrow t)) = \inf f f^{-1}(\uparrow t) = \inf \uparrow t = t$ . Thus  $s \in f^{-1}(\uparrow t)$  and hence  $f^{-1}(\uparrow t) = \uparrow s$ . Therefore  $f^{-1}(\uparrow t)$  is closed. Now let  $V \in \sigma_Z(Q)$ . We shall show that  $f^{-1}(V) \in \sigma_Z(P)$ . Since  $f$  is a monotone map,  $f^{-1}(V)$  is an upper set. Let  $S$  be a  $Z$ -set in  $P$  such that  $\sup S \in f^{-1}(V)$ . Then  $f(\sup S) \in V$  and, since  $f$  is  $Z$ -continuous,  $\sup f(S) \in V$ . Since  $f(S)$  is a  $Z$ -set in  $Q$  and since  $V \in \sigma_Z(Q)$ , there exists  $x \in S$  such that  $f(x) \in V$ ; that is,  $x \in f^{-1}(V)$ . Thus  $f^{-1}(V) \in \sigma_Z(P)$ . This completes the proof that  $f$  is continuous.  $\square$

The following lemma was proved in [BE].

**Lemma 1.4.** *Let  $P, Q$  be  $Z$ -continuous posets, and let  $(g, d)$  be a Galois connection from  $P$  to  $Q$ . If  $g$  is  $Z$ -continuous, then  $d$  preserves the waybelow relation.*

A subposet of a  $Z$ -continuous poset is called a subalgebra if the inclusion map is an upper adjoint which preserves the sups of  $Z$ -sets. It was shown in [V1] that a subalgebra of a  $Z$ -continuous poset is  $Z$ -continuous.

**Proposition 1.5.** *Every subalgebra of a strongly  $Z$ -continuous poset  $P$  is a closed subspace of  $P$ .*

PROOF: Let  $j$  be the lower adjoint of the inclusion map  $i : S \rightarrow P$ . Let  $x \in P \setminus S$ . We want to find an open set containing  $x$  and contained in  $P \setminus S$ . Note that  $ij(x) \geq x$  which implies that  $j(x) > x$ . Then there exists  $y \in P$  such that  $y \not\leq x$  and  $y \ll j(x)$ . Therefore  $x \in P \setminus \uparrow y = V_1$ . Since  $j$  preserves sups,  $y \ll_P j(x) = j(\sup_P \downarrow x) = \sup_S j(\downarrow x)$ . Then by the above lemma,  $j(y) \leq_S j(x) = \sup_S j(\downarrow x)$  and hence there exists  $z \ll x$  such that  $j(y) \leq j(z)$ . Therefore  $x \in \uparrow z = V_2$ . Let  $V = V_1 \cap V_2$ . We claim  $S \cap V = \emptyset$ . Indeed, if  $r \in S \cap V$ , then  $y \not\leq r$  and  $z \ll r$ . Then  $y \leq j(y) \leq j(z) \leq j(r) = r$ . This contradiction proves the claim. This completes the proof of the proposition.  $\square$

A subposet  $B$  of a  $Z$ -continuous poset  $P$  is called a basis if, for all  $x \in P$ ,

- (i)  $\downarrow x \cap B \in I_Z(P)$  and (ii)  $x = \sup \downarrow x \cap B$  ([V1]).

**Proposition 1.6.** *If  $P$  is a  $Z$ -continuous poset with a countable basis, then  $P$  is metrizable.*

PROOF: Let  $B$  be a countable basis of  $P$ . We shall show that  $\{P \setminus \uparrow b : b \in B\} \cup \{\uparrow b : b \in B\}$  is a subbasis of the topology. Let  $V \in \sigma_Z(P)$  and let  $x \in V$ . Since  $\sup(\downarrow x \cap B) = x$  and  $\downarrow x \cap B \in I_Z(P)$ , there exists  $y \in V$  such that  $y \in \downarrow x \cap B$ . Then  $x \in \uparrow y \subseteq V$ . Now let  $P \setminus \uparrow x \in \omega_Z(P)$ . Then  $P \setminus \uparrow x = P \setminus \uparrow \sup(\downarrow x \cap B) = P \setminus (\bigcap_{b \in \downarrow x \cap B} \uparrow b) = \bigcup_{b \in \downarrow x \cap B} P \setminus \uparrow b$ . This proves the claim, and the proposition follows from Urysohn's Metrization Theorem.  $\square$

## REFERENCES

- [BE] Bandelt H.J., Erné M., *The category of  $Z$ -continuous posets*, J. Pure Appl. Algebra **30** (1983), 219–226.
- [COMP] Gierz G., Hofmann K.H., Keimel K., Lawson J.D., Mislove M., Scott D.S., *A Compendium of Continuous Lattices*, Springer-Verlag, Berlin, Heidelberg, and New York, 1980.
- [M] Martinez J., *Unique factorization in partially ordered sets*, Proc. Amer. Math. Soc. **33** (1972), 213–220.
- [N] Novak D., *Generalization of continuous posets*, Trans. Amer. Math. Soc. **272** (1982), 645–667.
- [R] Raney G., *A subdirect-union representation for completely distributive lattices*, Proc. Amer. Math. Soc. **4** (1953), 518–522.
- [V1] Venugopalan P.,  *$Z$ -continuous posets*, Houston J. Math. **12** (1986), 275–294.
- [V2] Venugopalan P., *Union complete subset system*, Houston J. Math. **14** (1988), 583–600.
- [WWT] Wright J.B., Wagner E.G., Thatcher J.W., *A uniform approach to inductive posets and inductive closure*, Theor. Computer Science **7** (1978), 57–77.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STAMFORD, CONNECTICUT  
06903, USA

(Received January 19, 1996)