

W. Feng; Jeffrey Ronald Leslie Webb

Surjectivity results for nonlinear mappings without oddness conditions

Commentationes Mathematicae Universitatis Carolinae, Vol. 38 (1997), No. 1, 15--28

Persistent URL: <http://dml.cz/dmlcz/118899>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1997

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Surjectivity results for nonlinear mappings without oddness conditions

W. FENG, J.R.L. WEBB

Abstract. Surjectivity results of Fredholm alternative type are obtained for nonlinear operator equations of the form $\lambda T(x) - S(x) = f$, where T is invertible, and T, S satisfy various types of homogeneity conditions. We are able to answer some questions left open by Fučík, Nečas, Souček, and Souček. We employ the concept of an a -stably-solvable operator, related to nonlinear spectral theory methodology. Applications are given to a nonlinear Sturm-Liouville problem and a three point boundary value problem recently studied by Gupta, Ntouyas and Tsamatos.

Keywords: (K, L, a) homeomorphism, a -homogeneous operator, a -stably solvable map

Classification: 47H15, 47H12, 34B10

1. Introduction

The authors of [1] studied various surjectivity results for nonlinear operator equations of the form

$$(1.1) \quad \lambda T(x) - S(x) = f$$

when T is invertible. They considered various types of homogeneity conditions, in particular a key assumption was that T was a so-called (K, L, a) -homeomorphism. The precise definition of this and other concepts mentioned in the introduction is given a little later.

In their paper [1], the authors gave theorems of Fredholm alternative type under the assumptions that T is an odd (K, L, a) -homeomorphism and $S : X \rightarrow Y$ is an odd compact (completely continuous) operator. Furthermore, they established existence of a solution of the equation (1.1) for each $f \in Y$ provided $\lambda \neq 0$ if T is an odd a -homogeneous map and S is an odd b -strongly quasihomogeneous map with $a > b$. In the case $a < b$ they proved the same assertion in finite dimensional spaces but the infinite-dimensional case was an unsolved problem.

In this paper, we employ different methods which allow us to answer some of their open questions. By introducing the concept of an a -stably-solvable operator, we obtain some surjectivity results for $\lambda T - S$ under weaker conditions. One of the theorems generalizes the result of existence of a solution of (1.1) in case $a < b$ to the infinite-dimensional case. These results seem not to be able to be proven by their methods.

It is possible to give simple examples that show that our results are real extensions of the earlier ones, but we prefer to give more substantial applications. We discuss a nonlinear Sturm-Liouville problem on the half line following work by Toland [4]. He studied eigenvalues and asymptotic bifurcation points whereas we obtain surjectivity when λ is not one of these eigenvalues.

We also discuss existence of solutions to a three point boundary value problem recently studied by Gupta, Ntouyas and Tsamatos, in [6], [7], [8]. The boundary conditions are of the type $x(0) = 0$, $x(1) = \alpha x(\eta)$. Those authors assume that $\alpha < 1/\eta$ but we suppose only that $\alpha \neq 1/\eta$. We obtain a different criterion for existence which improves on Theorem 4 of [6] in some cases but is less good in others.

2. Prerequisites

We will make use of the class of k -set contractive maps and of the theory of degree for $I - f$ where f is k -set contractive, see for example [5]. We give some notations and definitions that we shall use.

Given any continuous map f from a subset $D(f)$ of a complex Banach space X into a Banach space Y written $f : D(f) \subseteq X \rightarrow Y$, let $\alpha(\Omega)$ denote the measure of noncompactness of the bounded set Ω (see for example [5]), and let

$$\begin{aligned} \alpha(f) &= \inf\{k \geq 0 : \alpha(f(\Omega)) \leq k\alpha(\Omega) \text{ for every bounded } \Omega \subset D(f)\}, \\ \omega(f) &= \sup\{k \geq 0 : \alpha(f(\Omega)) \geq k\alpha(\Omega) \text{ for every bounded } \Omega \subset D(f)\}, \\ d(f) &= \liminf_{\|x\| \rightarrow \infty, x \in D(f)} \frac{\|f(x)\|}{\|x\|}, \quad |f| = \limsup_{\|x\| \rightarrow \infty, x \in D(f)} \frac{\|f(x)\|}{\|x\|}. \end{aligned}$$

Here $|f|$ is called the quasinorm of f and f is said to be quasibounded if $|f| < \infty$. Maps with $\alpha(f) < 1$ are k -set contractive (also condensing) with $k = \alpha(f)$. Note that a map f satisfies $\alpha(f) = 0$ if and only if f is compact, that is, $\overline{f(\Omega)}$ is compact for every bounded set Ω .

We shall also use of some notions employed by Furi, Martelli and Vignoli [2] in their theory of spectrum of nonlinear operators. We recall some of these.

Definition 2.1. Let $f : X \rightarrow Y$ be a continuous map from a Banach space X into a Banach space Y . The map f is said to be *stably-solvable* if the equation

$$f(x) = h(x)$$

has a solution $x \in X$ for any continuous compact map $h : X \rightarrow Y$ with quasinorm $|h| = 0$.

f is said to be *regular* if it is stably-solvable and $d(f)$ and $\omega(f)$ are both positive. When $Y = X$, the resolvent set of f is the set

$$\rho(f) = \{\lambda \in \mathbb{C}, \lambda I - f \text{ is regular}\}$$

and the spectrum of f is $\sigma(f) = \mathbb{C} \setminus \varrho(f)$.

If f is invertible then $\alpha(f^{-1}) = 1/\omega(f)$, so regular invertible maps have k -set contractive inverses.

We will consider a generalization of the concept of stably-solvable maps below.

3. Surjectivity theorems

We begin with a result which generalizes Theorem 1.2 and Corollary 1.1 in Chapter II of [1]. Those authors studied operators T that are (K, L, a) -homeomorphisms, where a (not necessarily linear) map $T : X \rightarrow Y$ is said to be a (K, L, a) -homeomorphism if

- (a) T is a homeomorphism of X onto Y , and
- (b) there exists real numbers $K > 0, a > 0, L > 0$ such that

$$L\|x\|^a \leq \|T(x)\| \leq K\|x\|^a \quad \text{for each } x \in X.$$

We do not assume so much.

Theorem 3.1. *Let $T : D(T) \subseteq X \rightarrow Y$ be an operator satisfying the following conditions:*

1. T is one to one, onto and $T^{-1} : Y \rightarrow D(T)$ is continuous;
2. there exist real numbers $L > 0, a > 0$ and $b > 0$ such that

$$\|T(x)\| \geq L\|x\|^a - b \quad \text{for every } x \in D(T);$$

3. T is bounded, that is, maps bounded sets into bounded sets.

Let $S : X \rightarrow Y$ be bounded, continuous and suppose that

$$\limsup_{x \in D(T), \|x\| \rightarrow \infty} \frac{\|S(x)\|}{\|x\|^a} = A.$$

Then $\lambda T - S$ maps $D(T)$ onto Y under any one of the following conditions:

1. $|\lambda| > \max\{\frac{A}{L}, \frac{\alpha(S)}{\omega(T)}\}$;
2. S is compact, and $|\lambda| > \frac{A}{L}$;
3. Y is a finite dimensional space, and $|\lambda| > \frac{A}{L}$;
4. S is compact, $A = 0$, and $\lambda \neq 0$.

PROOF: Clearly it suffices to prove case 1. Also it is clear that $\lambda T - S$ maps $D(T)$ onto Y if $I - F$ maps Y onto Y where $F : Y \rightarrow Y$ is defined by $F(y) = ST^{-1}(y/\lambda)$.

For any bounded set $\Omega \in Y$, we have

$$\begin{aligned} \alpha(F(\Omega)) &= \alpha(ST^{-1}(\Omega/\lambda)) \\ &\leq \alpha(ST^{-1})\alpha(\Omega/\lambda) \\ &\leq \frac{1}{|\lambda|} \frac{\alpha(S)}{\omega(T)} \alpha(\Omega). \end{aligned}$$

Therefore,

$$\alpha(F) \leq \frac{1}{|\lambda|} \frac{\alpha(S)}{\omega(T)} < 1.$$

[If S is compact or Y is finite dimensional, then $\alpha(F) = 0$.]

Also we have,

$$\begin{aligned} |F| &= \limsup_{\|y\| \rightarrow \infty} \|F(y)\| / \|y\| \\ &= \limsup_{\|y\| \rightarrow \infty} \|ST^{-1}(y/\lambda)\| / \|y\|. \end{aligned}$$

Writing $x = T^{-1}(y/\lambda)$, we have $Tx = y/\lambda$, and we obtain

$$\begin{aligned} |F| &= \limsup_{\|Tx\| \rightarrow \infty} \frac{\|S(x)\|}{|\lambda| \|Tx\|} \\ &= \limsup_{x \in D(T), \|x\| \rightarrow \infty} \frac{\|S(x)\|}{|\lambda| \|Tx\|} \\ &\leq \limsup_{x \in D(T), \|x\| \rightarrow \infty} \frac{\|S(x)\|}{|\lambda| (L\|x\|^a - b)} \\ &= \frac{A}{|\lambda|L} < 1. \end{aligned}$$

Hence, by the results of [2], $1 \in \varrho(F)$, in particular $I - F$ maps Y onto Y . \square

Remark 3.2. A result similar to Theorem 3.1 was obtained in [3], where a different method was used.

Remark 3.3. Theorem 1.2 of [1] requires that T is a (K, L, a) -homeomorphism and that T, S are both odd with S compact, but allow either $|\lambda| > \frac{A}{L}$ or $|\lambda| < \frac{A}{K}$, $\lambda \neq 0$.

The following simple example shows that for $0 \neq \lambda \in \mathbb{C}$, even when T is a (K, L, a) -homeomorphism there is no result similar to Theorem 3.1 in the case $|\lambda| < \frac{A}{K}$ without some extra hypothesis (such as oddness of the maps).

Example 3.4. Let T and $S : \mathbb{C} \rightarrow \mathbb{C}$ be defined by

$$T(z) = z, \quad S(x + iy) = |x| + iy,$$

and let $\lambda = 1/2$. Then $a = 1, K = L = 1, T$ is odd, S is not odd. Also $A = 1, S, T$ are compact maps, but $\lambda T - S$ is not onto since $z/2 - S(z) = 1$ has no solution.

We recall the following concepts from [1].

Definition 3.5. Suppose that $a > 0$.

(a) A map $F_0 : X \rightarrow Y$ is called *a-homogeneous* if $F_0(tu) = t^a F_0(u)$ for every $t \geq 0$ and $u \in X$.

(b) $F : X \rightarrow Y$ is said to be *a-quasihomogeneous* relative to F_0 if $F_0 : X \rightarrow Y$ is *a-homogeneous* and

$$t_n \searrow 0, u_n \rightharpoonup u_0, t_n^a F(u_n/t_n) \rightarrow g \in Y$$

together imply that $g = F_0(u_0)$. [Here $u_n \rightharpoonup u_0$ denotes weak convergence.]

(c) $F : X \rightarrow Y$ is said to be *a-strongly quasihomogeneous* relative to F_0 if

$$t_n \searrow 0, u_n \rightharpoonup u_0 \text{ imply that } t_n^a F(u_n/t_n) \rightarrow F_0(u_0) \in Y.$$

It is known ([1]) that in case (c) F_0 is *a-homogeneous* and also must be strongly continuous, that is $u_n \rightharpoonup u_0$ implies $F_0 u_n \rightarrow F_0 u_0$.

By applying Theorem 3.1 instead of Corollary 1.1 of [1], we obtain the following generalization of Theorem 4.1 of [1], where we can dispense with the assumption that T, S are odd maps.

Theorem 3.6. *Let X be reflexive and let T satisfy the conditions of Theorem 3.1. Let $S : X \rightarrow Y$ be a compact b -strongly quasihomogeneous operator relative to S_0 and suppose that $a > b$. Then for $\lambda \neq 0$, $\lambda T - S$ maps $D(T)$ onto Y .*

PROOF: By Theorem 3.1, part 4, it suffices to show that

$$\lim_{\|x\| \rightarrow \infty, x \in D(T)} \frac{\|S(x)\|}{\|x\|^a} = 0.$$

This was proved in Theorem 4.1 of [1] but we include the proof for completeness. If this is false, there is a sequence $\{x_n\}$ with $\|x_n\| \rightarrow \infty$ and $\varepsilon > 0$ such that $\|Sx_n\|/\|x_n\|^a \geq \varepsilon$, for all sufficiently large n . Letting $u_n = x_n/\|x_n\|$ and $t_n = 1/\|x_n\|$ we have, for a subsequence, that

$$S(x_n)/\|x_n\|^b \rightarrow S_0(u_0).$$

Since $a > b$ this gives $S(x_n)/\|x_n\|^a \rightarrow 0$, a contradiction. \square

Remark 3.7. The authors of [1] say that the case $a < b$ seems to be unsolved in the infinite dimensional case. We shall give an answer below, see Theorem 3.12.

We introduce the following extension of the concept of stably solvable maps which is appropriate to our needs.

Definition 3.8. A continuous map $f : D(f) \subseteq X \rightarrow Y$ is said to be *a-stably-solvable* for some $a > 0$ if the equation

$$f(x) = h(x)$$

has a solution $x \in D(f)$ for any continuous compact map $h : X \rightarrow Y$ with

$$|h|_a := \limsup_{\|x\| \rightarrow \infty} \frac{\|h(x)\|}{\|x\|^a} = 0.$$

Lemma 3.9. Suppose $T : D(T) \subseteq X \rightarrow Y$ is as in Theorem 3.1. Then T is *a-stably-solvable*.

PROOF: Let $h : X \rightarrow Y$ be a compact map with $|h|_a = 0$. Then $\alpha(T^{-1}h) = 0$, and

$$\begin{aligned} \limsup_{\|x\| \rightarrow \infty} \frac{\|T^{-1}h(x)\|}{\|x\|} &= \limsup_{\|x\| \rightarrow \infty} \frac{\|T^{-1}h(x)\|}{\|h(x)\|^{\frac{1}{a}}} \left(\frac{\|h(x)\|}{\|x\|^a} \right)^{\frac{1}{a}} \\ &\leq \limsup_{\|x\| \rightarrow \infty} \left(\frac{1}{L} \right)^{\frac{1}{a}} \left(\frac{\|h(x)\|}{\|x\|^a} \right)^{\frac{1}{a}} \rightarrow 0. \end{aligned}$$

Therefore, $|T^{-1}h| = 0$. This implies that $1 \in \varrho(T^{-1}h)$, so that $I - T^{-1}h$ is onto, that is, there exists $x \in D(T)$ such that $x = T^{-1}h(x)$, that is, $Tx = hx$. \square

Lemma 3.10 (The Continuation Principle for *a-stably-solvable* maps).

Let $f : D(f) \subseteq X \rightarrow Y$ be *a-stably-solvable*, $h : X \times [0, 1] \rightarrow Y$ be continuous, compact and such that $h(x, 0) = 0$ for all $x \in D(f)$. Let

$$U = \{x \in D(f), f(x) = h(x, t) \text{ for some } t \in [0, 1]\}.$$

Then, if $f(U)$ is bounded, the equation

$$f(x) = h(x, 1)$$

has a solution.

PROOF: Let $B_r = \{y \in Y, \|y\| < r\}$, and let $r > 0$ be chosen so that $\overline{f(U)} \subset B_r$. Let $\varphi : X \rightarrow [0, 1]$ be continuous and such that

$$\varphi(y) = \begin{cases} 1, & \text{for } y \in \overline{f(U)}, \\ 0, & \text{for } \|y\| \geq r, \end{cases}$$

and let π be the radial retraction of Y onto $\overline{B_r}$. Then the equation

$$f(x) = \pi h(x, \varphi(f(x)))$$

has a solution $x_0 \in D(f)$ since πh is compact and

$$|\pi h|_a = \lim_{\|x\| \rightarrow \infty} \frac{\|(\pi h)(x)\|}{\|x\|^a} = 0.$$

If $\|f(x_0)\| = r$, then $\varphi(f(x_0)) = 0$, and $f(x_0) = \pi h(x_0, 0) = 0$, a contradiction. Thus $\|f(x_0)\| < r$, and $f(x_0) = h(x_0, \varphi(f(x_0)))$, which shows that $x_0 \in U$ and therefore $f(x_0) = h(x_0, 1)$. \square

Theorem 3.1 of [1] gave theorems of Fredholm alternative type for the couple (T, S) when T, S were both odd. Recall that λ is said to be an eigenvalue for the couple T_0, S_0 if there is $x_0 \neq 0$ such that $\lambda T_0 x_0 - S_0 x_0 = 0$. Using Lemmas 3.9 and 3.10 we can give the following result when neither T nor S is odd.

Theorem 3.11. *Let X be a reflexive Banach space, and let T be as in Theorem 3.1 with $D(T) = X$ and also a -quasihomogeneous relative to T_0 . Let $S : X \rightarrow Y$ be a compact a -strongly-quasihomogeneous operator relative to S_0 . If $\lambda \neq 0$, and for every $t \in (0, 1]$, λ/t is not an eigenvalue for the couple (T_0, S_0) , then $\lambda T - S$ maps X onto Y .*

PROOF: For arbitrary $y \in Y$, let

$$U = \{x \in X, \lambda T(x) = h(x, t) = t[S(x) + y], t \in [0, 1]\}.$$

We show that U is bounded. If not, there exists $x_n \in U$, $\|x_n\| \rightarrow \infty$, such that

$$\lambda T(x_n) = t_n[S(x_n) + y], \quad t_n \in [0, 1],$$

so that

$$\begin{aligned} \frac{\lambda T(x_n)}{\|x_n\|^a} &= t_n \left(\frac{S(x_n)}{\|x_n\|^a} + \frac{y}{\|x_n\|^a} \right) \\ &= t_n \frac{1}{\|x_n\|^a} S \left(\frac{x_n/\|x_n\|}{1/\|x_n\|} \right) + t_n \frac{y}{\|x_n\|^a}. \end{aligned}$$

Without loss of generality we assume that $x_n/\|x_n\| \rightarrow x_0$, $t_n \rightarrow t_0 \in [0, 1]$. Then there exists a subsequence $\{x_{n_k}\}$ such that

$$\begin{aligned} t_{n_k} \frac{1}{\|x_{n_k}\|^a} S \left(\frac{x_{n_k}/\|x_{n_k}\|}{1/\|x_{n_k}\|} \right) &\rightarrow t_0 S_0(x_0), \\ \lim_{n \rightarrow \infty} \frac{\lambda T(x_{n_k})}{\|x_{n_k}\|^a} &= t_0 S_0(x_0). \end{aligned}$$

Since T is a -quasihomogeneous relative to T_0 , we obtain

$$\lambda T_0(x_0) = t_0 S_0(x_0).$$

However,

$$\frac{\|\lambda T(x_{n_k})\|}{\|x_{n_k}\|^a} \geq |\lambda|L - \frac{|\lambda|b}{\|x_{n_k}\|^a} > |\lambda|L/2,$$

for n_k sufficiently large so that $\|t_0 S_0(x_0)\| > 0$. Hence $t_0 \neq 0$, and $S_0(x_0) \neq 0$.

From the definition of a -strongly-quasihomogeneous operator it is easy to show that $S_0(0) = 0$. Thus $x_0 \neq 0$, and λ/t_0 is an eigenvalue of (T_0, S_0) , a contradiction. Thus U is bounded. By Lemma 3.9, $\lambda T : X \rightarrow Y$ is a -stably-solvable. So by Lemma 3.10, the equation $\lambda T(x) = S(x) + y$ has a solution $x \in X$, that is $\lambda T - S$ is onto. \square

The next two results extend Theorem 4.2 of [1] to the infinite dimensional case.

Theorem 3.12. *Let X be a reflexive Banach space. Let T be a bounded, odd mapping satisfying the following conditions:*

1. $T : D(T) \subseteq X \rightarrow Y$ is one to one, onto and $T^{-1} : Y \rightarrow D(T)$ is continuous;
2. there exist real numbers $K > 0, a > 0$ and q such that

$$\|T(x)\| \leq K\|x\|^a + q \text{ for every } x \in D(T).$$

Suppose that S is odd, continuous and b -strongly quasihomogeneous relative to S_0 , and that $\inf_{\{\|x\|=1\}} \|S_0(x)\| > 0$. If $a < b$, then for every λ with $|\lambda| > \alpha(S)/\omega(T)$, $\lambda T - S$ is a -stably-solvable.

PROOF: First we show that there exists $R > 0$ such that $\lambda x - T^{-1}Sx \neq 0$ whenever $\|x\| \geq R$. If there exists $\{x_n\} \subset X$, $\|x_n\| \rightarrow \infty$ such that

$$\lambda x_n - T^{-1}S(x_n) = 0$$

we may assume that $\frac{x_n}{\|x_n\|} \rightarrow x_0$. Then we have

$$\frac{\|S(x_n)\|}{\|x_n\|^b} = \frac{\lambda T(x_n)}{\|x_n\|^b} \leq \frac{|\lambda|K\|x_n\|^a + q}{\|x_n\|^b} \rightarrow 0.$$

Since S is b -strongly quasihomogeneous relative to S_0 , we have

$$\frac{1}{\|x_n\|^b} S(x_n) = \frac{1}{\|x_n\|^b} S\left(\frac{x_n/\|x_n\|}{1/\|x_n\|}\right) \rightarrow S_0(x_0).$$

As S_0 is strongly continuous we also have $S_0\left(\frac{x_n}{\|x_n\|}\right) \rightarrow S_0(x_0)$. Since $\inf_{\|x\|=1} \|S_0(x)\| > 0$ it follows that $S_0(x_0) \neq 0$, this contradicts the above. Let $B_r(0) = \{x \in X, \|x\| < r\}$, where $r > R$. Then $\alpha(T^{-1}S) < |\lambda|$ and the topological degree $d(I - T^{-1}S/\lambda, B_r(0), 0)$ is odd, hence nonzero (see, for example, [5]). For a compact operator $h : X \rightarrow Y$ with $h = 0$ for $\|x\| = r$,

$$d\left(I - T^{-1}S/\lambda - T^{-1}h/\lambda, B_r(0), 0\right) \neq 0$$

because of boundary value dependence of degree.

For each $n \in \mathbb{N}$ let σ_n be continuous and such that

$$\sigma_n(x) = \begin{cases} 1 & \text{for } \|x\| \leq n, \\ 0 & \text{for } \|x\| \geq 2n. \end{cases}$$

Then, if $h : X \rightarrow Y$ is a compact operator, with $|h|_a = 0$, for every $n > R/2$, the equation

$$\lambda T(x) - S(x) = \sigma_n(x)h(x)$$

has a solution $x_n \in D(T)$. If for all n , we have $\|x_n\| > n$, then

$$\frac{\lambda T(x_n) - S(x_n)}{\|x_n\|^b} = \frac{\sigma_n(x_n)h(x_n)}{\|x_n\|^b}.$$

Assume that $x_n/\|x_n\| \rightarrow x_0$. Then from

$$\frac{\lambda T(x_n) - S(x_n)}{\|x_n\|^b} \rightarrow -S_0(x_0) \neq 0 \quad (n \rightarrow \infty),$$

and

$$\frac{\sigma_n(x_n)h(x_n)}{\|x_n\|^b} = \sigma_n(x_n) \frac{h(x_n)}{\|x_n\|^a} \frac{\|x_n\|^a}{\|x_n\|^b} \rightarrow 0 \quad (n \rightarrow \infty),$$

we reach a contradiction. Hence there exists n , such that $\|x_n\| \leq n$, and then

$$\lambda T(x_n) - S(x_n) = h(x_n),$$

and we are done. □

Theorem 3.13. *Let X be a reflexive Banach space, $T, T_1 : D(T) \rightarrow Y$ and $S, S_1 : X \rightarrow Y$ be of the form $T = T_1 + R$, $S = S_1 + R'$, where T_1 satisfies the same conditions as T in Theorem 3.12, S_1 is odd, continuous and b -strongly quasihomogeneous relative to S_0 , and $R, R' : X \rightarrow Y$ are compact operators with $|R|_a = |R'|_a = 0$. Suppose that $a < b$, and that $\inf_{\{\|x\|=1\}} \|S_0(x)\| > 0$. Then $\lambda T - S$ maps $D(T)$ onto Y for every λ with $|\lambda| > \alpha(S)/\omega(T)$.*

PROOF: For $y \in Y$, let $h(x) = -\lambda R(x) + R'(x) + y$, so that h is compact and $|h|_a = 0$. By Theorem 3.12, the equation

$$\lambda T_1(x) - S_1(x) = h(x)$$

has a solution $x_0 \in D(T)$. Hence

$$\lambda T(x_0) - S(x_0) = y,$$

that is $\lambda T - S$ is onto. □

4. Applications

The following applications are examples of situations that can be settled by the above theorems but apparently cannot be handled by the results in [1].

Example 4.1. We consider a nonlinear Sturm-Liouville problem on an unbounded domain, namely the following nonlinear differential equation:

$$(4.1) \quad \begin{aligned} -(p(x)u'(x))' + q(x)u(x) &= \lambda\{u(x) + g(x)f(u(x))\}, \\ \text{for } x \in (0, \infty), \text{ and } u(0) &= 0. \end{aligned}$$

In [4] it was shown that certain eigenvalues λ are asymptotic bifurcation points. Under the same assumptions we will show that if v is continuous, the equation

$$(4.2) \quad \begin{aligned} -(p(x)u'(x))' + q(x)u(x) &= \lambda\{u(x) + g(x)f(u(x))\} + v(x) \\ \text{for } x \in (0, \infty), \text{ and } u(0) &= 0 \end{aligned}$$

has a solution when λ is not one of these eigenvalues.

We recall the assumptions made in [4].

1. $p : [0, \infty) \rightarrow \mathbb{R}$ is continuous and continuously differentiable on $(0, \infty)$, with p' bounded and $0 < P_1 \leq p(x) \leq P_2 < \infty$ for all $x \in [0, \infty)$.
2. $q : [0, \infty) \rightarrow \mathbb{R}$ is continuous with $0 < Q_1 \leq q(x) \leq Q_2 < \infty$ for all $x \in [0, \infty)$.
3. f is a continuously differentiable function from \mathbb{R} into itself, and there exist positive real numbers P and K such that $|f(p)| \leq K|p|^r$ for all $p \geq P$, for some $r < 1$.
4. $g \in H_0^1(0, \infty)$.

For $u : [0, \infty) \rightarrow \mathbb{R}$ and $x \in [0, \infty)$ let H be defined by $(Hu)(x) = g(x)f(u(x))$. Let $A : H_0^1 \cap W^{2,2} \rightarrow L^2$ be the self-adjoint extension of the operator A_0 defined by $A_0u = -(p(x)u'(x))' + q(x)u(x)$ with domain the set of twice continuously differentiable functions with compact support in $(0, \infty)$. Then, ([4]), A is a positive self-adjoint operator in L^2 and its positive square root $A^{\frac{1}{2}}$ is a linear homeomorphism of H_0^1 onto L^2 , where H_0^1 is the closure of C_0^∞ in $W^{1,2}$ and C_0^∞ is the linear space of all infinitely differentiable, real-valued functions with compact support in $(0, \infty)$.

We claim (and will show below) that for $0 < |\lambda| < Q := \liminf_{x \rightarrow \infty} q(x)$, and λ not an eigenvalue of A , the operator

$$u \mapsto u - \lambda A^{-1}u + \lambda A^{-1/2}HA^{-1/2}u$$

from $L^2 \rightarrow L^2$ is onto. Assuming this, it follows that the equation

$$Au = \lambda u + \lambda Hu + v$$

has a solution $u \in H_0^1 \cap W^{2,2}$ for any $v \in L^2$ ([4, Lemma 4.18]). Hence if v is continuous, using the same arguments as in Lemma 4.20 of [4] it follows that the equation (4.2) has a solution.

We now establish the claim made above. Let $\mu = 1/\lambda$, and let $T, S : L^2 \rightarrow L^2$ be defined by

$$Tu = \mu u - A^{-1}u, \quad Su = A^{-1/2}HA^{-1/2}u.$$

Suppose that $|\mu| > \alpha(A^{-1}) = 1/Q$ ([4, Theorem 4.23]), and that μ is not an eigenvalue of A^{-1} . Then T is a bounded linear operator, which is one to one, onto, and has a continuous inverse. So it is a $(K, L, 1)$ -homeomorphism of L^2 onto L^2 . Furthermore, T is 1-quasihomogeneous relative to T since it has continuous inverse. It has been shown that S is a compact operator and the quasinorm $|S| = 0$ in the space L^2 ([4, Lemma 4.17]). Assume that there exist $u_n \in L^2$ with $u_n \rightharpoonup u_0$, $t_n \searrow 0$ such that

$$t_n S(u_n/t_n) > \varepsilon_0 > 0.$$

Then $\{\|u_n/t_n\|_2\}$ is unbounded. If $\|u_{n_k}/t_{n_k}\|_2 \rightarrow \infty$, ($n_k \rightarrow \infty$), then we have

$$\|t_{n_k} S(u_{n_k}/t_{n_k})\|_2 = \frac{\|S(u_{n_k}/t_{n_k})\|_2}{\|u_{n_k}\|_2/t_{n_k}} \|u_{n_k}\|_2 \rightarrow 0,$$

a contradiction. Thus we have shown that S is a 1-strongly quasihomogeneous operator relative to $S_0 = 0$ in the space L^2 . For any $t \in (0, 1]$,

$$(1/t)(\mu I - A^{-1})(u) = 0 \implies u = 0,$$

so $1/t$ is not an eigenvalue of the couple $(T, 0)$. By Theorem 3.11, $T - S$ maps L^2 onto L^2 . Thus we have reached the conclusion.

The following second-order m -point nonlinear boundary value problem (BVP) has been studied recently by Gupta, Ntouyas and Tsamatos ([6], [7], [8]):

$$(4.3) \quad \begin{aligned} x''(t) &= f(t, x(t), x'(t)) + e(t) \quad 0 < t < 1, \\ x(0) &= 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i). \end{aligned}$$

It was shown that the problem of existence of a solution for the BVP (4.3) can be studied via the three boundary value problem

$$(4.4) \quad \begin{aligned} x''(t) &= f(t, x(t), x'(t)) + e(t) \quad 0 < t < 1, \\ x(0) &= 0, \quad x(1) = \alpha x(\eta), \end{aligned}$$

where $\eta \in (0, 1)$ and $\alpha \in \mathbb{R}$.

Some conditions for the existence of a solution for the BVP (4.4) were obtained in [6] using the Leray-Schauder continuation theorem. Their results suppose that $\alpha < 1/\eta$. By using Theorem 3.1, we obtain the following result which gives a different condition for the existence of a solution for (4.4) under the more general hypothesis $\alpha \neq 1/\eta$.

Theorem 4.2. *Let $f: [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function satisfying Carathéodory's conditions. Assume that there exist functions $p(t)$, $q(t)$, $r(t)$ in $L^1(0, 1)$ such that*

$$|f(t, x_1, x_2)| \leq p(t)|x_1| + q(t)|x_2| + r(t)$$

for a.e. $t \in [0, 1]$ and all $(x_1, x_2) \in \mathbb{R}^2$. Also let $\eta \in (0, 1)$, $\alpha \geq 0$, $\alpha \neq 1/\eta$ be given. Then for any given $e \in L^1(0, 1)$ the boundary value problem (4.4) has at least one solution in $C^1[0, 1]$ provided that

$$(4.5) \quad \|p\|_1 + \|q\|_1 < \begin{cases} (1 - \alpha\eta)/2, & \text{if } \alpha\eta < 1, \\ (\alpha\eta - 1)/2\alpha\eta, & \text{if } \alpha\eta > 1. \end{cases}$$

PROOF: Let X denote the Banach space $C^1[0, 1]$ with the norm

$$\|x\| = \max\{\|x\|_\infty, \|x'\|_\infty\}.$$

Let Y denote the Banach space $L^1(0, 1)$ with its usual norm.

The linear operator $L: D(L) \subset X \rightarrow Y$ is defined by setting

$$D(L) = \{x \in W^{2,1}(0, 1) : x(0) = 0, x(1) = \alpha x(\eta)\},$$

and for $x \in D(L)$,

$$Lx = x''.$$

For $x \in X$, let

$$(Nx)(t) = f(t, x(t), x'(t)), \quad t \in [0, 1].$$

Then N is a bounded map from X into Y . It can be shown that $L: D(L) \subset X \rightarrow Y$ is one to one and onto when $\alpha \neq 1/\eta$. In fact, $L^{-1} = K$, where K is the linear operator $K: Y \rightarrow D(L) \subset X$ defined by

$$(Ky)(t) = \int_0^t (t-s)y(s) ds + \frac{\alpha t}{1-\alpha\eta} \int_0^\eta (\eta-s)y(s) ds - \frac{t}{1-\alpha\eta} \int_0^1 (1-s)y(s) ds.$$

For $y \in Y$, we have

$$\|Ky\|_\infty \leq \left(1 + \frac{\alpha\eta + 1}{|1 - \alpha\eta|}\right) \|y\|_1,$$

where $\|y\|_1$ is the norm of y in the space $L^1(0, 1)$. Also

$$\|(Ky)'\|_\infty \leq \left(1 + \frac{\alpha\eta + 1}{|1 - \alpha\eta|}\right) \|y\|_1.$$

Thus we have

$$\|Ky\| \leq \left(1 + \frac{\alpha\eta + 1}{|1 - \alpha\eta|}\right) \|y\|_1.$$

Let $T = I$ and $S = KN$. Then $\alpha(S) = 0$ by the Arzela-Ascoli theorem. Also we have

$$\begin{aligned}
 A &= \limsup_{\|x\| \rightarrow \infty} \frac{\|S(x)\|}{\|x\|} \\
 &= \limsup_{\|x\| \rightarrow \infty} \frac{\|KN(x)\|}{\|x\|} \\
 &\leq \limsup_{x \rightarrow \infty} \left(1 + \frac{\alpha\eta + 1}{|1 - \alpha\eta|} \right) \frac{\|N(x)\|_1}{\|x\|} \\
 &\leq \left(1 + \frac{\alpha\eta + 1}{|1 - \alpha\eta|} \right) \limsup_{\|x\| \rightarrow \infty} \frac{\|p\|_1 \|x\|_\infty + \|q\|_1 \|x'\|_\infty + \|r\|_1}{\|x\|} \\
 &\leq \left(1 + \frac{\alpha\eta + 1}{|1 - \alpha\eta|} \right) \limsup_{\|x\| \rightarrow \infty} \frac{(\|p\|_1 + \|q\|_1) \|x\| + \|r\|_1}{\|x\|} \\
 &= \left(1 + \frac{\alpha\eta + 1}{|1 - \alpha\eta|} \right) (\|p\|_1 + \|q\|_1) \\
 &= \begin{cases} \frac{2}{1 - \alpha\eta} (\|p\|_1 + \|q\|_1) & \text{for } \alpha\eta < 1 \\ \frac{2\alpha\eta}{\alpha\eta - 1} (\|p\|_1 + \|q\|_1) & \text{for } \alpha\eta > 1. \end{cases}
 \end{aligned}$$

By the assumption (4.5) we see that $A < 1$. Hence, from Theorem 3.1, the operator $I - S = I - KN$ maps X onto X .

Hence, given any $e \in L^1(0, 1)$, there exists $x \in C^1[0, 1]$ such that

$$x(t) - (KNx)(t) = Ke(t).$$

Thus $x = KNx + Ke \in D(L)$ and

$$Lx - Nx = e.$$

This proves that the BVP (4.4) has at least one solution in $C^1[0, 1]$. \square

Remark 4.3. When $\alpha\eta < 1$, the condition (4.5) gives a better result than Theorem 4 of [6] in case $\alpha(1 - \eta) > 2$ since their condition demands $\|p\|_1 + \|q\|_1 < \frac{1 - \alpha\eta}{\alpha(1 - \eta)}$, but is worse in the case $\alpha(1 - \eta) < 2$. Also our result can apply when $\alpha\eta > 1$.

REFERENCES

- [1] Fučík S., Nečas J., Souček J., Souček V., *Spectral Analysis of Nonlinear Operators*, Lecture Notes in Mathematics **346**, Springer-Verlag, Berlin, Heidelberg, New York, 1973.
- [2] Furi M., Martelli M., Vignoli A., *Contributions to the spectral theory for nonlinear operators in Banach spaces*, Ann. Mat. Pura. Appl. (IV) **118** (1978), 229–294.
- [3] Webb J.R.L., *On degree theory for multivalued mappings and applications*, Boll. Un. Mat. It. (4) **9** (1974), 137–158.

- [4] Toland J.F., *Topological Methods for Nonlinear Eigenvalue Problems*, Battelle Advanced Studies Centre, Geneva, Mathematics Report No. 77, 1973.
- [5] Deimling K., *Nonlinear Functional Analysis*, Springer Verlag, Berlin, 1985.
- [6] Gupta C.P., Ntouyas S.K., Tsamatos P.Ch., *On an m -point boundary-value problem for second-order ordinary differential equations*, *Nonlinear Analysis, Theory, Methods & Applications* **23** (1994), 1427–1436.
- [7] Gupta C.P., Ntouyas S.K., Tsamatos P.Ch., *Solvability of an m -point boundary value problem for second order ordinary differential equations*, *J. Math. Anal. Appl.* **189** (1995), 575–584.
- [8] Gupta C.P., *A note on a second order three-point boundary value problem*, *J. Math. Anal. Appl.* **186** (1994), 277–281.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GLASGOW, GLASGOW G12 8QW, SCOTLAND, UNITED KINGDOM

(Received October 24, 1995)