

Jarmila Ranošová

Sets of determination for solutions of the Helmholtz equation

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 38 (1997), No. 2, 309--328

Persistent URL: <http://dml.cz/dmlcz/118929>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1997

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Sets of determination for solutions of the Helmholtz equation

JARMILA RANOŠOVÁ

*Abstract.* Let  $\alpha > 0$ ,  $\lambda = (2\alpha)^{-1/2}$ ,  $S^{n-1}$  be the  $(n - 1)$ -dimensional unit sphere,  $\sigma$  be the surface measure on  $S^{n-1}$  and  $h(x) = \int_{S^{n-1}} e^{\lambda\langle x,y \rangle} d\sigma(y)$ .

We characterize all subsets  $M$  of  $\mathbb{R}^n$  such that

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for every positive solution  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ . A closely related problem of representing functions of  $L_1(S^{n-1})$  as sums of blocks of the form  $e^{\lambda\langle x_k, \cdot \rangle} / h(x_k)$  corresponding to points of  $M$  is also considered. The results provide a counterpart to results for classical harmonic functions in a ball, and for parabolic functions on a slab, see References.

*Keywords:* Helmholtz equation, set of determination, decomposition of  $L^1$

*Classification:* 35J05, 31B10

### Preliminaries

In this paper the following notation is used: Small letters, such as  $x, y$ , will denote points in  $\mathbb{R}^n$ ,  $S^{n-1}$  the  $(n - 1)$ -dimensional unit sphere and  $\sigma$  the surface measure on  $S^{n-1}$ .

Consider, for  $\alpha > 0$  fixed, the Helmholtz equation

$$\Delta u - 2\alpha u = 0 \quad \text{on } \mathbb{R}^n.$$

**Theorem A.** *A function  $u$  on  $\mathbb{R}^n$  is a difference of two positive solutions of the Helmholtz equation if and only if there is a signed measure  $\mu_u$  on  $S^{n-1}$  such that for all  $x \in \mathbb{R}^n$*

$$\int_{S^{n-1}} e^{\lambda\langle x,y \rangle} d|\mu_u|(y) < \infty$$

and

$$u(x) = \int_{S^{n-1}} e^{\lambda\langle x,y \rangle} d\mu_u(y),$$

---

Support of the Charles University Grant Agency (GAUK 186/96) is gratefully acknowledged

where  $\lambda = (2\alpha)^{-1/2}$ .

The solution  $u$  is positive if and only if  $\mu_u$  is a measure.

PROOF: This representation theorem can be proved by means of Martin boundary, see [8]. For a different proof, see [6]. □

The solution corresponding to  $\sigma$  will be denoted by  $h$ .

For  $\nu \in \mathbb{R}$  the function  $I_\nu$  is “the Bessel function with an imaginary argument” of the order  $\nu$  regular at zero. (For details see any book about Bessel functions, for example [14, p. 17].)

Then

$$h(x) = C\lambda^{(2-n)/2}\|x\|^{(2-n)/2}I_{(n-2)/2}(\lambda\|x\|),$$

with  $C$  chosen so that  $h(0) = \omega_n$ , the area of the unit sphere in  $\mathbb{R}^n$ . (See [6, p. 261].)

For  $f, g$  two functions on  $\mathbb{R}^n$ ,  $f \sim g$  will mean that  $\lim_{\|x\| \rightarrow \infty} \frac{f(x)}{g(x)} = 1$ .

As  $I_\nu(\|x\|) \sim (2\pi\|x\|)^{-1/2}e^{\|x\|}$  (see for example [14, pages 17 and 203]), we have that

$$\lim_{\|x\| \rightarrow \infty} \frac{h(x)\|x\|^{(n-1)/2}}{e^{\lambda\|x\|}} = C\lambda^{(2-n)/2}(2\pi)^{-1/2},$$

this constant will be denoted by  $\kappa$ .

A solution  $u$  of the Helmholtz equation will be called  $h$ -bounded if there exist real constants  $c_1$  and  $c_2$  such that  $c_1h(x) \leq u(x) \leq c_2h(x)$  for all  $x \in \mathbb{R}^n$ .

Moreover, a solution  $u$  of the Helmholtz equation will be called simple if there exists a  $\sigma$ -measurable subset  $A$  of  $S^{n-1}$  such that  $u(x) = \int_A e^{\lambda\langle x,y \rangle} d\sigma(y)$  for any  $x \in \mathbb{R}^n$ .

**Definition.** For  $y \in S^{n-1}$ ,  $b \in \mathbb{R}^+$ ,  $k \in \mathbb{R}^+$  define the admissible region  $A(y, b)$  to be

$$\{x \in \mathbb{R}^n; \|x - \|x\|y\| < b\|x\|^{1/2}\}$$

and the truncated admissible region  $A^k(y, b)$  to be

$$A(y, b) \cap \{x \in \mathbb{R}^n; \|x\| > k\}.$$

Let  $M \subset \mathbb{R}^n$  and  $y \in S^{n-1}$ . The point  $y$  will be called a  $b$ -admissible limit point of  $M$  if for any  $k \in \mathbb{R}^+$  the set  $M \cap A^k(y, b)$  is not empty. The point  $y$  will be called an admissible limit point of  $M$  if there exists  $b \in \mathbb{R}^+$  such that  $y$  is a  $b$ -admissible limit point of  $M$ .

A function  $f$  on  $\mathbb{R}^n$  is said to converge admissibly at  $y$  if, for all  $b > 0$ ,  $f$  restricted to  $A(y, b)$  has a limit at  $\infty$ .

We will write  $A\text{-}\lim_{x \rightarrow y} f(x)$ .

The space  $\mathbb{R}^n$  endowed with the sheaf of solutions of the Helmholtz equation is a strong harmonic space in the sense of Bauer, see [2, p. 86].

Terms as harmonic functions, superharmonic functions and reduced functions are related to this harmonic space and have a standard meaning.

This harmonic space satisfies conditions (1)–(10) in [13], see [13], and so minimal thinness at points of  $S^{n-1}$  is well defined and the Fatou-Naïm-Doob theorem holds. For the reader’s convenience the basic facts are presented here.

**Definition.** Let  $M \subset \mathbb{R}^n$ ,  $v$  positive superharmonic function on  $D$ . The reduction of  $v$  on  $M$  is defined as

$$R_v^M = \inf\{u; u \geq v \text{ on } M, u \text{ is positive superharmonic function on } \mathbb{R}^n\}.$$

Let  $M \subset \mathbb{R}^n$  and  $y \in S^{n-1}$ . The set  $M$  is minimal thin at  $y$  if

$$R_{e^{\lambda(\cdot, y)}}^M \neq e^{\lambda(\cdot, y)}.$$

The minimal fine filter at  $y$  is filter:  $\mathcal{F}(y) = \{M \subset \mathbb{R}^n; \mathbb{R}^n \setminus M \text{ is minimal thin at } y\}$ .

A function  $f$  converging along  $\mathcal{F}(y)$  is said to have a minimal fine limit at  $y$ . This limit will be denoted  $\text{mf-lim } f(x)$ .

**Theorem B** (Limit theorems). Let  $u$  be a positive solution and  $v$  be a strictly positive solution of the Helmholtz equation defined on all  $\mathbb{R}^n$  and  $\mu_u, \mu_v$  be their representing measures on  $S^{n-1}$ .

Then the following equalities hold:

$$A\text{-}\lim_{x \rightarrow y} \frac{u(x)}{v(x)} = \frac{d\mu_u}{d\mu_v}(y)$$

for  $\mu_v$ -almost all points  $y$  of  $S^{n-1}$  (admissible convergence);

$$\text{mf-lim}_{x \rightarrow y} \frac{u(x)}{v(x)} = \frac{d\mu_u}{d\mu_v}(y)$$

for  $\mu_v$ -almost all points  $y$  of  $S^{n-1}$  (the Fatou-Naïm-Doob limit theorem).

PROOF: See [9, p. 85] and [13]. □

*Remark.* For  $v = h$ , the admissible convergence follows from the minimal fine convergence (even in a more general situation); see [9, p. 84].

Let  $x \in \mathbb{R}^n$ ,  $b, c, k \in \mathbb{R}^+$  and  $M \subset \mathbb{R}^n$ . In this paper, the following subsets of  $\mathbb{R}^n$  will be of special interest:

$$\begin{aligned}
 B(x, c) &= \{z \in \mathbb{R}^n; \|z - x\| \leq c\}, \\
 S(x, b, k) &= \{z \in \mathbb{R}^n; \|z\| = k\|x\| \text{ and } \|z - kx\| < k^{\frac{1}{2}}b\|x\|^{\frac{1}{2}}\}, \\
 M_{S,b,k} &= \cup_{x \in M} S(x, b, k), \\
 S(x, b) &= S(x, b, 1), \\
 M_{S,b} &= \cup_{x \in M} S(x, b), \\
 cM &= \{z \in \mathbb{R}^n; \text{there exists } x \in M \text{ such that } z = cx\}.
 \end{aligned}$$

Let  $x, y \in \mathbb{R}^n$ ,  $\alpha_{x,y}$  will denote the angle between  $x$  and  $y$ .

**The main results**

**Theorem.** *Let  $M \subset \mathbb{R}^n$ . Then the following statements are equivalent:*

(i)

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all simple solutions  $u$  of the Helmholtz equation;

(ii)

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all  $h$ -bounded solutions  $u$  of the Helmholtz equation;

(iii)

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all positive solutions  $u$  of the Helmholtz equation;

(iv) the set of points of  $S^{n-1}$  which are not admissible limit points of  $M$  has  $\sigma$ -measure zero;

(v) for any  $b \in \mathbb{R}^+$ , the set of points of  $S^{n-1}$  which are not  $b$ -admissible limit points of  $M$  has  $\sigma$ -measure zero;

(vi) there exist  $b, k \in \mathbb{R}^+$ , such that the set of points of  $S^{n-1}$  at which  $M_{S,b,k}$  is minimal thin has  $\sigma$ -measure zero;

(vii) for any  $b, k \in \mathbb{R}^+$ , the set of points of  $S^{n-1}$  at which  $M_{S,b,k}$  is minimal thin has  $\sigma$ -measure zero;

(viii) if  $\nu$  is a countably finite Borel measure with  $\text{supp}(\nu) = \overline{M}$ , then for every  $f \in L_1(S^{n-1})$  there exists  $\Phi \in L_1(\nu)$  such that

$$(1) \quad f(y) = \int_{\mathbb{R}^n} \Phi(x) \frac{e^{\lambda \langle x, y \rangle}}{h(x)} d\nu(x)$$

for  $\sigma$ -almost all  $y$  and

$$\|f\|_{L_1(S^{n-1})} = \inf \{ \|\Phi\|_{L_1(\nu)}; (1) \text{ holds for some } \Phi \in L_1(\nu) \};$$

(ix) for every  $f \in L_1(S^{n-1})$ , there is a sequence  $\{x_k\}$ ,  $x_k \in M$  and  $\{\lambda_k\} \in l_1$  such that

$$(2) \quad f(y) = \sum_{k=1}^{\infty} \lambda_k \frac{e^{\lambda \langle x_k, y \rangle}}{h(x_k)}$$

for  $\sigma$ -almost all  $y$  and

$$\|f\|_{L_1(S^{n-1})} = \inf \left\{ \sum_{k=1}^{\infty} |\lambda_k|; (2) \text{ holds for some } \{x_k\} \text{ in } M \right\};$$

(x) if  $\nu$  is a countably finite Borel measure with  $\text{supp}(\nu) = \overline{M}$ , then for every  $f \in L_1(S^{n-1})$  there exists  $\Phi \in L_1(\nu)$  such that

$$(3) \quad f(y) = \kappa^{-1} \int_{\mathbb{R}^n} \Phi(x) e^{\lambda \|x\| (\cos \alpha_{x,y} - 1)} \|x\|^{(n-1)/2} d\nu(x)$$

for  $\sigma$ -almost all  $y$ ;

moreover for any  $c \in \mathbb{R}^+$  there exists a function  $\Phi$  satisfying (3), such that  $\Phi = 0$  on  $B(0, c)$ , and

$$\|f\|_{L_1(S^{n-1})} = \inf \{ \|\Phi\|_{L_1(\nu)}; (3) \text{ holds for some } \Phi \in L_1(\nu), \Phi = 0 \text{ on } B(0, c) \};$$

(xi) for every  $f \in L_1(S^{n-1})$  and for any  $c \in \mathbb{R}^+$ , there is a sequence  $\{x_k\}$ ,  $x_k \in M$ ,  $\|x_k\| > c$  and  $\{\lambda_k\} \in l_1$  such that

$$(4) \quad f(y) = \kappa^{-1} \sum_{k=1}^{\infty} \lambda_k e^{\lambda \|x_k\| (\cos \alpha_{x,y} - 1)} \|x_k\|^{(n-1)/2}$$

for  $\sigma$ -almost all  $y$ ; such that

$$\|f\|_{L_1(S^{n-1})} = \inf \left\{ \sum |\lambda_k|; (4) \text{ holds for some } \{x_k\} \text{ in } M \setminus B(0, c) \right\}.$$

*Remark.* A set satisfying the condition (i) will be called a set of determination.

### Proof of Theorem

We will need the following theorem:

**Theorem 1.** Let  $u$  be a positive solution of the Helmholtz equation on  $\mathbb{R}^n$  and  $\mu_u$  its representing measure on  $S^{n-1}$ . Then

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \text{ess inf}_{y \in S^{n-1}} \frac{d\mu_u}{d\sigma}(y).$$

If  $u$  is an  $h$ -bounded function then

$$\sup_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \operatorname{ess\,sup}_{y \in S^{n-1}} \frac{d\mu_u}{d\sigma}(y) \quad \text{and} \quad \sup_{x \in \mathbb{R}^n} \frac{|u(x)|}{h(x)} = \operatorname{ess\,sup}_{y \in S^{n-1}} \left| \frac{d\mu_u}{d\sigma}(y) \right|.$$

PROOF: By the Lebesgue-Radon-Nikodym theorem the existence of measures  $\mu_a$  and  $\mu_s$ , such that  $\mu_u = \mu_a + \mu_s$ ,  $\mu_a \leq \sigma$  and  $\mu_s \perp \sigma$ , is guaranteed.

Let  $f_u = \frac{d\mu_u}{d\sigma}$ . Denote  $k_1 = \inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)}$  and  $k_2 = \operatorname{ess\,inf}_{y \in S^{n-1}} f_u(y)$ .

Obviously,

$$u(x) = \int_{S^{n-1}} e^{\lambda\langle x,y \rangle} d(f_u\sigma + \mu_s)(y) = \int_{S^{n-1}} f_u(y)e^{\lambda\langle x,y \rangle} d\sigma(y) + \int_{S^{n-1}} e^{\lambda\langle x,y \rangle} d\mu_s(y)$$

and, as the last term is positive,

$$u(x) \geq \int_{S^{n-1}} f_u(y)e^{\lambda\langle x,y \rangle} d\sigma(y) \geq k_2 \int_{S^{n-1}} e^{\lambda\langle x,y \rangle} d\sigma(y) = k_2 h(x)$$

for all  $x \in \mathbb{R}^n$ . This gives  $k_1 \geq k_2$ .

On the other hand  $u(x) - k_1 h(x)$  is a positive solution of the Helmholtz equation and thus  $\mu_u - k_1\sigma$  is a measure, so  $(f_u - k_1)\sigma + \mu_s$  is a measure. Since  $\mu_s \perp \sigma$ ,  $(f_u - k_1)\sigma$  is a measure and consequently  $\operatorname{ess\,inf}_{y \in S^{n-1}} f_u(y) \geq k_1$ , or  $k_2 \geq k_1$ .

The proof of the rest of the theorem is analogous. □

**Proof of equivalence of (i), (ii), (iii), (iv) and (v).**

As the implications (v)⇒(iv), (ii)⇒(i) and (iii)⇒(ii) are trivial (in the last implication just take  $u - c_1 h$  instead of  $u$ ), we will prove (iv)⇒(iii) and (i)⇒(v).

**Theorem 2.** *Let  $M$  be a subset of  $\mathbb{R}^n$  and  $\sigma$ -almost every point  $y \in S^{n-1}$  be an admissible limit point of  $M$ . Then*

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for every positive solution  $u$  of the Helmholtz equation on  $\mathbb{R}^n$  and

$$\sup_{x \in \mathbb{R}^n} \frac{|u(x)|}{h(x)} = \sup_{x \in M} \frac{|u(x)|}{h(x)}$$

for every  $h$ -bounded solution  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ .

PROOF: The assertion follows immediately from the previous theorem and the limit theorem. □

**Lemma 1.** *Let  $b$  be a positive number and  $x \in \mathbb{R}^n$ . Denote  $C(x, b)$  the set of all  $y \in S^{n-1}$  such that  $x \in A(y, b)$ . Then*

$$C(x, b) = \left\{ y \in S^{n-1}; \left\| y - \frac{x}{\|x\|} \right\| < \frac{b}{\sqrt{\|x\|}} \right\}$$

and there exists a positive number  $c$  such that

$$\int_{C(x,b)} e^{\lambda \langle x, y \rangle} d\sigma(y) \geq c \cdot h(x),$$

whenever  $x \in \mathbb{R}^n \setminus \{0\}$ .

PROOF: See [9, p. 84]. □

**Theorem 3.** *Let  $M \subset \mathbb{R}^n$  and  $b \in \mathbb{R}^+$ . If*

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all simple solutions of the Helmholtz equation, then  $\sigma$ -almost every point  $y \in S^{n-1}$  is a  $b$ -admissible limit point of  $M$ .

PROOF: Suppose that it is not true.

Denote the set  $M \cap \{x \in \mathbb{R}^n; \|x\| > k\}$  by  $M^k$  and the set of all  $b$ -admissible limit points of  $M$  by  $M_b$ . As  $M_b = \bigcap_{k \in \mathbb{N}} (\bigcup_{x \in M^k} C(x, b))$  is a  $G_\delta$  set, it is a  $\sigma$ -measurable subset of  $S^{n-1}$ . Then its complement  $M'_b$  is also measurable and by our assumption  $\sigma(M'_b) > 0$ .

Recall that for  $k \in \mathbb{N}$  and  $y \in S^{n-1}$ ,  $A^k(y, b)$  denotes the truncated admissible region  $A(y, b) \cap \{x \in \mathbb{R}^n; \|x\| > k\}$ . Then, for every  $y \in M'_b$ , there is  $k_y \in \mathbb{N}$  such that  $A^{k_y}(y, b) \cap M$  is empty. Denote by  $D_k$  the set of  $y \in M'_b$  for which  $A^k(y, b) \cap M$  is empty.

As  $D_k$  is a complement of  $\bigcup_{x \in M^k} C(x, b)$ , it is a  $\sigma$ -measurable subset of  $S^{n-1}$ .

Since  $\bigcup_{k=1}^\infty D_k = M'_b$ , the Lebesgue measure of at least one of the sets  $D_k$ , say of  $D_{k_0}$ , is strictly positive. Denote this set by  $D$  and its complement  $(S^{n-1}) \setminus D$  by  $D'$ .

It is clear that  $C(x, b) \subset D'$ , whenever  $x \in M^k$ .

For any measurable set  $A \subset S^{n-1}$  we define

$$u_A(x) = \int_A e^{\lambda \langle x, y \rangle} d\sigma(y), \quad x \in \mathbb{R}^n.$$

So  $u_A$  is a simple solution of the Helmholtz equation. By Theorem 1 we get that if  $\sigma(A) > 0$ , then  $\sup_{x \in \mathbb{R}^n} \frac{u_A(x)}{h(x)} = 1$  and if  $\sigma(A') > 0$ , then  $\inf_{x \in \mathbb{R}^n} \frac{u_A(x)}{h(x)} = 0$ .



The set  $D$  has a positive measure, so the function  $u_{D'}$  is a simple solution of the Helmholtz equation and

$$\inf_{x \in \mathbb{R}^n} \frac{u_{D'}(x)}{h(x)} = 0.$$

But  $C(x, b)$  is a subset of  $D'$  for every  $x \in M^k$ . Now from the above lemma there exists a constant  $c$  such that

$$\frac{u_{D'}(x)}{h(x)} \geq \frac{u_{C(x,b)}(x)}{h(x)} \geq c$$

for every  $x \in M^k$ .

We arrive at

$$\inf_{x \in M^k} \frac{u_{D'}(x)}{h(x)} \geq c.$$

Now it will be shown that

$$\inf_{x \in M \setminus M^k} \frac{u_{D'}(x)}{h(x)} > 0.$$

As  $h$  is positive and continuous and  $B(0, k)$  is compact, there exists  $c_1 \in \mathbb{R}^+$  such that  $h(x) \leq c_1$  for all  $x \in B(0, k)$ .

It follows

$$\begin{aligned} u_{D'}(x) &= \int_{D'} e^{\lambda \langle x, y \rangle} d\sigma(y) \geq \int_{D'} e^{-\lambda \|x\| \cdot \|y\|} d\sigma(y) = \\ &= \int_{D'} e^{-\lambda \|x\|} d\sigma(y) = \sigma(D') \cdot e^{-\lambda \|x\|} \geq \sigma(D') \cdot e^{-\lambda k}. \end{aligned}$$

Let us denote this positive constant by  $c_2$ .

Thus

$$\inf_{x \in B(0, k)} \frac{u_{D'}(x)}{h(x)} \geq \frac{c_2}{c_1}.$$

Consequently,

$$\inf_{x \in M} \frac{u_{D'}(x)}{h(x)} \geq \min(c, \frac{c_2}{c_1}) > 0,$$

contradicting our assumption. □

### Proof of (vi) and (vii).

The implication (vii) $\Rightarrow$ (vi) is trivial. Now it will be proved, that (vi) $\Rightarrow$ (iii) and (v) $\Rightarrow$ (vii).

**Theorem 4.** Let  $n \in \mathbb{N}$ ,  $b \in \mathbb{R}^+$ . Then there exists a positive constant  $c$ , such that for every  $x \in \mathbb{R}^n$ , for every  $z \in S(x, b, \frac{1}{2})$  and for every positive solution  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ ,

$$\frac{u(z)}{h(z)} \geq c \frac{u(x)}{h(x)},$$

and for any  $M \subset \mathbb{R}^n$

$$\inf_{x \in M_{S,b,\frac{1}{2}}} \frac{u(x)}{h(x)} \geq c \inf_{x \in M} \frac{u(x)}{h(x)}.$$

PROOF: For the first part, see [9, p. 83]. The second part immediately follows. □

**Theorem 5.** Let  $M \subset \mathbb{R}^n$  such that the set of points of  $S^{n-1}$  at which  $M$  is minimal thin is of  $\sigma$ -measure zero.

Then

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all positive solutions  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ .

PROOF: It follows from Theorem 1 and from the Fatou-Naim-Doob limit theorem. □

**Theorem 6.** Let  $M \subset \mathbb{R}^n$  and  $b \in \mathbb{R}^+$  such that the set of points of  $S^{n-1}$  at which  $M_{S,b,\frac{1}{2}}$  is minimal thin is of  $\sigma$ -measure zero.

Then there exists a constant  $c$  depending only on  $b$  and  $n$  such that

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} \geq c \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all positive solutions  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ .

PROOF: This theorem is obtained by combining Theorems 4 and 5. □

**Theorem 7.** Let  $M \subset \mathbb{R}^n$ . Then the following statements are equivalent:

(i)

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all positive solutions  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ ;

(ii) there exists  $c > 0$  such that

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} \geq c \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all positive solutions  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ .

PROOF: (i)  $\Rightarrow$  (ii) is clear, put  $c = 1$ .

(ii)  $\Rightarrow$  (i) Let us suppose that there exists a set  $M$  satisfying (ii), but not (i). Then  $c$  in (ii) belongs to  $(0, 1)$ .

Let  $u$  be a positive solution of the Helmholtz equation for which (i) is not true.

$$\text{Denote } \inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = c_1 \quad \text{and} \quad \inf_{x \in M} \frac{u(x)}{h(x)} = c_2.$$

Thus by our assumptions,  $c_2 > c_1 \geq c \cdot c_2$ .

Let  $v(x) = u(x) - c_1 h(x)$  for  $x \in \mathbb{R}^n$ .

Then  $v$  is a positive solution of the Helmholtz equation and

$$\inf_{x \in \mathbb{R}^n} \frac{v(x)}{h(x)} = c_2 - c_1 = 0, \quad \text{and} \quad \inf_{x \in M} \frac{v(x)}{h(x)} = c_2 - c_1 > 0,$$

which is a contradiction with (ii). □

**Theorem 8.** *Let  $M \subset \mathbb{R}^n$  and  $b \in \mathbb{R}^+$  such that the set of points of  $S^{n-1}$  at which  $M_{S,b,\frac{1}{2}}$  is minimal thin has  $\sigma$ -measure zero.*

Then

$$\inf_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \inf_{x \in M} \frac{u(x)}{h(x)}$$

for all positive solution  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ .

PROOF: The result is obtained by combining two previous theorems. □

**Theorem 9.** *Let  $M \subset \mathbb{R}^n$ ,  $y \in S^{n-1}$  and  $b \in \mathbb{R}^+$ . If  $y$  is an admissible limit point of  $M$ , then  $M_{S,b}$  is not minimal thin at  $y$ .*

PROOF: Let  $\{x_k\}$  be a sequence of points of  $M$  converging to  $y$  admissibly — it means that there exists  $b_1 \in \mathbb{R}^+$  such that  $\{x_k\}$  converges  $b_1$ -admissibly.

Then a straightforward calculation gives that  $S(x_k, b) \subset A(y, b_1 + b)$ .

Since the Helmholtz equation is invariant with respect to linear isometries of  $\mathbb{R}^n$ , the harmonic measure  $\mu_0$  (for the notion of the harmonic measure, see [2, p. 120]) on  $\partial B(0, r)$  corresponding to 0, is invariant with respect to isometries of  $\partial B(0, r)$  and hence it is a multiple of the surface measure  $\sigma_n$  on  $\partial B(0, r)$ .

As  $\mu_0(\partial B(0, r)) = \frac{h(0)}{h(r \cdot e_1)}$  and  $h(0) = \omega_n$  we have that for any  $\sigma$ -measurable subset  $E$  of  $\partial B(0, r)$

$$\mu_0(E) = \frac{h(0)\sigma_n(E)}{h(r \cdot e_1)\omega_n r^{n-1}} = \frac{\sigma(r^{-1}E)}{h(r \cdot e_1)}.$$

The proof of the theorem is finished in the same way as the proof of Proposition 2.2 in [9, p. 82]; for the reader's convenience it is given here.

Let us denote  $u_k$  the solution of the Dirichlet problem on  $B(0, \|x_k\|)$  with boundary value 1 on  $S(x_k, b)$  and 0 on the rest of the boundary.

Hence  $u_k(0) = h(\|x_k\|)^{-1} \sigma(\|x_k\|^{-1} S(x_k, b)) \sim b^{(n-1)/2} \|x_k\|^{-(n-1)/2} h(\|x_k\|)^{-1}$ .

As  $h(x) \sim \frac{\kappa e^{\lambda \|x\|}}{\|x\|^{(n-1)/2}}$  (see Preliminaries),

$$u_k(0) \sim \kappa b^{(n-1)/2} e^{-\lambda \|x_k\|}.$$

Now denote  $v_k$  the solution of the Dirichlet problem on  $B(0, \|x_k\|)$  with boundary value  $e^{\lambda \langle x, y \rangle}$  on  $S(x_k, b)$  and 0 on the rest of the boundary.

For any  $b_0 \in \mathbb{R}^+$  there is a positive constant  $c_1$  such that for all  $x \in A(y, b_0)$   $c_1^{-1} e^{\lambda \|x\|} \leq e^{\lambda \langle x, y \rangle} \leq c_1 e^{\lambda \|x\|}$  whenever  $x \in A(y, b_0)$ .

(Indeed,  $0 \leq \lambda(\|x\| - \langle x, y \rangle) = \lambda \|x\|(1 - \langle x', y \rangle) = \frac{1}{2} \lambda \|x\| \|x' - y\|^2 \leq \frac{1}{2} \lambda b_0^2$ , where  $x' = \frac{x}{\|x\|}$ .)

As  $S(x_k, b) \subset A(y, b_1 + b)$ , for the boundary values of  $u_k$  and  $v_k$  holds

$$c_1^{-1} e^{\lambda \|x_k\|} u_k(x) \leq v_k(x) \leq c_1 e^{\lambda \|x_k\|} u_k(x)$$

for  $x \in \partial B(0, \|x_k\|)$  and hence for any  $x \in B(0, \|x_k\|)$ .

Namely this is true for 0 and so, using the above relation for  $u_k(0)$ , the existence of a positive constant  $c_2$  such that

$$c_2^{-1} \leq v_k(0) \leq c_2$$

for any  $k \in \mathbb{N}$  is guaranteed.

Let  $S = \cup_{k \in \mathbb{N}} S(x_k, b)$ . The Perron-Wiener-Brelot method of solving the Dirichlet problem shows that, for any  $k \in \mathbb{N}$ , the inequality  $v_k \leq R_{e^{\lambda \langle \cdot, y \rangle}}^S$  holds on  $B(0, \|x_k\|)$ . As  $\{v_k\}$  is bounded in 0, it has by virtue of the Harnack inequality a converging subsequence. Denoting its limit by  $v$ , it is easy to see that  $v$  is a positive solution of the Helmholtz equation,  $v(0) \geq c_2^{-1}$  and  $v \leq R_{e^{\lambda \langle \cdot, y \rangle}}^S$ . Hence its representing measure  $\mu_v \leq \delta_y$  and thus  $R_{e^{\lambda \langle \cdot, y \rangle}}^S = e^{\lambda \langle \cdot, y \rangle}$ , it means that  $S$  is not minimal thin at  $y$  and hence  $M_{S,b}$  is not minimal thin at  $y$ .  $\square$

So far we have proved the implication (vi)  $\Rightarrow$  (iii) for  $k = \frac{1}{2}$  and the implication (v)  $\Rightarrow$  (vii) for  $k = 1$ . The conditions for  $k$  will be removed using the following lemma.

**Lemma 2.** *Let  $M \subset \mathbb{R}^n$ ,  $c \in \mathbb{R}^+$ ,  $y \in S^{n-1}$ . The point  $y$  is an admissible limit point of the set  $M$  if and only if  $y$  is a admissible limit point of  $cM$ .*

Let  $x \in \mathbb{R}^n$  and  $b, k \in \mathbb{R}^+$ . Then

$$S(x, b, k) = S(kx, b) = S(2kx, b, \frac{1}{2})$$

and

$$M_{S,b,k} = (kM)_{S,b} = (2kM)_{S,b,\frac{1}{2}}.$$

PROOF: A straightforward calculation. □

Now, it is easy to finish the proof of (vi) and (vii).

Let  $k \in \mathbb{R}^+$ . Using the first part of the lemma and equivalence of (i) and (v) it follows that  $M$  is a set of determination if and only if  $kM$  is a set of determination.

From that and from  $(kM)_{S,b} = M_{S,b,k}$  it immediately follows that (vii) is true for any positive  $k$ .

From  $M_{S,b,k} = (2kM)_{S,b,\frac{1}{2}}$  it follows that if (vi) holds for some  $k$  then  $2kM$  is a set of determination, so  $M$  is a set of determination.

**Proof of (viii) and (ix).**

The implication (viii) $\Rightarrow$ (ix) is trivial. (Take a countable subset of  $M$  and the counting measure on it.) We will prove (v) $\Rightarrow$ (viii) and (ix) $\Rightarrow$ (ii).

**Theorem 10.** *Let  $M$  be a subset of  $\mathbb{R}^n$  and  $\nu$  be a countably-finite measure on  $\mathbb{R}^n$  such that  $\text{supp}(\nu) = \overline{M}$ . Let*

$$\sup_{x \in \mathbb{R}^n} \frac{|u(x)|}{h(x)} = \sup_{x \in M} \frac{|u(x)|}{h(x)}$$

for every  $h$ -bounded solution  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ .

Then, for any  $f$  in  $L_1(S^{n-1})$ , there exists  $\Phi$  in  $L_1(\nu)$  such that

$$(1) \quad f = \int_{\mathbb{R}^n} \Phi(x) \frac{e^{\lambda \langle x, \cdot \rangle}}{h(x)} d\nu(x)$$

$\sigma$ -almost everywhere and

$$\|f\|_{L_1(S^{n-1})} = \inf \{ \|\Phi\|_{L_1(\nu)}; (1) \text{ holds for some } \Phi \in L_1(\nu) \}.$$

We will need the following version of the closed range theorem (see [12, p. 97]). Let  $\mathcal{X}$  and  $\mathcal{Y}$  be Banach spaces,  $T$  a bounded linear mapping of  $\mathcal{X}$  into  $\mathcal{Y}$ . If there exists a constant  $c > 0$  such that  $\|T^*y^*\| \geq c\|y^*\|$  for all  $y^* \in \mathcal{Y}^*$  then  $T\mathcal{X} = \mathcal{Y}$ . In our situation,  $\mathcal{X} = L_1(\nu)$ ,  $\mathcal{Y} = L_1(S^{n-1})$  and for  $\Phi \in L_1(\nu)$  we define

$$T_\nu \Phi = \int_{S^{n-1}} \Phi(x) \frac{e^{\lambda \langle x, \cdot \rangle}}{h(x)} d\nu(x).$$

**Lemma 3.** *The mapping  $T_\nu$  is a bounded linear mapping of  $L_1(\nu)$  into  $L_1(S^{n-1})$ ,  $\|T_\nu\| = 1$ ;  $T_\nu^*$  is the bounded mapping  $L_\infty(S^{n-1})$  into  $L_\infty(\nu)$  such that*

$$T_\nu^*g(x) = \frac{1}{h(x)} \int_{S^{n-1}} e^{\lambda\langle x,y \rangle} g(y) d\sigma(y).$$

PROOF: Using the Fubini theorem we arrive at

$$\begin{aligned} \|T_\nu\Phi\|_{L_1(S^{n-1})} &= \int_{S^{n-1}} |T_\nu\Phi| d\sigma = \int_{S^{n-1}} \left| \int_{\mathbb{R}^n} \Phi(x) \frac{e^{\lambda\langle x,y \rangle}}{h(x)} d\nu(x) \right| d\sigma(y) \leq \\ &\int_{S^{n-1}} \left( \int_{\mathbb{R}^n} |\Phi(x)| \frac{e^{\lambda\langle x,y \rangle}}{h(x)} d\nu(x) \right) d\sigma(y) = \int_{\mathbb{R}^n} \left( \int_{S^{n-1}} |\Phi(x)| \frac{e^{\lambda\langle x,y \rangle}}{h(x)} d\sigma(y) \right) d\nu(x) = \\ &\int_{\mathbb{R}^n} \frac{|\Phi(x)|}{h(x)} \left( \int_{S^{n-1}} e^{\lambda\langle x,y \rangle} d\sigma(y) \right) d\nu(x) = \int_{\mathbb{R}^n} \frac{|\Phi(x)|}{h(x)} h(x) d\nu(x) = \|\Phi\|_{L_1(\nu)}. \end{aligned}$$

So the first part of Lemma is proved.

Let  $g \in L_\infty(S^{n-1})$  and  $\Phi \in L_1(\nu)$ . Using again the Fubini theorem we have

$$\begin{aligned} [\Phi, T_\nu^*g] &= [T_\nu\Phi, g] = \int_{S^{n-1}} g \cdot T_\nu\Phi d\sigma = \int_{S^{n-1}} g(y) \left( \int_{\mathbb{R}^n} \Phi(x) \frac{e^{\lambda\langle x,y \rangle}}{h(x)} d\nu(x) \right) d\sigma(y) = \\ &\int_{\mathbb{R}^n} \frac{\Phi(x)}{h(x)} \left( \int_{S^{n-1}} g(y) e^{\lambda\langle x,y \rangle} d\sigma(y) \right) d\nu(x) = \left[ \Phi, \frac{1}{h} \int_{S^{n-1}} e^{\lambda\langle \cdot, y \rangle} g(y) d\sigma(y) \right]. \end{aligned}$$

□

**Proof of Theorem.** We shall prove the existence of a constant  $c > 0$  such that  $\|T_\nu^*g\|_{L_\infty(\nu)} \geq c\|g\|_{L_\infty(S^{n-1})}$  for all  $g \in L_\infty(S^{n-1})$  and the first part of the theorem will be proved.

The function  $h \cdot (T_\nu^*g)$  is an  $h$ -bounded solution of the Helmholtz equation on  $\mathbb{R}^n$ . Then, by hypothesis,

$$\sup_{x \in M} |(T_\nu^*g)(x)| = \sup_{x \in \mathbb{R}^n} |(T_\nu^*g)(x)| = \|g\|_{L_\infty(S^{n-1})}.$$

Since  $T_\nu^*g$  is a continuous function on  $\mathbb{R}^n$  and  $\text{supp}(\nu) = \overline{M}$ ,

$$\|T_\nu^*g\|_{L_\infty(\nu)} = \sup_{x \in M} |(T_\nu^*g)(x)|.$$

Consequently,

$$\|T_\nu^*g\|_{L_\infty(\nu)} = \|g\|_{L_\infty(S^{n-1})}.$$

So we can take  $c = 1$ . The first part of Theorem is proved.

To prove the other part define the space

$$\mathcal{Z} = L_1(\nu) / \ker T_\nu.$$

For  $z \in \mathcal{Z}$  and  $\Phi \in z$  put  $Sz = T_\nu \Phi$ .

Then  $S$  is an invertible bounded linear mapping of  $\mathcal{Z}$  into  $L_1(S^{n-1})$  and so its adjoint  $S^*$  is an invertible bounded linear mapping of  $L_\infty(S^{n-1})$  into  $\mathcal{Z}^*$  (see [12, p. 94]).

Let  $z \in \mathcal{Z}$ ,  $\Phi \in z$  and  $g \in L_\infty(S^{n-1})$ . Then we have

$$(S^*g)(z) = [Sz, g] = [T_\nu \Phi, g] = [\Phi, T_\nu^*g].$$

If  $\varepsilon > 0$ , there exists  $\Phi_0 \in L_1(\nu)$  with  $\|\Phi_0\|_{L_1(\nu)} = 1$  and

$$|[\Phi_0, T_\nu^*g]| > \|T_\nu^*g\|_{L_\infty(\nu)} - \varepsilon.$$

Let  $z_0$  denote the coset of  $\Phi_0$  in  $\mathcal{Z}$ . Then

$$|(S^*g)(z_0)| > \|T_\nu^*g\|_{L_\infty(\nu)} - \varepsilon$$

and

$$\|z_0\|_{\mathcal{Z}} \leq \|\Phi_0\|_{L_1(\nu)} = 1.$$

Therefore, the norm of the functional  $S^*g$  satisfies

$$\|S^*g\|_{\mathcal{Z}^*} > \|T_\nu^*g\|_{L_\infty(\nu)} - \varepsilon = \|g\|_{L_\infty(S^{n-1})} - \varepsilon.$$

Since  $\varepsilon$  was arbitrary, we proved that

$$\|S^*g\|_{\mathcal{Z}^*} \geq \|g\|_{L_\infty(S^{n-1})}$$

for any  $g \in L_\infty(S^{n-1})$ , and so, using the fact that the norm of any operator is the same as the norm of its adjoint (see [1, p. 93]) and the obvious fact that  $(S^*)^{-1} = (S^{-1})^*$ , we have

$$\|S^{-1}\| = \|(S^*)^{-1}\| \leq 1.$$

Fix  $f \in L_1(S^{n-1})$  and put  $z = S^{-1}f$ . Then

$$\|z\|_{\mathcal{Z}} \leq \|f\|_{L_1(S^{n-1})},$$

that is

$$\inf\{\|\Phi\|_{L_1(\nu)}; T_\nu \Phi = f\} \leq \|f\|_{L_1(S^{n-1})}.$$

By Lemma we have

$$\|f\|_{L_1(S^{n-1})} = \|T_\nu \Phi\|_{L_1(S^{n-1})} \leq \|T_\nu\| \cdot \|\Phi\|_{L_1(\nu)} = \|\Phi\|_{L_1(\nu)}.$$

So the opposite inequality holds as well. □

**Theorem 11.** *Let  $\nu$  be a countably finite measure on  $\mathbb{R}^n$  and  $\text{supp}(\nu) = \overline{M}$ . Assume that for every function  $f \in L_1(S^{n-1})$  there exists  $\Phi$  in  $L_1(\mathbb{R}^n)$  such that*

$$(1) \quad f = \int_{\mathbb{R}^n} \Phi(x) \frac{e^{\lambda\langle x, \cdot \rangle}}{h(x)} d\nu(x)$$

$\sigma$ -almost everywhere and

$$\|f\|_{L_1(S^{n-1})} = \inf \{ \|\Phi\|_{L_1(\nu)}; (1) \text{ holds for some } \Phi \text{ in } L_1(\nu) \}.$$

Then

$$\sup_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \sup_{x \in M} \frac{u(x)}{h(x)}$$

for any  $h$ -bounded positive solution  $u$  of the Helmholtz equation on  $\mathbb{R}^n$ .

PROOF: Put  $c = \sup_{x \in M} \frac{u(x)}{h(x)}$ . We have  $c < \infty$ .

Let  $\varepsilon > 0$ . If we fix  $x_0 \in \mathbb{R}^n$ , then  $e^{\lambda\langle x_0, \cdot \rangle} \in L_1(S^{n-1})$  and  $\|e^{\lambda\langle x_0, \cdot \rangle}\|_{L_1(S^{n-1})} = h(x_0)$ . By our assumptions there is a function  $\Phi \in L_1(\nu)$  such that

$$e^{\lambda\langle x_0, \cdot \rangle} = \int_{\mathbb{R}^n} \Phi(x) \frac{e^{\lambda\langle x, \cdot \rangle}}{h(x)} d\nu(x) \leq \int_{\mathbb{R}^n} |\Phi(x)| \frac{e^{\lambda\langle x, \cdot \rangle}}{h(x)} d\nu(x)$$

and

$$\|\Phi\|_{L_1(\nu)} < h(x_0) + \varepsilon.$$

As  $u$  is an  $h$ -bounded positive solution of the Helmholtz equation, we can integrate the first inequality with respect to  $f_u d\sigma$ . Using the Fubini theorem and the fact that  $u \leq ch$  on  $\text{supp}(\nu)$ , we have

$$\begin{aligned} u(x_0) &= \int_{S^{n-1}} e^{\lambda\langle x_0, y \rangle} f_u(y) d\sigma(y) \leq \int_{S^{n-1}} \left( \int_{\mathbb{R}^n} |\Phi(x)| e^{\lambda\langle x, y \rangle} d\nu(x) \right) f_u(y) d\sigma(y) = \\ & \int_{\mathbb{R}^n} |\Phi(x)| \left( \int_{S^{n-1}} e^{\lambda\langle x, y \rangle} f_u(y) d\sigma(y) \right) d\nu(x) = \int_{S^{n-1}} |\Phi(x)| u(x) d\nu(x) \leq \\ & \int_{S^{n-1}} c |\Phi(x)| d\nu(x) = c \|\Phi\|_{L_1(\nu)} \leq c(h(x_0) + \varepsilon). \end{aligned}$$

Since  $x_0$  and  $\varepsilon$  were arbitrary, we have  $\sup_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = c$ . □

Of course, the following special form of Theorem 11 holds:



**Theorem 12.** Let  $M$  be a subset of  $\mathbb{R}^n$ . Assume that for every function  $f \in L_1(S^{n-1})$  there exist  $\{\lambda_k\}_{k=1}^\infty \in l_1$  and a sequence  $\{x_k\}_{k=1}^\infty$  of points in  $M$  such that

$$(2) \quad f = \sum_{k=1}^\infty \lambda_k \frac{e^{\lambda \langle x_k, \cdot \rangle}}{h(x_k)}$$

$\sigma$ -almost everywhere and

$$\|f\|_{L_1(S^{n-1})} = \inf \left\{ \sum_{k=1}^\infty |\lambda_k|; (2) \text{ holds for some } \{x_k\} \text{ in } M \right\}.$$

Then

$$\sup_{x \in \mathbb{R}^n} \frac{u(x)}{h(x)} = \sup_{x \in M} \frac{u(x)}{h(x)}$$

for any bounded positive solution  $u$  of the Helmholtz equation.

**Proof of the conditions (x) and (xi).**

We will prove the equivalence of (viii) and (x). The equivalence of (ix) and (xi) is just a special form of it.

**Proof of (viii)  $\Rightarrow$  (x)**

Let us denote

$$K_1(x, y) = \frac{e^{\lambda \langle x, y \rangle}}{h(x)} \quad \text{and} \quad K_2(x, y) = \frac{e^{\lambda \langle x, y \rangle} \|x\|^{(n-1)/2}}{\kappa e^{\lambda \|x\|}}.$$

Then we have

$$\|K_1(x, \cdot)\|_{L_1(S^{n-1})} = \int_{S^{n-1}} \left| \frac{e^{\lambda \langle x, y \rangle}}{h(x)} \right| d\sigma(y) = 1$$

and

$$\begin{aligned} \|K_1(x, \cdot) - K_2(x, \cdot)\|_{L_1(S^{n-1})} &= \int_{S^{n-1}} \left| \frac{e^{\lambda \langle x, y \rangle}}{h(x)} - \frac{e^{\lambda \langle x, y \rangle} \|x\|^{(n-1)/2}}{\kappa e^{\lambda \|x\|}} \right| d\sigma(y) = \\ &= \int_{S^{n-1}} e^{\lambda \langle x, y \rangle} \left| \frac{1}{h(x)} - \frac{\|x\|^{(n-1)/2}}{\kappa e^{\lambda \|x\|}} \right| d\sigma(y) = \\ &= \left| \frac{1}{h(x)} - \frac{\|x\|^{(n-1)/2}}{\kappa e^{\lambda \|x\|}} \right| \int_{S^{n-1}} e^{\lambda \langle x, y \rangle} d\sigma(y) = \left| 1 - \frac{h(x) \|x\|^{(n-1)/2}}{\kappa e^{\lambda \|x\|}} \right|, \end{aligned}$$

from the asymptotic behaviour of the function  $h$  (see Preliminaries) it follows, that to every positive  $\varepsilon$ , there exists a positive number  $c_\varepsilon$  such that

$$\|K_1(x, \cdot) - K_2(x, \cdot)\|_{L_1(S^{n-1})} < \varepsilon$$

and

$$\|K_2(x, \cdot)\|_{L_1(S^{n-1})} < 1 + \varepsilon,$$

whenever  $\|x\| > c_\varepsilon$ .

Let  $f \in L_1(S^{n-1})$  and  $c > 1$ . Then there exists  $\Phi_0 \in L_1(\nu)$ , such that

$$f = \int_{\mathbb{R}^n} \Phi_0(x) K_1(x, \cdot) d\nu(x), \quad \text{and} \quad \|f\|_{L_1(S^{n-1})} \leq \|\Phi_0\|_{L_1(\nu)} \leq c \|f\|_{L_1(S^{n-1})},$$

and moreover, as (viii) is equivalent to (v) and (v) holds for  $M$ , if and only if it holds for  $M \setminus B(0, c_\varepsilon)$ ,  $\Phi_0$  can be chosen to be zero on  $B(0, c_\varepsilon)$ .

Put  $f_0 = f$ . Now, functions  $f_k \in L_1(S^{n-1})$  and  $\Phi_k \in L_1(\nu)$  for any  $k = 1, 2, \dots$ , will be defined.

$$f_{k+1} = f_k - \int_{\mathbb{R}^n} \Phi_k(x) K_2(x, \cdot) d\nu(x), \quad \text{for } k = 0, 1, \dots;$$

$\Phi_{k+1}$  is, for  $k = 0, 1, \dots$ , a function for which

$$f_{k+1} = \int_{\mathbb{R}^n} \Phi_{k+1}(x) K_1(x, \cdot) d\nu(x),$$

$$\|f_{k+1}\|_{L_1(S^{n-1})} \leq \|\Phi_{k+1}\|_{L_1(\nu)} \leq c \|f_{k+1}\|_{L_1(S^{n-1})}$$

and  $\Phi_{k+1}$  is zero on  $B(0, c_\varepsilon)$ .

We have  $f_0 \in L_1(S^{n-1})$  and  $\Phi_0 \in L_1(\nu)$  and above relations are satisfied. Suppose, it is true for  $0, 1, \dots, k$ , and prove it for  $k + 1$ :

$$\|f_{k+1}\|_{L_1(S^{n-1})} = \|f_k - \int_{\mathbb{R}^n} \Phi_k(x) K_2(x, \cdot) d\nu(x)\|_{L_1(S^{n-1})} =$$

$$\| \int_{\mathbb{R}^n} \Phi_k(x) K_1(x, y) d\nu(x) - \int_{\mathbb{R}^n} \Phi_k(x) K_2(x, y) d\nu(x) \|_{L_1(S^{n-1})} \leq$$

$$\int_{S^{n-1}} \int_{\mathbb{R}^n} |\Phi_k(x)(K_1(x, y) - K_2(x, y))| d\nu(x) d\sigma(y)$$

using Fubini theorem

$$= \int_{\mathbb{R}^n} |\Phi_k(x)| \int_{S^{n-1}} |K_1(x, y) - K_2(x, y)| d\sigma(y) \leq \varepsilon \|\Phi_k\|_{L_1(\nu)}.$$

So  $f_{k+1} \in L_1(S^{n-1})$  and by this fact and (v) and (viii) the existence of a function  $\Phi_{k+1}$  with required properties is guaranteed.

Combining the above estimates for  $\|\Phi_k\|_{L_1(\nu)}$  and  $\|f_{k+1}\|_{S^{n-1}}$  we obtain

$$\|f_{k+1}\|_{S^{n-1}} \leq c\varepsilon \|f_k\|_{L_1(S^{n-1})} \text{ for all } k = 0, 1, 2, \dots,$$

and from that

$$\|f_k\|_{S^{n-1}} \leq (c\varepsilon)^k \|f_0\|_{L_1(S^{n-1})} \text{ for all } k = 1, 2, \dots$$

Put  $\Phi = \sum_{k=0}^{\infty} \Phi_k$ . From the previous estimates it follows

$$\begin{aligned} \|\Phi\|_{L_1(\nu)} &\leq \sum_{k=0}^{\infty} \|\Phi_k\|_{L_1(\nu)} \leq \sum_{k=0}^{\infty} c \|f_k\|_{L_1(S^{n-1})} \leq \\ &c \|f_0\|_{L_1(S^{n-1})} + \sum_{k=1}^{\infty} (c\varepsilon)^k \|f_0\|_{L_1(S^{n-1})} = (c + \frac{c\varepsilon}{1 - c\varepsilon}) \|f_0\|_{L_1(S^{n-1})}. \end{aligned}$$

The constant  $(c + \frac{c\varepsilon}{1 - c\varepsilon})$  can be chosen arbitrarily close to 1.

We have proved that  $\Phi \in L_1(\nu)$  and the required relation between  $\|f\|_{L_1(S^{n-1})}$  and  $\|\Phi\|_{L_1(\nu)}$ , and we have proved as well that  $\sum_{k=1}^{\infty} |\Phi_k| \in L_1(\nu)$ .

As  $\Phi_k = 0$  on  $B(0, c_\varepsilon)$  for any  $k = 0, 1, \dots$ , the same is true for  $\Phi$  (what was to be proved) and  $\sum_{k=1}^{\infty} |\Phi_k|$ .

From these facts and the fact that  $\|K_2(x, \cdot)\|_{L_1(S^{n-1})} < 1 + \varepsilon$  whenever  $\|x\| > c_\varepsilon$  we get (using the Fubini theorem) that

$$\int_{\mathbb{R}^n} (\sum_{k=0}^{\infty} |\Phi_k(x)|) K_2(x, \cdot) d\nu(x) \in L_1(S^{n-1}).$$

From here it follows that for  $\sigma$ -almost all  $y$

$$\sum_{k=0}^{\infty} |\Phi_k(\cdot)| K_2(\cdot, y) \in L_1(\nu).$$

Using the Lebesgue Dominated Convergence Theorem with the above sum as dominating function we arrive to

$$\int_{\mathbb{R}^n} \Phi(x) K_2(x, y) d\nu(x) = \int_{\mathbb{R}^n} \left( \sum_{k=0}^{\infty} \Phi_k(x) \right) \cdot K_2(x, y) d\nu(x) =$$

$$\sum_{k=0}^{\infty} \int_{\mathbb{R}^n} \Phi_k(x) K_2(x, y) d\nu(x) = \sum_{k=0}^{\infty} (f_k(y) - f_{k+1}(y)) = f_0(y) = f(y)$$

for  $\sigma$ -almost all  $y \in S^{n-1}$ .

So

$$f = \int_{\mathbb{R}^n} \Phi(x) K_2(x, \cdot) d\nu(x)$$

and the proof is finished.  $\square$

The implication (x) $\Rightarrow$ (viii) can be proved in the same way.

### Remark

Similar problems have been recently investigated for classical harmonic functions on a ball in [3], [4], [5], [7] and for more general domains in [1], and for parabolic functions on a slab in [10] and [11]. In the present paper methods of proofs adopted in [7] and [5] turned out to be useful.

### REFERENCES

- [1] Aikawa H., *Sets of determination for harmonic functions in an NTA domains*, J. Math. Soc. Japan, to appear.
- [2] Bauer H., *Harmonische Räume und ihre Potentialtheorie*, Springer-Verlag, Berlin-Heidelberg-New York, 1966.
- [3] Bonsall F.F., *Decomposition of functions as sums of elementary functions*, Quart J. Math. Oxford (2) **37** (1986), 129–136.
- [4] Bonsall F.F., *Domination of the supremum of a bounded harmonic function by its supremum over a countable subset*, Proc. Edinburgh Math. Soc. **30** (1987), 441–477.
- [5] Bonsall F.F., *Some dual aspects of the Poisson kernel*, Proc. Edinburgh Math. Soc. **33** (1990), 207–232.
- [6] Caffarelli L.A., Littman W., *Representation formulas for solutions to  $\Delta u - u = 0$  in  $\mathbb{R}^n$* , Studies in Partial Differential Equations, Ed. W. Littman, MAA Studies in Mathematics 23, MAA (1982).
- [7] Gardiner S.J., *Sets of determination for harmonic function*, Trans. Amer. Math. Soc. **338** (1993), 233–243.
- [8] Korányi A., *A survey of harmonic functions on symmetric spaces*, Proc. Symposia Pure Math. XXV, part 1 (1979), 323–344.
- [9] Korányi A., Taylor J.C., *Fine convergence and parabolic convergence for the Helmholtz equation and the heat equation*, Illinois J. Math. **27.1** (1983), 77–93.
- [10] Ranošová J., *Sets of determination for parabolic functions on a half-space*, Comment. Math. Univ. Carolinae **35** (1994), 497–513.

- [11] Ranošová J., *Characterization of sets of determination for parabolic functions on a slab by coparabolic (minimal) thinness*, Comment. Math. Univ. Carolinae **37** (1996), 707–723.
- [12] Rudin W., *Functional analysis*, McGraw-Hill Book Company, 1973.
- [13] Taylor J.C., *An elementary proof of the theorem of Fatou-Naïm-Doob*, 1980 Seminar on Harmonic Analysis (Montreal, Que., 1980), CMS Conf. Proc., vol.1, Amer. Math. Soc., Providence, R.I (1981), 153–163.
- [14] Watson G.N., *Theory of Bessel functions*, 2nd ed., Cambridge Univ. Press, Cambridge, 1944.

MATHEMATICAL INSTITUTE, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY,  
SOKOLOVSKÁ 83, 186 00 PRAHA 8, CZECH REPUBLIC

*E-mail:* ranosova@karlin.mff.cuni.cz

(Received April 11, 1996)