

Roman Hric

Topological sequence entropy for maps of the circle

Commentationes Mathematicae Universitatis Carolinae, Vol. 41 (2000), No. 1, 53--59

Persistent URL: <http://dml.cz/dmlcz/119140>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2000

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Topological sequence entropy for maps of the circle

ROMAN HRIC

Abstract. A continuous map f of the interval is chaotic iff there is an increasing sequence of nonnegative integers T such that the topological sequence entropy of f relative to T , $h_T(f)$, is positive ([FS]). On the other hand, for any increasing sequence of nonnegative integers T there is a chaotic map f of the interval such that $h_T(f) = 0$ ([H]). We prove that the same results hold for maps of the circle. We also prove some preliminary results concerning topological sequence entropy for maps of general compact metric spaces.

Keywords: chaotic map, circle map, topological sequence entropy

Classification: Primary 26A18, 54H20, 58F13

Introduction

Let (X, ρ) be a compact metric space; denote by $C(X)$ the space of all continuous maps of this space into itself. We will pay a special attention to the case when X is the circle $\mathbb{S} = \{z \in \mathbb{C}; |z| = 1\}$; the metric on \mathbb{S} is given by $\|x, y\| = \text{dist}(\Pi^{-1}x, \Pi^{-1}y)$ where Π denotes the natural projection of the real line \mathbb{R} onto \mathbb{S} , i.e., $\Pi(x) = e^{2\pi ix}$. By \mathbb{N} we denote the set of all positive integers. If $T = (t_i)_{i=1}^\infty$ is an arbitrary sequence of nonnegative integers then the (T, f, n) -trajectory of $x \in X$ is the sequence $(f^{t_i}x)_{i=1}^n$. The set of all periodic points of f is denoted by $\text{Per}(f)$ and the set of periods of all periodic points of f by $P(f)$. A set $A \subseteq X$ is called a *retract* of X if there is a map $r : X \rightarrow A$ such that $r(a) = a$ for every $a \in A$.

Definition. Let (X, ρ) be a compact metric space. A map $f \in C(X)$ is said to be *chaotic* if there are points $x, y \in X$ such that

$$\begin{aligned} \limsup_{i \rightarrow \infty} \rho(f^i x, f^i y) &> 0, \\ \liminf_{i \rightarrow \infty} \rho(f^i x, f^i y) &= 0. \end{aligned}$$

(The set $\{x, y\}$ is called a *scrambled set*.) A map is called *nonchaotic* if it is not chaotic.

Remark. This definition of a chaotic map is equivalent to the original one by Li and Yorke in [LY] for maps of the interval (see [KuS]) and for maps of the circle (see [Ku]).

The author has been supported by the Slovak grant agency, grant number 1/4015/97.

Definition. Let (X, ρ) be a compact metric space, $f \in C(X)$ and $T = (t_i)_{i=1}^{\infty}$ be an increasing sequence of nonnegative integers. We say that a set $A \subseteq X$ (T, f, ε, n) -spans a set $B \subseteq X$ if for any $x \in B$ there is $y \in A$ such that $\rho(f^{t_i}x, f^{t_i}y) < \varepsilon$ for all $1 \leq i \leq n$. (We also say that the point y spans the point x .)

Definition ([G]). Let (X, ρ) be a compact metric space, $f \in C(X)$ and $T = (t_i)_{i=1}^{\infty}$ be an increasing sequence of nonnegative integers.

A set $A \subseteq X$ is said to be (T, f, ε, n) -separated if for any $x, y \in A$, $x \neq y$ there is an index i , $1 \leq i \leq n$, such that $\rho(f^{t_i}x, f^{t_i}y) > \varepsilon$. Let $\text{Sep}(T, f, \varepsilon, n)$ denote the largest of cardinalities of all (T, f, ε, n) -separated sets. Put

$$\text{Sep}(T, f) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Sep}(T, f, \varepsilon, n).$$

A subset of X is said to be a (T, f, ε, n) -span if it (T, f, ε, n) -spans X . Let $\text{Span}(T, f, \varepsilon, n)$ denote the smallest of cardinalities of all (T, f, ε, n) -spans. Put

$$\text{Span}(T, f) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Span}(T, f, \varepsilon, n).$$

Then $\text{Sep}(T, f) = \text{Span}(T, f)$ (see [G]) and we define the *topological sequence entropy* of f relative to T , $h_T(f)$, to be $\text{Sep}(T, f)$.

Remark. If $t_i = i - 1$, $i = 1, 2, \dots$ then $h_T(f)$ is the topological entropy $h(f)$ of f . Topological sequence entropy can be viewed as the topological entropy of the nonautonomous dynamical system given on the space X by the sequence of maps $f^{t_1}, f^{t_2-t_1}, f^{t_3-t_2}, \dots$ (see [KS]).

In [FS] Franzová and Smítal proved that a continuous map f of the interval is chaotic if and only if there is an increasing sequence of nonnegative integers T such that $h_T(f) > 0$. A natural question arose whether there is some universal sequence which characterizes chaos. This is not the case as it was proved in [H] — for any increasing sequence of nonnegative integers T there is a chaotic map f with $h_T(f) = 0$. The main aim of this paper is to prove the same results for maps of the circle.

Theorem 1. *A map $f \in C(\mathbb{S})$ is chaotic if and only if there is an increasing sequence of nonnegative integers T such that $h_T(f) > 0$.*

Remark. Theorem 1 does not hold in general, even for triangular maps of the square. There is a nonchaotic triangular map with positive topological sequence entropy relative to a suitable sequence (see [FPS, Theorem 2]) and, on the other hand, there is a chaotic triangular map with zero topological sequence entropy relative to any sequence (see [FPS, Theorem 3]).

Theorem 2. *Let X be a compact metric space containing a homeomorphic image of an interval and let T be an increasing sequence of nonnegative integers. Then there is a chaotic map $f \in C(X)$ such that $h_T(f) = 0$.*

Remark. The analysis of the proof of Theorem 6 in [H] shows that Theorem 2 holds also when X is a Cantor set.

Corollary 3. *Let T be an increasing sequence of nonnegative integers. Then there is a chaotic map $f \in C(\mathbb{S})$ such that $h_T(f) = 0$.*

Preliminary results

Let (X, ρ) and (Y, σ) be compact metric spaces, $f \in C(X)$, $g \in C(Y)$, and let $\pi : X \rightarrow Y$ be a continuous map such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & X \\ \pi \downarrow & & \downarrow \pi \\ Y & \xrightarrow{g} & Y \end{array}$$

commutes. In this situation we have the following

Lemma 4. *Let T be an increasing sequence of nonnegative integers. Then*

- (i) *if π is injective then $h_T(f) \leq h_T(g)$;*
- (ii) *if π is surjective then $h_T(f) \geq h_T(g)$;*
- (iii) *if π is bijective then $h_T(f) = h_T(g)$.*

PROOF:

(ii) and (iii). See [G, p. 332].

(i). We have that π is a homeomorphism between X and πX . Thus, by (iii), $h_T(f) = h_T(g|_{\pi X})$. Now let $E \subseteq \pi X$ be $(T, g|_{\pi X}, \varepsilon, n)$ -separated. Trivially, it is also (T, g, ε, n) -separated which gives $h_T(g|_{\pi X}) \leq h_T(g)$. \square

It is known that some of the properties of topological entropy are not satisfied by topological sequence entropy. For example, contrary to the formula $h(f^k) = k \cdot h(f)$, an analogous formula for topological sequence entropy does not hold — it is even possible that $h_S(f) < h_T(f)$ for a subsequence S of T ([L]). In this case the following result can be useful.

Theorem 5. *Let (X, ρ) be a compact metric space, $f \in C(X)$, T be an increasing sequence of nonnegative integers and k be a positive integer. Then there is an increasing sequence of nonnegative integers S such that $h_S(f^k) \geq h_T(f)$.*

PROOF: Since X is compact, f, f^2, \dots, f^{k-1} are equicontinuous, i.e., for any $\varepsilon > 0$ there is $\delta = \delta(\varepsilon) > 0$ such that if $x, y \in X$ and $\rho(x, y) \leq \delta$ then $\rho(f^i x, f^i y) < \varepsilon$ for $i = 1, \dots, k-1$. We may assume that $\delta \leq \varepsilon$.

Let $T = (t_i)_{i=1}^\infty$. Define $S = (s_i)_{i=1}^\infty$ as follows. Put $s_1 = [\frac{t_1}{k}]$ (where $[\cdot]$ stands for the integer part) and for any m let s_{m+1} will be the first $[\frac{t_i}{k}]$ greater than s_m .

Let $E \subseteq X$ be an (T, f, ε, n) -separated set. We are going to show that E is a (S, f^k, δ, m) -separated set where m is such that $s_m = [\frac{t_n}{k}]$. To this end let $x, y \in E$, $x \neq y$. Then for some $i \in \{1, 2, \dots, n\}$, $\rho(f^{t_i} x, f^{t_i} y) > \varepsilon$. Take j with $s_j = [\frac{t_i}{k}]$. Then $j \leq m$ and from the definition of δ we have $\rho(f^{k \cdot s_j} x, f^{k \cdot s_j} y) > \delta$. Thus E is an (S, f^k, δ, m) -separated set. From this we have $\text{Sep}(T, f, \varepsilon, n) \leq$

$\text{Sep}(S, f^k, \delta, m)$. Now, $h_T(f) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Sep}(T, f, \varepsilon, n) \leq$
 $\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Sep}(S, f^k, \delta, m) \leq \lim_{\delta \rightarrow 0} \limsup_{m \rightarrow \infty} \frac{1}{m} \log \text{Sep}(S, f^k, \delta, m) = h_S(f^k)$. \square

Corollary 6. *Let X be a compact metric space, $f \in C(X)$ and k be a positive integer. Then the following two conditions are equivalent:*

- (i) *there is an increasing sequence T of nonnegative integers such that $h_T(f) > 0$;*
- (ii) *there is an increasing sequence T of nonnegative integers such that $h_T(f^k) > 0$.*

In the sequel we will discuss the space of maps of the circle. The space $C(\mathbb{S})$ can be decomposed into the following classes (see [ALM, Chapter 3], cf. also [Ku, p. 384]):

$$\begin{aligned} C_1 &= \{f \in C(\mathbb{S}); f \text{ has no periodic point}\}; \\ C_2 &= \{f \in C(\mathbb{S}); P(f^n) = \{1\} \text{ or } P(f^n) = \{1, 2, 2^2, \dots\} \text{ for some } n \in \mathbb{N}\}; \\ C_3 &= \{f \in C(\mathbb{S}); P(f^n) = \mathbb{N} \text{ for some } n \in \mathbb{N}\}. \end{aligned}$$

According to this we will distinguish three different cases.

Maps without periodic points

In all of this section we assume $f \in C(\mathbb{S})$ to have no periodic point. We are going to show that Theorem 1 holds for such maps. Since, by [Ku, Theorem B], f is not chaotic, we need only to show that $h_T(f) = 0$ for any increasing sequence T . So fix T . If f is a homeomorphism then $h_T(f) = 0$ by [KS, Theorem D]. Otherwise, by [AK, Theorem 1 and Theorem 2], there is a nowhere dense perfect set E which is the only ω -limit set of f , all (closed) contiguous intervals are wandering, the preimage of any contiguous interval is a contiguous interval, the image of any contiguous interval is either a contiguous interval or a point from E . Moreover, $f|_E$ is monotone. By linear extension of $f|_E$ we obtain a monotone map $g \in C(\mathbb{S})$. By [KS, Theorem D], $h_T(g) = 0$. By Lemma 4(i), $h_T(f|_E) \leq h_T(g)$. Hence, $\limsup_{n \rightarrow \infty} \frac{1}{n} \log \text{Span}(T, f|_E, \varepsilon, n) = 0$ for any $\varepsilon > 0$.

Now fix an arbitrary $\varepsilon > 0$. We are going to estimate $\text{Span}(T, f, \varepsilon, n)$. Let I_1, \dots, I_k be all contiguous intervals longer than $\frac{\varepsilon}{2}$. Let A be a $(T, f|_E, \frac{\varepsilon}{2}, n)$ -span. Take any point x whose (T, f, n) -trajectory lies in $\mathbb{S} \setminus \bigcup_{i=1}^k I_i$. If $x \in E$ then x is (T, f, ε, n) -spanned by A . For $x \notin E$ put y to be an endpoint of the contiguous interval which contains x . Then $\|f^{ti}x, f^{ti}y\| \leq \frac{\varepsilon}{2}$ for all $1 \leq i \leq n$. Since $y \in E$ is $(T, f, \frac{\varepsilon}{2}, n)$ -spanned by a point $z \in A$, the set A obviously (T, f, ε, n) -spans all such points x .

So it remains to consider those points whose (T, f, n) -trajectories meet $\bigcup_{i=1}^k I_i$. Fix $N \in \mathbb{N}$ such that $N > \frac{1}{\varepsilon}$. We are going to show that there is a set of cardinality at most $n \cdot k \cdot N^k$ which (T, f, ε, n) -spans all considered points. It is sufficient to show that there is a set with cardinality at most N^k which (T, f, ε, n) -spans

the set $I(t_i, I_j) = \{x \in \mathbb{S}; f^{t_i}x \in I_j\}$ (for fixed $1 \leq i \leq n$ and $1 \leq j \leq k$). First, it is obvious that $I(t_i, I_j)$ is a contiguous interval. Consider its (T, f, n) -trajectory $(f^{t_1}I(t_i, I_j), \dots, f^{t_n}I(t_i, I_j))$. Each element in this trajectory is either a contiguous interval or a point from E . At most k of them have lengths greater than or equal to ε — cut each of such elements to N segments shorter than ε . All the other elements of the trajectory will be considered to be segments themselves. To each $x \in I(t_i, I_j)$ assign its code — the sequence $(S_1(x), \dots, S_n(x))$ where $S_l(x)$ is the segment containing $f^{t_l}x$. We have at most N^k different codes and all points with the same code can be (T, f, ε, n) -spanned by one point.

From what has been said above we see that $\text{Span}(T, f, \varepsilon, n) \leq \text{Span}(T, f|_E, \frac{\varepsilon}{2}, n) + n \cdot k \cdot N^k$ which finishes the proof of Theorem 1 for maps without periodic points.

Maps with periodic points

We will first deal with the case C_2 . We know that for any $n \in \mathbb{N}$ f is chaotic if and only if f^n is chaotic. Taking into account Corollary 6 we can without loss of generality assume that $P(f) = \{1\}$ or $P(f) = \{1, 2, 2^2, \dots\}$. Since f has a fixed point, by [Ku, Lemma 2.5] there is a lifting F and an F -invariant compact interval J longer than 1. In the following discussion of the case C_2 we will write F and Π instead of $F|_J$ and $\Pi|_J$, respectively, as in the next commutative diagram

$$\begin{array}{ccc} J & \xrightarrow{F} & J \\ \Pi \downarrow & & \downarrow \Pi \\ \mathbb{S} & \xrightarrow{f} & \mathbb{S} \end{array}$$

Note that if $x, y \in J$ then $\|\Pi x, \Pi y\| \leq |x - y|$ with the equality whenever $|x - y| \leq \frac{1}{2}$.

Lemma 7. *F is chaotic if and only if f is chaotic.*

PROOF: Let F be chaotic. Then there are two points $u, v \in J$ such that

$$(1) \quad \limsup_{i \rightarrow \infty} |F^i u - F^i v| = \gamma > 0;$$

$$(2) \quad \liminf_{i \rightarrow \infty} |F^i u - F^i v| = 0.$$

We claim that $\Pi u, \Pi v$ form a scrambled set for f . From (2), $\liminf_{i \rightarrow \infty} \|f^i \Pi u, f^i \Pi v\| = 0$. Now put $\eta = \min\{\gamma, \frac{1}{2}\}$. Take $0 < \delta < \eta$ such that $|x - y| < \delta$ implies $|Fx - Fy| < \eta$. From this and (1) and (2) we have that $|F^i u - F^i v| \in [\delta, \eta]$ infinitely many times. Since $\eta \leq \frac{1}{2}$, the same holds for $\|f^i \Pi u, f^i \Pi v\|$. Thus $\limsup_{i \rightarrow \infty} \|f^i \Pi u, f^i \Pi v\| \geq \delta > 0$.

Let F be nonchaotic. Then by [JS, Theorem 3] every trajectory of F is approximable by cycles, i.e. for any $\varepsilon > 0$ and any $x \in J$ there is some periodic point $p \in \text{Per}(F)$ such that

$$(3) \quad |F^i x - F^i p| < \varepsilon \quad \text{for all } i = 0, 1, 2, \dots$$

Fix any $z \in \mathbb{S}$. Take any of its preimages $x \in \Pi^{-1}z$. Let $\varepsilon > 0$ be arbitrary, $p \in \text{Per}(F)$ such that (3) is satisfied. Clearly, Πp is a periodic point for f and $\|f^i z, f^i \Pi p\| < \varepsilon$ for all $i = 0, 1, 2, \dots$. Thus f is not chaotic by [Ku, Theorem A]. \square

Lemma 8. *Let F be chaotic. Then there is an increasing sequence T such that $h_T(f) > 0$.*

PROOF: If F has a periodic point of period $k \cdot 2^m$ where $k > 1$ is odd then, by Sharkovsky theorem, it has also a periodic point of period $k' \cdot 2^m$ where $k' > \text{diam } J + 1$ is odd. Since $\Pi|_J$ is at most $[\text{diam } J] + 1$ to one, f has a periodic point of period $k'' \cdot 2^{m'}$ where $k'' > 1$ is odd. This is a contradiction since $P(f)$ is $\{1\}$ or $\{1, 2, 2^2, \dots\}$. So F is of type 2^∞ , chaotic. By [S] there is an orbit of periodic intervals of period $p > \text{diam } J$ such that F^p is chaotic on each of them. At least one interval K in this orbit is shorter than 1. Then $\Pi|_K$ is injective and so $F^p|_K$ is topologically conjugate with $f^p|_{\Pi K}$. By [FS, Theorem] there is an increasing sequence of nonnegative integers S such that $h_S(F^p|_K) > 0$. Since $h_{p \cdot S}(f) = h_S(f^p)$ it is sufficient to use Lemma 4(iii) and (i) to get $h_{p \cdot S}(f) \geq h_S(f^p|_{\pi K}) = h_S(F^p|_K) > 0$. \square

We are going to show that Theorem 1 holds for maps from the class C_2 . Let $f \in C_2$ be chaotic. Then we obtain the required result using Lemma 7 and Lemma 8.

Now let $f \in C_2$ and let there be an increasing sequence of nonnegative integers T such that $h_T(f) > 0$. Theorem 4(ii) then implies that $h_T(F) > 0$ where F has the same meaning as above. By [FS, Theorem] F is chaotic. Lemma 7 finishes the proof.

Finally we will discuss the situation for maps from the remaining class C_3 . So let $P(f^n) = \mathbb{N}$ for some n . By [BC, Theorem IX.28(i) and (ii)] we have that $h(f^n)$ is positive and so is $h(f)$. By the same theorem, conditions (ii) and (iii), we have that $f^{m \cdot n}$ is strictly turbulent for a suitable $m \in \mathbb{N}$ which implies that f is chaotic for the same reason as in the interval case. This finishes the proof of Theorem 1.

Proof of Theorem 2

The space X contains a homeomorphic image J of the interval $[0, 1]$. The set J is a retract of X by [HY, Theorem 2-34]. Let $r : X \rightarrow J$ be a corresponding retraction. By [H, Theorem 6] there is a chaotic onto map $g \in C([0, 1])$ such that $h_T(g) = 0$. Let $\tilde{g} \in C(J)$ be a map topologically conjugate with g . Define $f \in C(X)$ by $f = \tilde{g} \circ r$. Since $\bigcap_{i=0}^{\infty} f^i X = fX = J$, we have that $h_T(f) = h_T(f|_J) = 0$ by [BCJ, Proposition 1].

Acknowledgment. The author thanks Professor Lubomír Snoha for contributing ideas useful for this paper and for his helpful comments during the last stage of its preparation.

REFERENCES

- [ALM] Alsedà L., Llibre J., Misiurewicz M., *Combinatorial Dynamics and Entropy in Dimension One*, World Scientific Publ., Singapore, 1993.
- [AK] Auslander J., Katznelson Y., *Continuous maps of the circle without periodic points*, Israel. J. Math. **32** (1979), 375–381.
- [BCJ] Balibrea F., Cánovas J.S., Jiménez López V., *Commutativity and noncommutativity of topological sequence entropy*, preprint.
- [BC] Block L.S., Coppel W.A., *Dynamics in One Dimension*, Lecture Notes in Math., vol. 1513, Springer, Berlin, 1992.
- [FS] Franzová N., Smítal J., *Positive sequence entropy characterizes chaotic maps*, Proc. Amer. Math. Soc. **112** (1991), 1083–1086.
- [G] Goodman T.N.T., *Topological sequence entropy*, Proc. London Math. Soc. **29** (1974), 331–350.
- [HY] Hocking J.G., Young G.S., *Topology*, Dover, New York, 1988.
- [H] Hric R., *Topological sequence entropy for maps of the interval*, Proc. Amer. Math. Soc. **127** (1999), 2045–2052.
- [JS] Janková K., Smítal J., *A characterization of chaos*, Bull. Austral. Math. Soc. **34** (1986), 283–292.
- [KS] Kolyada S., Snoha L., *Topological entropy of nonautonomous dynamical systems*, Random and Comp. Dynamics **4** (1996), 205–233.
- [Ku] Kuchta M., *Characterization of chaos for continuous maps of the circle*, Comment. Math. Univ. Carolinae **31** (1990), 383–390.
- [KuS] Kuchta M., Smítal J., *Two point scrambled set implies chaos*, European Conference on Iteration Theory ECIT'87, World Sci. Publishing Co., Singapore.
- [L] Lemańczyk M., *The sequence entropy for Morse shifts and some counterexamples*, Studia Math. **52** (1985), 221–241.
- [LY] Li T Y., Yorke J.A., *Period three implies chaos*, Amer. Math. Monthly **82** (1975), 985–992.
- [S] Smítal J., *Chaotic functions with zero topological entropy*, Trans. Amer. Math. Soc. **297** (1986), 269–281.

DEPARTMENT OF MATHEMATICS, FACULTY OF NATURAL SCIENCES, MATEJ BEL UNIVERSITY,
TAJOVSKÉHO 40, SK-974 01 BANSKÁ BYSTRICA, SLOVAK REPUBLIC

E-mail: hric@fpv.umb.sk

(Received March 31, 1999)