

Juan Carlos Ferrando

On the convergence of certain sums of independent random elements

Commentationes Mathematicae Universitatis Carolinae, Vol. 43 (2002), No. 1, 77--81

Persistent URL: <http://dml.cz/dmlcz/119301>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On the convergence of certain sums of independent random elements

J.C. FERRANDO

Abstract. In this note we investigate the relationship between the convergence of the sequence $\{S_n\}$ of sums of independent random elements of the form $S_n = \sum_{i=1}^n \varepsilon_i x_i$ (where ε_i takes the values ± 1 with the same probability and x_i belongs to a real Banach space X for each $i \in \mathbb{N}$) and the existence of certain weakly unconditionally Cauchy subseries of $\sum_{n=1}^{\infty} x_n$.

Keywords: independent random elements, copy of c_0 , Pettis integrable function, perfect measure space

Classification: 46B15, 46B09

1. Preliminaries

Our notation is standard ([1], [3], [4], [9]). Throughout this note Δ will denote the Cantor space $\{-1, 1\}^{\mathbb{N}}$, Σ the σ -algebra of subsets of Δ generated by the n -cylinders of Δ for each $n \in \mathbb{N}$, and ν the Borel probability $\otimes_{i=1}^{\infty} \nu_i$ on Σ , where $\nu_i : 2^{\{-1, 1\}} \rightarrow [0, 1]$ is defined by $\nu_i(\emptyset) = 0$, $\nu_i(\{-1\}) = \nu_i(\{1\}) = 1/2$ and $\nu_i(\{-1, 1\}) = 1$ for each $i \in \mathbb{N}$. In what follows X will be a real Banach space and $L_0(\nu, X)$ will stand for the (F) -space over \mathbb{R} of all [classes of] ν -measurable X -valued functions equipped with the (F) -norm

$$\|f\|_0 = \int_{\Delta} \frac{\|f(\varepsilon)\|}{1 + \|f(\varepsilon)\|} d\nu(\varepsilon)$$

of the convergence in probability. We shall represent by $P_1(\nu, X)$ the (real) normed space consisting of all those [classes of] ν -measurable X -valued Pettis integrable functions f defined on Δ provided with the semivariation norm

$$\|f\|_{P_1(\nu, X)} = \sup \left\{ \int_{\Delta} |x^* f(\omega)| d\nu(\omega) : x^* \in X^*, \|x^*\| \leq 1 \right\}.$$

As it is well known, $P_1(\nu, X)$ is not a Banach space whenever X is infinite-dimensional. In the sequel we shall shorten by wuC the sentence ‘weakly unconditionally Cauchy’.

Supported by DGESIC PB97-0342 and Presidencia de la Generalitat Valenciana.

In [5] we have shown that if a series of independent random elements of the form $\sum_{n=1}^{\infty} f_n$, with $f_n(\omega) = \omega_n x_n$ for $\omega \in \Delta$ and $\{x_n\} \subseteq X$, converges ν -almost surely in X , then $\sum_{n=1}^{\infty} x_n$ has a subseries which is unconditionally convergent in norm. In this note we continue the investigation on the relationship among the convergence of the functional series $\sum_{n=1}^{\infty} f_n$ under different topologies and the existence of certain wuC subseries of $\sum_{n=1}^{\infty} x_n$.

2. On certain weakly unconditionally Cauchy subseries

Lemma 2.1. *If there are a closed set A in Δ with $\nu(A) > 1/2$ and a nonempty set $S \subseteq X^*$ such that $\sum_{i=1}^{\infty} x^* f_i(\omega)$ converges for $\omega \in A$ and $x^* \in S$, then there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $\sum_{i=1}^{\infty} |x^* x_{n_i}| < \infty$ for each $x^* \in S$.*

PROOF: The following fact is contained in the proof of [8, Proposition] (see also [5, Claim]). We shall denote by $C_{i_1 i_2 \dots i_k}$ or $C_{i_1 i_2 \dots i_k}(\varepsilon)$ any rectangle of Δ with fixed coordinates i_1, i_2, \dots, i_k , i.e., $C_{i_1 i_2 \dots i_k}(\varepsilon) = \{\omega \in \Delta : \omega_{i_j} = \varepsilon_j, 1 \leq j \leq k\}$ for some $\varepsilon \in \Delta$. On the other hand, given a strictly increasing sequence $Q = \{n_i : i \in \mathbb{N}\}$ of positive integers, for each $\omega \in \Delta$ we shall design by ω' (as in [8]) the element of Δ defined by $\omega'_i = \omega_i$ if $i \in Q$ and $\omega'_i = -\omega_i$ if $i \notin Q$.

Fact. *Let $A \in \Sigma$. If $\nu(A) > 1/2$, there is a strictly increasing sequence $\{n_i\}$ of positive integers such that $A \cap A' \cap C_{n_1 n_2 \dots n_k} \neq \emptyset$ for each $C_{n_1 n_2 \dots n_k}$ and each $k \in \mathbb{N}$.*

By hypothesis there is a closed set A in Δ with $\nu(A) > 1/2$ such that $\sum_{n=1}^{\infty} \omega_n x^* x_n$ converges for $\omega \in A$ and $x^* \in S$. According to the preceding fact there exists a strictly increasing sequence $Q = \{n_i\}$ of positive integers such that, given $\varepsilon \in \Delta$, then $A \cap A' \cap C_{n_1 n_2 \dots n_k}(\varepsilon) \neq \emptyset$ for each $k \in \mathbb{N}$. Since $\{A \cap A' \cap C_{n_1 n_2 \dots n_k}(\varepsilon) : k \in \mathbb{N}\}$ is a decreasing sequence of nonempty closed sets in the compact space Δ , there is a point ζ (which depends of ε) in Δ which belongs to the intersection $\bigcap_{k=1}^{\infty} A \cap A' \cap C_{n_1 n_2 \dots n_k}(\varepsilon)$. Hence, for each $x^* \in S$ and each pair (r, s) of positive integers, with $s > r$, one has

$$\left| \sum_{i=r+1}^s \varepsilon_i x^* x_{n_i} \right| = \left| \sum_{i=r+1}^s \zeta_{n_i} x^* x_{n_i} \right| \leq \frac{1}{2} \left(\left| \sum_{i=n_r+1}^{n_s} x^* f_i(\zeta) \right| + \left| \sum_{i=n_r+1}^{n_s} x^* f_i(\zeta') \right| \right).$$

Since $\zeta, \zeta' \in A$ and $x^* \in S$, both series $\sum_{i=1}^{\infty} x^* f_i(\zeta)$ and $\sum_{i=1}^{\infty} x^* f_i(\zeta')$ are convergent. So, for a given $\epsilon > 0$ there is a $k \in \mathbb{N}$ such that $\left| \sum_{i=n_r+1}^{n_s} x^* f_i(\zeta) \right| < \epsilon$ and $\left| \sum_{i=n_r+1}^{n_s} x^* f_i(\zeta') \right| < \epsilon$ for $s > r \geq k$, which implies that $\left| \sum_{i=r+1}^s \varepsilon_i x^* x_{n_i} \right| \leq \epsilon$ for $s > r \geq k$. Hence the numerical series $\sum_{i=1}^{\infty} \varepsilon_i x^* x_{n_i}$ converges. Given that this is true for each $\varepsilon \in \Delta$, it follows that $\sum_{i=1}^{\infty} |x^* x_{n_i}| < \infty$ for each $x^* \in S$ and we are done. \square

Theorem 2.2. *Assume that $\|x_n\| = 1$ for each $n \in \mathbb{N}$ and X has a dual unit ball with countably many extreme points. If*

$$\sup_{n \in \mathbb{N}} \int_{\Delta} |x^* S_n(\omega)| \, d\nu(\omega) < \infty$$

for each $x^* \in \text{Ext } B_{X^*}$, then X contains a copy of c_0 .

PROOF: By hypothesis, for each $x^* \in \text{Ext } B_{X^*}$ there exists $C_{x^*} > 0$ such that

$$(2.1) \quad \sup_{n \in \mathbb{N}} \int_{\Delta} \left| \sum_{i=1}^n x^* f_i(\omega) \right| \, d\nu(\omega) < C_{x^*}.$$

Hence, given $x^* \in \text{Ext } B_{X^*}$, as a consequence of (2.1) and of Khinchine's inequalities there exists a $K > 0$ such that

$$(2.2) \quad \left\{ \sum_{i=1}^n \sigma^2(x^* f_i) \right\}^{1/2} = \left\{ \sum_{i=1}^n (x^* x_i)^2 \right\}^{1/2} \\ \leq K \int_{\Delta} \left| \sum_{i=1}^n x^* f_i(\omega) \right| \, d\nu(\omega) < KC_{x^*}$$

for each $n \in \mathbb{N}$. Considering that the sequence $\{x^* f_i\}$ consists of independent random variables such that

$$\mathbf{E}(x^* f_i) = \int_{\Delta} x^* f_i(\omega) \, d\nu(\omega) = 0$$

for each $i \in \mathbb{N}$, according to [7, Section 46, Theorem B] equation (2.2) ensures that $\sum_{i=1}^{\infty} x^* f_i(\omega)$ converges almost surely for $\omega \in \Delta$. Since $\text{Ext } B_{X^*}$ is countable, it follows that there exists a ν -null set N such that $\sum_{i=1}^{\infty} x^* f_i(\omega)$ converges for each $\omega \in \Delta - N$ and each $x^* \in \text{Ext } B_{X^*}$. So, using inner regularity we may choose a closed set A with $A \subseteq \Delta - N$ and $\nu(A) > 1/2$ such that $\sum_{i=1}^{\infty} x^* f_i(\omega)$ converges for each $\omega \in A$ and each $x^* \in \text{Ext } B_{X^*}$. On the basis of Lemma 2.1, this implies that there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $\sum_{i=1}^{\infty} |x^* x_{n_i}| < \infty$ for each $x^* \in \text{Ext } B_{X^*}$. Since $\sum_{n=1}^{\infty} x_n$ diverges, Elton's theorem guarantees that X contains a copy of c_0 . \square

Proposition 2.3. *If the sums $\{S_n\}$ are bounded inside of a complete linear subspace L of $P_1(\nu, X)$, then $\sum_{n=1}^{\infty} x_n$ has a wuC subseries.*

PROOF: Since $\{S_n\}$ is bounded inside of a complete linear subspace L of $P_1(\nu, X)$ and given that the canonical inclusion map from $P_1(\nu, X)$ into $L_0(\nu, X)$ has closed graph ([6, Lemma 4]), then Banach-Schauder's theorem guarantees that $\{S_n\}$ is stochastically bounded. So, according to [9, Section 5.2.3, Theorem 2.2] the sums $\{S_n\}$ are bounded almost surely, i.e. $\nu(\{\omega \in \Delta : \sup_{n \in \mathbb{N}} \|\sum_{i=1}^n f_i(\omega)\| = \infty\}) = 0$. Hence Kwapien's theorem [8, Proposition] assures the existence of a wuC subseries of $\sum_{n=1}^{\infty} x_n$. \square

Corollary 2.4. *Assume that $\{f_n\}$ is a basic sequence in $\widehat{P_1(\nu, X)}$ equivalent to the unit vector basis of c_0 . If $[f_n]$ is contained in $P_1(\nu, X)$, then there exists a subsequence $\{f_{n_i}\}$ such that $[f_{n_i}]$ is isomorphic to a complemented copy of c_0 .*

PROOF: Since the series $\sum_{i=1}^{\infty} f_i$ is wuC in $P_1(\nu, X)$, there is $K > 0$ such that

$$\sup_{n \in \mathbb{N}} \left\| \sum_{i=1}^n \xi_i f_i \right\|_{P_1(\nu, X)} < K \|\xi\|_{\infty}$$

for each $\xi \in \ell_{\infty}$. Hence the sums $\{S_n\}$ are bounded in the complete linear subspace $[f_i]$ of $P_1(\nu, X)$ and Proposition 2.3 guarantees that $\sum_{n=1}^{\infty} x_n$ has a wuC subseries. Since $\|x_n\| = \|f_n\|_{P_1(\nu, X)}$ for each $n \in \mathbb{N}$, then $\inf_{n \in \mathbb{N}} \|x_n\| > 0$ and the classic Bessaga-Pelczyński allows us to conclude that $\{x_n\}$ contains a subsequence $\{x_{n_i}\}$ equivalent to the unit vector basis of c_0 . Therefore, there exists a bounded sequence $\{y_i^*\}$ in X^* such that $y_i^* x_{n_j} = \delta_{ij}$ for each $i, j \in \mathbb{N}$. Assuming without loss of generality that $y_i \in B_{X^*}$, set $g_i(\varepsilon) = \varepsilon_i y_i^*$ for each $i \in \mathbb{N}$ and define

$$\langle g_i, f \rangle = \int_{\Delta} \varepsilon_i y_i^* f(\varepsilon) d\nu(\varepsilon)$$

for each $f \in P_1(\nu, X)$. So we have $\langle g_i, f_{n_j} \rangle = \delta_{ij}$ for each $i, j \in \mathbb{N}$. On the other hand, denoting by C_n the rectangle of Δ formed by all those $\varepsilon \in \Delta$ with $\varepsilon_n = 1$ and noting that $\nu(E \cap C_n) \rightarrow \nu(E)/2$ for all $E \in \Sigma$, it follows that

$$\mathbf{E}_{C_n}(\varphi) = \frac{1}{\nu(C_n)} \int_{C_n} \varphi d\nu \rightarrow \int_{\Delta} \varphi d\nu = \mathbf{E}(\varphi)$$

for each ν -simple function $\varphi : \Delta \rightarrow \mathbb{R}$. This implies that $\mathbf{E}_{C_n}(\varphi) \rightarrow \mathbf{E}(\varphi)$ for each $\varphi \in L_1(\nu)$, which leads to $\int_{\Delta} \varepsilon_i \varphi(\varepsilon) d\nu \rightarrow 0$ for each $\varphi \in L_1(\nu)$. Since, in addition, (Δ, Σ, ν) is a perfect measure space, it can be shown as in [2] that $\langle g_i, f \rangle \rightarrow 0$ for each $f \in P_1(\nu, X)$. Consequently the map $P : P_1(\nu, X) \rightarrow P_1(\nu, X)$ defined by

$$Pf = \sum_{i=1}^{\infty} \langle g_i, f \rangle f_{n_i}$$

is a bounded linear projection operator from the barreled space $P_1(\nu, X)$ onto $[f_{n_i}]$. \square

Proposition 2.5. *If there exists a complete linear subspace L in $P_1(\nu, X)$ such that $\{f_i\} \subseteq L$ and $\sum_{i=1}^{\infty} f_i$ converges in $P_1(\nu, X)$ to some separably-valued $f \in L$, then there exists a subseries of $\sum_{i=1}^{\infty} x_i$ which is unconditionally convergent in X .*

PROOF: Given that $\sum_{i=1}^{\infty} f_i = f$ in $P_1(\nu, X)$ and L is complete, then $\sum_{i=1}^{\infty} f_i = f$ in L . Then, using the fact that the inclusion map from $P_1(\nu, X)$ into $L_0(\nu, X)$ has closed graph together with the Banach-Schauder theorem, we get that $\sum_{i=1}^{\infty} f_i = f$ in probability. Since the range of f is separable in norm, then [9, Section 5.2.3, Theorem 2.1] guarantees that the series $\sum_{i=1}^{\infty} f_i(\omega)$ converges in X to $f(\omega)$ almost surely for $\omega \in \Delta$. Hence [5, Theorem 2.1] establishes the existence of a subseries of $\sum_{n=1}^{\infty} x_n$ which is unconditionally convergent in X . \square

Question. *We do not know whether the statement of Theorem 2.2 is true without the assumption that B_{X^*} has countable many extreme points.*

REFERENCES

- [1] Cembranos P., Mendoza J., *Banach Spaces of Vector-Valued Functions*, LNM **1676**, Springer, 1997.
- [2] Díaz S., Fernández A., Florencio M., Paúl P.J., *Complemented copies of c_0 in the space of Pettis integrable functions*, Quaestiones Math. **16** (1993), 61–66.
- [3] Diestel J., *Sequences and series in Banach spaces*, GTM **92**, Springer-Verlag, New York-Berlin-Heidelberg-Tokyo, 1984.
- [4] Diestel J., Uhl J., *Vector measures*, Math Surveys **15**, Amer. Math. Soc., Providence, 1977.
- [5] Ferrando J.C., *On a theorem of Kwapien*, Quaestiones Math. **24** (2001), 51–54.
- [6] Freniche F.J., *Embedding c_0 in the space of Pettis integrable functions*, Quaestiones Math. **21** (1998), 261–267.
- [7] Halmos P.R., *Measure Theory*, GTM **18**, Springer, New York-Berlin-Heidelberg-Barcelona, 1950.
- [8] Kwapien S., *On Banach spaces containing c_0* , Studia Math. **52** (1974), 187–188.
- [9] Vakhania N.N., Tarieladze V.I., Chobanian S.A., *Probability Distributions on Banach Spaces*, D. Reidel Publishing Company, Dordrecht, 1987.

CENTRO DE INVESTIGACIÓN OPERATIVA, UNIVERSIDAD MIGUEL HERNÁNDEZ, E-03202 ELCHE (ALICANTE), SPAIN

E-mail: jc.ferrando@umh.es

(Received June 27, 2001)