

Giovanni Lo Faro; Giorgio Nardo; Jack R. Porter
On minimal- α -spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 44 (2003), No. 4, 727--740

Persistent URL: <http://dml.cz/dmlcz/119427>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2003

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On minimal- α -spaces

G. LO FARO, G. NORDO*, J.R. PORTER

Abstract. An α -space is a topological space in which the topology is generated by the family of all α -sets (see [N]). In this paper, minimal- $\alpha\mathcal{P}$ -spaces (where \mathcal{P} denotes several separation axioms) are investigated. Some new characterizations of α -spaces are also obtained.

Keywords: α -space, αT_i -space, minimal- αT_i space, T_2 -closed space, minimal- T_2 space, ψ -space

Classification: Primary 54A10, 54D25; Secondary 54A05, 54D80

1. Introduction

The family of all topologies on a set X is a complete atomic lattice. There has been a considerable amount of interest in topologies which are minimal or maximal in this lattice with respect to certain topological properties.

Given a topological property \mathcal{P} , we say that a topology on a set X is \mathcal{P} -*minimal* if every weaker topology on X does not possess property \mathcal{P} .

Throughout this paper, the word “space” will mean topological space, the topology on a space X is denoted by $\tau(X)$, int_τ and cl_τ (or int_X and cl_X when no confusion is possible about the topology on X) will denote respectively the interior and the closure operators with respect to $\tau(X)$ and if σ is a topology on the underlying set of X , then σ is called an *expansion* (respectively a *compression*) of $\tau(X)$ if $\tau(X) \subseteq \sigma$ (resp. $\sigma \subseteq \tau(X)$).

A subset R of a space X is called *regular open* if $\text{int}_\tau(\text{cl}_\tau(R)) = R$. The family of all regular open sets of X is denoted by $RO(X)$ and forms a base for a topology $\tau_s(X)$ on X which is a compression of $\tau(X)$ and it is called the *semiregularization* of X . We say that X is *semiregular* if $\tau_s(X) = \tau(X)$.

The notion of an α -set was introduced in 1965 by Njåstad [N]. Given a space X , we say that $A \subseteq X$ is an α -*set* if $A \subseteq \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A)))$. It is easy to prove that the family $\alpha\tau(X)$ of all α -sets of X is a topology on X which will be called the α -*topology* induced by $\tau(X)$ and it consists of all the subsets A of X such that there exists some open set $U \in \tau(X)$ such that $U \subseteq A \subseteq \text{int}_\tau(\text{cl}_\tau(U))$. The members of $\alpha\tau(X)$ will be called the α -*open* sets of X while their complements

* This research was supported by a grant of cofin. 2001 “Strutture Geometriche, Combinatorie e loro Applicazioni” and from the C.N.R. through I.N.D.A.M. (Italy).

will be called the α -closed sets of X . Evidently, every open set is an α -set and hence the α -topology is an expansion of $\tau(X)$. We say that X is an α -space if $\tau(X) = \tau_\alpha(X)$.

Obviously, the partial ordered set of all α -topologies contain both a maximum (the discrete topology) and a minimum (the trivial topology) element. If \mathcal{P} is a topological property, we denote by $\alpha\mathcal{P}$ the class of α -spaces which satisfy property \mathcal{P} . In this paper, we investigate the minimal- $\alpha\mathcal{P}$ property with particular regard to the properties $\mathcal{P} = T_0, T_1, T_2$ and $T_{2\frac{1}{2}}$.

2. Basic facts on α -spaces

The α -topology $\alpha\tau(X)$ of a space $(X, \tau(X))$ has some interesting similarities with the notion of *semiregularization* $\tau_s(X)$ (see, for example, [PW]).

Lemma 1 ([N]). *For any α -open set A and any α -closed set C of a space X , we have:*

- (1) $\text{cl}_{\alpha\tau}(A) = \text{cl}_\tau(A)$;
- (2) $\text{int}_{\alpha\tau}(C) = \text{int}_\tau(C)$;
- (3) $\text{int}_{\alpha\tau}(\text{cl}_{\alpha\tau}(A)) = \text{int}_\tau(\text{cl}_\tau(A))$.

Proposition 2 ([N]). *For any space X , we have $\alpha(\alpha\tau(X)) = \alpha\tau(X)$.*

PROOF: Evidently $\alpha\tau(X) \subseteq \alpha(\alpha\tau(X))$. Let $B \in \alpha(\alpha\tau(X))$. Then, there is some $A \in \alpha\tau(X)$ such that $A \subseteq B \subseteq \text{int}_{\alpha\tau}(\text{cl}_{\alpha\tau}(A))$. So, there exists some $U \in \tau(X)$ such that $U \subseteq A \subseteq \text{int}_\tau(\text{cl}_\tau(U))$. Hence, $\text{int}_\tau(\text{cl}_\tau(U)) = \text{int}_\tau(\text{cl}_\tau(A))$ and, by Lemma 1(2), we have

$$U \subseteq A \subseteq B \subseteq \text{int}_{\alpha\tau}(\text{cl}_{\alpha\tau}(A)) = \text{int}_\tau(\text{cl}_\tau(A)) = \text{int}_\tau(\text{cl}_\tau(U)).$$

This proves that $B \in \alpha\tau(X)$. □

Definition 1 ([N]). A space X is called an α -space if $\tau(X) = \alpha\tau(X)$ or, equivalently, if $\tau(X) = \alpha(\sigma(X))$ for some topology $\sigma(X)$ on X .

Proposition 3. *Let $(X, \tau(X))$ be a space. The following are equivalent:*

- (1) X is an α -space;
- (2) $\tau(X) = \{U \cup \{p\} : U \in \tau(X), p \in \text{int}(\text{cl}(U))\}$;
- (3) $\{U \cup \{p\} : U \in \tau(X) \text{ is dense in } X \text{ and } p \in X\} \subseteq \tau(X)$.

PROOF: (1) \Rightarrow (2) Evidently, $\tau(X) \subseteq \{U \cup \{p\} : U \in \tau(X), p \in \text{int}_\tau(\text{cl}_\tau(U))\}$ (it suffices to take $p \in U$). Conversely, let $U \in \tau(X)$ and $p \in \text{int}_\tau(\text{cl}_\tau(U))$. Then $U \subseteq U \cup \{p\} \subseteq U \cup \text{int}_\tau(\text{cl}_\tau(U)) = \text{int}_\tau(\text{cl}_\tau(U))$. Since X is an α -space, $U \cup \{p\} \in \tau(X)$.

(2) \Rightarrow (3) Obvious because, when $\text{cl}_\tau(U) = X$, every $p \in \text{int}_\tau(\text{cl}_\tau(U)) = X$.

(3) \Rightarrow (1) Suppose that $\{U \cup \{p\} : U \in \tau(X) \text{ is dense in } X \text{ and } p \in X\} \subseteq \tau(X)$ and let $A \in \alpha\tau(X)$, i.e. $A \subseteq \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A)))$. The set $D = \text{int}_\tau(A) \cup \text{int}_\tau(X \setminus \text{int}_\tau(A))$ is open and dense because

$$\begin{aligned} \text{cl}(D) &= \text{cl}_\tau(\text{int}_\tau(A)) \cup \text{cl}_\tau(\text{int}_\tau(X \setminus \text{int}_\tau(A))) \\ &= \text{cl}_\tau(\text{cl}_\tau(\text{int}_\tau(A))) \cup \text{cl}_\tau(X \setminus \text{cl}_\tau(\text{int}_\tau(A))) \\ &= \text{cl}_\tau(\text{cl}_\tau(\text{int}_\tau(A)) \cup (X \setminus \text{cl}_\tau(\text{int}_\tau(A)))) \\ &= \text{cl}_\tau(X) \\ &= X. \end{aligned}$$

So, for any $p \in A$, by hypothesis we have that $D \cup \{p\} \in \tau(X)$.

Now, we consider the open set:

$$\begin{aligned} W_p &= (D \cup \{p\}) \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A))) \\ &= (\text{int}_\tau(A) \cup \text{int}_\tau(X \setminus \text{int}_\tau(A)) \cup \{p\}) \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A))) \\ &= (\text{int}_\tau(A) \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A)))) \\ &\quad \cup (\text{int}_\tau(X \setminus \text{int}_\tau(A)) \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A)))) \\ &\quad \cup (\{p\} \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A)))) \\ &= \text{int}_\tau(A) \cup \{p\} \end{aligned}$$

as

$$\begin{aligned} &(\text{int}_\tau(X \setminus \text{int}_\tau(A)) \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A)))) \\ &= (X \setminus \text{cl}_\tau(\text{int}_\tau(A))) \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A))) \\ &\subseteq (X \setminus \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A)))) \cap \text{int}_\tau(\text{cl}_\tau(\text{int}_\tau(A))) = \emptyset. \end{aligned}$$

Thus, for every $p \in A$, $W_p = \text{int}_\tau(A) \cup \{p\} \in \tau(X)$ and so, also the union $\bigcup_{p \in A} W_p = \text{int}_\tau(A) \cup A = A$ is an open set of X . This proves that $\alpha\tau(X) \subseteq \tau(X)$ and hence that X is an α -space. \square

In [N] it is proved the following:

Proposition 4. *A space X is an α -space if and only if all the nowhere dense sets are closed sets.*

The following proposition improves the previous one.

Proposition 5. *A space (X, τ) is an α -space if and only if every nowhere dense set is discrete.*

PROOF: Suppose that X is an α -space and that N is a nowhere dense subset of X . By Proposition 4, N is a closed set and so $X \setminus N$ is an open dense set. For any $p \in N$, by Proposition 3(3), the set $U = (X \setminus N) \cup \{p\}$ is open in X . Since $U \cap N = \{p\}$, it follows that N is discrete.

Conversely, let U be an open dense set of X and $p \in X$. Then $X \setminus U$ is a nowhere dense set and by hypothesis, it is discrete. So, if $p \in X \setminus U$, there exists some $V \in \tau$ such that $V \cap (X \setminus U) = \{p\}$. Hence $U \cup \{p\} = U \cup V \in \tau(X)$. Since the case when $p \in U$ is trivial, by Proposition 3(3), it is proved that X is an α -space. □

It is shown in [Lo] that the operator α is not monotonic, i.e. that, in general, for two topologies $\tau(X)$ and $\sigma(X)$ on a set X , $\tau(X) \subseteq \sigma(X)$ does not imply $\alpha\tau(X) \subseteq \alpha\sigma(X)$. However, we have the following:

Lemma 6. *Let $\tau(X)$ and $\sigma(X)$ be topologies on a set X such that $\tau(X) \subseteq \sigma(X)$ and $\tau_s(X) = \sigma_s(X)$. Then $\alpha\tau(X) \subseteq \alpha\sigma(X)$.*

PROOF: Let $A \in \alpha\tau(X)$. Then there exists some $U \in \tau(X)$ such that $U \subseteq A \subseteq \text{int}_\tau(\text{cl}_\tau(U))$. So, being

$$\text{int}_\sigma(\text{cl}_\sigma(U)) = \text{int}_{\sigma_s}(\text{cl}_{\sigma_s}(U)) = \text{int}_{\tau_s}(\text{cl}_{\tau_s}(U)) = \text{int}_\tau(\text{cl}_\tau(U)),$$

we have $U \subseteq A \subseteq \text{int}_\sigma(\text{cl}_\sigma(U))$ with $U \in \tau \subseteq \sigma$, that is $A \in \alpha\sigma(X)$. □

Let us recall that a space $(X, \tau(X))$ is called:

- T_2 -closed (resp. $T_{2\frac{1}{2}}$ -closed) if it is closed in every Hausdorff (resp. $T_{2\frac{1}{2}}$) space containing X as a subspace,
- minimal- T_2 (resp. minimal- $T_{2\frac{1}{2}}$) if it is a T_2 - (resp. $T_{2\frac{1}{2}}$ -) space and there is no strictly coarser T_2 (resp. $T_{2\frac{1}{2}}$) topology on the same set X .

The following properties are well-known and will be used later.

Proposition 7. *A Hausdorff space is T_2 -closed if and only if every open ultrafilter on X is fixed.*

Proposition 8. *A space $(X, \tau(X))$ is Hausdorff if and only if its semiregularization $(X, \tau_s(X))$ is Hausdorff.*

Corollary 9. *A space $(X, \tau(X))$ is T_2 -closed if and only if its semiregularization $(X, \tau_s(X))$ is minimal- T_2 .*

Proposition 10. *Every regular closed subspace of a T_2 -closed space is T_2 -closed.*

Proposition 11. *A Hausdorff space X is minimal- T_2 if and only if it is semi-regular and T_2 -closed.*

Proposition 12. *If $\sigma(X)$ and $\tau(X)$ are two topologies on a set X such that $\tau_s(X) \subseteq \sigma(X) \subseteq \tau(X)$ then $\tau_s(X) = \sigma_s(X)$.*

PROOF: Let R be a regular open set of $(X, \tau(X))$. Then there exists some $U \in \tau(X)$ such that $R = \text{int}_\tau(\text{cl}_\tau(U))$. So, $R \in \tau_s(X) \subseteq \sigma(X)$. Since $\tau_s(X) \subseteq \sigma(X)$ and using a well-known property of the closure of the semiregularization (see [PW]), we have that

$$\text{cl}_\sigma(R) \subseteq \text{cl}_{\tau_s}(R) = \text{cl}_\tau(R)$$

while, being $\sigma(X) \subseteq \tau(X)$, it follows

$$\text{int}_\sigma(\text{cl}_\sigma(R)) \subseteq \text{int}_\tau(\text{cl}_\tau(R)) = R.$$

Obviously, being $R \in \sigma(X)$, we also have that $R \subseteq \text{int}_\sigma(\text{cl}_\sigma(R))$ and hence that $R = \text{int}_\sigma(\text{cl}_\sigma(R))$. Since $R \in \sigma(X)$, this means that R is a regular open set of $(X, \sigma(X))$. On the other hand, let S be a regular open set of $(X, \sigma(X))$. Then there exists some $V \in \sigma(X)$ such that $S = \text{int}_\sigma(\text{cl}_\sigma(V))$. Since $S \in \sigma(X) \subseteq \tau(X)$, we have that

$$\text{cl}_\tau(S) \subseteq \text{cl}_\sigma(S)$$

Hence, being $\tau_s(X) \subseteq \sigma(X)$ and by some well-known properties of the interior of the semiregularization it follows that

$$\text{int}_\tau(\text{cl}_\tau(S)) = \text{int}_{\tau_s}(\text{cl}_\tau(S)) \subseteq \text{int}_\sigma(\text{cl}_\sigma(S)) = S.$$

Obviously, being $S \in \tau(X)$, we also have that $S \subseteq \text{int}_\tau(\text{cl}_\tau(S))$ and hence that $S = \text{int}_\tau(\text{cl}_\tau(S))$, that is S is a regular open set of $(X, \tau(X))$. Thus, the topologies generated by these families of regular open sets, i.e. the semiregularization of $(X, \tau(X))$ and $(X, \sigma(X))$ coincide and we can conclude that $\tau_s(X) = \sigma_s(X)$. \square

Proposition 13. *Let \mathcal{U} be a free open ultrafilter on a Hausdorff space X and p be a fixed point in X . Then, there exists a Hausdorff topology $\tau_{\mathcal{U}}$ on X such that $\alpha_{\mathcal{U}}(X) \not\subseteq \alpha\tau(X)$.*

PROOF: Let us consider the family $\tau_{\mathcal{U}}(X) = \{U \in \tau(X) : p \in U \Rightarrow U \in \mathcal{U}\}$. It is a simple routine to verify that $\tau_{\mathcal{U}}(X)$ forms a topology on X such that $\tau_{\mathcal{U}}(X) \subseteq \tau(X)$.

The space $(X, \tau_{\mathcal{U}}(X))$ is T_2 . In fact, for every $x \neq y$ in X , since $(X, \tau(X))$ is Hausdorff, there are $U, V \in \tau(X)$ such that $x \in U, y \in V$ and $U \cap V = \emptyset$. If $p \notin U \cup V, U, V \in \tau_{\mathcal{U}}(X)$ and we are done. Otherwise, if, for example, $p \in U$, we have $p \notin V$ and $V \in \tau(X)$. Furthermore, since \mathcal{U} is free with respect to $(X, \tau(X))$, there exist some $N \in \tau(X)$ and some $W \in \mathcal{U}$ such that $y \in N$ and $W \cap N = \emptyset$.

Thus $W \subseteq U \cup W$ implies $U \cup W \in \mathcal{U}$ and hence $U \cup W \in \tau_{\mathcal{U}}(X)$, while being $p \notin V \cap N \in \tau(X)$ it follows that $V \cap N \in \tau_{\mathcal{U}}(X)$. So, $U \cup W$ and $V \cap N$ are two open neighborhoods of x and y respectively in $\tau_{\mathcal{U}}(X)$ such that

$$\begin{aligned} (U \cup W) \cap (V \cap N) &= (U \cap (V \cap N)) \cup (W \cap (V \cap N)) \\ &\subseteq (U \cap V) \cup (W \cap N) \\ &= \emptyset \cup \emptyset = \emptyset \end{aligned}$$

and this proves that the space $(X, \tau_{\mathcal{U}}(X))$ is Hausdorff.

Since it is immediate to see that every neighborhood of p in $\tau_{\mathcal{U}}(X)$ belongs to \mathcal{U} and $(X, \tau_{\mathcal{U}}(X))$ is T_2 , it follows that p is the unique convergence point of \mathcal{U} with respect to $\tau_{\mathcal{U}}(X)$.

In order to show that $\alpha\tau_{\mathcal{U}}(X) \subseteq \alpha\tau(X)$, we observe first that, for every $U \in \tau_{\mathcal{U}}(X)$, it results:

$$(1) \quad \text{cl}_{\tau_{\mathcal{U}}}(U) = \begin{cases} \text{cl}_{\tau}(U) & \text{if } U \notin \mathcal{U}, \\ \text{cl}_{\tau}(U) \cup \{p\} & \text{if } U \in \mathcal{U}. \end{cases}$$

In fact, $\tau(X) \subseteq \tau_{\mathcal{U}}(X)$ implies, in any case, $\text{cl}_{\tau_{\mathcal{U}}}(U) \subseteq \text{cl}_{\tau}(U)$.

Now, consider the case $U \notin \mathcal{U}$ and suppose, by contradiction that there is some $x \in \text{cl}_{\tau_{\mathcal{U}}}(U)$ such that $x \notin \text{cl}_{\tau}(U)$. So, there exists some neighborhood N of x in $\tau(X)$ such that $N \cap U = \emptyset$.

Since \mathcal{U} is an open ultrafilter on $(X, \tau(X))$, $U \notin \mathcal{U}$ implies that $X \setminus \text{cl}_{\tau}(U) \in \mathcal{U}$. Now, $W = (X \setminus \text{cl}_{\tau}(U)) \cup N$ is an open neighborhood of x with respect to $\tau_{\mathcal{U}}(X)$ (because $W \in \tau(X)$, $x \in N \subseteq W$, and $X \setminus \text{cl}_{\tau}(U) \subseteq W$ implies $W \in \mathcal{U}$) but it results

$$W \cap U = \left((X \setminus \text{cl}_{\tau}(U)) \cup N \right) \cap U = \left((X \setminus \text{cl}_{\tau}(U)) \cap U \right) \cup (N \cap U) = \emptyset \cup \emptyset = \emptyset$$

which is a contradiction to $x \in \text{cl}_{\tau_{\mathcal{U}}}(U)$.

Let us consider the case $U \in \mathcal{U}$. Evidently $p \in \text{cl}_{\tau_{\mathcal{U}}}(U)$ as for every open neighborhood N of p in $\tau_{\mathcal{U}}(X)$, it follows that $N \in \mathcal{U}$ and hence $N \cap U \in \mathcal{U}$ implies $N \cap V \neq \emptyset$. Conversely, suppose, by contradiction, that there is some $x \in \text{cl}_{\tau_{\mathcal{U}}}(U)$ such that $x \neq p$ and $x \notin \text{cl}_{\tau}(U)$. Then, there exists some open neighborhood N of p in $\tau(X)$ such that $N \cap U = \emptyset$ and, it must be $N \notin \tau_{\mathcal{U}}(X)$, that is $p \in N \notin \mathcal{U}$. Since $(X, \tau(X))$ is T_2 and $x \neq p$, there is some open neighborhood G of $x \in \tau(X)$ such that $p \notin G$. Thus, $p \notin N \cap G \in \tau(X)$ implies that $N \cap G \in \tau_{\mathcal{U}}(X)$. So, $N \cap G$ is an open neighborhood of x in $\tau_{\mathcal{U}}(X)$ such that $(N \cap G) \cap U \subseteq N \cap U = \emptyset$. A contradiction to $x \in \text{cl}_{\tau_{\mathcal{U}}}(U)$. Applying the usual duality rules to formula (1), we also obtain that, for every $U \in \tau_{\mathcal{U}}(X)$, it results:

$$(2) \quad \text{int}_{\tau_{\mathcal{U}}}(\text{cl}_{\tau_{\mathcal{U}}}(U)) = \begin{cases} \text{int}_{\tau}(\text{cl}_{\tau}(U)) \setminus \{p\} & \text{if } U \notin \mathcal{U}, \\ \text{int}_{\tau}(\text{cl}_{\tau}(U)) & \text{if } U \in \mathcal{U}. \end{cases}$$

In fact, if $U \notin \mathcal{U}$, as \mathcal{U} is an open ultrafilter on $(X, \tau(X))$, $X \setminus \text{cl}_\tau(U) \in \mathcal{U}$ and so we have that:

$$\begin{aligned} \text{int}_{\tau_{\mathcal{U}}}(\text{cl}_{\tau_c U}(U)) &= \text{int}_{\tau_{\mathcal{U}}}(\text{cl}_\tau(U)) \\ &= X \setminus \text{cl}_{\tau_{\mathcal{U}}}(X \setminus (\text{cl}_\tau(U))) \\ &= X \setminus \text{cl}_\tau(X \setminus (\text{cl}_\tau(U) \cup \{p\})) \\ &= (X \setminus \text{cl}_\tau(X \setminus \text{cl}_\tau(U))) \cap (X \setminus \{p\}) \\ &= \text{int}_\tau(\text{cl}_\tau(U)) \setminus \{p\}. \end{aligned}$$

If $U \in \mathcal{U}$, since \mathcal{U} is an open ultrafilter on $(X, \tau(X))$, it necessarily results $X \setminus \text{cl}_\tau(U) \notin \mathcal{U}$ and we have:

$$\begin{aligned} \text{int}_{\tau_{\mathcal{U}}}(\text{cl}_{\tau_{\mathcal{U}}}(U)) &= X \setminus \text{cl}_{\tau_{\mathcal{U}}}(X \setminus \text{cl}_{\tau_{\mathcal{U}}}(U)) \\ &= X \setminus \text{cl}_\tau(X \setminus \text{cl}_{\tau_{\mathcal{U}}}(U)) \\ &= \text{int}_\tau(\text{cl}_{\tau_{\mathcal{U}}}(U)) \\ &= \text{int}_\tau(\text{cl}_\tau(U) \cup \{p\}) \\ &= \text{int}_\tau(\text{cl}_\tau(U)), \end{aligned}$$

where the last equality is due to the fact that $p \in \text{cl}_\tau(U)$ (because for every neighborhood N of p in $\tau(X)$, we have $N \in \mathcal{U}$ and so, being $U \in \mathcal{U}$, it follows that $N \cap U \in \mathcal{U}$ and $N \cap U \neq \emptyset$).

Now, for every $A \in \alpha\tau_{\mathcal{U}}(X)$, there exists some $U \in \tau_{\mathcal{U}}(X)$ such that $U \subseteq A \subseteq \text{int}_{\tau_{\mathcal{U}}}(\text{cl}_{\tau_{\mathcal{U}}}(U))$ and by formula (2), it immediately follows, in both cases, that $U \subseteq A \subseteq \text{int}_\tau(\text{cl}_\tau(U))$ with $U \in \alpha\tau(X) \subseteq \tau(X)$ and so that $A \in \alpha\tau(X)$. Thus $\alpha\tau_{\mathcal{U}}(X) \subseteq \alpha\tau(X)$. To finish the proof, we will show that $\alpha\tau_{\mathcal{U}}(X) \neq \alpha\tau(X)$. Since \mathcal{U} is free, p is not an adherent point for \mathcal{U} and so there exist some neighborhood V of p in $\tau(X)$ and some $U \in \mathcal{U}$ such that $U \cap V = \emptyset$. Thus $V \notin \mathcal{U}$ and, being $p \in V$, by definition of $\tau_{\mathcal{U}}(X)$, it follows that $V \notin \tau_{\mathcal{U}}(X)$. Evidently $V \in \alpha\tau(X)$ (as $V \in \tau(X)$) but $V \notin \alpha\tau_{\mathcal{U}}(X)$. Suppose, by contradiction, that there exists some $W \in \tau_{\mathcal{U}}(X)$ such that $W \subseteq V \subseteq \text{int}_{\tau_{\mathcal{U}}}(\text{cl}_{\tau_{\mathcal{U}}}(W))$. Since $V \notin \mathcal{U}$, it follows a fortiori that $W \notin \mathcal{U}$ and so, by formula (2), we have that $W \subseteq V \subseteq \text{int}_\tau(\text{cl}_\tau(W)) \setminus \{p\}$ and thus that $p \notin V$. A contradiction. This proves that $\alpha\tau_{\mathcal{U}}(X) \subsetneq \alpha\tau(X)$ and concludes our proof. \square

3. Minimal- αT_0 -spaces

Definition 2. Let \mathcal{P} be a topological property. A space X is said to be an $\alpha\mathcal{P}$ -space if it is an α -space and property \mathcal{P} holds.

Definition 3. Let \mathcal{P} be a topological property. A space X is called *minimal- $\alpha\mathcal{P}$* if it is an $\alpha\mathcal{P}$ -space and there is no strictly coarser $\alpha\mathcal{P}$ -topology on the same set.

Since the unique T_1 -minimal-space is the cofinite topology which is an α -space (by Proposition 3), it is evident that the class of minimal- αT_1 -spaces coincides with the well-known class of minimal- T_1 .

One would suspect that if X is minimal- T_0 , then $(X, \alpha\tau(X))$ is minimal- αT_0 . Here is a counterexample.

Example 14. Consider \mathbb{R} with $\tau(\mathbb{R}) = \{(-\infty, a) : a \in \mathbb{R}\} \cup \{\emptyset, \mathbb{R}\}$. Then \mathbb{R} is minimal- T_0 . If $\emptyset \neq U \in \tau(\mathbb{R})$, $\text{cl}_{\mathbb{R}}(U) = \mathbb{R}$, then $\text{int}_{\mathbb{R}}(\text{cl}_{\mathbb{R}}(U)) = \mathbb{R}$. So, if $a < b$ and $c = a - 1$, then $(-\infty, c) \cup \{a\}, (-\infty, c) \cup \{b\} \in \alpha\tau(\mathbb{R})$. In particular, it follows that $(\mathbb{R}, \alpha\tau(\mathbb{R}))$ is T_1 . If σ is the cofinite topology on \mathbb{R} , then $\sigma \subset \tau(\mathbb{R})$.

Lemma 15. Let X be a set, $p \in X$, and $\tau(X) = \{U : p \in U \text{ and } X \setminus U \text{ is finite}\} \cup \{\emptyset\}$. Then X is minimal- αT_0 .

PROOF: Clearly, X is T_0 . If $\emptyset \neq U \in \tau(X)$, then $p \in U$, $\text{cl}_X U = X$, and $\text{int}_X \text{cl}_X U = X$. Now $U \cup \{q\}$ is open for all $q \in X$, by Proposition 3, since X is αT_0 . Let $\sigma \subseteq \tau(X)$ and (X, σ) be αT_0 . If $\emptyset \neq U \in \tau(X)$, then $p \in U$, and $X \setminus U = \{q_1, \dots, q_n\}$ is finite. Since $\emptyset \neq V \in \sigma \subseteq \tau(X)$ implies $p \in V$, then for each $1 \leq i \leq n$, there is a $V_i \in \sigma$ such that $p \in V_i \subseteq X \setminus \{q_i\}$. For $T = \bigcap \{V_i : 1 \leq i \leq n\}$, $p \in T \in \sigma$ and $T \subseteq U$. As (X, σ) is αT_0 and $\text{cl}_{\sigma} T = X$, then $U = \bigcup \{T \cup \{q\} : q \in U \setminus T\} \in \sigma$. □

In the topology just considered, every point different from p is closed. Now, let us consider the case in which some point of X is not closed.

Lemma 16. Let X be an αT_0 -space, and $p \in X$ such that $\text{cl}_X \{p\} \neq \{p\}$. Then $\text{cl}_X(\{p\})$ is a regular-closed set and $p \in \text{int}_X(\text{cl}_X(\{p\}))$.

PROOF: Let $U = X \setminus \text{cl}_X(\{p\})$. It suffices to show that U is regular-open. If $p \in \text{cl}_X(U)$, then $\text{cl}_X(U) = X$ and $\text{int}_X(\text{cl}_X(U)) = X$. So, for $q \in \text{cl}_X(\{p\}) \setminus \{p\}$, $U \cup \{q\}$ is open and $(U \cup \{q\}) \cap \{p\} = \emptyset$. That is, $q \notin \text{cl}_X(\{p\})$, a contradiction. So, $p \notin \text{cl}_X(U)$. For $V = X \setminus \text{cl}_X(U)$, we have that $p \in V \subseteq \text{cl}_X(\{p\})$. Thus, $\text{cl}_X(V) = \text{cl}_X(\{p\})$. □

In order to obtain a characterization of minimal- T_0 -spaces, we need some other lemmas.

Lemma 17. Let X be an αT_0 -space, and $p, q \in X$ such that $\text{cl}_X(\{p\}) \neq \{p\}$, $\text{cl}_X(\{q\}) \neq \{q\}$, and $p \neq q$. Then $p \notin \text{cl}_X(\{q\})$ and $q \notin \text{cl}_X(\{p\})$.

PROOF: Since X is T_0 , then $\text{cl}_X(\{p\}) \neq \text{cl}_X(\{q\})$. Assume that $q \in \text{cl}_X(\{p\})$. Then $p \notin \text{cl}_X(\{q\})$ as $\text{cl}_X(\{p\}) \neq \text{cl}_X(\{q\})$. So, $\text{cl}_X(\{q\}) \subseteq \text{cl}_X(\{p\})$. By Lemma 16, $q \in \text{int}_X(\text{cl}_X(\{q\})) \subseteq \text{int}_X(\text{cl}_X(\{p\})) \subseteq \text{cl}_X(\{p\})$. As $p \notin \text{cl}_X(\{q\})$, it follows that $p \notin \text{int}_X(\text{cl}_X(\{q\}))$. Therefore, $q \notin \text{cl}_X(\{p\})$, a contradiction. □

Lemma 18. *Let X be a minimal- αT_0 -space, and $p, q \in X$ such that $\text{cl}_X(\{p\}) \neq \{p\}$, $\text{cl}_X(\{q\}) \neq \{q\}$, and $p \neq q$. Then $\text{int}_X(\text{cl}_X(\{p\})) = \{p\}$ and $\text{int}_X(\text{cl}_X(\{q\})) = \{q\}$.*

PROOF: Let $\tau = \tau(X)$ be the topology on X , $r \in \text{int}_X(\text{cl}_X(\{p\})) \setminus \{p\}$ and consider the topology $\sigma = \{U \in \tau : r \in U \text{ implies } q \in U\}$ on X . Clearly, $\sigma \subseteq \tau$. There are a number of cases to verify that (X, σ) is T_0 but each case is straightforward. Next we show that (X, σ) is an α -space. Let U be open and dense in (X, σ) and $t \in X \setminus U$. We want to show that $U \cup \{t\} \in \sigma$. If $r \in U$, then U is dense and open in (X, τ) and hence, $U \cup \{t\} \in \sigma$. Suppose that $r \notin U$. Our first goal is to show that U is dense in τ . If $\text{cl}_\tau(U) \neq \text{cl}_\sigma(U)$, then $r \in \text{cl}_\sigma(U) \setminus \text{cl}_\tau(U)$. Then $p \notin U$ and $\text{cl}_\tau(\{p\}) \cap U = \emptyset$. In particular, $\text{int}_\tau(\text{cl}_\tau(\{p\})) \cap U = \emptyset$. There is an $V \in \tau$ such that $p \in V$ and $r \notin V$. Now $p \in \text{int}_\tau(\text{cl}_\tau(\{p\})) \cap V \in \sigma$ and $\text{int}_\tau(\text{cl}_\tau(\{p\})) \cap V \cap U = \emptyset$. So, U is not dense in (X, σ) , a contradiction. Thus, U is dense in τ . Also the above proof shows that $p \in U$. If $q \notin U$, then $\text{cl}_\tau(\{q\}) \cap U = \emptyset$. In particular, $\text{int}_\tau(\text{cl}_\tau(\{q\})) \cap U = \emptyset$. But $q \in \text{int}_\tau(\text{cl}_\tau(\{q\})) \in \sigma$, a contradiction. Thus, $q \in U$. With both $p, q \in U$, we have that $U \cup \{r\} \in \sigma$. For $t \neq r$, then $U \cup \{t\} \in \sigma$ as $U \cup \{t\} \in \tau$ and $r \notin U \cup \{t\}$. □

Proposition 19. *Let X be a minimal- αT_0 -space, and $P = \{p \in X : \text{cl}_X(\{p\}) \neq \{p\}\}$ such that $|P| \geq 2$. Then P is dense in X and if $V \in \tau(X)$ and $V \setminus P \neq \emptyset$, then $P \subseteq V$.*

PROOF: Let $\tau = \tau(X)$ be the topology of the space X . First we show that $Q = \{q \in X : \{q\} \in \tau\}$ is dense. Clearly, $Q \supseteq P$. Fix $r \in X \setminus \text{cl}_X(Q)$. Note that the topology $\sigma = \{U \in \tau : r \in U \text{ implies } Q \subseteq U\} \subseteq \tau$. If we show that (X, σ) is αT_0 , it will follow that Q is dense. Since $\{q\} \in \sigma$ for all $q \in Q$, q can be T_0 -separated from all $p \in X \setminus \{q\}$ in σ . As $(X \setminus \text{cl}_X(Q)) \cup Q \in \sigma$ and $(X \setminus \text{cl}_X(Q)) \cup Q \cup \{t\} \in \sigma$ for $t \in \text{cl}_X(Q) \setminus Q$, a point $t \in \text{cl}_X(Q) \setminus Q$ can be T_0 -separated from all $p \in X \setminus \{t\}$ in σ . Let $s, t \in X \setminus \text{cl}_X(Q)$. As τ is T_0 , there is some $V \in \tau$ such that $s \in V$ and $t \notin V$ or vice versa. Now, $s \in V \cup Q \in \sigma$ and $t \notin V \cup Q$. This completes the proof that (X, σ) is T_0 . Next we show that (X, σ) is an α -space. Let U be an open and dense subset of (X, σ) . Note that $Q \subseteq U$. If $\text{cl}_\tau(U) \neq \text{cl}_\sigma(U)$, then $r \in \text{cl}_\sigma(U) \setminus \text{cl}_\tau(U)$. There is an $V \in \tau$ such that $r \in V$ and $V \cap Q = \emptyset$. As $r \notin P$, $\text{cl}_\tau(\{r\}) = \{r\}$, $\emptyset \neq V \setminus \{r\} \in \sigma$ and $V \setminus \{r\} \cap U = \emptyset$, a contradiction as U is dense in (X, σ) . This shows that $\text{cl}_\tau(U) = X$. As τ is an α topology, for $t \in X, U \cup \{t\} \in \tau$. As $Q \subseteq U, U \cup \{t\} \in \sigma$. This completes the proof that (X, σ) is an αT_0 -space. Finally, we show that $Q = P$. Assume that $q \in Q \setminus P$. The topology $\sigma = \{U \in \tau : q \in U \text{ implies } P \subseteq U\} \subseteq \tau$. Similar to the above, it is straightforward to show that (X, σ) is T_0 and an α -space, a contradiction as τ is minimal- αT_0 . Thus, $P = Q$. □

Proposition 20. *Let X be a minimal- αT_0 -space, and $P = \{p \in X : \text{cl}_X(\{p\}) \neq \{p\}\}$ such that $|P| \geq 2$. Then $\tau(X)$ is generated by the base $\{P \cup \{q\} : q \in X \setminus P\} \cup \{\{p\} : p \in P\}$.*

PROOF: This is an obvious consequence of Proposition 19. □

Proposition 21. *Let X be a set and $P \subsetneq X$ such that $|P| \geq 2$. Let $\tau(X)$ be generated by the base $\{P \cup \{q\} : q \in X \setminus P\} \cup \{\{p\} : p \in P\}$. Then X is a minimal- αT_0 -space.*

PROOF: Let $\tau = \tau(X)$ be the topology on X and $\sigma \subseteq \tau$ be an αT_0 topology. For $p \in P, (X \setminus P) \cup \{p\} = \text{cl}_\tau(\{p\}) \subseteq \text{cl}_\sigma(\{p\})$. Thus, for all $p \in P, \text{cl}_\sigma(\{p\}) \neq \{p\}$. By Lemma 17, for $q \in P \setminus \{p\}, q \notin \text{cl}_\sigma(\{p\})$. Hence $\text{cl}_\sigma(\{p\}) = (X \setminus P) \cup \{p\} = \text{cl}_\tau(\{p\})$. Thus, $P \setminus \{p\} \in \sigma$ for all $p \in P$. By Lemma 16, $p \in \text{int}_\sigma(\text{cl}_\sigma(\{p\}))$. As $|P| \geq 2$, let $q \in P \setminus \{p\}$. Then $\text{int}_\sigma(\text{cl}_\sigma\{q\}) \cap P \setminus \{p\} = \{q\} \in \sigma$ and $P \in \sigma$. As $X = \text{cl}_\tau(P) \subseteq \text{cl}_\sigma(P), X = \text{int}_\sigma(\text{cl}_\sigma(P))$. Since (X, σ) is an αT_0 -space, it follows that for $q \in X \setminus P, P \cup \{q\} \in \sigma$. This shows that $\sigma = \tau$. □

Finally, we obtain the characterization of αT_0 -spaces.

Theorem 22. *Let X be an αT_0 -space and $P = \{p \in X : \text{cl}_X(\{p\}) \neq \{p\}\}$. Then X is minimal- αT_0 iff $P \neq \emptyset$ and*

- (i) *if $P = \{p\}$, then $\tau(X) = \{U : p \in U \text{ and } X \setminus U \text{ is finite}\} \cup \{\emptyset\}$, or*
- (ii) *if $|P| \geq 2$, then $\tau(X)$ is generated by the base $\{P \cup \{q\} : q \in X \setminus P\} \cup \{\{p\} : p \in P\}$.*

PROOF: (\Leftarrow) It follows from Lemma 18 and Proposition 20. □
 (\Rightarrow) It follows from Lemma 15 and Proposition 21. □

4. Minimal- αT_2 -spaces

Proposition 23. *Let X be an αT_2 -space. Then X is a minimal- αT_2 -space if and only if $(X, \tau_s(X))$ is minimal- T_2 and $\tau(X) = \alpha \tau_s(X)$.*

PROOF: (\Rightarrow) Suppose that X is a minimal- αT_2 -space. Thus $\alpha \tau(X) = \tau(X)$. Then the space $(X, \tau(X))$ is T_2 -closed. In fact, if, by contradiction, it is not, by Proposition 7, there exists some free open ultrafilter \mathcal{U} on $(X, \tau(X))$ and by Proposition 13, there exists a strictly coarser αT_2 topology $\tau_{\mathcal{U}}$. A contradiction to the α -minimality of $(X, \tau(X))$. Hence, by Corollary 9, $(X, \tau_s(X))$ is minimal- T_2 . Furthermore, being $\tau_s(X) \subseteq \tau(X)$ and, obviously, $(\tau_s(X))_s = \tau_s(X)$, by Lemma 6, we have that $\alpha \tau_s(X) \subseteq \alpha \tau(X)$, i.e. $\alpha \tau_s(X) \subseteq \tau(X)$ where $\alpha \tau_s(X)$ is Hausdorff by Proposition 11 as $\tau(X)$ is Hausdorff and the T_2 axioms are expansive. Since $(X, \tau(X))$ is minimal- αT_2 , we conclude that $\alpha \tau_s(X) = \tau(X)$.

(\Leftarrow) Let us suppose that $(X, \tau_s(X))$ is minimal- T_2 and $\tau(X) = \alpha \tau_s(X)$. Then $(X, \tau(X))$ is H -closed by Corollary 9. Now, let $\sigma(X)$ be an αT_2 topology on X such that

$$\sigma(X) \subseteq \tau(X).$$

For every R regular open set of $(X, \tau(X))$, $X \setminus R$ is a regular closed set and hence, by Proposition 10, it is a T_2 -closed subspace of the Hausdorff space $(X, \sigma(X))$. Thus $X \setminus R$ is a closed set of $(X, \sigma(X))$ and $R \in \sigma(X)$. This proves that

$$\tau_s(X) \subseteq \sigma(X).$$

Hence, by Proposition 12, $\tau_s(X) = \sigma_s(X)$ and so, applying Lemma 6 to $\tau_s(X) \subseteq \sigma(X)$, we have that $\alpha\tau_s(X) \subseteq \alpha\sigma(X)$. Since $\sigma(X)$ is an α topology and, by hypothesis $\alpha\tau_s(X) = \tau(X)$, it follows that $\tau(X) \subseteq \sigma(X)$ and so that $\tau(X) = \sigma(X)$. This shows that $(X, \tau(X))$ is a minimal- αT_2 -space and concludes our proof. \square

Corollary 24. *Let $(X, \tau(X))$ be a space with a dense set D of isolated points. Then:*

- (1) *the α -topology $\alpha\tau(X)$ coincides with the topology generated by $\tau(X) \cup \{D \cup \{x\} : x \in X\}$, i.e. it is a simple extension of the subspace D ;*
- (2) *X is an α -space if and only if $X \setminus D$ is discrete;*
- (3) *if X is a semiregular α -space, it results:*

$$\{\sigma : \sigma \text{ is a topology on } X \text{ such that } \sigma_s = \tau(X)\} = \{\tau(X)\}.$$

PROOF: Straightforward applications of Propositions 3, 5 and 6. \square

Example 25. Let us consider the set $X = \mathbb{R} \times [0, +\infty[$. It is easy to verify that

$$\tau_s(X) = \{U \subseteq X : (x, 0) \in U \Rightarrow \exists \epsilon > 0 \text{ such that }]x - \epsilon, x + \epsilon[\times]0, \epsilon[\subseteq U\}$$

defines a Tychonoff topology on X and that $D = \mathbb{R} \times]0, +\infty[$ is a dense set of isolated points. Now, if we consider another topology on X ,

$$\begin{aligned} \tau(X) = \{ & U \subseteq X : (x, 0) \in U \setminus \mathbb{Q} \times \{0\} \Rightarrow \exists \epsilon > 0 \\ & \text{such that }]x - \epsilon, x + \epsilon[\times]0, \epsilon[\subseteq U \\ & \text{and } (x, 0) \in U \cap \mathbb{Q} \times \{0\} \Rightarrow \exists \epsilon > 0 \text{ such that} \\ &]x - \epsilon, x + \epsilon[\times]0, \epsilon[\cup (]x - \epsilon, x + \epsilon[\cap \mathbb{Q} \times \{0\}) \subseteq U \}, \end{aligned}$$

it is a simple routine to check that $\tau_s(X)$ is the semiregularization of $\tau(X)$ and that $\tau(X) \neq \tau_s(X)$. Furthermore, since $C = X \setminus (\mathbb{R} \times]0, +\infty[)$ is a closed nowhere dense subset with respect to both $\tau(X)$ and $\tau_s(X)$, by Corollary 24 follows that $\alpha\tau(X) = \alpha\tau_s(X)$ coincides with the topology generated by $\tau(X) \cup \{(\mathbb{R} \times]0, +\infty[) \cup \{(x, 0)\} : x \in \mathbb{R}\}$ and so that $\tau(X) \neq \alpha\tau(X)$. Thus, we have that

$$\tau_s(X) \subsetneq \tau(X) \subsetneq \alpha\tau(X) = \alpha\tau_s(X).$$

Example 26 (A Tychonoff α -space which is not minimal- T_2). Let us recall that two sets are *almost disjoint* if their intersection is a finite set. It is a simple routine to show (by using Zorn's Lemma) that there exists a maximal almost disjoint family \mathcal{M} of subsets of \mathbb{N} . The space generated on the set $\psi = \mathbb{N} \cup \mathcal{M}$ by the base

$$\mathcal{B} = \left\{ \{n\} : n \in \mathbb{N} \right\} \cup \left\{ \{M\} \cup S : M \in \mathcal{M} \text{ and } S \text{ is a cofinite subset of } M \right\}$$

is a 0-dimensional (and hence Tychonoff) but not normal (and hence not minimal- T_2). This space is known in literature as the ψ -space (see 1N, [PW]). Since, every closed nowhere dense set of ψ is discrete, by Proposition 5, it is evident that ψ is an α -space.

Example 27 (A minimal- T_2 , α -space). Let us consider the set

$$Z = \left\{ \left(\frac{1}{n}, 0 \right) : n \in \mathbb{N} \right\} \cup \left\{ \left(\frac{1}{n}, \frac{1}{m} \right) : n \in \mathbb{N}, m \in \mathbb{Z} \right\}$$

with the topology $\tau(Z)$ induced by the usual topology on \mathbb{R}^2 . Let $X = Z \cup \{a, b\}$ and define a topology $\tau(X)$ on X by saying that a subset $U \subset Z$ is open if $U \cap Z \in \tau(Z)$ and if $a \in U$ (respectively, $b \in U$) there exists some $k \in \mathbb{N}$ such that $\left\{ \left(\frac{1}{n}, \frac{1}{m} \right) : n \in \mathbb{N}, m \geq k \right\} \subseteq U$ (respectively, $\left\{ \left(\frac{1}{n}, -\frac{1}{m} \right) : n \in \mathbb{N}, m \geq k \right\} \subseteq U$). It is well-known (see 4.8(d), [PW]) that the space X is Urysohn, not compact and minimal- T_2 . Furthermore, since every its nowhere dense subset (namely, $\left\{ \left(\frac{1}{n}, 0 \right) : n \in \mathbb{N} \right\} \cup \{a, b\}$) is discrete, it follows from Proposition 5 that X is an α -space.

Let us note that, by 24(3), in both of the above spaces, we have that $\{\sigma : \sigma \text{ is a topology on } X \text{ such that } \sigma_s = \tau(X)\} = \{\tau(X)\}$. Thus, the question remains: *is there some semiregular α -space $(X, \tau(X))$ such that $\{\tau(X)\} \subsetneq \{\sigma : \sigma \text{ is a topology on } X \text{ such that } \sigma_s = \tau(X)\}$?*

We now provide an example of a Tychonoff α -space X such that

$$\{\tau(X)\} \subsetneq \{\sigma : \sigma \text{ is a topology on } X \text{ such that } \sigma_s = \tau(X)\}.$$

Example 28. Recall that a measurable subset A of \mathbb{R} has density 1 if

$$A = \left\{ x \in \mathbb{R} : \lim_{h \rightarrow 0} \frac{m(A \cap [x - h, x + h])}{2h} = 1 \right\}.$$

The set of $\{A \subseteq \mathbb{R} : A \text{ measurable with density } 1\}$ forms a topology $\delta(\mathbb{R})$, called the density topology, on the set \mathbb{R} . The space $(\mathbb{R}, \delta(\mathbb{R}))$ is a Tychonoff space without isolated points, strictly finer than the usual topology $\tau(\mathbb{R})$, and has the property that every nowhere dense subset is closed and discrete (see 2.7 in [T]). In

particular, $(\mathbb{R}, \delta(\mathbb{R}))$ is an α -space by Proposition 4. We need one additional fact about $(\mathbb{R}, \delta(\mathbb{R}))$. Note that for $x \in U \in \delta(\mathbb{R})$, $U \cap (x, \infty) \neq \emptyset$ and $U \cap (-\infty, x) \neq \emptyset$. In particular, the $\delta(\mathbb{R})$ open neighborhood filter is contained in two distinct $\delta(\mathbb{R})$ open ultrafilters on \mathbb{R} . Before continuing with this example, we need a result about absolutes of spaces.

For a Hausdorff space X and let $EX = \{\mathcal{U} : \mathcal{U} \text{ is a convergent, open ultrafilter on } X\}$. For $U \in \tau(X)$, let $O(U) = \{\mathcal{U} \in EX : U \in \mathcal{U}\}$. For $U, V \in \tau(X)$, it is easy to verify (see [PW]) that $O(\emptyset) = \emptyset$, $O(X) = EX$, $O(U \cap V) = O(U) \cap O(V)$, $O(U \cup V) = O(U) \cup O(V)$, $EX \setminus O(U) = O(X \setminus \text{cl}_X(U))$, and $O(U) = O(\text{int}_X \text{cl}_X(U))$. EX with the topology generated by $\{O(U) : U \in \tau(X)\}$ is an extremally disconnected Tychonoff space, called the *absolute* of X . The function $k : EX \rightarrow X$ defined by $k(\mathcal{U})$ is the unique convergent point of \mathcal{U} is called a covering function and has the properties that k is irreducible, θ -continuous, perfect and onto. If X is regular, then k is also continuous. If $D \subseteq EX$ such that $k[D] = X$, then D is dense in EX .

Proposition 29. *Let X be a regular α -space and Y a subspace of EX such that for each $x \in X$, $|k^{\leftarrow}(\{x\})| = 2$. Then Y is an α -space.*

PROOF: Let $\bar{k} = k|_Y$. The function $\bar{k} = k|_Y : Y \rightarrow X$ is continuous and onto. So, Y is dense in EX and extremally disconnected. Let N be a nowhere dense subset of Y . Suppose $U = \text{int}_{EX}(\text{cl}_{EX}(N)) \neq \emptyset$. Then, as Y is dense, $\emptyset \neq U \cap Y \subseteq \text{cl}_{EX}(N) \cap Y = \text{cl}_Y(N)$, contradicting that N is nowhere dense in Y . Thus, N is nowhere dense in EX . By 6.5d(2) in [PW], $k[N] = \bar{k}[N]$ is nowhere dense in X and hence discrete in X . Let $p \in N$. There is an open set V in X such that $V \cap \bar{k}[N] = \{k(p)\}$. So, $N \cap \bar{k}^{\leftarrow}[V] = \bar{k}^{\leftarrow}(\{p\})$. But $\bar{k}^{\leftarrow}(\{p\})$ has only two points and there is an open set W in Y such that $N \cap \bar{k}^{\leftarrow}[V] \cap W = \{p\}$. This shows that N is discrete in Y and Y is an α -space. \square

We are ready to apply Proposition 29 to the regular α -space $(\mathbb{R}, \delta(\mathbb{R}))$. First note that the covering function $k : E\mathbb{R} \rightarrow (\mathbb{R}, \delta(\mathbb{R}))$ has the property that $|k^{\leftarrow}(r)| \geq 2$ for each $r \in \mathbb{R}$ since each open neighborhood filter is contained in two distinct $\delta(\mathbb{R})$ open ultrafilters on \mathbb{R} . Let $Y \subseteq E\mathbb{R}$ have the property that $|k^{\leftarrow}(r) \cap Y| = 2$ for each $r \in \mathbb{R}$, and let $Z \subset Y$ be such that $|k^{\leftarrow}(r) \cap Z| = 1$ for each $r \in \mathbb{R}$. Now, both Y and Z are dense in $E\mathbb{R}$. By Proposition 29, Y is a Tychonoff α -space. Let σ be the topology on Y generated by $\tau(Y) \cup \{Z\}$. Clearly, $\tau(Y) \subsetneq \sigma$, and it is straightforward to show that $\sigma_s = \tau(Y)$. The space Y is the desired space.

5. A problem

Since the property to be $T_{2\frac{1}{2}}$ -closed does not pass to the regular open subspaces, we cannot use a result like Proposition 10 to prove a proposition similar to Proposition 23. So, we leave the following as an unsolved problem:

Conjecture. Let X be an $\alpha T_{2\frac{1}{2}}$ -space. Then X is a minimal- $\alpha T_{2\frac{1}{2}}$ -space if and only if $(X, \tau_s(X))$ is semiregular, $T_{2\frac{1}{2}}$ -closed and $\tau(X) = \alpha\tau_s(X)$.

REFERENCES

- [D] Dontchev J., *Survey on pre-open sets*, preprint, 1999.
- [E] Engelking R., *General Topology*, Heldermann, Berlin, 1989.
- [L] Larson R., *Minimal T_0 -spaces and minimal T_D -spaces*, Pacific J. Math. **31** (1969), 451–458.
- [Le] Levine N., *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly **70** (1963), 36–41.
- [Lo] Lo Faro G., *Su alcune proprietà degli insiemi α -aperti*, Atti Sem. Mat. Fis. Univ. Modena **XXIX** (1980), 242–252 (in Italian).
- [N] Njåstad O., *On some classes of nearly open sets*, Pacific J. Math. **15** (1965), no. 3, 961–970.
- [PW] Porter J.R., Woods R.G., *Extensions and Absolutes of Hausdorff Spaces*, Springer, New York, 1988.
- [T] Tall F.D., *The density topology*, Pacific J. Math. **62** (1976), 275–284.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MESSINA, CONTRADA PAPARDO,
SALITA SPERONE 31, 98166 SANT'AGATA, MESSINA, ITALY

E-mail: lofaro@unime.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MESSINA, CONTRADA PAPARDO,
SALITA SPERONE 31, 98166 SANT'AGATA, MESSINA, ITALY

E-mail: nardo@dipmat.unime.it

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, 405 SNOW HALL,
1460 JAYHAWK BLVD., LAWRENCE, KANSAS 66045-7567, USA

E-mail: porter@math.ukans.edu

(Received October 7, 2002)