

Pierre Matet

Club-guessing, good points and diamond

Commentationes Mathematicae Universitatis Carolinae, Vol. 48 (2007), No. 2, 211--216

Persistent URL: <http://dml.cz/dmlcz/119651>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2007

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Club-guessing, good points and diamond

PIERRE MATET

Abstract. Shelah’s club-guessing and good points are used to show that the two-cardinal diamond principle $\diamond_{\kappa,\lambda}$ holds for various values of κ and λ .

Keywords: $P_\kappa(\lambda)$, diamond principle

Classification: 03E05

Let κ be a regular cardinal greater than or equal to ω_2 , and λ be a cardinal greater than κ . In [4] Jech introduced the following notions. $P_\kappa(\lambda)$ denotes the collection of all subsets of λ of size less than κ . A subset C of $P_\kappa(\lambda)$ is *closed unbounded* if (a) it is cofinal in the partially ordered set $(P_\kappa(\lambda), \subseteq)$, and (b) for any infinite ordinal $\theta < \kappa$ and any sequence $\langle a_\alpha : \alpha < \theta \rangle$ of elements of C such that $a_\beta \subseteq a_\alpha$ whenever $\beta < \alpha < \theta$, $\bigcup_{\alpha < \theta} a_\alpha \in C$. A subset S of $P_\kappa(\lambda)$ is *stationary* if $S \cap C \neq \emptyset$ for every closed unbounded subset C of $P_\kappa(\lambda)$. The principle $\diamond_{\kappa,\lambda}$ asserts the existence of a sequence $\langle s_a : a \in P_\kappa(\lambda) \rangle$ with $s_a \subseteq a$ such that for any $X \subseteq \lambda$, $\{a : s_a = X \cap a\}$ is a stationary subset of $P_\kappa(\lambda)$. Jech showed that $\diamond_{\kappa,\lambda}$ could be introduced by forcing. Moreover, he proved that $\diamond_{\kappa,\lambda}$ holds in the constructible universe L . It was shown in [2] that if $2^{<\kappa} < \lambda$, then $\diamond_{\kappa,\lambda}$ holds. In this paper we show that if $2^{<\kappa} \leq \mu^+$ for some cardinal μ such that $\omega < \text{cf}(\mu) < \kappa < \mu \leq \lambda$, then $\diamond_{\kappa,\lambda}$ holds. So if either $2^{<\kappa} = \lambda$ and λ is the successor of a cardinal of uncountable cofinality less than κ , or $2^{<\kappa} = \lambda^+$ and $\omega < \text{cf}(\lambda) < \kappa$, then $\diamond_{\kappa,\lambda}$ holds (and hence the nonstationary ideal on $P_\kappa(\lambda)$ is not 2^λ -saturated). Our result is proved by modifying the argument used by Foreman and Magidor in [3] to show that if $\text{cf}(\lambda) < \kappa$, then there is a family of λ^{++} stationary subsets of $P_\kappa(\lambda)$ such that any two of them have nonstationary intersection.

We need a few lemmas.

Lemma 1 (Solovay [9]). *Let ρ be a regular uncountable cardinal. Then every stationary subset of ρ is the union of ρ disjoint stationary sets.*

Given two regular infinite cardinals $\theta < \rho$, E_θ^ρ denotes the set of all infinite limit ordinals $\alpha < \rho$ such that $\text{cf}(\alpha) = \theta$.

Lemma 2 (Shelah see [5]). *Let $\rho > \omega_1$ be a regular cardinal, and S be a stationary subset of E_ω^ρ . Then one can find a cofinal, order-type ω subset c_γ of γ for each $\gamma \in S$ so that $\{\gamma \in S : c_\gamma \subseteq C\}$ is stationary in ρ for any closed unbounded subset C of ρ .*

Our source for the following notions and facts is [1]. Let $\mu > \kappa$ be a singular cardinal of uncountable cofinality $\nu < \kappa$. Suppose $\langle \mu_i : i < \nu \rangle$ is an increasing sequence of regular cardinals such that $\kappa < \mu_0$ and $\sup\{\mu_i : i < \nu\} = \mu$. Given $f, g \in \prod_{i < \nu} \mu_i$, $f <^* g$ means that $|\{i : f(i) \geq (i)\}| < \nu$. Similarly, $f \leq^* g$ means that $|\{i : f(i) > g(i)\}| < \nu$. By a *scale of length ξ* , ξ an ordinal, we mean a sequence $\langle f_\alpha : \alpha < \xi \rangle$ of elements of $\prod_{i < \nu} \mu_i$ such that (a) $f_\beta <^* f_\alpha$ whenever $\beta < \alpha < \xi$, and (b) for every $g \in \prod_{i < \nu} \mu_i$, there is $\alpha < \xi$ with $g <^* f_\alpha$. Shelah proved that the μ_i can be chosen so that there exists a scale of length μ^+ . Let $\langle f_\alpha : \alpha < \mu^+ \rangle$ be such a scale. Given an infinite limit ordinal $\alpha < \mu^+$, an *exact upper bound* for the sequence $\langle f_\beta : \beta < \alpha \rangle$ is an element g of $\prod_{i < \nu} \mu_i$ such that (i) $f_\beta <^* g$ for any $\beta < \alpha$, and (ii) for every $h \in \prod_{i < \nu} \mu_i$ with $h <^* g$, there is $\beta < \alpha$ with $h <^* f_\beta$. By a *good point*, an infinite limit ordinal $\alpha < \mu^+$ is meant such that $\text{cf}(\alpha) > \nu$ and there exists an exact upper bound g_α for $\langle f_\beta : \beta < \alpha \rangle$ with the property that for any $i < \nu$, $g_\alpha(i)$ is an infinite limit ordinal of cofinality $\text{cf}(\alpha)$. Letting S denote the set of good points α such that $\text{cf}(\alpha) = \kappa$, S is stationary in μ^+ . Now consider the sequence $\langle h_\alpha : \alpha < \mu^+ \rangle$ defined by: $h_\alpha = f_\alpha$ if $\alpha \notin S$, and $h_\alpha = g_\alpha$ otherwise. Then $\langle h_\alpha : \alpha < \mu^+ \rangle$ is a scale. Moreover, for each $\alpha \in S$, h_α is an exact upper bound for $\langle h_\beta : \beta < \alpha \rangle$. Let us sum it up all in the following.

Lemma 3 (Shelah see [1]). *Let $\mu > \kappa$ be a singular cardinal of uncountable cofinality $\nu < \kappa$. Then one can find sequences $\langle \mu_i : i < \nu \rangle$ and $\langle h_\alpha : \alpha < \mu^+ \rangle$ and a set S such that (a) $\langle \mu_i : i < \nu \rangle$ is an increasing sequence of regular cardinals such that $\kappa < \mu_0$ and $\sup\{\mu_i : i < \nu\} = \mu$, (b) $\langle h_\alpha : \alpha < \mu^+ \rangle$ is a scale of length μ^+ in $\prod_{i < \nu} \mu_i$, (c) S is a stationary subset of $E_\kappa^{\mu^+}$, and (d) for each $\alpha \in S$, $\text{ran}(h_\alpha) \subseteq E_\kappa^{\mu^+}$ and h_α is an exact upper bound for $\langle h_\beta : \beta < \alpha \rangle$.*

Suppose μ is a cardinal greater than κ . For $n < \omega$, let R_n^μ be the set of all increasing functions from n to $E_\kappa^{\mu^+}$. Let \mathcal{T}_μ be the collection of all nonempty subsets T of $\bigcup_{n < \omega} R_n^\mu$ such that for any $n < \omega$ and any $t \in T \cap R_n^\mu$, $\{t \upharpoonright \ell : \ell < n\} \subseteq T$ and $\{\alpha \in E_\kappa^{\mu^+} : t \cup \{n, \alpha\} \in T\}$ is stationary in μ^+ .

Lemma 4 (Shioya [8]). *Suppose that $\mu > \kappa$ is a cardinal, $T \in \mathcal{T}_\mu$, $n < \omega$ and $\varphi : T \cap (\bigcup_{n < q < \omega} R_q^\mu) \rightarrow \mu^+$ is such that for every $t \in \text{dom}(\varphi)$, (a) $\varphi(t) \in t(n)$, and (b) $\varphi(t \upharpoonright q) \leq \varphi(t)$ for $n < q < \text{dom}(t)$. Then one can find $T' \in \mathcal{T}_\mu \cap P(T)$ and $\psi : T \cap R_n^\mu \rightarrow \mu^+$ so that (i) $T' \cap R_n^\mu = T \cap R_n^\mu$ and (ii) $\varphi(t) \leq \psi(t \upharpoonright n)$ for any $t \in T' \cap (\bigcup_{n < q < \omega} R_q^\mu)$.*

For $A \subseteq P_\kappa(\lambda)$, $G_{\kappa,\lambda}(A)$ denotes the following two-person game lasting ω moves. Player I makes the first move. I and II alternately pick members of $P_\kappa(\lambda)$, thus building a sequence $\langle a_n : n < \omega \rangle$ with the condition that $a_0 \subseteq a_1 \subseteq \dots$. II wins the game just in case $\bigcup_{n < \omega} a_n \in A$. Let $NG_{\kappa,\lambda}$ be the set of all $B \subseteq P_\kappa(\lambda)$ such that II has a winning strategy in $G_{\kappa,\lambda}(P_\kappa(\lambda) \setminus B)$.

Lemma 5 (Matet [6]). $NG_{\kappa,\lambda}$ is a normal ideal on $P_\kappa(\lambda)$.

Proposition 6. Suppose $2^{<\kappa} \leq \mu^+$ for some cardinal μ such that $\kappa < \mu \leq \lambda$ and $\omega < \text{cf}(\mu) < \kappa$. Then there is a sequence $\langle s_a : a \in P_\kappa(\lambda) \rangle$ with $s_a \subseteq a$ such that for any $X \subseteq \lambda$, $\{a : s_a = X \cap a\} \in NG_{\kappa,\lambda}^+$.

PROOF: Let μ be a fixed cardinal such that $\omega < \text{cf}(\mu) < \kappa < \mu \leq \lambda$ and $2^{<\kappa} \leq \mu^+$. Fix a stationary subset H of $E_\omega^{\mu^+}$. Using Lemma 2, select an increasing function $\bar{\gamma}$ from ω into γ for each $\gamma \in H$ so that $\text{ran}(\bar{\gamma})$ is cofinal in γ for every $\gamma \in H$, and $\{\gamma \in H : \text{ran}(\bar{\gamma}) \subseteq C\}$ is stationary in μ^+ for any closed unbounded subset C of μ^+ .

Set $\nu = \text{cf}(\mu)$ and let $\langle \mu_i : i < \nu \rangle$, $\langle h_\alpha : \alpha < \mu^+ \rangle$ and S be as in the statement of Lemma 3. For $\alpha < \mu^+$, set $x_\alpha = \text{ran}(h_\alpha)$. For $b \in P_\kappa(\mu)$, define $g_b \in \prod_{i < \nu} \mu_i$ by $g_b(i) = \sup(b \cap \mu_i)$. Define $\rho : P_\kappa(\mu) \rightarrow \mu^+$ by $\rho(b) =$ the least $\beta < \mu^+$ such that $g_b \leq^* h_\beta$. For $\alpha \in S$ and $b \in P_\kappa(\mu)$, define $g_b^\alpha \in \prod_{i < \nu} h_\alpha(i)$ by $g_b^\alpha(i) = \sup(b \cap h_\alpha(i))$. For $\alpha \in S$, define $\rho_\alpha : P_\kappa(\mu) \rightarrow \alpha$ by $\rho_\alpha(b) =$ the least $\beta < \mu^+$ such that $g_b^\alpha \leq^* h_\beta$. Note that given any sequence $\langle b_n : n < \omega \rangle$ of elements of $P_\kappa(\mu)$, $\rho(\bigcup_{n < \omega} b_n) = \sup\{\rho(b_n) : n < \omega\}$. Moreover, for every $\alpha \in S$, $\rho_\alpha(\bigcup_{n < \omega} b_n) = \sup\{\rho_\alpha(b_n) : n < \omega\}$.

We will prove that there is a sequence $\langle s_a : a \in P_\kappa(\lambda) \rangle$ such that for any $X \subseteq \lambda$,

$$\{a : \rho(\mu \cap a) \in H \text{ and } s_a = X \cap a\} \in NG_{\kappa,\lambda}^+.$$

For $n < \omega$ and $0 < \zeta < \kappa$, let F_n^ζ be the set of all $(n+1)$ -tuples (f_0, \dots, f_n) of functions from ζ to 2. By Lemma 1, S can be partitioned into disjoint stationary subsets Z_n , $n < \omega$. Again by Lemma 1, for each n , Z_n can be decomposed into disjoint stationary subsets $Z(f_0, \dots, f_n)$, $(f_0, \dots, f_n) \in \bigcup_{0 < \zeta < \kappa} F_n^\zeta$.

For $b \subseteq \lambda$, let $e(b) : \text{o.t.}(b) \rightarrow b$ be the function that enumerates the elements of b in increasing order. For $a, b \in P_\kappa(\lambda)$ with $a \subseteq b$, let $\chi(a, b) : \text{o.t.}(b) \rightarrow 2$ be defined by $(\chi(a, b))(\delta) = 1$ if and only if $(e(b))(\delta) \in a$.

The proof will go as follows. Given $A \in NG_{\kappa,\lambda}^*$ and $X \subseteq \lambda$, we will construct a_n and α_n for $n < \omega$, f_n^i for $i \leq n < \omega$, and a and γ so that (a) $a_0, a_1, \dots \in P_\kappa(\lambda)$ and $a_0 \subseteq a_1 \subseteq \dots$, (b) $f_n^i = \chi(a_i, a_n)$ for $i < n$ and $f_n^n = \chi(X \cap a_n, a_n)$, (c) $a = \bigcup_{n < \omega} a_n$, $a \in A$ and $\gamma = \rho(a \cap \mu)$, (d) $\alpha_{n+1} =$ the least α such that $\bar{\gamma}(n) < \alpha < \mu^+$ and $\text{ran}(h_\alpha) \subseteq a$, and (e) $\alpha_{n+1} \in Z(f_n^0, f_n^1, \dots, f_n^n)$.

The guessing sequence $\langle s_a : a \in P_\kappa(\lambda) \rangle$ is now defined in the obvious way. Given $a \in P_\kappa(\lambda)$, put $\xi = \text{o.t.}(a)$ and $\gamma = \rho(a \cap \mu)$. Let $(*)$ assert that $\gamma \in H$ and there exist ζ_n and α_{n+1} for $n < \omega$ and f_n^i for $i \leq n < \omega$ such that (0) $0 < \zeta_n < \kappa$, (1) f_n^i is a function from ζ_n to 2, (2) $\alpha_{n+1} =$ the least $\alpha < \mu^+$ such that $\alpha > \bar{\gamma}(n)$ and $\text{ran}(h_\alpha) \subseteq a$, and (3) $\alpha_{n+1} \in Z(f_n^0, f_n^1, \dots, f_n^n)$. s_a can be any subset of a if $(*)$ does not hold. Now suppose that $(*)$ holds. By induction on θ , define a_n^θ for $\theta < \xi$ and $n < \omega$ as follows. Put $a_n^0 = \phi$ for every $n < \omega$. If θ is an infinite limit ordinal, set $a_n^\theta = \bigcup_{\eta < \theta} a_n^\eta$ for all $n < \omega$. Assuming a_n^θ has been defined for all n , look for a $j < \omega$ such that (α) for $j < n < \omega$, $\text{o.t.}(a_n^\theta) \in \text{dom}(f_n^j)$ and $f_n^j(\text{o.t.}(a_n^\theta)) = 1$, and (β) for $\ell < j \leq n < \omega$, $\text{o.t.}(a_n^\theta) \in \text{dom}(f_n^\ell)$ and $f_n^\ell(\text{o.t.}(a_n^\theta)) = 0$. If there is no such j , set $a_n^{\theta+1} = a_n^\theta$ for every $n < \omega$. If there is one, it must be unique. Set $a_n^{\theta+1} = a_n^\theta$ for $n < j$, and $a_n^{\theta+1} = a_n^\theta \cup \{(e(a))(\theta)\}$ for $j \leq n < \omega$. Finally, set $s_a = \bigcup_{n < \omega} s_n$, where $s_n = \{(e(a_n))(\eta) : \eta \in \text{dom}(f_n^n) \cap \text{o.t.}(a_n) \text{ and } f_n^n(\eta) = 1\}$.

Now fix $A \in NG_{\kappa,\lambda}^*$ and $X \subseteq \lambda$. We must find $a \in A$ such that $s_a = X \cap a$. Let τ be a winning strategy for player II in the game $G_{\kappa,\lambda}(A)$. Define $k : \bigcup_{n < \omega} R_{n+1}^\mu \rightarrow P_\kappa(\lambda)$ as follows. Set $k(t) = \tau(x_{t(0)})$ for any $t \in R_1^\mu$. Given $0 < n < \omega$ and $t \in R_{n+1}^\mu$, define a_m and b_m for $m \leq n$ by: $a_0 = x_{t(0)}$, $a_m = b_{m-1} \cup x_{t(m)}$ for $m > 0$, and $b_m = \tau(a_0, \dots, a_m)$, and set $k(t) = b_n$.

Define W_n for $n < \omega$ by induction as follows. Set $W_0 = R_0^\mu$, $W_1 = R_1^\mu$ and

$$W_2 = \{t \in R_2^\mu : t(1) \in Z(\chi(X \cap k(t \upharpoonright 1), k(t \upharpoonright 1)))\}.$$

For $n \geq 2$, let W_{n+1} be the set of all $t \in R_{n+1}^\mu$ such that $t \upharpoonright n \in W_n$ and $t(n)$ belongs to $Z(f_0, \dots, f_{n-1})$, where $f_i = \chi(k(t \upharpoonright (i+1)), k(t \upharpoonright n))$ for $i < n-1$, and $f_{n-1} = \chi(X \cap k(t \upharpoonright n), k(t \upharpoonright n))$. Put $T_0 = \bigcup_{n < \omega} W_n$. For $0 < r < \omega$, define $\varphi_r : T_0 \cap (\bigcup_{r < q < \omega} R_q^\mu) \rightarrow \mu^+$ by $\varphi_r(t) = \rho_{t(r)}(\mu \cap k(t))$. Using Lemma 4, select $T_r \in \mathcal{T}_\mu$ and $\psi_r : T_r \cap R_r^\mu \rightarrow \mu^+$ for $0 < r < \omega$ so that $T_r \subseteq T_{r-1}$, $T_r \cap R_r^\mu = T_{r-1} \cap R_r^\mu$, and $\varphi_r(t) \leq \psi_r(t \upharpoonright r)$ for every $t \in T_r \cap (\bigcup_{r < q < \omega} R_q^\mu)$. Set $T = \bigcap_{r < \omega} T_r$.

Let C be the set of all γ with $\kappa < \gamma < \mu^+$ such that for any r with $0 < r < \omega$, and any $t \in T \cap R_r^\mu$ with $\text{ran}(t) \subseteq \gamma$, $\rho(\mu \cap k(t)) < \gamma$, $\psi_r(t) < \gamma$ and $\{\alpha < \gamma : t \cup \{(r, \alpha)\} \in T\}$ is cofinal in γ . Since C is a closed unbounded subset of μ^+ , there is $\gamma \in H$ such that $\text{ran}(\bar{\gamma}) \subseteq C$. Pick $y : \omega \rightarrow \mu^+$ so that $\{y \upharpoonright m : m < \omega\} \subseteq T$ and $y(0) < \bar{\gamma}(0) < y(1) < \bar{\gamma}(1) < \dots$. Set $a_n = k(y \upharpoonright (n+1))$ for $n < \omega$, and $a = \bigcup_{n < \omega} a_n$. Then for each $m < \omega$,

$$\bar{\gamma}(m) < y(m+1) \leq \rho(\mu \cap a_{m+1}) < \bar{\gamma}(m+1)$$

since $x_{y(m+1)} \subseteq a_{m+1}$ and $\text{ran}(y \upharpoonright (m+2)) \subseteq \bar{\gamma}(m+1)$. Hence

$$\rho(\mu \cap a) = \sup\{\rho(\mu \cap a_{m+1}) : m < \omega\} = \gamma.$$

For $0 < r < \omega$,

$$\rho_{y(r)}(\mu \cap a) = \sup\{\rho_{y(r)}(\mu \cap a_q) : r < q < \omega\} \leq \bar{\gamma}(r - 1)$$

since $\rho_{y(r)}(\mu \cap a_q) = \varphi_r(y \upharpoonright q) \leq \psi_r(y \upharpoonright r) < \bar{\gamma}(r - 1)$ whenever $r < q < \omega$. It follows that $y(r) =$ the least $\alpha < \mu^+$ such that $\alpha > \bar{\gamma}(r - 1)$ and $x_\alpha \subseteq a$, since $\delta \leq \rho_{y(r)}(a \cap \mu)$ for any $\delta < y(r)$ such that $x_\delta \subseteq a$. Define f_n^j for $j \leq n < \omega$ by $f_n^j = \chi(a_j, a_n)$ if $j < n$, and $f_n^n = \chi(X \cap a_n, a_n)$. Then $y(n + 1) \in Z(f_n^0, f_n^1, \dots, f_n^n)$ for all $n < \omega$. Finally, $s_a = \bigcup_{n < \omega} \{(e(a_n))(\eta) : \eta \in \text{o.t.}(a_n) \text{ and } f_n^n(\eta) = 1\} = \bigcup_{n < \omega} (X \cap a_n) = X \cap a$. \square

In the case when κ is the successor of a cardinal of cofinality ω , the assumption of Proposition 6 can be weakened.

Let $\nu > 0$ be a cardinal. For $A \subseteq P_\kappa(\lambda)$, the game $G_{\kappa,\lambda}^\nu(A)$ is defined similarly to $G_{\kappa,\lambda}(A)$, where now the choices are made from $P_\nu(\lambda)$.

Lemma 7 (Matet [7]). *Suppose κ is the successor of a cardinal ν of cofinality ω . Then for any $A \subseteq P_\kappa(\lambda)$, $A \in NG_{\kappa,\lambda}^*$ if and only if II has a winning strategy in the game $G_{\kappa,\lambda}^\nu(A)$.*

It is now straightforward to modify the proof of Proposition 6 so as to get the following.

Proposition 8. *Suppose that κ is the successor of a cardinal ν of cofinality ω , and $2^{<\nu} \leq \mu^+$ for some cardinal μ such that $\omega < \text{cf}(\mu) < \kappa < \mu \leq \lambda$. Then there is a sequence $\langle s_a : a \in P_\kappa(\lambda) \rangle$ with $s_a \subseteq a$ such that for any $X \subseteq \lambda$, $\{a : s_a = X \cap a\} \in NG_{\kappa,\lambda}^+$.*

REFERENCES

- [1] Cummings J., Foreman M., Magidor M., *Canonical structure in the universe of set theory, part one*, Ann. Pure Appl. Logic **129** (2004), 211–243.
- [2] Donder H.-D., Matet P., *Two cardinal versions of diamond*, Israel J. Math. **83** (1993), 1–43.
- [3] Foreman M., Magidor M., *Mutually stationary sequences of sets and the non-saturation of the non-stationary ideal on $P_\kappa(\lambda)$* , Acta Math. **186** (2001), 271–300.
- [4] Jech T.J., *Some combinatorial problems concerning uncountable cardinals*, Ann. Math. Logic **5** (1973), 165–198.
- [5] Kojman M., *The A, B, C of pcf: a companion to pcf theory, part I*, 1995, unpublished.
- [6] Matet P., *Concerning stationary subsets of $[\lambda]^{<\kappa}$* , in: Set Theory and its Applications (J. Steprāns and S. Watson, eds.), Lecture Notes in Mathematics 1401, Springer, Berlin, 1989, pp. 119–127.
- [7] Matet P., *Game ideals*, preprint.
- [8] Shioya M., *Splitting $P_\kappa\lambda$ into maximally many stationary sets*, Israel J. Math. **114** (1999), 347–357.

- [9] Solovay R.M., *Real-valued measurable cardinals*, in: Axiomatic Set Theory (D.S. Scott, ed.), Proceedings of Symposia in Pure Mathematics, vol. 13, part 1, American Mathematical Society, Providence, 1971, pp. 397–428.

UNIVERSITÉ DE CAEN - CNRS, LABORATOIRE DE MATHÉMATIQUES, BP 5186,
14032 CAEN CEDEX, FRANCE

E-mail: matet@math.unicaen.fr

(*Received* May 9, 2006, *revised* February 1, 2007)