

Yanzheng Duan; Bor-Luh Lin

Characterizations of L^1 -predual spaces by centerable subsets

Commentationes Mathematicae Universitatis Carolinae, Vol. 48 (2007), No. 2, 239--243

Persistent URL: <http://dml.cz/dmlcz/119654>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2007

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Characterizations of L^1 -predual spaces by centerable subsets

YANZHENG DUAN, BOR-LUH LIN

Abstract. In this note, we prove that a real or complex Banach space X is an L^1 -predual space if and only if every four-point subset of X is centerable. The real case sharpens Rao's result in [*Chebyshev centers and centerable sets*, Proc. Amer. Math. Soc. **130** (2002), no. 9, 2593–2598] and the complex case is closely related to the characterizations of L^1 -predual spaces by Lima [*Complex Banach spaces whose duals are L_1 -spaces*, Israel J. Math. **24** (1976), no. 1, 59–72].

Keywords: Chebyshev radius, centerable subsets and L^1 -predual spaces

Classification: Primary 41A65; Secondary 46B20

1. Introduction

Let X be a Banach space. For $a \in X$, $r > 0$, let $B[a, r] = \{x \in X : \|a - x\| \leq r\}$ and let $S(X) = \{x \in X : \|x\| = 1\}$. If A is a bounded subset of X , let

$$r(A, x) = \sup_{a \in A} \|x - a\|$$

and let

$$r(A) = \inf_{x \in X} r(A, x)$$

denote the Chebyshev radius of A . Let

$$\delta(A) = \sup\{\|a - b\| : a, b \in A\}$$

denote the diameter of A . Then $\delta(A) \leq 2r(A)$ for every bounded subset A of X .

Definition 1.1 ([4]). Let X be a Banach space and A a bounded subset of X . If $\delta(A) = 2r(A)$, then A is said to be centerable.

Definition 1.2 ([6]). A Banach space X whose dual X^* is isometrically isomorphic to $L^1(\mu)$ for some positive measure μ is called an L^1 -predual space.

L^1 -predual spaces are also called Lindenstrauss spaces. In [9], Lindenstrauss gave several characterizations of real L^1 -predual spaces using intersection properties of balls. In [7], Lima gave several characterizations of complex L^1 -predual spaces using intersection properties of balls.

A Banach space X is called a \mathcal{P}_1 space if X is norm-one complemented in every Banach space Z containing X . Let X be a real or complex Banach space. By Theorem 6.1 of [9] and Theorem 4.1 of [7], X is an L^1 -predual space if and only if X^{**} is a \mathcal{P}_1 space. By [4, p. 193], every bounded subset of a \mathcal{P}_1 space is centerable. In 1977, W.J. Davis [2] proved that the converse is true.

In 2002, Rao [10] proved that a real Banach space X is an L^1 -predual space if and only if every finite subset of X is centerable. In this note, we prove that if every four-point subset of a real or complex Banach space X is centerable, then X is an L^1 -predual space. The result for the real case sharpens Rao's result in [10] and our proof is different from Rao's. The result for the complex case is a new form of characterizations of L^1 -predual spaces. We also point out that it cannot be sharpened anymore, i.e., that every three-point subset of a real or complex Banach space X is centerable does not imply that X is an L^1 -predual space.

2. Main results

We first give a characterization of n -point subsets of Banach spaces to be centerable.

Proposition 2.1. *Let X be a real or complex Banach space and $n \geq 3$ be an integer. Then every n -point subset of X is centerable if and only if for every $r > 0$ and every family of pairwise intersecting closed balls $\{B[a_i, r]\}_{i=1}^n$ in X , $\bigcap_{i=1}^n B[a_i, r + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$.*

PROOF: \Rightarrow . Let $A = \{a_1, a_2, \dots, a_n\}$. Since $\{B[a_i, r]\}_{i=1}^n$ are pairwise intersecting, $\|a_i - a_j\| \leq 2r$ for all i, j . Hence, $2r(A) = \delta(A) \leq 2r$. So $r(A) \leq r$. Therefore, for any $\varepsilon > 0$, there exists $x_0 \in X$ such that $r(A, x_0) \leq r + \varepsilon$, which implies that $x_0 \in \bigcap_{i=1}^n B[a_i, r + \varepsilon]$.

\Leftarrow . For any n -point subset $A = \{a_1, a_2, \dots, a_n\}$ of X , to prove that $\delta(A) = 2r(A)$, it suffices to prove that $\delta(A) \geq 2r(A)$. In fact, let $r = \frac{1}{2}\delta(A)$. Then the family $\{B[a_i, r]\}_{i=1}^n$ are pairwise intersecting. Hence, for any $\varepsilon > 0$, $\bigcap_{i=1}^n B[a_i, r + \varepsilon] \neq \emptyset$. Let $x_0 \in \bigcap_{i=1}^n B[a_i, r + \varepsilon]$, then $\|a_i - x_0\| \leq r + \varepsilon$, $i = 1, 2, \dots, n$, i.e., $r(A, x_0) \leq r + \varepsilon$. Hence, $r(A) = \inf_{x \in X} r(A, x) \leq r = \frac{1}{2}\delta(A)$. \square

Next theorem is due to Lindenstrauss [9].

Theorem 2.2 ([9]). *Let X be a real Banach space and $n \geq 3$ an integer. Then the following statements are equivalent.*

- (1) *For every $r > 0$ and every family of pairwise intersecting closed balls*

- $\{B[a_i, r]\}_{i=1}^n$ in X , $\bigcap_{i=1}^n B[a_i, r + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$.
- (2) For every $r > 0$ and every family of pairwise intersecting closed balls $\{B[a_i, r]\}_{i=1}^n$ in X , $\bigcap_{i=1}^n B[a_i, r] \neq \emptyset$.
- (3) For every family of pairwise intersecting closed balls $\{B[a_i, r_i]\}_{i=1}^n$ in X , $\bigcap_{i=1}^n B[a_i, r_i] \neq \emptyset$.
- (4) For every family of pairwise intersecting closed balls $\{B[a_i, r_i]\}_{i=1}^n$ in X , $\bigcap_{i=1}^n B[a_i, r_i + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$.

Combining Proposition 2.1 and Theorem 2.2, we have the following theorem.

Theorem 2.3. *Let X be a real Banach space and $n \geq 3$ an integer. Then every n -point subset of X is centerable if and only if one of the four conditions in Theorem 2.2 holds.*

The following theorem is a special case of Lindenstrauss’s Theorem 4.1 in [9].

Theorem 2.4 ([9]). *Let X be a real Banach space and $n \geq 3$ an integer. Then the following statements are equivalent.*

- (1) For every family of pairwise intersecting closed balls $\{B[a_i, r_i]\}_{i=1}^4$ in X , $\bigcap_{i=1}^4 B[a_i, r_i] \neq \emptyset$.
- (2) For every family of pairwise intersecting closed balls $\{B[a_i, r_i]\}_{i=1}^n$ in X , $\bigcap_{i=1}^n B[a_i, r_i] \neq \emptyset$.

Following Theorem 2.3 and Theorem 2.4, we have the following.

Theorem 2.5. *Let X be a real Banach space. Then every four-point subset of X is centerable if and only if every finite subset of X is centerable.*

In [1], P. Bandyopadhyay and T.S.S.R.K. Rao proved the following result.

Theorem 2.6 ([1, Theorem 3.9]). *Let X be a real or complex L^1 -predual space. Then any compact subset A of X is centerable.*

Now we are ready to give characterizations of real L^1 -predual spaces by centerable subsets.

Theorem 2.7. *Let X be a real Banach space. Then the following statements are equivalent.*

- (1) X is an L^1 -predual space.
- (2) Every four-point subset of X is centerable.
- (3) Every finite subset of X is centerable.
- (4) Every compact subset of X is centerable.

PROOF: (4) \Rightarrow (3) \Rightarrow (2). Trivial. (2) \Rightarrow (1). Following Theorem 2.3, Theorem 2.5 and Theorem 6.1 in [9]. (1) \Rightarrow (4). Following Theorem 2.6. □

In order to give a similar characterization of complex L^1 -predual spaces by centerable subsets, we need Lima’s results [7], [8] about characterizations of complex L^1 -predual spaces.

Definition 2.8 ([5]). A family of closed balls $\{B[a_i, r_i]\}_{i \in I}$ in a complex (real) Banach space X is said to have the weak intersection property if for any $f \in S(X^*)$, $\bigcap_{i \in I} B[f(a_i), r_i] \neq \emptyset$ in $\mathbb{C}(\mathbb{R})$.

Definition 2.9 ([7]). Let $n \geq 3$ be an integer. We say that a real or complex Banach space X is an $E(n)$ -space if for every family $\{B[a_i, r_i]\}_{i=1}^n$ of n closed balls in X with the weak intersection property, $\bigcap_{i=1}^n B[a_i, r_i] \neq \emptyset$.

Theorem 2.10 ([8, Corollary 2.5]). *Let X be a real or complex Banach space and let $n \geq 3$ be an integer. Then the following statements are equivalent.*

- (1) For every $r > 0$ and every family of n closed balls $\{B[a_i, r]\}_{i=1}^n$ in X such that any three of them have nonempty intersection, $\bigcap_{i=1}^n B[a_i, r + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$.
- (2) For every family of n closed balls $\{B[a_i, r_i]\}_{i=1}^n$ in X such that any three of them have nonempty intersection, $\bigcap_{i=1}^n B[a_i, r_i + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$.

Theorem 2.11 ([7, Corollary 4.3]). *Let X be a complex Banach space. If for every family of 4 closed balls $\{B[a_i, r_i]\}_{i=1}^4$ in X such that any three of them have nonempty intersection, $\bigcap_{i=1}^4 B[a_i, r_i + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$, then X is an $E(n)$ -space for all $n \geq 3$.*

Theorem 2.12 ([7, Theorem 4.1]). *Let X be a complex Banach space. Then the following statements are equivalent.*

- (1) X is an L^1 -predual space.
- (2) X is an $E(n)$ -space for all $n \geq 3$.

Theorem 2.13. *Let X be a complex Banach space. Then the following statements are equivalent.*

- (1) X is an L^1 -predual space.
- (2) Every four-point subset of X is centerable.
- (3) Every finite subset of X is centerable.
- (4) Every compact subset of X is centerable.

PROOF: (1) \Rightarrow (4). Following Theorem 2.6. (4) \Rightarrow (3) \Rightarrow (2). Trivial.

(2) \Rightarrow (1). Let $r > 0$ and let $\{B[a_i, r]\}_{i=1}^4$ be a family of four closed balls such that any three of them intersect. Since $\{B[a_i, r]\}_{i=1}^4$ is pairwise intersecting and $\{a_1, a_2, a_3, a_4\}$ is centerable, by Theorem 2.1, $\bigcap_{i=1}^4 B[a_i, r + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$. Hence, by Theorem 2.10, for every family of 4 closed balls $\{B[a_i, r_i]\}_{i=1}^4$ in X such that any three of them have nonempty intersection, $\bigcap_{i=1}^4 B[a_i, r_i + \varepsilon] \neq \emptyset$ for all $\varepsilon > 0$. Following Theorem 2.11, X is an $E(n)$ -space for all $n \geq 3$. Therefore by Theorem 2.12, X is an L^1 -predual space. \square

Remark 2.14. Let us show that centerability of all three-point subsets of X does not imply that X is an L^1 -predual space. Consider the real or complex space ℓ_1 . Since every three pairwise intersecting closed balls in \mathbb{R} or \mathbb{C} intersect (see [3, p.65]), the same holds also for ℓ_1 by Theorem 4.6(c) in [9]. Hence, by Theorem 2.3, every three-point set in ℓ_1 is centerable. On the other hand, ℓ_1 is not an L^1 -predual space since $\ell_1^* = \ell_\infty$.

Acknowledgment. The authors are very grateful to the referee for reading the paper very thoroughly and giving many useful suggestions.

REFERENCES

- [1] Bandyopadhyay P., Rao T.S.S.R.K., *Central subspaces of Banach spaces*, J. Approx. Theory **103** (2000), 206–222.
- [2] Davis W.J., *A characterization of \mathcal{P}_1 spaces*, J. Approx. Theory **21** (1977), 315–318.
- [3] Hanner O., *Intersections of translates of convex bodies*, Math. Scand. **4** (1956), 65–87.
- [4] Holmes R.B., *A course on optimization and best approximation*, Lecture Notes in Math., Vol. 257, Springer, Berlin, 1972.
- [5] Hustad O., *Intersection properties of balls in complex Banach spaces whose dual are L^1 -spaces*, Acta Math. **132** (1974), 282–313.
- [6] Lacey H.E., *The isometric theory of classical Banach spaces*, Grundlehren Math. Wiss., Band 208, Springer, New York-Heidelberg, 1974.
- [7] Lima A., *Complex Banach spaces whose duals are L_1 -spaces*, Israel J. Math. **24** (1976), no. 1, 59–72.
- [8] Lima A., *Intersection properties of balls and subspaces in Banach spaces*, Trans. Amer. Math. Soc. **227** (1977), 1–62.
- [9] Lindenstrauss J., *Extension of compact operators*, Mem. Amer. Math. Soc., Vol. 48, Providence, 1964.
- [10] Rao T.S.S.R.K., *Chebyshev centers and centerable sets*, Proc. Amer. Math. Soc. **130** (2002), no. 9, 2593–2598.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY, IA 52242-1419, U.S.A.

E-mail: yduan@math.uiowa.edu
bllin@math.uiowa.edu

(Received July 6, 2006, revised October 24, 2006)