

Arkady G. Leiderman; Igor V. Protasov

Cellularity of a space of subgroups of a discrete group

Commentationes Mathematicae Universitatis Carolinae, Vol. 49 (2008), No. 3, 519–522

Persistent URL: <http://dml.cz/dmlcz/119740>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2008

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Cellularity of a space of subgroups of a discrete group

A. LEIDERMAN, I.V. PROTASOV

Abstract. Given a discrete group G , we consider the set $\mathcal{L}(G)$ of all subgroups of G endowed with topology of pointwise convergence arising from the standard embedding of $\mathcal{L}(G)$ into the Cantor cube $\{0, 1\}^G$. We show that the cellularity $c(\mathcal{L}(G)) \leq \aleph_0$ for every abelian group G , and, for every infinite cardinal τ , we construct a group G with $c(\mathcal{L}(G)) = \tau$.

Keywords: space of subgroups, cellularity, Shanin number

Classification: 54B20, 54A25

Let G be a discrete group and $\exp G$ the set of all subsets of G . We identify $\exp G$ with the Cantor cube $\{0, 1\}^G$ and consider the set $\mathcal{L}(G)$ of all subgroups of G as a subspace of $\{0, 1\}^G$. Let $\mathcal{F}(G)$ be the family of all finite subsets of G . Given any $F, H \in \mathcal{F}(G)$, we put

$$[F, G \setminus H] = \{A \in \mathcal{L}(G) : F \subseteq A \subseteq G \setminus H\}.$$

Then the family $\{[F, G \setminus H] : F, H \in \mathcal{F}(G)\}$ forms a base of the topology in $\mathcal{L}(G)$. It is easy to see that $\mathcal{L}(G)$ is a zero-dimensional compact space.

We denote by $c(X)$ the *cellularity* of a topological space X . Remind that $c(X)$ is the supremum of cardinalities of disjoint families of open subsets of X .

We say that a topological space X has *Shanin number* ω (see [4] or [1, Problem 2.7.11]) if any uncountable family \mathcal{U} of non-empty open subsets of X has an uncountable subfamily $\mathcal{V} \subseteq \mathcal{U}$ with $\bigcap \mathcal{V} \neq \emptyset$. Evidently, if a space X has Shanin number ω then $c(X) \leq \aleph_0$.

Given any topological group G , we denote by $L(G)$ the space of all closed subgroups of G endowed with the Vietoris topology. By [3, Theorem 2], $c(L(G)) \leq \aleph_0$ for every compact group G .

Theorem 1. *For every discrete abelian group G , the space $\mathcal{L}(G)$ has Shanin number ω , in particular, $c(\mathcal{L}(G)) \leq \aleph_0$.*

PROOF: By [6, Theorem 24.1], G is a subgroup of some divisible group D . By [6, Theorem 23.1], D is a direct sum $\bigoplus_{\alpha \in I} G_\alpha$ of some family of countable groups.

The first author was partially supported by the Israel Science Foundation Grant no. 508/06.

Thus we suppose that G is a subgroup of $\bigoplus_{\alpha \in I} G_\alpha$. Let $\{U_\beta : \beta \in J\}$ be an uncountable family of non-empty open subsets of $\mathcal{L}(G)$. For every $\beta \in J$, we pick finite subsets F_β, H_β of G such that $\emptyset \neq [F_\beta, G \setminus H_\beta] \subseteq U_\beta$, and put

$$K_\beta = \{\alpha \in I : \text{pr}_\alpha g \neq 0 \text{ for some } g \in F_\beta \cup H_\beta\}.$$

Then $\{K_\beta : \beta \in J\}$ is an uncountable family of finite subsets of I . By the Δ -Lemma, there exist an uncountable subset $J_1 \subseteq J$ and a finite subset $K \subseteq I$ such that $K_\beta \cap K_\gamma = K$ for all distinct $\beta, \gamma \in J_1$. We put

$$G_K = \{g \in G : \text{pr}_\alpha g = 0 \text{ for every } \alpha \notin K\}.$$

For every subset X of G , we denote by $\langle X \rangle$ the subgroup of G generated by X . Every subgroup of a finitely generated abelian group is finitely generated, hence it follows that for every $\beta \in J_1$, the subgroup $\langle F_\beta \rangle \cap G_K$ is finitely generated. Since G_K is a countable group, there are countably many finitely generated subgroups of G_K , and therefore there exist an uncountable subset $J_2 \subseteq J_1$ and a subgroup $A \subseteq G_K$ such that $\langle F_\beta \rangle \cap G_K = A$ for every $\beta \in J_2$. We put $S = \langle \bigcup_{\beta \in J_2} F_\beta \rangle$ and prove that $S \in \bigcap_{\beta \in J_2} [F_\beta, G \setminus H_\beta]$. It suffices to show that $H_\beta \cap S = \emptyset$ for every $\beta \in J_2$.

Pick any element $g \in S$ and fix $\beta \in J_2$. Clearly, if $g = 0$ then $g \notin H_\beta$ because of $0 \in \langle F_\beta \rangle$ and $\emptyset \neq [F_\beta, G \setminus H_\beta]$. If $g \neq 0$, we can write g as

$$g = g_{\alpha_1} + \dots + g_{\alpha_n},$$

where $g_{\alpha_i} \in G_{\alpha_i} \setminus \{0\}$ for every $i \in \{1, \dots, n\}$. If $\alpha_i \notin K_\beta$ for some $i \in \{1, \dots, n\}$ then, by definition of K_β , we have $g \notin H_\beta$.

Now consider the possibility of $\{\alpha_1, \dots, \alpha_n\} \subseteq K_\beta$. We claim that in this case $g \in \langle F_\beta \rangle$. Let us assume the contrary, $g \notin \langle F_\beta \rangle$.

By the choice of J_2 , we have $\langle F_\beta \rangle \cap G_K = \langle F_\gamma \rangle \cap G_K$ for every distinct $\beta, \gamma \in J_2$, therefore by definition of S , there exist $\gamma \in J_2 \setminus \{\beta\}$ and $j \in \{1, \dots, n\}$ such that $\alpha_j \in K_\gamma \setminus K$. On the other hand, $K_\beta \cap K_\gamma = K$, hence $\alpha_j \notin K_\beta$, and we get a contradiction with $\{\alpha_1, \dots, \alpha_n\} \subseteq K_\beta$. So, $g \in \langle F_\beta \rangle$ which implies that $g \notin H_\beta$. □

Remind that a Hausdorff compact space X is called *dyadic* if X is a continuous image of some Cantor cube. If G is a compact abelian group, by [5, Theorems 3, 4], the space $L(G)$ is dyadic if and only if the weight $w(G) \leq \aleph_1$. A natural question arises to characterize discrete groups G for which $\mathcal{L}(G)$ is a dyadic space.

Theorem 2. *For a discrete abelian group G , the space $\mathcal{L}(G)$ is dyadic if and only if $|G| \leq \aleph_1$.*

PROOF: By [2, Theorem 3], for every compact abelian group G the space $L(G)$ is homeomorphic to $\mathcal{L}(\hat{G})$, where \hat{G} is the dual group to G . Therefore, by Pontryagin's duality and by above-mentioned [5, Theorems 3, 4], the space $\mathcal{L}(G)$ is dyadic if and only if $|G| \leq \aleph_1$. \square

Our last theorem shows that, in contrast to Theorem 1, there are non-abelian groups G with arbitrary large cellularity $c(\mathcal{L}(G))$.

Theorem 3. *For every infinite cardinal τ there exists a discrete group G such that $c(\mathcal{L}(G)) = |G| = \tau$.*

PROOF: Let F be a free group generated by the set $\{x_\alpha, y_\alpha : \alpha < \tau\}$ of free generators. We shall define a group G as the quotient $G = F/N$, where N is a certain normal subgroup of the free group F containing all the words $x_\alpha^2, y_\alpha^2, x_\beta x_\alpha x_\beta y_\alpha, \alpha < \beta < \tau$.

We fix $\alpha < \tau$ and put

$$A = \langle a_\alpha \rangle \times \langle b_\alpha \rangle, \quad B = \otimes_{\alpha < \beta < \tau} \langle c_\beta \rangle,$$

where $\langle a_\alpha \rangle, \langle b_\alpha \rangle, \langle c_\beta \rangle$ are the cyclic subgroups of order 2. Then we define a group G_α as the semidirect product $G_\alpha = A \rtimes B$ with $c_\beta a_\alpha c_\beta = b_\alpha$ for all $\alpha < \beta < \tau$. Let $f_\alpha : F \rightarrow G_\alpha$ be a homomorphism defined by the rule

$$\begin{aligned} f_\alpha(x_\lambda) &= f_\alpha(y_\lambda) = 1 \quad \text{for all } \lambda < \alpha, \\ f_\alpha(x_\alpha) &= a_\alpha, \quad f_\alpha(y_\alpha) = b_\alpha, \\ f_\alpha(x_\beta) &= f_\alpha(y_\beta) = c_\beta \quad \text{for all } \alpha < \beta < \tau. \end{aligned}$$

Finally, we define

$$N = \bigcap \{ \text{Ker } f_\alpha : \alpha < \tau \}, \quad \text{and } G = F/N.$$

Clearly, $f_\alpha(x_\lambda^2) = f_\alpha(y_\lambda^2) = 1$ for every $\lambda < \tau$, and it easily could be seen that $f_\alpha(x_\beta x_\lambda x_\beta) = f_\alpha(y_\lambda)$ for all $\lambda < \beta < \tau$. Hence, N is a normal subgroup of the free group F containing all the words $x_\alpha^2, y_\alpha^2, x_\beta x_\alpha x_\beta y_\alpha, \alpha < \beta < \tau$, but $x_\alpha, y_\alpha, x_\alpha y_\alpha^{-1} \notin N$. It follows that $x_\alpha N \neq 1, y_\alpha N \neq 1, x_\alpha N \neq y_\alpha N$ in the quotient group $G = F/N$. In the sequel we denote the cosets wN , i.e. the elements of the group G , simply by w .

In order to finish the proof of Theorem 3 we show that $c(\mathcal{L}(G)) = |G| = \tau$. Let us consider the family $\{[\{x_\alpha\}, G \setminus \{y_\alpha\}] : \alpha < \tau\}$ of non-empty open subsets of $\mathcal{L}(G)$, and show that this family is disjoint. We assume the contrary and choose α, β such that $\alpha < \beta < \tau$ and

$$[\{x_\alpha\}, G \setminus \{y_\alpha\}] \cap [\{x_\beta\}, G \setminus \{y_\beta\}] \neq \emptyset.$$

Then the subgroup $\langle x_\alpha, x_\beta \rangle$ generated by x_α, x_β is in the above intersection. On the other hand, $y_\alpha = x_\beta x_\alpha x_\beta$, so

$$y_\alpha \in \langle x_\alpha, x_\beta \rangle, \quad \langle x_\alpha, x_\beta \rangle \in [\{x_\beta\}, G \setminus \{y_\alpha\}]$$

and we get a contradiction. \square

Question. Does there exist a nilpotent group G such that $c(\mathcal{L}(G)) > \aleph_0$?

REFERENCES

- [1] Engelking R., *General Topology*, PWN, Warszawa, 1985.
- [2] Protasov I.V., *Dualisms of topological Abelian groups*, Ukrain. Mat. Zh. **31** (1979), 207–211; English translation: Ukrainian Math. J. **31** (1979), 164–166.
- [3] Protasov I.V., *The Suslin number of a space of subgroups of a locally compact group*, Ukrain. Mat. Zh. **40** (1988), 654–658; English translation: Ukrainian Math. J. **40** (1988), 559–562.
- [4] Shanin N.A., *On product of topological spaces*, Trudy Mat. Inst. Akad. Nauk SSSR **24** (1948), 1–112 (in Russian).
- [5] Tsybenko Yu.V., *Dyadicity of a space of subgroups of a topological group*, Ukrain. Mat. Zh. **38** (1986), 635–639.
- [6] Fuchs L., *Infinite Abelian Groups*, Vol. 1, Academic Press, New York and London, 1970.

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, BEER SHEVA, ISRAEL

E-mail: arkady@math.bgu.ac.il

TARAS SHEVCHENKO KYIV NATIONAL UNIVERSITY, UKRAINE

E-mail: protasov@unicyb.kiev.ua

(Received September 20, 2007, revised March 21, 2008)