

Botir Zakirov

Abstract characterization of Orlicz-Kantorovich lattices associated with an L_0 -valued measure

Commentationes Mathematicae Universitatis Carolinae, Vol. 49 (2008), No. 4, 595--610

Persistent URL: <http://dml.cz/dmlcz/119748>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2008

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Abstract characterization of Orlicz-Kantorovich lattices associated with an L_0 -valued measure

BOTIR ZAKIROV

Abstract. An abstract characterization of Orlicz-Kantorovich lattices constructed by a measure with values in the ring of measurable functions is presented.

Keywords: Orlicz-Kantorovich lattice, vector-valued measure, Orlicz function

Classification: 46B42, 46E30, 46G10

1. Introduction

The development of the theory of integration for measures with values in the algebra L_0 of all real measurable functions has inspired the study of Banach L_0 -modules of measurable functions. The theory of L_p -spaces associated with a vector-valued measure is given in monographs [7], [10]. Precise description of Orlicz-Kantorovich spaces $L_M(\nabla, m)$ associated with a complete Boolean algebra ∇ , an N -function M and an L_0 -valued measure m defined on ∇ is given in [13], [14], [15]. Spaces $L_M(\nabla, m)$ are important examples of Banach-Kantorovich spaces (see, for example, [7], [8], [4] for definition and basic properties).

The abstract characterization of Banach lattices isomorphic to L_p -spaces is well known (see, for example, [9]). The same is done for Orlicz spaces in [2]. One can expect similar results for Banach L_0 -modules $L_p(\nabla, m)$ and $L_M(\nabla, m)$. This problem was considered in [7] for $L_p(\nabla, m)$. Here we solve this problem for $L_M(\nabla, m)$.

We use terminology and notations from the theory of Boolean algebras from [11], the theory of vector lattices from [12], [5], the theory of vector integration from [10], [8], the theory of lattice-normed spaces from [7], [8], and also terminology for Orlicz-Kantorovich lattices from [13], [14].

2. Preliminaries

Let E be a vector lattice, E_+ be the set of all non-negative elements from E . Any element $x \in E$ can be uniquely decomposed as $x = x_+ - x_-$, where $x_+, x_- \in E_+$ and $x_+ \wedge x_- = 0$. The element $|x| = x_+ + x_-$ is called the absolute value of x , and elements x_+ and x_- are called the positive and negative parts of x , respectively. Elements $x, y \in E$ are disjoint iff $|x| \wedge |y| = 0$.

Let $u \in E_+$. If no non-zero element is disjoint with u , then u is called a weak order unit. Fix some weak order unit (if it exists) \mathbf{I} . An element $e \in E_+$ is called a *unitary element* if $e \wedge (\mathbf{I} - e) = 0$. The set $\nabla(E)$ of all unitary elements from E is a Boolean algebra with respect to the order induced from E . A complement in $\nabla(E)$ is given as $\mathbf{I} - e$.

A vector lattice is called *complete* (σ -*complete*) if $\sup A$ and $\inf A$ exist for every (countable) bounded subset A .

Let E be a σ -complete vector lattice with weak unit \mathbf{I} . For every $x \in E$, the element $e_x := \sup\{\mathbf{I} \wedge (n|x|) : n \in \mathbb{N}\}$ is unitary. It is called the *support* of x . Define $e_t^x := e_{(t\mathbf{I}-x)_+}$. The set $\{e_t^x\}_{t \in \mathbb{R}}$ is called a family of *spectral unitary elements* of x . If $x_n \in E$, $x = \inf x_n$, then $e_t^x = \sup_{n \geq 1} e_t^{x_n}$ for all $t \in \mathbb{R}$ (see [12, Lemma IV.10.2]).

Suppose that a σ -complete vector lattice E is of countable type, i.e. every set of non-zero mutually disjoint elements from E is at most countable. Then E is order complete. Moreover, for every bounded set $A \subset E$, there exists a subset $\{x_n\}_{n=1}^\infty \subset A$, such that $\sup A = \sup_{n \geq 1} x_n$.

A Boolean algebra ∇ is called *complete* (σ -*complete*) if $\sup A$ exists for every (countable) subset $A \subset \nabla$. Let E be a complete (σ -complete) vector lattice with a weak unit. Then, the Boolean algebra $\nabla(E)$ (see above) is complete (σ -complete). Evidently, the operation \sup is the same in E and $\nabla(E)$. The decomposition of a unit in Boolean algebra is an arbitrary set $(e_\alpha)_{\alpha \in A}$ satisfying $\sup_{\alpha \in A} e_\alpha = \mathbf{I}$, $e_\alpha \neq 0$, $e_\alpha \wedge e_\beta = 0$, $\alpha \neq \beta$, $\alpha, \beta \in A$.

Let (Ω, Σ, μ) be a σ -finite measurable space. Let $L_0 = L_0(\Omega)$ be the algebra of all real measurable functions on (Ω, Σ, μ) (functions equal a.e. are identified). L_0 is a complete vector lattice with respect to the natural order ($x \geq y$ if $x(\omega) \geq y(\omega)$ for almost all ω). The weak order unit is $\mathbf{1}(\omega) \equiv 1$. The set $\nabla(\Omega)$ of all idempotents in L_0 is a complete Boolean algebra.

The support e_x of an element $x \in L_0$ is also denoted by $s(x)$. It is clear that $s(x) = \chi_{\{|x|>0\}}$. Also, $xs(x) = x$. If $xy = 0$ then $s(x)y = 0$. In particular, $|x| \wedge |y| = 0$ if and only if $s(x)s(y) = 0$.

Let $e = \chi_A \in \nabla(\Omega)$. Set $e\Omega = (A, \Sigma_A, \mu)$, where $\Sigma_A = \{B \cap A : B \in \Sigma\}$. The rings $L_0(e\Omega)$ and $eL_0(\Omega)$ can be canonically identified. The Boolean algebras $\nabla(e\Omega)$ and $e\nabla(\Omega) = \{g \in \nabla(\Omega) : g \leq e\}$ can also be identified canonically. Define the map $\mu : \nabla(\Omega) \rightarrow [0, \infty]$ as $\mu(e) = \mu(A)$ if $e = \chi_A \in \nabla(\Omega)$. Obviously, μ is a strongly positive (i.e. $\mu(e) > 0$ for $e \neq 0$) countably additive σ -finite measure on $\nabla(\Omega)$.

A sequence $\{x_n\} \subset L_0$ converges locally with respect to a measure μ to the element $x \in L_0$ (notation: $x_n \xrightarrow{l, \mu} x$) if for any $A \in \Sigma$ with $\mu(A) < \infty$ the sequence $x_n \chi_A$ converges with respect to the measure to $x \chi_A$. If $\mu(\Omega) < \infty$, then local convergence with respect to the measure coincides with convergence with respect to the measure. There exists a countable set of non-zero disjoint idempotents

$\{e_n\} \subset \nabla(\Omega)$ such that $\sup_{n \geq 1} e_n = \mathbf{1}$ and $\mu(e_n) < \infty$. The algebra $L_0(\Omega)$ is canonically identified with the direct product $\prod_{n=1}^{\infty} L_0(e_n\Omega)$. Local convergence with respect to the measure is now identified with convergence of each coordinate with respect to the measure. $L_0(\Omega)$ with this topology is a complete metrizable topological vector lattice.

Now we define a Banach-Kantorovich space for an L_0 -valued norm.

Let E be a vector space over the field \mathbb{R} . A mapping $\|\cdot\| : E \rightarrow L_0$ is said to be a *vector (L_0 -valued) norm* if it satisfies the following axioms:

1. $\|x\| \geq 0$, and $\|x\| = 0 \Leftrightarrow x = 0$ ($x \in E$);
2. $\|\lambda x\| = |\lambda| \|x\|$ ($\lambda \in \mathbb{R}, x \in E$);
3. $\|x + y\| \leq \|x\| + \|y\|$ ($x, y \in E$).

A norm $\|\cdot\|$ is called *decomposable* or *Kantorovich* if the following property holds:

Property 1. If $e_1, e_2 \geq 0$ and $\|x\| = e_1 + e_2$, then there exist $x_1, x_2 \in E$ such that $x = x_1 + x_2$ and $\|x_k\| = e_k$ ($k = 1, 2$).

If property 1 is valid only for disjoint elements $e_1, e_2 \in L_0$, the norm is called *disjointly decomposable* or, briefly, *d-decomposable*.

A pair $(E, \|\cdot\|)$ is called a *lattice-normed space* (shortly, LNS). If the norm $\|\cdot\|$ is decomposable (*d-decomposable*), then so is the space $(E, \|\cdot\|)$.

A sequence $\{x_n\} \subset E$ (*bo*)-converges to $x \in E$ if the sequence $\{\|x_n - x\|\}$ (*o*)-converges to 0 in L_0 . A sequence $\{x_n\}$ is said to be a (*bo*)-*Cauchy sequence* if $\sup_{n, k \geq m} \|x_n - x_k\| \xrightarrow{(o)} 0$ as $m \rightarrow \infty$. An LNS is called (*bo*)-*complete* if any (*bo*)-Cauchy sequence (*bo*)-converges. A *Banach-Kantorovich space* (shortly, BKS) is a *d-decomposable* (*bo*)-complete LNS. It is well known that every BKS is a decomposable LNS.

Suppose that $(E, \|\cdot\|)$ is an LNS and a vector lattice simultaneously. The norm $\|\cdot\|$ is called *monotone* if $|x| \leq |y|$ implies that $\|x\| \leq \|y\|$. BKS with a monotone norm is called a *Banach-Kantorovich lattice*.

Let E be an L_0 -module. It is called a *normal L_0 -module* if

1. for any non-zero $e \in \nabla(\Omega)$, there exists $x \in E$ such that $ex \neq 0$;
2. for any decomposition of unit $\{e_n\}_{n=1}^{\infty} \subset \nabla(\Omega)$ and any $\{x_n\}_{n=1}^{\infty} \subset E$, there exists $x \in E$ such that $e_n x = e_n x_n$ for all n ;
3. if $x \in E$ and $\{e_n\} \in \nabla(\Omega)$ is a disjoint sequence, then $e_n x = 0$ for all n implies that $(\sup_{n \geq 1} e_n)x = 0$.

An ordered normal L_0 -module E is called an *L_0 -vector lattice* if for any $x, y, z \in E, \lambda \in L_0, \lambda \geq 0$, the inequality $x \leq y$ implies $x + z \leq y + z$ and $\lambda x \leq \lambda y$. The simplest example of an L_0 -vector lattice is L_0 itself considered as a module over L_0 .

Lemma 2.1. *Let E be an L_0 -vector lattice, $x, y \in E, x \geq 0, y \geq 0, e, g \in \nabla(\Omega), eg = 0$. Then the elements ex and gy are disjoint.*

PROOF: Let $z = ex \wedge gy$. Since $ex \geq 0, gy \geq 0$, we have $z \geq 0$ and it follows that $0 \leq ez \leq egy = 0$, i.e. $ez = 0$. Further, $0 \leq (1 - e)z \leq (1 - e)ex = 0$, and therefore $(1 - e)z = 0$, i.e. $z = ez = 0$. \square

Remark 2.2. If $x, z \in E, e \in \nabla(\Omega)$ and $0 \leq z \leq ex$, then $z = ez$.

Lemma 2.3. Let E be an L_0 -vector lattice with a weak order unit \mathbf{I} . Then

- (i) $\lambda\mathbf{I} \neq 0$ for any non-zero $\lambda \in L_0$;
- (ii) $(\lambda\mathbf{I}) \vee 0 = \lambda_+\mathbf{I}$ for any $\lambda \in L_0$.

PROOF: (1) Let $\lambda \in L_0, \lambda \geq 0, \lambda \neq 0$. Then $\lambda \geq \varepsilon e$ for some $e \in \nabla(\Omega), e \neq 0, \varepsilon > 0$. Hence, $\lambda\mathbf{I} \geq \varepsilon e\mathbf{I}$. Let us show that $e\mathbf{I} \neq 0$. Select $x \in E$ such that $ex \neq 0$. Let $x = x_+ - x_-$. Either $ex_+ \neq 0$ or $ex_- \neq 0$. Let $ex_+ \neq 0$. Set $z = (ex_+) \wedge \mathbf{I} \neq 0$. If $e\mathbf{I} = 0$, then by Remark 2.2, $0 \leq ez = z \leq e\mathbf{I} = 0$, i.e. $z = 0$. Therefore, $e\mathbf{I} \neq 0$ and $\lambda\mathbf{I} \neq 0$. Let now λ be an arbitrary element from L_0 , and $\lambda = \lambda_+ - \lambda_-$, moreover $\lambda_- \neq 0$. Suppose $\lambda_+\mathbf{I} - \lambda_-\mathbf{I} = 0$. Then $\lambda_-\mathbf{I} = s(\lambda_-)\lambda_-\mathbf{I} = s(\lambda_-)\lambda_+\mathbf{I} = 0$, which is not the case.

(2) It is clear that $\lambda_+\mathbf{I} \geq 0$ and $\lambda_+\mathbf{I} - \lambda\mathbf{I} = \lambda_-\mathbf{I} \geq 0$, i.e. $\lambda_+\mathbf{I} \geq \lambda\mathbf{I} \vee 0$. On the other hand, if $a = \lambda\mathbf{I} \vee 0$, then

$$a \geq s(\lambda_+)a \geq s(\lambda_+)\lambda\mathbf{I} = \lambda_+\mathbf{I}.$$

Hence, $\lambda_+\mathbf{I} = (\lambda\mathbf{I}) \vee 0$. \square

Submodules and morphisms are defined in a usual way.

Proposition 2.4. Let E be an L_0 -vector lattice and \mathbf{I} be a weak order unit in E . Then $N = \{\lambda\mathbf{I} : \lambda \in L_0\}$ is a normal L_0 -submodule in E and a vector sublattice in E , canonically isomorphic to L_0 . Moreover, $N(\Omega) = \{e\mathbf{I} : e \in \nabla(\Omega)\}$ is a σ -Boolean subalgebra in $\nabla(E)$.

PROOF: Only the second assertion needs to be proved. It follows from Lemma 2.1 that $N(\Omega)$ is a Boolean subalgebra of ∇ .

Let $\{e_n\} \subset \nabla(\Omega)$ and $e = \sup e_n$. If $g \in \nabla$ and $g \geq e_n\mathbf{I}$, then $\mathbf{I} - g \leq (1 - e_n)\mathbf{I}$, and therefore $e_n(\mathbf{I} - g) \leq e_n(1 - e_n)\mathbf{I} = 0$. Hence, $e_n(\mathbf{I} - g) = 0$. Then $e(\mathbf{I} - g) = 0$ because E is normal. Hence, $e\mathbf{I} = \sup_{n \geq 1} e_n\mathbf{I}$. This means that $N(\Omega)$ is a σ -subalgebra in $\nabla(E)$. \square

Proposition 2.5. Let E be a σ -complete L_0 -vector lattice, \mathbf{I} a weak order unit in E and let $\{\alpha_n\} \subset L_0$ be bounded from above (below). Then $\sup_{n \geq 1} (\alpha_n\mathbf{I}) = (\sup_{n \geq 1} \alpha_n)\mathbf{I}$ ($\inf_{n \geq 1} (\alpha_n\mathbf{I}) = (\inf_{n \geq 1} \alpha_n)\mathbf{I}$, respectively).

PROOF: First, let us show that the equality

$$e_{\alpha\mathbf{I}} := \sup_{n \geq 1} (\mathbf{I} \wedge n|\alpha|\mathbf{I}) = s(\alpha)\mathbf{I}$$

holds for any $\alpha \in L_0$. One can assume that $\alpha \geq 0$. Let $g_n = \{\alpha \geq \frac{1}{n}\}$ be a spectral idempotent for α in L_0 . It is obvious that $g_n \uparrow s(\alpha)$ and by Proposition 2.4, $g_n \mathbf{I} \uparrow s(\alpha) \mathbf{I}$.

Let $f_n = s(\alpha) - g_n$ and $\beta_n = n\alpha f_n$, $n = 1, 2, \dots$. It is clear that $0 \leq \beta_n \leq f_n \leq \mathbf{1}$ and $\beta_n g_i = 0$ for all $i = 1, 2, \dots, n$. Hence, $0 \leq \beta_n \mathbf{I} \leq f_n \mathbf{I} \leq f_i \mathbf{I} \leq \mathbf{I}$ as $n \geq i$. Let $a_n = \sup_{k \geq n} \beta_k \mathbf{I}$ and $a = \inf_{n \geq 1} a_n$. Since $a_n \leq f_n \mathbf{I}$, we have $a \leq f_n \mathbf{I}$ for all $n = 1, 2, \dots$. We thus have $0 \leq g_n a \leq g_n f_n \mathbf{I} = 0$, i.e. $g_n a = 0$, $n = 1, 2, \dots$. Hence, $s(\alpha)a = (\sup_{n \geq 1} g_n)a = 0$. On the other hand, $a \leq f_n \mathbf{I} \leq s(\alpha) \mathbf{I}$. By Remark 2.2 we obtain $a = s(\alpha)a$, and so $a = 0$. Thus, $\beta_n \mathbf{I} \xrightarrow{(o)} 0$. Since $\mathbf{1} \wedge n\alpha = g_n + \beta_n$, it follows that $\mathbf{I} \wedge (n\alpha) \mathbf{I} = (\mathbf{1} \wedge n\alpha) \mathbf{I} = g_n \mathbf{I} + \beta_n \mathbf{I}$. Hence, $e_{\alpha \mathbf{I}} = (o)\text{-}\lim(\mathbf{I} \wedge (n\alpha) \mathbf{I}) = (o)\text{-}\lim g_n \mathbf{I} + (o)\text{-}\lim \beta_n \mathbf{I} = s(\alpha) \mathbf{I}$. Now let us show that $\inf_{n \geq 1} (\alpha_n \mathbf{I}) = (\inf_{n \geq 1} \alpha_n) \mathbf{I}$ for any bounded from below sequence (α_n) in L_0 . Let $\alpha = \inf_{n \geq 1} \alpha_n$, $x = \inf_{n \geq 1} \alpha_n \mathbf{I}$.

Consider in E the families $\{e_t^x\}_{t \in \mathbb{R}}$ and $\{e_t^{\alpha_n \mathbf{I}}\}_{t \in \mathbb{R}}$ of spectral unitary elements for x and $\alpha_n \mathbf{I}$, respectively. By Lemma 2.3(ii) we have

$$e_t^{\alpha_n \mathbf{I}} = e_{(t\mathbf{I} - \alpha_n \mathbf{I})_+} = e_{((t\mathbf{1} - \alpha_n) \mathbf{I})_+} = e_{(t\mathbf{1} - \alpha_n)_+} \mathbf{I} = s((t\mathbf{1} - \alpha_n)_+) \mathbf{I}.$$

This together with Proposition 2.4 and [11, Lemma IV.10.2] imply that

$$\begin{aligned} e_t^x &= \sup_{n \geq 1} e_t^{\alpha_n \mathbf{I}} = \sup_{n \geq 1} (s((t\mathbf{1} - \alpha_n)_+) \mathbf{I}) \\ &= \left(\sup_{n \geq 1} s((t\mathbf{1} - \alpha_n)_+) \right) \mathbf{I} = s((t\mathbf{1} - \alpha)_+) \mathbf{I} = g_t^\alpha, \end{aligned}$$

where $\{g_t^\alpha\}_{t \in \mathbb{R}}$ is the family of spectral idempotents for α in L_0 . Similarly, for the family of spectral idempotents $\{e_t^{\alpha \mathbf{I}}\}_{t \in \mathbb{R}}$ we have

$$e_t^{\alpha \mathbf{I}} = e_{(t\mathbf{I} - \alpha \mathbf{I})_+} = s((t\mathbf{1} - \alpha)_+) \mathbf{I} = g_t^\alpha \mathbf{I}.$$

Hence, $e_t^x = e_t^{\alpha \mathbf{I}}$ for all $t \in \mathbb{R}$.

It follows from the spectral theorem for σ -complete vector lattices [11, Theorem IV.10.1] that $x = \alpha \mathbf{I}$, i.e. $\inf_{n \geq 1} (\alpha_n \mathbf{I}) = (\inf_{n \geq 1} \alpha_n) \mathbf{I}$. If $\{\alpha_n\}$ is a bounded from above sequence from L_0 , then passing to the sequence $\{-\alpha_n\}$, we obtain $\sup_{n \geq 1} (\alpha_n \mathbf{I}) = (\sup_{n \geq 1} \alpha_n) \mathbf{I}$. □

Remark 2.6. Let E be a σ -complete L_0 -vector lattice with a weak order unit. Then L_0 can be identified with the normal L_0 -submodule N in E . In addition, operations \sup and \inf are identical in L_0 and N . The Boolean algebra $\nabla(\Omega)$ is a σ -subalgebra in $\nabla(E)$.

3. Banach L_0 -vector lattices

Let E be a normal L_0 -module. An L_0 -valued norm $\|\cdot\| : E \rightarrow L_0$ is said to be *compatible with the structure of the L_0 -module E* (shortly, *L_0 -norm*) if $\|\lambda x\| = |\lambda|\|x\|$ for any $x \in E$ and $\lambda \in L_0$. Then, the pair $(E, \|\cdot\|)$ is called a *normed L_0 -module*.

Let E be a normed L_0 -module. Let t be the topology of local convergence with respect to the measure in L_0 . A sequence $\{x_n\} \subset E$ *t -converges* to $x \in E$ if $\|x_n - x\| \xrightarrow{t} 0$. Cauchy sequences are defined as usual. A normed L_0 -module E is called *Banach (t -Banach)* if any (bo)-Cauchy (t -Cauchy, respectively) sequence in E (bo)-converges (t -converges, respectively). E is a Banach L_0 -module if and only if it is a t -Banach L_0 -module.

Let E be a BKS over L_0 . It is possible to define a structure of L_0 -module on E . This structure makes E a Banach L_0 -module. Vice versa, any Banach L_0 -module E is a BKS over L_0 .

If E is a normed L_0 -module and simultaneously an L_0 -vector lattice with a monotone norm, then E is called a *normed L_0 -vector lattice*. Any norm complete L_0 -vector lattice is called a *Banach L_0 -vector lattice*. The class of Banach L_0 -vector lattices coincides with the class of Banach-Kantorovich lattices over L_0 .

Let us give examples of Banach L_0 -vector lattices.

Suppose ∇ is a complete Boolean algebra. Denote by $X(\nabla)$ the Stone compactification of ∇ . Let $L_0(\nabla)$ be the set of all continuous functions $x : X(\nabla) \rightarrow [-\infty, +\infty]$ such that $x^{-1}(\{\pm\infty\})$ is a nowhere dense subset of $X(\nabla)$ (see [10, V, §2]). Evidently, $L_0(\nabla)$ is a ring and an order complete vector lattice. The function $\mathbf{1}$, equal to 1 identically on $X(\nabla)$, is a weak order unit in $L_0(\nabla)$. The order ideal generated by the element $\mathbf{1}$ coincides with the space $C(X(\nabla))$ of all continuous real functions on $X(\nabla)$.

A mapping $m : \nabla \rightarrow L_0$ is called an *L_0 -valued measure* on ∇ if

1. $m(e) \geq 0$ for any $e \in \nabla$,
2. $m(e \vee g) = m(e) + m(g)$ if $e, g \in \nabla$ and $e \wedge g = 0$,
3. if $e_n \downarrow 0$, $e_n \in \nabla$, then $m(e_n) \downarrow 0$.

A measure m is called *strongly positive* if $m(e) = 0$, $e \in \nabla$ implies $e = 0$. Using Lebesgue construction, one can obtain an integral $I_m : x \rightarrow \int x dm$ for every strongly positive L_0 -valued measure m (see [10], [8]). There exists the greatest order ideal $L := L_1(\nabla, m)$ in $L_0(\nabla)$ containing ∇ with the following properties:

1. $I_m e = m(e)$ for any $e \in \nabla$,
2. $I_m(ax + by) = aI_m x + bI_m y$, $x, y \in L$, $a, b \in \mathbb{R}$,
3. if $x_n, x \in L$ and $x_n \uparrow x$ then $I_m x_n \xrightarrow{(o)} I_m x$.

The mapping I_m satisfying the above properties is uniquely defined. The norm on $L_1(\nabla, m)$ is defined as $\|x\|_1 = \int |x| dm$. Now, $(L_1(\nabla, m), \|\cdot\|_1)$ is a (bo)-complete

LNS over L_0 (see [10]).

We suppose that $\nabla(\Omega)$ is a regular Boolean subalgebra in ∇ , i.e. $\sup A \in \nabla(\Omega)$ for every $A \subset \nabla(\Omega)$. We can always obtain this by considering the complete tensor product $\nabla \otimes \nabla(\Omega)$ of the Boolean algebras ∇ and $\nabla(\Omega)$ (see [2, VII, §7.2]). One can canonically identify $L_0(\Omega)$ with a subalgebra in $L_0(\nabla)$. It is also a regular vector sublattice in $L_0(\nabla)$. Moreover, sup and inf operations in $L_0(\Omega)$ and $L_0(\nabla)$ coincide. Hence, $L_0(\nabla)$ becomes an L_0 -vector lattice (multiplication of elements from $L_0(\nabla)$ by elements from L_0 coincides with the natural multiplication in $L_0(\nabla)$).

From now on, we require the measure $m : \nabla \rightarrow L_0$ to be compatible with the module structure, i.e. $m(ge) = gm(e)$ for all $e \in \nabla, g \in \nabla$. In this case, $L_1(\nabla, m)$ becomes a BKS over L_0 . In addition, the following property holds:

Let $x \in L_1(\nabla, m)$ and $\alpha \in L_0$. Then, $\alpha x \in L_1(\nabla, m)$ and $\int \alpha x dm = \alpha \int x dm$. In particular, $L_0 \subset L_1(\nabla, m)$ and $\int \alpha dm = \alpha m(\mathbf{1})$ for all $\alpha \in L_0$ (see [6, 6.1.10]).

Let $p > 1$. Set

$$L_p(\nabla, m) := \{x \in L_0(\nabla) : |x|^p \in L_1(\nabla, m)\}.$$

Then $L_p(\nabla, m)$ is a normal L_0 -module and a Banach L_0 -vector lattice with respect to the norm $\|x\|_p := (\int |x|^p dm)^{1/p}$ (see [1, 4.2.2], or [2, VIII, §8.2]).

Now we give examples of L_0 -valued measures compatible with the module structure.

Example 1. Let (Ω, Σ, μ) be a σ -finite complete measure space. Let $\mathcal{A} \subset \Sigma$ be a σ -subalgebra. Denote by $m(e) = E(e|\mathcal{A})$ the conditional expectation. It is clear that m is a strongly positive $L_0(\Omega, \mathcal{A}, \mu)$ -valued measure on $\nabla(\Omega, \Sigma, \mu)$ compatible with the module structure.

Example 2. Let (Ω, Σ, μ) be the same space as in Example 1, X be another complete Boolean algebra with a strongly positive scalar measure ν . Step mappings $u : (\Omega, \Sigma, \mu) \rightarrow X$ are defined in the usual way. Let $\Gamma(X)$ be the set of all step mappings $u : (\Omega, \Sigma, \mu) \rightarrow X$. A mapping $u : (\Omega, \Sigma, \mu) \rightarrow X$ is said to be measurable if there exists a sequence $\{u_n\} \subset \Gamma(X)$ such that $\nu(u(\omega)\Delta u_n(\omega)) \rightarrow 0$ as $n \rightarrow \infty$ for a.e. $\omega \in \Omega$. Here, $e\Delta g = (e \wedge Cg) \vee (Ce \wedge g)$, $e, g \in X$. Let $\mathcal{L}_0(\Omega, X)$ be the set of all measurable maps from (Ω, Σ, μ) into X . For arbitrary $u, v \in \mathcal{L}_0(\Omega, X)$ we set $u \leq v$ if $u(\omega) \leq v(\omega)$ for all $\omega \in \Omega$. Then, $\mathcal{L}_0(\Omega, X)$ becomes a Boolean algebra. Its unit is $\mathbf{1}(\omega) \equiv \mathbf{1}_X$. Its zero is $\mathbf{0}(\omega) = \mathbf{0}_X$. The complement is defined as $(Cu)(\omega) = C(u(\omega))$. Moreover $(u \vee v)(\omega) = u(\omega) \vee v(\omega)$, $(u \wedge v)(\omega) = u(\omega) \wedge v(\omega)$, $\omega \in \Omega$.

Consider the ideal $J = \{u \in \mathcal{L}_0(\Omega, X) : u(\omega) = 0 \text{ a.e.}\}$. Define $L_0(\Omega, X)$ as a Boolean factor-algebra $\mathcal{L}_0(\Omega, X)/J$. $L_0(\Omega, X)$ is a complete Boolean algebra (see [1]). $\nabla(\Omega) = \{u \in L_0(\Omega, X) : u = \chi_A, A \in \Sigma\}$ is a regular Boolean subalgebra in $L_0(\Omega, X)$. If $u \in \Gamma(X)$, then the scalar function $\nu \circ u \in L_0(\Omega)$. Hence, for any $v \in L_0(\Omega, X)$, the function $\nu(v(\omega)) = \lim_{n \rightarrow \infty} \nu(v_n(\omega)) \in L_0(\Omega)$. Here $v_n \in \Gamma(X)$,

$\nu(v(\omega)\Delta v_n(\omega)) \rightarrow 0$. So, we defined a mapping $\nu : L_0(\Omega, X) \rightarrow L_0(\Omega)$. It is an L_0 -valued strongly positive measure on $L_0(\Omega, X)$ compatible with the module structure (see [1]).

Let $(E, \|\cdot\|)$ be a normed L_0 -vector lattice. A norm in E is called *order continuous* if for any $\{x_n\} \subset E_+, x_n \downarrow 0$ implies $\|x_n\| \xrightarrow{t} 0$.

The following order and topological properties of normed L_0 -vector lattices can be proved in the same way as in the case of normed lattices.

Theorem 3.1. *Let $(E, \|\cdot\|)$ be a normed L_0 -vector lattice. Then*

1. *if $\{x_n\} \subset E$ is an increasing t -converging sequence, then*

$$\lim_{n \rightarrow \infty} x_n = \sup_n x_n.$$

2. (Amemiya theorem). *The following conditions are equivalent:*
 - (a) *E is a Banach L_0 -vector lattice;*
 - (b) *if $\{x_n\}$ is a (bo)-Cauchy increasing sequence from E_+ , then $\{x_n\}$ (bo)-converges in E ;*
 - (c) *if $\{x_n\}$ is a (bo)-Cauchy increasing sequence from E_+ , then there exists $x = (\sup_{n \geq 1} x_n) \in E$.*
3. *Let $(E, \|\cdot\|)$ be a σ -complete normed L_0 -vector lattice with an order continuous norm. Then E is of countable type. Therefore E is an order complete vector lattice.*
4. *Let E be a Banach L_0 -vector space. The following conditions are equivalent:*
 - (a) *E is an order complete lattice and $\|\cdot\|$ is order continuous.*
 - (b) *Any bounded sequence of positive mutually disjoint elements t -converges to zero.*

4. Orlicz-Kantorovich lattices associated with Orlicz L_0 -modulators

Let us start with some definitions.

Definition. $\psi : [0, \infty) \rightarrow \mathbb{R}$ is called an *Orlicz function* if it is a convex non-negative function such that $\psi(0) = 0$ and $\psi(t) > 0$ for $t > 0$. An additional requirement is the so called (δ_2, Δ_2) -condition, i.e. $\psi(2t) \leq c\psi(t)$ for all $t \geq 0$ and a constant $c > 0$.

Let $x \in L_0(\nabla)$. By definition, $G = \{t \in X(\nabla) : |x(t)| < \infty\} \subset X(\nabla)$ is an open and dense subset. Hence, we can define $y \in L_0(\nabla)$ as $y = \psi \circ |x| := \psi(|x|)$. Define

$$L_\psi := L_\psi(\nabla, m) := \{x \in L_0(\nabla) : \psi(|x|) \in L_1(\nabla, m)\}.$$

It is clear that L_ψ is a normal L_0 -submodule and a vector sublattice in $L_0(\nabla)$.

Let $\mathcal{P}(L_0) = \{\lambda \geq 0 \in L_0 : s(\lambda) = \mathbf{1}\}$. Obviously, for any $\lambda \in \mathcal{P}(L_0)$ there exists $\lambda^{-1} \in \mathcal{P}(L_0)$.

Lemma 4.1. *Let $x \in L_\psi$. There exists $\lambda \in \mathcal{P}(L_0)$ such that*

$$\int \psi(\lambda^{-1}|x|) dm \leq \mathbf{1}.$$

PROOF: Let $\lambda_0 = \int \psi(|x|) dm + \mathbf{1}$. It is clear that $\lambda_0 \in \mathcal{P}(L_0)$ and $0 \leq \lambda_0^{-1} \leq \mathbf{1}$. Since $\psi(st) \leq s\psi(t)$ for all $s \in [0, 1]$, we are done. \square

Hence, we can define an L_0 -valued function

$$\|x\|_{(\psi)} = \inf \left\{ \lambda \in \mathcal{P}(L_0) : \int \psi(\lambda^{-1}|x|) dm \leq \mathbf{1} \right\}.$$

Theorem 4.2. *$(L_\psi, \|\cdot\|_{(\psi)})$ is a Banach L_0 -vector lattice.*

We need some lemmas to prove Theorem 4.2.

Lemma 4.3. *Let $x_n, x \in L_0(\nabla)$, $0 \leq x_n \uparrow x$. Then $\psi(x_n) \uparrow \psi(x)$.*

The proof of this lemma is similar to that of Lemma 2.4 from [14].

Lemma 4.4. *$\|x\|_{(\psi)}$ is a monotone L_0 -norm on L_ψ , i.e. $(L_\psi, \|\cdot\|_{(\psi)})$ is a normed L_0 -vector lattice.*

PROOF: Obviously, $\|\cdot\|$ is monotone, convex and positive. Assume now that $\|x\| = 0$ for some $x \in L_\psi$. Consider $\lambda \in \mathcal{P}(L_0)$ such that $\int \psi(\lambda^{-1}|x|) dm \leq \mathbf{1}$. Then, $\lambda \wedge \mathbf{1} \in \mathcal{P}(L_0)$ and $0 \leq \lambda \wedge \mathbf{1} \leq \mathbf{1}$. Obviously, $(\lambda \wedge \mathbf{1})^{-1}|x| = \lambda^{-1}|x|\{\lambda < \mathbf{1}\} + |x|\{\lambda \geq \mathbf{1}\}$. Hence, $\psi((\lambda \wedge \mathbf{1})^{-1}|x|) = \psi(\lambda^{-1}|x|)\{\lambda < \mathbf{1}\} + \psi(|x|)\{\lambda \geq \mathbf{1}\}$. Therefore,

$$\begin{aligned} \int \psi((\lambda \wedge \mathbf{1})^{-1}|x|) dm &= \{\lambda < \mathbf{1}\} \int \psi(\lambda^{-1}|x|) dm + \{\lambda \geq \mathbf{1}\} \int \psi(|x|) dm \\ &\leq \{\lambda < \mathbf{1}\} + \int \psi(|x|) dm \leq \mathbf{1} + \int \psi(|x|) dm. \end{aligned}$$

However, $\psi((\lambda \wedge \mathbf{1})^{-1}|x|) \geq (\lambda \wedge \mathbf{1})^{-1}\psi(x)$. Therefore,

$$\int \psi(|x|) dm \leq (\lambda \wedge \mathbf{1}) \left(\mathbf{1} + \int \psi(|x|) dm \right).$$

Now, one can take infimum over all such λ and obtain $\int \psi(|x|) dm = 0$. Hence, $x = 0$. \square

Lemma 4.5. *Let $x \in L_\psi$, $e = \mathbf{1} - s(\|x\|_{(\psi)})$. Then*

$$\int \psi \left(\left(\|x\|_{(\psi)} + e \right)^{-1} |x| \right) dm \leq \mathbf{1}.$$

PROOF: Obviously, $s(x) = s(\|x\|_{(\psi)})$. Hence $(\mathbf{1} - e)|x| = |x|$. Since L_0 has countable type, then there exists a sequence $\{\lambda_n\} \subset \mathcal{P}(L_0)$, such that $\int \psi(\lambda_n^{-1}|x|) \leq \mathbf{1}$ and $\lambda_n \downarrow = \|x\|_{(\psi)}$. Set $\alpha_n = \lambda_n(\mathbf{1} - e) + e$, $n = 1, 2, \dots$. Then $\alpha_n \downarrow (\|x\|_{(\psi)} + e)$ and $\alpha_n^{-1} = (\lambda_n^{-1}(\mathbf{1} - e) + e) \uparrow (\|x\|_{(\psi)} + e)^{-1}$. Hence, $\psi(\alpha_n^{-1}|x|) \uparrow \psi((\|x\|_{(\psi)} + e)^{-1}|x|)$ (see Lemma 4.3). By the monotone convergence theorem (see [6, 6.1.5]), we have

$$\begin{aligned} \int \psi \left(\left(\|x\|_{(\psi)} + e \right)^{-1} |x| \right) dm &= \sup_{n \geq 1} \int \psi \left(\alpha_n^{-1}|x| \right) dm \\ &= \sup_{n \geq 1} \int \psi \left(\left(\lambda_n^{-1}(\mathbf{1} - e) + e \right) |x| \right) dm \\ &= \sup_{n \geq 1} \int \psi(\lambda_n^{-1}|x|) dm \leq \mathbf{1}. \end{aligned}$$

□

PROOF OF THEOREM 4.2: Consider a (bo)-Cauchy increasing sequence $\{x_n\} \in (L_\psi)_+$. Obviously, the sequence $\|x_n\|_{(\psi)}$ is a (o)-Cauchy sequence in L_0 . That is, $\|x_n\|_{(\psi)} \uparrow \alpha$. Set $e_n = \mathbf{1} - s(x_n)$ and $\alpha_n = \|x_n\|_{(\psi)} + e_n$. Then $0 \leq \alpha_n \leq \alpha + \mathbf{1}$. By Lemma 4.5, $\int \psi(\alpha_n^{-1}x_n) \leq \mathbf{1}$. Therefore, $\int \psi((\alpha + \mathbf{1})^{-1}x_n) \leq \mathbf{1}$. The sequence $\psi((\alpha + \mathbf{1})^{-1}x_n) \in L_1$ is monotone and L_1 -bounded. Hence, $\psi((\alpha + \mathbf{1})^{-1}x_n) \uparrow y \in L_1$. Therefore, $x_n \uparrow (\alpha + \mathbf{1})\psi^{-1}(y) \in L_\psi$. □

A Banach L_0 -vector lattice $(L_\psi, \|\cdot\|_{(\psi)})$ is called *the Orlicz-Kantorovich space*. See examples after Theorem 5.1.

Denote $\Phi(x) = \int \psi(|x|) dm$. It is easy to see that the mapping $\Phi : L_\psi \rightarrow L_0$ satisfies the following properties:

1. $\Phi(x) \geq 0$ and $\Phi(x) = 0 \Leftrightarrow x = 0$;
2. $\Phi(x) \leq \Phi(y)$ if $|x| \leq |y|$;
3. $\Phi(\alpha x + (\mathbf{1} - \alpha)y) \leq \alpha\Phi(x) + (\mathbf{1} - \alpha)\Phi(y)$, $\alpha \in L_0$, $0 \leq \alpha \leq \mathbf{1}$;
4. $\Phi(2x) \leq c\Phi(x)$ for some constant $c > 0$;
5. $\Phi(x + y) = \Phi(x) + \Phi(y)$ if $x \wedge y = 0$;
6. $\Phi(ex) = e\Phi(x)$ for all $e \in \nabla(\Omega)$;
7. $\Phi(t\mathbf{1}) = \varphi(t)\Phi(\mathbf{1})$ for all $t \geq 0$, where $\varphi : [0, \infty) \rightarrow [0, \infty)$ is a scalar function.

Now we define an Orlicz L_0 -lattice. Let E be an L_0 -vector lattice with a weak order unit $\mathbf{1}$. A map $\Phi : E \rightarrow L_0$ is called an Orlicz L_0 -modulator if Φ satisfies properties 1-7. Obviously, $\Phi(x) = \Phi(|x|)$ and $\Phi(\alpha x) \leq \alpha\Phi(x)$ for $\alpha \in L_0, 0 \leq \alpha \leq \mathbf{1}$. The element $\Phi(\mathbf{1})$ is invertible in L_0 . Indeed, let $e = s(\Phi(\mathbf{1}))$. Then $\Phi((\mathbf{1} - e)\mathbf{1}) = (\mathbf{1} - e)\Phi(\mathbf{1}) = 0$. Hence, $(\mathbf{1} - e)\mathbf{1} = 0$ and $e = \mathbf{1}$. Properties 1-7 imply that φ is an Orlicz function satisfying the (δ_2, Δ_2) -condition.

Set $B(x) = \{\lambda \in \mathcal{P}(L_0) : \Phi(\lambda^{-1}x) \leq \mathbf{1}\}$. If $\lambda = \Phi(x) + \mathbf{1}$, then $\Phi(\lambda^{-1}x) \leq \lambda^{-1}\Phi(x) \leq \mathbf{1}$. Hence $B(x)$ is a non-empty set. For any $x \in E$, set $\|x\|_\Phi = \inf\{\lambda : \lambda \in B(x)\}$.

Proposition 4.6. *$(E, \|\cdot\|_\Phi)$ is a normed L_0 -vector lattice.*

PROOF: Obviously, $\|\cdot\|_\Phi$ is monotone, convex and positive. If $\|x\|_\Phi = 0$, then repeating the proof of Lemma 4.4 and using properties of the Orlicz L_0 -modulator Φ , we obtain $x = 0$. Let $x, y \in E, \lambda_1 \in B(x), \lambda_2 \in B(y)$. Then

$$\begin{aligned} \Phi((\lambda_1 + \lambda_2)^{-1}(x + y)) &= \Phi(\lambda_1(\lambda_1 + \lambda_2)^{-1}\lambda_1^{-1}x + \lambda_2(\lambda_1 + \lambda_2)^{-1}\lambda_2^{-1}y) \\ &\leq \lambda_1(\lambda_1 + \lambda_2)^{-1}\Phi(\lambda_1^{-1}x) + \lambda_2(\lambda_1 + \lambda_2)^{-1}\Phi(\lambda_2^{-1}y) \leq \mathbf{1}, \end{aligned}$$

i.e. $\lambda_1 + \lambda_2 \in B(x + y)$. This means that $B(x) + B(y) \subseteq B(x + y)$, and so

$$\|x + y\|_\Phi \leq \|x\|_\Phi + \|y\|_\Phi.$$

Let us now show that $\|ex\|_\Phi = e\|x\|_\Phi$ for any idempotent $e \in L_0$ and $x \in E$. Take $\lambda, \beta \in \mathcal{P}(L_0)$ such that $\Phi(\lambda^{-1}x) \leq \mathbf{1}, \Phi(\beta^{-1}xe) \leq \mathbf{1}$. Then $\gamma = \beta e + \lambda(\mathbf{1} - e) \in \mathcal{P}(L_0)$, in addition $\gamma^{-1} = \beta^{-1}e + \lambda^{-1}(\mathbf{1} - e)$ and

$$\begin{aligned} \Phi(\gamma^{-1}x) &= \Phi(\gamma^{-1}xe) + \Phi(\gamma^{-1}x(\mathbf{1} - e)) \\ &= \Phi(\beta^{-1}xe) + \Phi(\lambda^{-1}x(\mathbf{1} - e)) \\ &= e\Phi(\beta^{-1}xe) + (\mathbf{1} - e)\Phi(\lambda^{-1}x) \\ &\leq e + (\mathbf{1} - e) = \mathbf{1}. \end{aligned}$$

Hence, $\|x\|_\Phi \leq \gamma$ and therefore $e\|x\|_\Phi \leq \|ex\|_\Phi$.

Since $|ex| \leq |x|$, we have $\|ex\|_\Phi \leq \|x\|_\Phi$. That is why $e\|x\|_\Phi \leq \|ex\|_\Phi \leq e\|x\|_\Phi$, i.e. $e\|x\|_\Phi = \|ex\|_\Phi$.

Further, if $\lambda \in \mathcal{P}(L_0)$ and $\Phi(\lambda^{-1}ex) \leq \mathbf{1}$, then $\Phi(\beta^{-1}ex) = \Phi(\lambda^{-1}ex) \leq \mathbf{1}$ for $\beta = \lambda e + \varepsilon(\mathbf{1} - e)$. Hence $\|ex\|_\Phi(\mathbf{1} - e) = 0$ and $\|ex\|_\Phi = e\|ex\|_\Phi = e\|x\|_\Phi$.

Let now α be an invertible element from L_0 . Then

$$\begin{aligned} \|\alpha x\|_\Phi &= \inf \left\{ \lambda \in \mathcal{P}(L_0) : \Phi(\lambda^{-1}\alpha x) \leq \mathbf{1} \right\} \\ &= \inf \left\{ |\alpha|\gamma : \Phi(\gamma^{-1}x) \leq \mathbf{1}, \gamma = \lambda|\alpha|^{-1} \in \mathcal{P}(L_0) \right\} = |\alpha|\|x\|_\Phi. \end{aligned}$$

If α is an arbitrary non-zero element from L_0 , $e = \mathbf{1} - s(\alpha)$, then $\alpha + e$ is invertible in L_0 , and therefore

$$\begin{aligned} \|\alpha x\|_{\Phi} &= \|(\alpha + e)(\mathbf{1} - e)x\|_{\Phi} = (|\alpha| + e)\|(\mathbf{1} - e)x\|_{\Phi} \\ &= (|\alpha| + e)\|x\|_{\Phi} = |\alpha|\|x\|_{\Phi}. \end{aligned}$$

Thus, $(E, \|\cdot\|_{\Phi})$ is a normed L_0 -vector lattice. □

Definition. A norm-complete L_0 -vector lattice $(E, \|\cdot\|_{\Phi})$ is called an *Orlicz L_0 -lattice*.

The Orlicz-Kantorovich space $(L_{\psi}, \|\cdot\|_{(\psi)})$ is a good example of Orlicz L_0 -lattices.

Theorem 4.7. *The Orlicz L_0 -lattice $(E, \|\cdot\|_{\Phi})$ is an order complete lattice, and the L_0 -norm $\|\cdot\|_{\Phi}$ is order continuous.*

PROOF: Consider a disjoint bounded sequence $\{x_n\} \subset E_+$. Since $x_n \leq x \in E_+$, we have $\sum_{i=1}^n x_i \leq x$. Using property 5, we obtain $\sum_{i=1}^n \Phi(x_i) \leq \Phi(x)$. Hence, $\Phi(x_n) \xrightarrow{(o)} 0$. For any fixed $i = 1, 2, \dots$, $\Phi(2^i x_n) \xrightarrow{(o)} 0$. The element $\lambda = \Phi(x) + \mathbf{1} \in B(x)$. Hence, $\|x\|_{\Phi} \leq \Phi(x) + \mathbf{1}$. Therefore, $\|x_n\|_{\Phi} \leq 2^{-i}\Phi(2^i x_n) + 2^{-i}\mathbf{1}$. Thus, $(o)\text{-}\lim \|x_n\|_{\Phi} \leq 2^{-i}\mathbf{1}$ for any i . Hence, $(o)\text{-}\lim \|x_n\|_{\Phi} = 0$. By Theorem 3.1.4, we are done. □

Lemma 4.8. *Let $\|x\|_{\Phi} \leq \mathbf{1}$ and $\{\|x\|_{\Phi} = \mathbf{1}\} = 0$. Then $\Phi(x) \leq \|x\|_{\Phi}$.*

PROOF: As in Proposition 2.7 from [13], one can choose $\lambda_n \in B(x)$ such that $\lambda_n \downarrow \|x\|_{\Phi}$. Let $\lambda \in L_0$, $\lambda \geq 0$, $\|x\|_{\Phi} \leq \lambda \leq \mathbf{1}$ and $\{\lambda = \|x\|_{\Phi}\} = 0$. Then, λ is invertible. Set $f_n = \{\lambda < \lambda_n\}$. Obviously, $f_n \downarrow 0$. We have

$$\begin{aligned} \Phi(\lambda^{-1}x) &= \Phi\left(\left(\lambda_n^{-1}\lambda_n\lambda^{-1}\right)x\right) \\ &= f_n\Phi\left(\lambda_n^{-1}x\lambda_n\lambda^{-1}\right) + (\mathbf{1} - f_n)\Phi\left(\lambda_n^{-1}x\left(\lambda_n\lambda^{-1}(\mathbf{1} - f_n)\right)\right) \\ &\leq f_n\Phi\left(\lambda_n^{-1}x\lambda_n\lambda^{-1}\right) + (\mathbf{1} - f_n)\Phi(\lambda_n^{-1}x) \\ &\leq f_n\Phi(\lambda_n^{-1}x\lambda_n\lambda^{-1}) + (\mathbf{1} - f_n). \end{aligned}$$

Since $f_n \downarrow 0$, $f_n\Phi(\lambda_n^{-1}x\lambda_n\lambda^{-1}) \xrightarrow{(o)} 0$. After switching to (o) -limit, we obtain $\Phi(\lambda^{-1}x) \leq \mathbf{1}$. Since $\lambda \leq \mathbf{1}$, we have $\lambda^{-1}\Phi(x) \leq \Phi(\lambda^{-1}x) \leq \mathbf{1}$.

Let $\alpha_n = \|x\|_{\Phi} + n^{-1}(\mathbf{1} - \|x\|_{\Phi})$. Then $\|x\|_{\Phi} \leq \alpha_n \leq \mathbf{1}$ and $\{\|x\|_{\Phi} = \alpha_n\} = 0$. Hence $\Phi(x) \leq \alpha_n$, $n = 1, 2, \dots$ and $\Phi(x) \leq \|x\|_{\Phi}$. □

Proposition 4.9. *Let $(E, \|\cdot\|_\Phi)$ be an Orlicz L_0 -lattice, $y_n \in E$. Then $\|y_n\|_\Phi \xrightarrow{(o)} 0$ if and only if $\Phi(y_n) \xrightarrow{(o)} 0$.*

PROOF: Let $\Phi(y_n) \xrightarrow{(o)} 0$. Then, $\|y_n\|_\Phi \xrightarrow{(o)} 0$ (see the proof of Theorem 4.7).

Set $g_n = \{\|y_n\|_\Phi < \mathbf{1}\}$. Since $\|y_n\|_\Phi \xrightarrow{(o)} 0$, we have $g_n \xrightarrow{(o)} \mathbf{1}$. Obviously, $\|g_n y_n\|_\Phi = g_n \|y_n\|_\Phi \leq \mathbf{1}$ and $\{g_n \|y_n\|_\Phi = \mathbf{1}\} = \emptyset$. By Lemma 4.8, $\Phi(g_n y_n) \leq \|g_n y_n\|_\Phi = g_n \|y_n\|_\Phi \xrightarrow{(o)} 0$. Since $(\mathbf{1} - g_n) \xrightarrow{(o)} 0$, we have $(\mathbf{1} - g_n)\Phi(y_n) \xrightarrow{(o)} 0$. Hence, $\Phi(y_n) = \Phi(g_n y_n) + \Phi((\mathbf{1} - g_n)y_n) \xrightarrow{(o)} 0$. □

Proposition 4.10. *Let $x_n \uparrow x$. Then $\Phi(x_n) \uparrow \Phi(x)$.*

PROOF: Obviously, $\sup_{n \geq 1} \Phi(x_n) \leq \Phi(x)$. Further, for any number $a \in (0, 1]$, we have $x = (1 - a)x_n + a(x_n + a^{-1}(x - x_n))$. Using properties of Φ , we obtain

$$\begin{aligned} \Phi(x) &\leq (1 - a)\Phi(x_n) + a\Phi\left(x_n + a^{-1}(x - x_n)\right) \\ &\leq \Phi(x_n) + 2^{-1}ac\left(\Phi(x_n) + \Phi\left(a^{-1}(x - x_n)\right)\right). \end{aligned}$$

By Theorem 4.7, $\|a^{-1}(x - x_n)\|_\Phi \xrightarrow{(o)} 0$. By Proposition 4.9, $\Phi(a^{-1}(x - x_n)) \downarrow 0$. Hence,

$$\begin{aligned} \Phi(x) &\leq (o)\text{-}\limsup_{n \rightarrow \infty} \left(\Phi(x_n) + 2^{-1}ac\left(\Phi(x_n) + \Phi\left(a^{-1}(x - x_n)\right)\right)\right) \\ &= \left(1 + \frac{1}{2}ac\right) \sup_{n \geq 1} \Phi(x_n). \end{aligned}$$

Since a is arbitrary, we obtain $\Phi(x) \leq \sup_{n \geq 1} \Phi(x_n)$. □

5. Abstract characterization of Orlicz-Kantorovich L_0 -spaces

Definition (compare with [2]). An Orlicz L_0 -lattice $(E, \|\cdot\|_\Phi)$ is called *component-invariant* if

$$\Phi(te) = \Phi(e)\Phi^{-1}(\mathbf{I})\Phi(t\mathbf{I})$$

for all $t \geq 0, e \in \nabla$.

The Orlicz-Kantorovich space $(L_\psi(\nabla, m), \|\cdot\|_{(\psi)})$ is a component-invariant Orlicz L_0 -lattice. The reverse assertion is proved in Theorem 5.1. This can be considered as an abstract characterization of Orlicz-Kantorovich spaces in the class of Banach L_0 -vector lattices.

Theorem 5.1. *Let $(E, \|\cdot\|_{\Phi})$ be a component-invariant Orlicz L_0 -lattice. There exists a strongly positive measure m on ∇ , with values in L_0 , such that $(E, \|\cdot\|_{\Phi})$ is isometrically isomorphic to the Orlicz-Kantorovich space $(L_{\psi}(\nabla, m), \|\cdot\|_{(\psi)})$. Here $\psi(t) \cdot \mathbf{1} = \Phi(t\mathbf{I})\Phi^{-1}(\mathbf{I})$.*

PROOF: E can be identified (see [12]) with a normal vector sublattice in $L_0(\nabla) = C_{\infty}(X(\nabla))$ so that \mathbf{I} coincides with the $f \equiv \mathbf{1}$. Moreover, $e \in \nabla$ if and only if e is a characteristic function of an open-closed set from $X(\nabla)$. For any $e \in \nabla$, set $m(e) = \Phi(e)$. Obviously, $m(e) \in L_0$, $m(e) \geq 0$. If $e \wedge g = 0$, $e, g \in \nabla$, then $m(e \vee g) = m(e) + m(g)$. Clearly, $m(e) = 0$ if and only if $e = 0$. Let $\{e_n\} \subset \nabla$ and $e_n \downarrow 0$. By Theorem 4.7, we have $\|e_n\|_{\Phi} \downarrow 0$. Proposition 4.9 implies $\Phi(e_n) \downarrow 0$. This means that m is a strongly positive measure on ∇ with values in L_0 . Obviously, $m(eg) = em(g)$. Hence, m is compatible with the module structure.

Let x be a positive simple element from $L_0(\nabla)$, i.e. $x = \sum_{i=1}^n \lambda_i g_i$. Here, $\lambda_i \geq 0$ and $g_i \in \nabla$ are mutually disjoint. $\sup g_i = \mathbf{I}$. Obviously, $x \in E$ and $x \in L_{\psi}(\nabla, m)$.

Using the component invariance of $(E, \|\cdot\|_{\Phi})$, we obtain

$$\begin{aligned} \Phi(x) &= \sum_{i=1}^n \Phi(\lambda_i g_i) = \sum_{i=1}^n \Phi(g_i)\Phi^{-1}(\mathbf{I})\Phi(\lambda_i \mathbf{I}) = \sum_{i=1}^n \psi(\lambda_i)m(g_i) \\ &= \int \sum_{i=1}^n \psi(\lambda_i)g_i \, dm = \int \psi\left(\sum_{i=1}^n \lambda_i g_i\right) \, dm = \int \psi(x) \, dm. \end{aligned}$$

Thus, $\|x\|_{\Phi} = \inf\{\lambda \in \mathcal{P}(L_0) : \Phi(\lambda^{-1}x) \leq \mathbf{I}\} = \inf\{\lambda \in \mathcal{P}(L_0) : \int \psi(\lambda^{-1}x) \, dm \leq \mathbf{I}\} = \|x\|_{(\psi)}$ for any positive simple element x from $L_0(\nabla)$.

However, simple elements are dense in E as well as in L_{ψ} . □

We now use Theorem 5.1 to construct examples of Orlicz-Kantorovich spaces.

Let $(\Omega, \Sigma, \mu), (X, \nu)$ be as in Example 2. Let $L_{\psi}(X, \nu)$ be an Orlicz space associated with (X, ν) and with the Orlicz function ψ satisfying the (δ_2, Δ_2) -condition. We denote by $\Gamma(L_{\psi}(X, \nu))$ the set of all step mappings $u : (\Omega, \Sigma, \mu) \rightarrow L_{\psi}(X, \nu)$ having the form $u = \sum_{i=1}^n x_i \chi_{A_i}$ where $x_i \in L_{\psi}(X, \nu)$, $A_i \in \Sigma$, $A_i \cap A_j = \emptyset$, $i \neq j$, $i, j = 1, \dots, n$, $n \in \mathbb{N}$.

A mapping $u : (\Omega, \Sigma, \mu) \rightarrow L_{\psi}(X, \nu)$, is called measurable if there exists a sequence $\{u_k\} \subset \Gamma(L_{\psi}(X, \nu))$ such that $\|u(\omega) - u_n(\omega)\|_{L_{\psi}(X, \nu)} \rightarrow 0$ as $n \rightarrow \infty$ for a.e. $\omega \in \Omega$. Let $\mathcal{L}_0(\Omega, L_{\psi}(X, \nu))$ be the set of all measurable mappings from (Ω, Σ, μ) into $L_{\psi}(X, \nu)$. Obviously, $\mathcal{L}_0(\Omega, L_{\psi}(X, \nu))$ is an $\mathcal{L}_0(\Omega)$ -module, in addition $\|u(\omega)\|_{L_{\psi}(X, \nu)}$ is a measurable function on (Ω, Σ, μ) for all $u \in \mathcal{L}_0(\Omega, L_{\psi}(X, \nu))$. Consider an $\mathcal{L}_0(\Omega)$ -submodule $J = \{u \in \mathcal{L}_0(\Omega, L_{\psi}(X, \nu)) : u(\omega) = 0 \text{ a.e.}\}$ and denote by $L_0(\Omega, L_{\psi}(X, \nu))$ the factor-module $\mathcal{L}_0(\Omega, L_{\psi}(X, \nu))/J$. Then $(L_0(\Omega, L_{\psi}(X, \nu)), \|\cdot\|)$ is a Banach L_0 -vector lattice [3], where $\|\tilde{u}\| = [\|u(\omega)\|_{L_{\psi}(X, \nu)}]_{\sim}$.

The norm in $L_\psi(X, \nu)$ is order continuous, and therefore $g_n, g \in X, \nu(g_n \Delta g) \rightarrow 0$ implies that $\|g_n - g\|_{L_\psi(X, \nu)} \rightarrow 0$. Hence, the complete Boolean algebra $L_0(\Omega, X)$ from Example 2 is a subset of $L_0(\Omega, L_\psi(X, \nu))$. Moreover, the Boolean algebra of unitary elements from $L_0(\Omega, L_\psi(X, \nu))$ with respect to the weak unit $\mathbf{1}(\omega) = \mathbf{1}_X, \omega \in \Omega$ coincides with $L_0(\Omega, X)$. It is clear that $m(\tilde{e}) = [\nu(e(\omega))]^\sim$ is a strongly positive L_0 -valued measure on $L_0(\Omega, X)$ and m is compatible with the module structure (see Example 2).

Theorem 5.2. *The Banach L_0 -vector lattices $L_0(\Omega, L_\psi(X, \nu))$ and $L_\psi(L_0(\Omega, X), m)$ are order and isometrically isomorphic.*

PROOF: Without loss of generality, one can assume that $\psi(1) = 1$. For any $u = \sum_{i=1}^n x_i \chi_{A_i} \in \Gamma(L_\psi(X, \nu))$, set

$$\Phi_0(u)(\omega) = \int \psi(|u(\omega)|) d\nu = \sum_{i=1}^n \left(\int \psi(|x_i|) d\nu \right) \chi_{A_i}(\omega), \omega \in \Omega.$$

Let $v \in L_0(\Omega, L_\psi(X, \nu))$ and $\{u_n\}$ be a sequence from $\Gamma(L_\psi(X, \nu))$ such that $\|v(\omega) - u_n(\omega)\|_{L_\psi(X, \nu)} \rightarrow 0$ as $n \rightarrow \infty$ for a.e. $\omega \in \Omega$. Fix $\omega \in \Omega$ for which $\|v(\omega) - u_n(\omega)\|_{L_\psi(X, \nu)} \rightarrow 0$. Let us show that $\lim_{n \rightarrow \infty} \Phi_0(u_n)(\omega) = \int \psi(|u(\omega)|) d\nu$. If not, then there exist $\varepsilon > 0$ and a sequence $\{u_{n_k}(\omega)\}$ such that

$$(1) \quad \left| \int \psi(|u(\omega)|) d\nu - \int \psi(|u_{n_k}(\omega)|) d\nu \right| > \varepsilon, \quad k = 1, 2, \dots$$

Choose a subsequence $a_s = u_{n_{k_s}}(\omega)$ which is (o) -converging to $u(\omega)$ in $L_\psi(X, \nu)$ [11, VII, §2]. Then the sequence $\{\psi(a_s)\}$ (o) -converges to $\psi(u(\omega))$ in $L_1(X, \nu)$, which contradicts (1).

Thus there exists a limit

$$\Phi_0(v)(\omega) := \int \psi(|v(\omega)|) d\nu = \lim_{n \rightarrow \infty} \int \psi(|u_n(\omega)|) d\nu = \lim_{n \rightarrow \infty} \Phi_0(u_n)(\omega),$$

for a.e. $\omega \in \Omega$, in particular, $\Phi_0(v) \in \mathcal{L}_0(\Omega)$. Let $\Phi(\tilde{u}) = [\Phi_0(u)]^\sim$. Clearly, Φ is a component-invariant L_0 -modulator on $L_0(\Omega, L_\psi(X, \nu))$, in addition $\Phi(t\mathbf{1}) = \psi(t)\Phi(\mathbf{1}), t \geq 0$.

If $\tilde{u} \in L_0(\Omega, L_\psi(X, \nu)), \lambda \in \mathcal{P}(L_0)$, then

$$\begin{aligned} \Phi(\lambda^{-1}\tilde{u}) \leq \mathbf{1} &\Leftrightarrow \int \psi(\lambda^{-1}(\omega)|u(\omega)|) d\nu \leq 1 \text{ a.e.} \Leftrightarrow \\ \|u(\omega)\|_{L_\psi(X, \nu)} \leq \lambda(\omega) \text{ a.e.} &\Leftrightarrow \|\tilde{u}\| \leq \lambda. \end{aligned}$$

Hence,

$$\|\tilde{u}\| = \inf\{\lambda \in \mathcal{P}(L_0) : \|\tilde{u}\| \leq \lambda\} = \inf\{\lambda \in \mathcal{P}(L_0) : \Phi(\lambda^{-1}\tilde{u}) \leq \mathbf{1}\} = \|\tilde{u}\|_\Phi.$$

Thus, $(L_0(\Omega, L_\psi(X, \nu)), \|\cdot\|)$ is a component-invariant Orlicz L_0 -lattice. In addition, $(L_0(\Omega, L_\psi(X, \nu)), \|\cdot\|)$ and $L_\psi(L_0(\Omega, X), m)$ are isometrically isomorphic (Theorem 5.1). □

Remark 5.3. If $\psi(t) = t^p$, $p \geq 1$, then $L_0(\Omega, L_p(X, \nu))$ is isometrically isomorphic to $L_p(L_0(\Omega, X), m)$.

Acknowledgment. The author would like to thank Prof. Vladimir Chilin for helpful discussions, and also the referee for useful remarks.

REFERENCES

- [1] Busakhla N.Yu., *Measurable bundles of Dedekind logics*, Uzbek. Mat. Zh. no. 3 (1999), 29–34 (Russian).
- [2] Claas W.J., Zaanen A.C., *Orlicz lattices*, Comment. Math. Special Issue **1** (1978), 77–93.
- [3] Ganiev I.G., *Measurable bundles of lattices and their applications*, Investigations in Functional Analysis and its Applications, Nauka, Moscow, 2006, pp. 9–49 (Russian).
- [4] Gutman A.E., *Banach bundles in the theory of lattice-normed spaces*, Order-compatible Linear Operators, Trudy Inst. Mat. 29 (1995), Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk, 1995, pp. 63–211 (Russian).
- [5] Kantorovich L.V., Akilov G.P., *Functional Analysis*, Nauka, Moscow, 1977 (Russian).
- [6] Krein S.G., Petunin Yu.T., Semenov E.M., *Interpolation of Linear Operators*, Nauka, Moscow, 1978 (Russian); English translation: Translations of Mathematical Monographs, vol. 54, American Mathematical Society, Providence, 1982.
- [7] Kusraev A.G., *Vector Duality and its Applications*, Nauka, Novosibirsk, 1985 (Russian).
- [8] Kusraev A.G., *Dominated Operators*, Mathematics and its Applications, 519, Kluwer Academic Publishers, Dordrecht, 2000.
- [9] Lacey H.E., *The Isometric Theory of Classical Banach Spaces*, Springer, New York-Heidelberg, 1974.
- [10] Sarymsakov T.A., *Topological Semifields and their Applications*, Fan, Tashkent, 1989 (Russian).
- [11] Vladimirov D.A., *Boolean Algebras*, Nauka, Moscow, 1969 (Russian).
- [12] Vulikh B.Z., *Introduction to the Theory of Partially Ordered Spaces*, Fizmatgiz, Moscow, 1961 (Russian); English translation: Wolters-Noordhoff, Groningen, 1967.
- [13] Zakirov B.S., *The Luxemburg norm in the Orlicz-Kantorovich space*, Uzbek. Mat. Zh. no. 2 (2007), 32–44 (Russian).
- [14] Zakirov B.S., *Orlicz-Kantorovich lattices associated with an L_0 -valued measure*, Uzbek. Mat. Zh. no. 4 (2007), 18–34 (Russian).
- [15] Zakirov B.S., *Analytical representation of L_0 -valued homomorphisms in Orlicz-Kantorovich modules*, Mat. Trudy **10** (2007), no. 2, 112–141 (Russian).

INSTITUTE OF MATHEMATICS AND INFORMATION TECHNOLOGIES, UZBEKISTAN ACADEMY OF SCIENCES, 29, F. KHODJAEV STR., TASHKENT 100125, UZBEKISTAN

E-mail: botirzakirov@list.ru

(Received November 20, 2007, revised June 6, 2008)