

Tran Quoc Chien

Medium distances of probability-fuzzy points and an application to linear programming

*Kybernetika*, Vol. 25 (1989), No. 6, 494--504

Persistent URL: <http://dml.cz/dmlcz/124667>

## Terms of use:

© Institute of Information Theory and Automation AS CR, 1989

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## MEDIUM DISTANCES OF PROBABILITY FUZZY-POINTS AND AN APPLICATION TO LINEAR PROGRAMMING

TRAN QUOC CHIEN

In the paper the notions of probability-fuzzy points and medium distances in metric space are defined. These concepts are then applied for solving degenerated linear programs.

### 1. PROBABILITY-FUZZY POINTS

It is well known that a fuzzy point  $a$  in a space  $X$  is defined by a characteristic function

$$f_a: X \rightarrow [0, 1].$$

Here  $f_a(x)$  presents the probability that  $x = a$ . If we accept the addition operation of such possibilities then it may happen that the possibility of  $a$  is greater than 1 for some  $x \in X$ . This fact somehow contradicts our normal 'probabilistic' thinking. Therefore we propose now a new concept, namely probability-fuzzy points, which eliminates the mentioned discrepancy.

In what follows we suppose that a  $\sigma$ -algebra  $\mathcal{X}$  of subsets in the space  $X$  is given. A probability-fuzzy point  $a$  in  $X$  is characterized by a probability measure  $\mu_a$  on  $(X, \mathcal{X})$  (it means, among others,  $\mu_a(X) = 1$ ). Now for all  $A \in \mathcal{X}$  we have

$$0 \leq \mu_a(A) \leq 1$$

what expresses the probability that  $a \in A$ .

A deterministic point  $a \in X$  has its characteristic probability measure  $\mu_a$  of special form

$$\mu_a(A) = \begin{cases} 1 & \text{if } a \in A \\ 0 & \text{if } a \notin A \end{cases}$$

The space  $X$  is thus embedded into the space of all probability-fuzzy points, denoted by  $X_p$ .

Now we suppose that a  $\sigma$ -finite Haar measure  $\mu$  is defined on  $(X, \mathcal{X})$ , where  $X$

is equipped with a group operation '+'. If there exist density functions  $f_a$  and  $f_b$  of the probability measures  $\mu_a$  and  $\mu_b$  with respect to  $\mu$ , we can easily define the sum  $a + b$  by means of convolutions (see [2, 3]). Indeed, the density function corresponding to the sum  $a + b$  is naturally defined as follows

$$\begin{aligned} f_{a+b}(x) &= \int_X f_a(z) \cdot f_b(x - z) \, d\mu(z) \\ &= \int_X f_a(x - y) \cdot f_b(y) \, d\mu(y) \end{aligned}$$

and

$$\mu_{a+b}(A) = \int_A f_{a+b}(x) \, d\mu(x).$$

In virtue of Fubini's Theorem on iterated integrals (cf. [1]) we have

$$\begin{aligned} \mu_{a+b}(X) &= \int_X f_{a+b}(x) \, d\mu(x) \\ &= \int_X \left( \int_X f_a(z) \cdot f_b(-z + x) \, d\mu(z) \right) d\mu(x) \\ &= \int_X \left( \int_X f_a(z) \cdot f_b(-z + x) \, d\mu(x) \right) d\mu(z) \\ &= \int_X f_a(z) \cdot \left( \int_X f_b(-z + x) \, d\mu(x) \right) d\mu(z) \\ &= \int_X f_a(z) \, d\mu(z) = 1. \end{aligned}$$

So  $\mu_{a+b}$  is in fact a probability measure where all integrals are allowable because the function  $f_{a+b}(x)$  is integrable and  $\mathcal{X}$ -measurable (see [3], Statement 2.1). It should be stressed that by means of probability-fuzzy points one eliminates the trouble appearing with the 'min' operation in the definition of sums of fuzzy quantities as it is done in [2].

## 2. MEDIUM DISTANCES

Now we suppose that the space  $X$  is provided with a metric  $\varrho(x, y)$  defined for deterministic points in  $X$ . We shall extend the metric  $\varrho(x, y)$  to the space  $X_p$  of probability-fuzzy points by means of the notion of medium distance proposed as follows.

Let  $a$  and  $b$  be probability-fuzzy points with characteristic measures  $\mu_a$  and  $\mu_b$  respectively. Let  $\mu_a \otimes \mu_b$  denote the product measure of  $\mu_a$  and  $\mu_b$  on the Cartesian product  $(X, \mathcal{X}) \times (X, \mathcal{X})$ , where  $\mathcal{X}$  contains all open subsets of  $X$ . Then the function  $\varrho(x, y)$  is integrable and the Lebesgue integral

$$\int_{X \times X} \varrho(x, y) \, d\mu_a \otimes \mu_b$$

will be called the medium distance of points  $a$  and  $b$  and denoted by  $d_m(a, b)$ .

It is easily seen that if  $a$  and  $b$  are deterministic points then  $d_m(a, b)$  coincides with  $\varrho(a, b)$ . The symmetry property

$$d_m(a, b) = d_m(b, a) \quad \forall a, b \in X_p$$

immediately follows from the definition. As for the triangle inequality it needs a detailed proof.

Let  $a, b$  and  $c$  be arbitrary probability-fuzzy points. We have

$$\begin{aligned}
 d_m(a, b) &= \int_{X \times X} \varrho(x, y) \, d\mu_a \otimes \mu_b(x, y) \\
 &= \int_{X \times X} \varrho(x, y) \, d\mu_a \otimes \mu_b(x, y) \cdot \int_X d\mu_c(z) \\
 &= \int_{X \times X \times X} \varrho(x, y) \, d\mu_a \otimes \mu_b \otimes \mu_c(x, y, z) \\
 &\leq \int_{X \times X \times X} [\varrho(x, z) + \varrho(z, y)] \, d\mu_a \otimes \mu_b \otimes \mu_c(x, y, z) \\
 &= \int_{X \times X \times X} \varrho(x, z) \, d\mu_a \otimes \mu_b \otimes \mu_c(x, y, z) + \\
 &\quad + \int_{X \times X \times X} \varrho(y, z) \, d\mu_a \otimes \mu_b \otimes \mu_c(x, y, z) \\
 &= \int_{X \times X} \varrho(x, z) \, d\mu_a \otimes \mu_c(x, z) \cdot \int_X d\mu_b(y) + \\
 &\quad + \int_{X \times X} \varrho(y, z) \, d\mu_b \otimes \mu_c(y, z) \cdot \int_X d\mu_a(x) \\
 &= d_m(a, c) + d_m(c, b).
 \end{aligned}$$

Now, let  $a \in X_p$  and  $B \subset X$ . The medium distance from  $a$  to  $B$  is similarly defined as

$$d_m(a, B) = \int_X \varrho(x, B) \, d\mu_a(x),$$

where

$$\varrho(x, B) = \inf \{ \varrho(x, b) : b \in B \}$$

is the usual distance from  $x$  to the set  $B$ .

*Remark.* The medium distance does not fulfill the property

$$d_m(a, b) = 0 \Leftrightarrow a = b$$

and it prevents  $d_m(a, b)$  to be a metric in the traditional sense. However, if we consider probability-fuzzy points as random points varying with respect to time, then the fact that the distance between a point and itself is positive is acceptable. Here it is in place to remind the famous statement of Heracleitos: 'One cannot enter twice into the same river'.

### 3. EXAMPLE

Let  $X = \mathbb{R}^n$  and

$$S_a = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : x_i \leq c \, \forall i \text{ and } \sum_{i=1}^n x_i = a\}$$

where  $c$  is a fixed positive constant and  $a$  is a parameter satisfying

$$(3.1) \quad |a| \leq c \cdot n.$$

Since  $S_a$  are parallel affine spaces of dimension  $n - 1$  there exists the same positive atomless measure  $\nu$ , on the  $\sigma$ -algebra of measurable sets, for all of them, similarly as in  $\mathbb{R}^{n-1}$ .

Further, we define  $p_a$  as a probability-fuzzy point with uniform probability distribution on  $S_a$ , i.e.  $p_a$  has the density function  $f_a$  on  $S_a$  such that

$$f_a(x) = 1/\nu(S_a) \quad \forall x \in S_a.$$

The characteristic measure  $\mu_a$  of  $p_a$  is then defined by

$$\mu_a(S) = v(S \cap S_a) / v(S_a) \quad \forall S \subset \mathbb{R}^n.$$

Let us denote  $\mathbb{R}_+^n = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : x_i \geq 0 \forall i = 1, \dots, n\}$ .

We shall consider the medium distance from  $p_a$  to  $\mathbb{R}_+^n$  and prove that, as a function of parameter  $a$ ,  $d_m(p_a, \mathbb{R}_+^n)$  is decreasingly monotone.

Putting

$$C = \underbrace{(c, \dots, c)}_{n\text{-times}} \in \mathbb{R}^n$$

then the following lemmas are easily verifiable.

**Lemma 3.1.** For all  $a \in [0, nc]$  we have

$$(3.2) \quad S_a - C = \lambda_a(S_0 - C),$$

where

$$(3.3) \quad \lambda_a = (nc - a) / nc.$$

**Lemma 3.2.** For all  $a \in [0, nc]$  we have

$$\begin{aligned} v(S_a) &= v(S_a - C) = \\ &= \lambda_a^{n-1} v(S_0 - C) = \lambda_a^{n-1} v(S_0) \end{aligned}$$

(see Fig. 1).




Fig. 1.

**Statement 3.1.** If  $nc \geq a > b \geq 0$ , then

$$d_m(p_a, \mathbb{R}_+^n) < d_m(p_b, \mathbb{R}_+^n).$$

**Proof.** We have

$$\begin{aligned} d_m(p_a, \mathbb{R}_+^n) &= \frac{1}{v(S_a)} \int_{S_a} \left[ \sum_{x_i < 0} x_i^2 \right]^{1/2} dv(x) = \frac{1}{\lambda_a^{n-1} v(S_0)} \int_{S_a - C} \left[ \sum_{y_i < -c} (y_i + c)^2 \right]^{1/2} dv(y) = \\ &\quad \text{(by Lemma 3.2 and substitution } y = x - C) \\ &= \frac{1}{\lambda_a^{n-1} v(S_0)} \int_{S_0 - C} \left[ \sum_{z_i < -c/\lambda_a} (\lambda_a z_i + c)^2 \right]^{1/2} \lambda_a^{n-1} dv(z) = \\ &\quad \text{(by substitution } y = \lambda_a \cdot z) \\ &= \frac{1}{v(S_0)} \int_{S_0 - C} \left[ \sum_{z_i < -c/\lambda_a} (\lambda_a z_i + c)^2 \right]^{1/2} dv(z). \end{aligned}$$

Analogically,

$$d_m(p_b, \mathbb{R}_+^n) = \frac{1}{v(S_0)} \int_{S_0 - c} \left[ \sum_{z_i < -c/\lambda_b} (\lambda_b z_i + c)^2 \right]^{1/2} dv(z).$$

Since  $a > b$  one has  $\lambda_a < \lambda_b$  (see (3.3)). Then

$$-c/\lambda_a < -c/\lambda_b$$

and hence

$$(\lambda_a \cdot z_i + c)^2 < (\lambda_b \cdot z_i + c)^2 \quad \text{for such } z_i \text{ that } z_i < -c/\lambda_a.$$

Consequently we obtain

$$d_m(p_a, \mathbb{R}_+^n) < d_m(p_b, \mathbb{R}_+^n)$$

and the proof is complete. □

Now consider the case  $-nc \leq a \leq 0$ . Obviously

**Lemma 3.3.** For all  $a \in [-nc, 0]$  we have

$$(3.4) \quad S_a + C = \beta_a(S_0 + C)$$

where

$$(3.5) \quad \beta_a = (nc + a)/nc.$$

**Lemma 3.4.** For all  $a \in [-nc, 0]$  we have

$$v(S_a) = v(S_a + C) = \beta_a^{n-1} v(S_0 + C) = \beta_a^{n-1} v(S_0)$$

(see Fig. 2).




Fig. 2.

**Statement 3.2.** If  $0 \geq a > b \geq -nc$ , then

$$d_m(p_a, \mathbb{R}_+^n) < d_m(p_b, \mathbb{R}_+^n).$$

**Proof.** We have

$$\begin{aligned} d_m(p_a, \mathbb{R}_+^n) &= \frac{1}{v(S_a)} \int_{S_a} \left[ \sum_{x_i < 0} x_i^2 \right]^{1/2} dv(x) = \\ &= \frac{1}{\beta_a^{n-1} v(S_0)} \int_{S_a + c} \left[ \sum_{y_i < c} (y_i - c)^2 \right]^{1/2} dv(x) = \\ &\quad \text{(by Lemma 3.4 and substitution } y = x + C) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\beta_a^{n-1} v(S_0)} \int_{S_0+c} \left[ \sum_{z_i < c/\beta_a} (\beta_a z_i - c)^2 \right]^{1/2} \beta_a^{n-1} dv(z) = \\
&\quad \text{(by substitution } y = \beta_a \cdot z \text{)} \\
&= \frac{1}{v(S_0)} \int_{S_0+c} \left[ \sum_{z_i < c/\beta_a} (\beta_a z_i - c)^2 \right]^{1/2} dv(z).
\end{aligned}$$

Analogously

$$d_m(p_b, \mathbb{R}_+^n) = \frac{1}{v(S_0)} \int_{S_0+c} \left[ \sum_{z_i < c/\beta_b} (\beta_b z_i - c)^2 \right]^{1/2} dv(z).$$

Since  $a > b$  one has  $\beta_a > \beta_b$  (see 3.5). Then

$$c/\beta_a < c/\beta_b$$

and hence

$$(\beta_a \cdot z_i - c)^2 < (\beta_b \cdot z_i - c)^2 \quad \text{for such } z_i \text{ that } z_i < c/\beta_a.$$

Consequently, we obtain

$$d_m(p_a, \mathbb{R}_+^n) < d_m(p_b, \mathbb{R}_+^n)$$

and the proof is complete.  $\square$

#### 4. AN APPLICATION TO LINEAR PROGRAMMING

First, we briefly recall the simplex method for solving linear programs. Consider the problem

$$(4.1) \quad f = c \cdot x \rightarrow \max$$

s.t.

$$(4.2) \quad A \cdot x = b$$

$$(4.3) \quad x \geq 0$$

where  $c = (c_1, \dots, c_n) \in \mathbb{R}^n$ ,  $b = (b_1, \dots, b_m)' \in \mathbb{R}^m$ ,  $A = (a_{ij})_{m \times n}$  and  $x = (x_1, \dots, x_n)' \in \mathbb{R}^n$ .

Suppose that  $m < n$  and  $\text{rank}(A) = m$ . A vector  $x \in \mathbb{R}^n$  is a feasible solution if it fulfills (4.2) and (4.3); an optimal solution if it fulfills (4.1), (4.2) and (4.3); a basic solution if it is a feasible solution and the columns  $a_j$  of  $A$  corresponding to positive components  $x_j > 0$  are linearly independent. A basis of a basic solution  $x$  is an arbitrary system of  $m$  linearly independent columns of  $A$  including all those corresponding to positive components of  $x$ . A basic solution is degenerated if it has less than  $m$  positive components. Problems (4.1)–(4.3) is degenerated if it has degenerated solutions.

Let us have a basic solution  $x = (x_1, \dots, x_n)$  of problems (4.1)–(4.3) with the basis

$$A_\sigma = \{a_j: a_j \text{ is the } j\text{th column of } A \text{ and } j \in \sigma\}$$

the basic variables

$$x_\sigma = \{x_j: j \in \sigma\}$$

and the nonbasic variables

$$x_\omega = \{x_j: j \in \omega = \{1, \dots, n\} \setminus \sigma\}$$

where  $\sigma$  denotes the set of indexes of basic variables.

We have then the corresponding simplex tableau

**Tableau 1.**

$x_\sigma$	— $x_\omega$	
	1	... — $x_j$ ...
$f$	$\beta_{00}$	... $\beta_{0j}$ ...
⋮	⋮	⋮
$x_i$	$\beta_{i0}$	... $\beta_{ij}$ ...
⋮	⋮	⋮

where

$$(4.4) \quad B = (\beta_{ij}) = A_\sigma^{-1} \cdot A_\omega, \quad \beta = (\beta_{i0}) = A_\sigma^{-1} \cdot b, \quad \Delta = (\beta_{0j}) = c_\sigma \cdot B - c_\omega, \\ \beta_{00} = c_\sigma \cdot \beta, \quad c_\sigma = \{c_j: j \in \sigma\} \quad \text{and} \quad c_\omega = \{c_j: j \in \omega\}.$$

The simplex iteration procedure consists of the following steps.

*Step 1. Testing of optimality:*

$$(4.5) \quad \Delta = (\beta_{0j})_{j \in \omega} \geq 0$$

If (4.5) holds then  $x$  is an optimal solution. Stop. Otherwise go to Step 2.

*Step 2. Testing of boundedness:*

$$(4.6) \quad \exists j \in \omega: \beta_{0j} < 0 \quad \text{and} \quad \beta_{ij} \leq 0 \quad \forall i \in \sigma.$$

If (4.6) holds then  $c \cdot x$  is not bounded from above. Stop. Otherwise go to Step 3.

*Step 3. Constructing of an improved solution:*

Choose a  $\beta_{0s} < 0$ , usually the smallest one, and determine

$$(4.7) \quad \theta = \min \{ \beta_{i0} / \beta_{is}: i \in \sigma \text{ and } \beta_{is} > 0 \}.$$

Let  $\beta_{i0} / \beta_{is} = \theta$ . Perform the Jordan-Gauss elimination procedure:

$$(4.8) \quad \begin{aligned} \beta'_{ij} &= \beta_{ij} - \frac{\beta_{is}\beta_{is}}{\beta_{is}} & \forall i \in (\sigma' \setminus \{s\}) \cup \{0\} \\ & & \forall j \in (\sigma' \setminus \{t\}) \cup \{0\} \\ \beta'_{sj} &= \beta_{tj} / \beta_{ts} & \forall j \in (\omega' \setminus \{t\}) \cup \{0\} \\ \beta'_{st} &= 1 / \beta_{ts} \\ \beta'_{it} &= -\beta_{is} / \beta_{ts} & \forall i \in (\sigma' \setminus \{t\}) \cup \{0\} \end{aligned}$$

where

$$\sigma' = (\sigma \setminus \{t\}) \cup \{s\}, \quad \omega' = (\omega \setminus \{s\}) \cup \{t\}.$$



We obtain a new simplex tableau, Tableau 2, and a new basic solution  $x'$  with improved values

$$(4.9) \quad \beta'_{00} = \beta_{00} - \theta \cdot \beta_{0s}.$$

**Tableau 2.**

$x_{\sigma'}$	$x_{\omega'}$	
	1	$\dots - x_j \dots$
$f$	$\beta'_{00}$	$\dots \beta'_{0j} \dots$
$\vdots$	$\vdots$	$\vdots$
$x_i$	$\beta_{i0}$	$\dots \beta_{ij} \dots$
$\vdots$	$\vdots$	$\vdots$

Go to Step 1.

If the problem (4.1)–(4.3) is degenerated, there may exist several basic variables  $x_t$  with  $\beta_{t0}/\beta_{ts} = \theta$  and one faces the question which of them to choose. In practice, the variable with the smallest index is usually chosen. In order to avoid cycling one can use the lexicographical rule.

However these rules do not take into consideration the optimality criterion and the iteration procedure for finding an optimum solution is in many cases unnecessarily prolonged.

Let us consider  $\beta'_{0j}$ ,  $j \in \omega'$ . We know that

$$\beta'_{0t} = -\beta_{0s}/\beta_{ts} > 0.$$

Hence the optimality criterion (4.5) in this iteration reduces to

$$(4.10) \quad \Delta_t = (\beta'_{0j})_{j \in \omega' \setminus \{t\}} \geq 0$$

which is equivalent to

$$(4.11) \quad d(\Delta_t, \mathbb{R}_+^{n-m-1}) = 0$$

where  $d(\cdot, \cdot)$  denotes the normal distance in  $\mathbb{R}^{n-m-1}$ .

If we know the exact value of  $\Delta_t$ , it is not a problem to verify condition (4.11). However, if the information about  $\Delta_t$  is not complete,  $\Delta_t$  becomes a non-exactly known vector and one naturally turns to ‘fuzzy theory’ for help. In what follows we shall present one such possible help.

Suppose we know only the sum of all components  $\Delta_t$

$$(4.12) \quad \delta_t = \sum_{j \in \omega' \setminus \{t\}} \beta'_{0j}$$

and the data are random enough and do not exceed some limit  $c > 0$ . Then we can assume that  $\Delta_t$  is uniformly distributed on the set

$$S_{\delta_t} = \left\{ z = (z_1, \dots, z_{n-m-1}) : \sum_{i=1}^{n-m-1} z_i = \delta_t \text{ and } |z_i| \leq c \ \forall i = 1, \dots, n-m-1 \right\}.$$

Now  $\Delta_t$  becomes a probability-fuzzy point in  $\mathbb{R}^{n-m-1}$  and one disposes of the medium distance  $d_m(\Delta_t, \mathbb{R}^{n-m-1})$  considered in the preceding section. Put

$$M = \{k: k \in \sigma \text{ and } \beta_{k0}/\beta_{ks} = \theta\}.$$

Then the optimality condition (4.11) is equivalent to

$$(4.13) \quad d(\Delta_t, \mathbb{R}_+^{n-m-1}) = \min \{d(\Delta_k, \mathbb{R}_+^{n-m-1}): k \in M\} = 0.$$

So it natural to choose  $t \in M$  that such

$$(4.14) \quad d(\Delta_t, \mathbb{R}_+^{n-m-1}) = \min_{k \in M} d(\Delta_k, \mathbb{R}_+^{n-m-1}).$$

However the  $\Delta_k$  are not exactly known. Therefore we logically use the medium distance and (4.14) becomes

$$(4.15) \quad d_m(\Delta_t, \mathbb{R}_+^{n-m-1}) = \min_{k \in M} d_m(\Delta_k, \mathbb{R}_+^{n-m-1}).$$

It follows from the results of Section 3 that (4.15) is equivalent to

$$(4.16) \quad \delta_t = \max \{\delta_k: k \in M\}.$$

By virtue of (4.8) and (4.12) we have

$$(4.17) \quad \delta_k = \sum_{j \in \omega \setminus \{s\}} \beta_{0j} - \beta_{0s} \left( \sum_{j \in \omega \setminus \{s\}} \beta_{kj} \right) / \beta_{ks}.$$

Since  $\sum_{j \in \omega \setminus \{s\}} \beta_{0j}$  is invariable and  $-\beta_{0s} > 0$ , condition (4.16) is equivalent to the following condition

$$(4.18) \quad \left( \sum_{j \in \omega \setminus \{s\}} \beta_{tj} \right) / \beta_{ts} = \max_{k \in M} \left( \sum_{j \in \omega \setminus \{s\}} \beta_{kj} \right) / \beta_{ks}$$

which is called the Maximum-Sum Rule.

The following examples show the advantage of the Maximum-Sum Rule as compared with the lexicographical rule.

**Example 1.** Consider the program

$$\begin{aligned} f &= -2x_1 - 3x_2 - 3x_3 - 8x_4 - 4x_5 \rightarrow \max \\ \text{s.t.} & \\ & 2x_2 + x_4 + 7x_5 = 14 \\ & 5x_2 + x_3 + 2x_5 = 4 \\ & x_1 + x_2 + 2x_5 = 8 \\ & x_j \geq 0 \quad \forall j \end{aligned}$$

Applying the lexicographical rule we get an optimal solution after three simplex tableaux, see Tableau 3, 4 and 5, and the comment attached to them.

In the passage from Tableau 3 to Tableau 4  $\frac{14}{7} = \frac{4}{2} = 2$  is the minimum in (4.7), hence according to the lexicographical rule  $x_4$  is excluded from the basis.

**Tableau 3.**

	1	$-x_2$	$-x_5$
$f$	-140	-30	-62
$x_4$	14	2	7
$x_3$	4	5	2
$x_1$	8	1	2

**Tableau 4.**

	1	$-x_2$	$-x_4$
$f$	-16	$-86/7$	$62/7$
$x_5$	2	$2/7$	$1/7$
$x_3$	0	$31/7$	$-2/7$
$x_1$	4	$3/7$	$-2/7$

**Tableau 5.**

	1	$-x_3$	$-x_4$
$f$	-16	$86/31$	$250/31$
$x_5$	2		
$x_2$	0		
$x_1$	4		

However the Maximum-Sum Rule enables to obtain the optimal solution only after two simplex tableaux as it is shown by Tableau 6 and 7 below. Here  $\frac{5}{2} > \frac{2}{7}$ ,

**Tableau 6.**

	1	$-x_2$	$-x_5$
$f$	-140	-30	-62
$x_4$	14	2	7
$x_3$	4	5	2
$x_1$	8	1	2

**Tableau 7.**

	1	$-x_2$	$-x_3$
$f$	-16	125	31
$x_4$	0		
$x_5$	2		
$x_1$	4		

hence  $\frac{5}{2}$  is the maximum in (4.18). So according to the Maximum-Sum Rule  $x_3$  is excluded from the basis.

**Example 2.** Let us consider the well known Beale's example (see Gass [4], Chapter 7)

$$\begin{aligned}
 & \frac{3}{4}x_1 - 150x_2 + \frac{1}{50}x_3 - 6x_4 \rightarrow \max \\
 & \text{s.t.} \\
 & \frac{1}{4}x_1 - 60x_2 - \frac{1}{25}x_3 + 9x_4 + x_5 = 0 \\
 & \frac{1}{2}x_1 - 90x_2 - \frac{1}{50}x_3 + 3x_4 + x_6 = 0 \\
 & \qquad \qquad \qquad x_3 + x_7 = 1 \\
 & \qquad \qquad \qquad x_j \geq 0 \quad \forall j
 \end{aligned}$$

This example is degenerated. If one chooses among the variables realizing the minimum in (4.7) the one with the smallest index, cycling will appear. In order to avoid cycling we can use the lexicographical rule and obtain the optimal solution after six simplex tableaux, see Gass [4], Chapter 7.

However, applying our Maximum-Sum Rule one gets the optimal solution much faster, after three simplex tableaux, as it is shown by Tableau 8, 9 and 10 below.

**Tableau 8.**

	1	$-x_1$	$-x_2$	$-x_3$	$-x_4$
$f$	0	$-3/4$	150	$-1/50$	6
$x_5$	0	$1/4$	$-60$	$-1/25$	9
$x_6$	0	$1/2$	$-90$	$-1/50$	3
$x_7$	1	0	0	1	0

**Tableau 9.**

	1	$-x_6$	$-x_2$	$-x_3$	$-x_4$
$f$	0	$3/2$	15	$-1/20$	$21/2$
$x_5$	0	$-1/2$	$-15$	$-3/100$	$15/2$
$x_1$	0	2	$-180$	$-1/25$	6
$x_7$	1	0	0	1	0

**Tableau 10.**

	1	$-x_6$	$-x_2$	$-x_7$	$-x_4$
$f$	$1/20$	$3/2$	15	$1/20$	$21/2$
$x_5$	$3/1000$				
$x_1$	$1/25$				
$x_3$	1				

In the passage from Tableau 8 to Tableau 9 we have  $(-60 - \frac{1}{25} + 9) : \frac{1}{4} = -204 \cdot 16$ ;  $(-90 - \frac{1}{50} + 3) : \frac{1}{2} = -174 \cdot 04$ . So according to the Maximum-Sum Rule  $x_6$  is excluded from the basis.

(Received January 25, 1988.)

#### REFERENCES

- [1] A. N. Kolmogorov and S. V. Fomin: Elements of Function Theory and Functional Analysis. Nauka, Moscow 1972. In Russian.
- [2] M. Mareš: How to handle fuzzy-quantities. *Kybernetika* 13 (1977), 1, 23–40.
- [3] M. Mareš: Fuzzy-quantities with real and integer values. *Kybernetika* 13 (1977), 1, 41–56.
- [4] S. I. Gass: Linear Programming – Methods and Applications. McGraw-Hill, New York 1969.

RNDr. Tran Quoc Chien, CSc., Department of Mathematics, Polytechnical School of Da-nang, Da-nang, Vietnam.