

Martin Kochol

Symmetrized and continuous generalization of transversals

*Mathematica Bohemica*, Vol. 121 (1996), No. 1, 95–106

Persistent URL: <http://dml.cz/dmlcz/125937>

## Terms of use:

© Institute of Mathematics AS CR, 1996

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SYMMETRIZED AND CONTINUOUS GENERALIZATION  
OF TRANSVERSALS

MARTIN KOCHOL,\* Bratislava

(Received September 19, 1994)

*Summary.* The theorem of Edmonds and Fulkerson states that the partial transversals of a finite family of sets form a matroid. The aim of this paper is to present a symmetrized and continuous generalization of this theorem.

*Keywords:* transversal, system of representatives, polymatroid

*AMS classification:* 05D15, 05B35, 52B40

## 1. INTRODUCTION

There are two classical results concerning both the transversal theory and the matroid theory. The first is the theorem of Rado [17], who established a necessary and sufficient condition for a finite family of sets to possess a transversal which is independent in a given matroid. The second result, stated by Edmonds and Fulkerson [4] (and also proved independently by Mirsky and Perfect [14]) states that the set of partial transversals of a finite family of sets form a matroid. There are many variations and generalizations of these two theorems. A comprehensive survey of this field can be found in the books of Mirsky [13] and Welsh [20].

In [8] and [9] we introduced  $\mathcal{M}$ -systems of representatives and  $\mathcal{M}$ -polytransversals. They present a new concept joining transversals and matroids. An  $\mathcal{M}$ -system of representatives of a finite family  $\mathcal{A} = (A_t : t \in T)$  of subsets of a finite set  $S$  is a family  $(x_t : t \in T)$  of elements of  $S$  such that  $x_t \in A_t$  for any  $t \in T$  and, for any  $s \in S$ , the set  $\{t \in T; x_t = s\}$  is independent in a given matroid  $M_s$ . Furthermore, the  $|S|$  dimensional vector  $(u_s : s \in S)$  where  $u_s = |\{t \in T; x_t = s\}|$  is called an

---

\*This research was partially supported by Grant of Slovak Academy of Sciences No. 2/1138/94 and by EC Cooperation Action IC 1000 "Algorithms for Future Technologies."

$\mathcal{M}$ -polytransversal of  $\mathcal{A}$ . In [9] we proved that the set of  $\mathcal{M}$ -polytransversals of  $\mathcal{A}$  forms the set of integral independent vectors of a polymatroid. This generalizes the theorem of Edmonds and Fulkerson.

Other generalizations of transversals and the theorem of Edmonds and Fulkerson are presented in [7] and [21].

Now we prove a symmetrized and continuous analogue of the results of [9]. As a motivation let us recall two theorems from transversal theory. We will express them in the language of bipartite graphs. A finite *bipartite graph*  $G = (S, T; E)$  consists of two finite disjoint vertex sets  $S, T$  and a set  $E$  of edges joining the vertex sets  $S$  and  $T$ . If  $X \subseteq S$  and  $Y \subseteq T$  we say that  $X = \{x_1, \dots, x_k\}$  can be *matched into*  $Y$  in  $G$  if there exists a set of edges joining each  $x_i$  to a distinct member of  $Y$  (in other words if the subgraph of  $G$  determined by  $X \cup Y$  has a matching which covers every vertex of  $X$ ). If  $X \subseteq S$  then  $\partial X$  is the set of vertices of  $T$  which are endpoints of an edge whose other endpoint is in  $X$ . The following theorem was proved by Brualdi [2]. Mirsky [13] calls it a symmetrized version of Rado's theorem.

**Theorem 1.** *Let  $G = (S, T; E)$  be a finite bipartite graph. Let  $M_1, M_2$  be matroids on  $S, T$  with rank functions  $\varrho_1, \varrho_2$ , respectively. Then there exists  $X \subseteq S$  with  $|X| = k$  such that  $X$  is independent in  $M_1$  and  $X$  can be matched into an independent set  $Y$  of  $M_2$ , if and only if for all  $X \subseteq S$ ,*

$$\varrho_1(S \setminus X) + \varrho_2(\partial X) \geq k.$$

The next theorem was proved by Perfect [15] (see also [20]) and generalizes the theorem of Edmonds and Fulkerson.

**Theorem 2.** *Let  $G = (S, T; E)$  be a finite bipartite graph. Let  $M$  be a matroid on  $T$  with rank function  $\varrho$ . Then the collection*

$$\{X: X \subseteq S, X \text{ can be matched in } G \text{ into an independent set of } M\}$$

*is the set of independent sets of a matroid  $M_1$  on  $S$  with rank function  $\varrho_1$  such that, for any  $X \subseteq S$ ,*

$$\varrho_1(X) = \min_{A \subseteq X} (\varrho(\partial A) + |X \setminus A|).$$

The aim of this paper is to show that symmetrized and continuous analogues of  $\mathcal{M}$ -polytransversals form a polymatroid. Our results generalize Theorems 1 and 2 but also the results from [7], [8], [9] and [21].

We assume familiarity with matroids and transversals. The main literature is the book of Welsh [20] where all basic results regarding matroids, polymatroids and transversals can be found. As other sources let us note [1], [5], [13] and [18].

## 2. PRELIMINARIES

Let  $\mathbb{R}_+$  ( $\mathbb{Z}_+$ ) denote the set of nonnegative real (integer) numbers. If  $S$  is a finite set, then denote by  $\mathbb{R}_+^S$  ( $\mathbb{Z}_+^S$ ) the space of real (integer) valued nonnegative vectors with coordinates indexed by  $S$ . Similarly, if also  $T$  is finite, then  $\mathbb{R}_+^{S \times T}$  ( $\mathbb{Z}_+^{S \times T}$ ) denotes the space of real (integer) valued nonnegative vectors with coordinates indexed by  $S \times T$ . For example

$$\begin{aligned}\mathbb{R}_+^S &= \{\mathbf{u} = (u_s : s \in S); u_s \in \mathbb{R}_+\}, \\ \mathbb{Z}_+^S &= \{\mathbf{u} = (u_s : s \in S); u_s \in \mathbb{Z}_+\}, \\ \mathbb{R}_+^{S \times T} &= \{\mathbf{u} = (u_{s,t} : s \in S, t \in T); u_{s,t} \in \mathbb{R}_+\}.\end{aligned}$$

For each  $\mathbf{x} \in \mathbb{R}_+^S$  and  $s \in S$  denote the  $s$ th coordinate of  $\mathbf{x}$  by  $x(s)$ . For  $\mathbf{x} \in \mathbb{R}_+^S$  and  $A \subseteq S$  we define  $\mathbf{x}(A) = \sum_{s \in A} x(s)$ , and  $\mathbf{x}|_A$  denotes the restriction of  $\mathbf{x}$  to  $A$ . We call the quantity  $|\mathbf{x}| = \mathbf{x}(S) = \sum_{s \in S} x(s)$  the *modulus* of  $\mathbf{x}$ .

A *polymatroid*  $\mathbb{P}$  (on  $S$ ) is a pair  $(S, \varrho)$  where  $S$ , the *ground set*, is a non-empty finite set and  $\varrho$ , the *ground set rank function*, is a function  $\varrho: 2^S \rightarrow \mathbb{R}_+$  such that

- (1)  $\varrho(\emptyset) = 0$ ,
- (2) if  $A \subseteq B \subseteq S$  then  $\varrho A \leq \varrho B$ ,
- (3) if  $A, B \subseteq S$  then  $\varrho A + \varrho B \geq \varrho(A \cup B) + \varrho(A \cap B)$ .

(Items (2) and (3) state that  $\varrho$  is *monotone* and *submodular*, respectively.) Then a vector  $\mathbf{u} \in \mathbb{R}_+^S$  such that  $\mathbf{u}(X) \leq \varrho X$  for all  $X \subseteq S$  is called an *independent vector* of  $\mathbb{P}$ .

If  $\varrho: 2^S \rightarrow \mathbb{Z}_+$  then  $\mathbb{P} = (S, \varrho)$  is called an *integral* polymatroid. Furthermore, if  $\varrho(\{s\}) = 0, 1$  for any  $s \in S$  then  $\mathbb{P}$  is called a *matroid*. One of the most important properties of polymatroids is expressed in the following theorem (see [3], [12]) known as the polymatroid intersection theorem of Edmonds.

**Theorem 3.** Let  $\mathbb{P}_1 = (S, \varrho_1)$  and  $\mathbb{P}_2 = (S, \varrho_2)$  be polymatroids and let  $k \in \mathbb{R}_+$ . Then there exists a vector  $\mathbf{u} \in \mathbb{R}_+^S$  independent in both  $\mathbb{P}_1$  and  $\mathbb{P}_2$  and with modulus at least  $k$  if and only if for all subsets  $X \subseteq S$

$$\varrho_1(X) + \varrho_2(S \setminus X) \geq k.$$

Furthermore, if both  $\mathbb{P}_1, \mathbb{P}_2$  are integral we may insist that the vector  $\mathbf{u}$  be integral.

If  $\mathbb{P} = (S, \varrho)$  is a polymatroid and  $k \in \mathbb{R}_+$ , then it is easy to check that  $\mathbf{P}^{(k)} = (S, \varrho^{(k)})$  such that  $\varrho^{(k)}(X) = \min\{k, \varrho X\}$  ( $X \subseteq S$ ) is polymatroid. We call  $\mathbf{P}^{(k)}$  the *truncation* of  $\mathbb{P}$  at  $k$ .

If  $\mathbf{P} = (S, \varrho)$  is a polymatroid and  $\emptyset \neq X \subseteq S$  then  $\mathbb{P}^{(X)} = (X, \varrho^{(X)})$  (where  $\varrho^{(X)}$  is the restriction of  $\varrho$  to  $X$ ) is a polymatroid. We call  $\mathbb{P}^{(X)}$  the *restriction* of  $\mathbf{P}$  to  $X$ .

Let  $I, S_i$  ( $i \in I$ ) be finite sets,  $S_i \cap S_j = \emptyset$  for any  $i \neq j$  and let  $\mathbf{P}_i = (S_i, \varrho_i)$  ( $i \in I$ ) be polymatroids. Let  $S = \bigcup_{i \in I} S_i$  and  $\varrho: 2^S \rightarrow \mathbb{R}_+$  be such that for any  $X \subseteq S$ ,

$$\varrho(X) = \sum_{i \in I} \varrho_i(X \cap S_i).$$

Then  $\mathbf{P} = (S, \varrho)$  is a polymatroid. We call  $\mathbf{P}$  the *product* of the polymatroids  $\mathbf{P}_i$  ( $i \in I$ ) and denote it by  $\prod_{i \in I} \mathbf{P}_i$ .

Clearly, if  $\mathbf{P}$  is integral and  $k \in \mathbb{Z}_+$  then  $\mathbb{P}^{(k)}$  and  $\mathbb{P}^{(X)}$  are integral. If  $\mathbf{P}_i$  ( $i \in I$ ) are integral then  $\prod_{i \in I} \mathbf{P}_i$  is integral.

Finally, if  $k \in \mathbb{R}_+$ , denote by  $\mathbf{U}_{k,S}$  the polymatroid  $(S, \varrho)$  such that  $\varrho X = k|X|$  for any  $X \subseteq S$ .

Now we introduce the main notions of this paper. Throughout the paper let  $S, T$  be two disjoint finite sets. Let  $\mathcal{P}_S = (\mathbb{P}_s = (T, \varrho_s): s \in S)$ ,  $\mathcal{P}_T = (\mathbb{P}_t = (S, \varrho_t): t \in T)$  be systems of polymatroids, let  $\mathbb{P}_2 = (T, \varrho_2)$  be a polymatroid and  $X \subseteq S, J \subseteq T$ .

A vector  $\mathbf{a} = (a_{s,t}: s \in X, t \in J) \in \mathbb{R}_+^{X \times J}$  is called an  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -*system of representatives* (in abbreviation  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -SR) if:

- the vector  $\mathbf{a}_s = (a_{s,t}: t \in J) \in \mathbb{R}_+^J$  is independent in  $\mathbb{P}_s$  for any  $s \in X$ ,
- the vector  $\mathbf{a}_t = (a_{s,t}: s \in X) \in \mathbb{R}_+^X$  is independent in  $\mathbb{P}_t$  for any  $t \in J$ ,
- the vector  $\mathbf{v} = (v_t = \sum_{s \in X} a_{s,t}: t \in J) \in \mathbb{R}_+^J$  is independent in  $\mathbb{P}_2$ .

Furthermore, the vector  $\mathbf{u} = (u_s = \sum_{t \in J} a_{s,t}: s \in X) \in \mathbb{R}_+^X$  is called an  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -*polytransversal*. In this case  $\mathbf{a}$  is called an  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -*origin* of  $\mathbf{u}$ .

Moreover, if  $\mathbf{P}_1 = (S, \varrho_1)$  is a polymatroid, then a vector  $\mathbf{a}' = (a'_{s,t}: s \in X, t \in J) \in \mathbb{R}_+^{X \times J}$  is called an  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbf{P}_1, \mathbb{P}_2)$ -*system of representatives* (in abbreviation  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbf{P}_1, \mathbb{P}_2)$ -SR) if:

- the vector  $\mathbf{a}'$  is an  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -SR,
- the vector  $\mathbf{u}' = (u'_s = \sum_{t \in J} a'_{s,t}: s \in X) \in \mathbb{R}_+^X$  is independent in  $\mathbf{P}_1$ .

The notions of transversals and systems of distinct representatives and also their generalizations introduced in [7], [8], [9], [10] and [20] are in fact integral  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversals and  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbf{P}_1, \mathbb{P}_2)$ -systems of representatives for special classes of  $X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbf{P}_1$  and  $\mathbb{P}_2$ . The main distinction introduced here is that we deal with vectors whose coordinates are from  $\mathbb{R}_+$  and not only from  $\mathbb{Z}_+$ . In this way we obtain a "continuous" analogue of transversals and systems of representatives. On the other hand our results presented in the next

section (Theorems 4 and 5) remain true if we deal only with integral vectors and integral polymatroids. Thus we generalize the results from [7], [8], [9], [10], [20] and [21].

At the end of this section we introduce another notation. Let  $Z \subseteq S \times T$ . Then denote

$$(4) \quad Z_{/s} = \{t \in T: (s, t) \in Z\} \text{ for any } s \in S,$$

$$(5) \quad Z_{/t} = \{s \in S: (s, t) \in Z\} \text{ for any } t \in T.$$

### 3. THE MAIN RESULTS

Primarily we generalize the operation product of polymatroids.

**Lemma 1.** *Let  $I, S_i$  ( $i \in I$ ) be finite sets,  $S_i \cap S_j = \emptyset$  for any  $i \neq j$ . Let  $\mathbb{P}_i = (S_i, \varrho_i)$  ( $i \in I$ ) and  $\mathbb{P}' = (I, \varrho')$  be (integral) polymatroids. Take  $S = \bigcup_{i \in I} S_i$  and  $\varrho: 2^S \rightarrow \mathbb{R}_+$  such that for any  $X \subseteq S$ ,*

$$(6) \quad \varrho(X) = \min_{L \subseteq I} \left( \varrho'(I \setminus L) + \sum_{i \in L} \varrho_i(X \cap S_i) \right).$$

*Then  $\mathbb{P} = (S, \varrho)$  is an (integral) polymatroid. Moreover, a vector  $\mathbf{a} = (a_s: s \in S) \in \mathbb{R}_+^S$  is independent in  $\mathbb{P}$  if and only if it is independent in  $\prod_{i \in I} \mathbb{P}_i$  and the vector  $\mathbf{u} = (u_i = \sum_{s \in S_i} a_s: i \in I) \in \mathbb{R}_+^I$  is independent in  $\mathbb{P}'$ . We will denote  $\mathbb{P}$  by  $\mathbb{P}^I \prod_{i \in I} \mathbb{P}_i$ .*

*Proof.* It is easy to check that  $\varrho$  is monotone and  $\varrho(\emptyset) = 0$ . Let  $X$  ( $X'$ ) be a subset of  $S$  and let  $L$  ( $L'$ ) be the subset of  $I$  for which the minimum occurs in (6). Then using the monotonicity and submodularity of  $\varrho', \varrho_i$  ( $i \in I$ ) we get

$$\begin{aligned} \varrho X + \varrho X' &= \varrho'(I \setminus L) + \sum_{i \in L} \varrho_i(X \cap S_i) + \varrho'(I \setminus L') + \sum_{i \in L'} \varrho_i(X' \cap S_i) \\ &\geq \varrho'(I \setminus (L \cap L')) + \varrho'(I \setminus (L \cup L')) \\ &\quad + \sum_{i \in L \cap L'} \varrho_i((X \cup X') \cap S_i) + \sum_{i \in L \cup L'} \varrho_i((X \cap X') \cap S_i) \\ &\geq \varrho(X \cup X') + \varrho(X \cap X'). \end{aligned}$$

Thus  $\varrho$  is submodular and  $\mathbb{P} = (S, \varrho)$  is a polymatroid.

Take  $\varphi: S \rightarrow I$  such that  $\varphi(x) = i$  iff  $x \in S_i$  ( $i \in I$ ). Let  $\varrho_1: 2^S \rightarrow \mathbb{R}_+$  be such that  $\varrho_1(X) = \varrho'(\varphi(X))$  for any  $X \subseteq S$  ( $\varphi(X) = \{\varphi(x); x \in X\}$ ). Then it is easy

to check that  $\mathbb{P}_1 = (S, \varrho_1)$  is a polymatroid and that  $\mathbf{a} = (a_s: s \in S) \in \mathbb{R}_+^S$  is independent in  $\mathbb{P}_1$  iff  $(\sum_{s \in S_i} a_s: i \in I) \in \mathbb{R}_+^I$  is independent in  $\mathbb{P}'$ . Finally, let  $\mathbb{P}_2 = (S, \varrho_2)$  denote the polymatroid  $\prod_{i \in I} \mathbb{P}_i$ . Then for any  $X \subseteq S$ ,

$$(7) \quad \varrho(X) = \min_{A \subseteq X} (\varrho_1(X \setminus A) + \varrho_2(A)).$$

Let  $\mathbf{a} \in \mathbb{R}_+^S$  be independent in  $\mathbb{P}$ . Then  $\mathbf{a}(X) \leq \varrho X$  and it follows from (7) that  $\mathbf{a}$  is independent in both  $\mathbb{P}_1$  and  $\mathbb{P}_2$ . On the other hand let  $\mathbf{a}$  be independent in both  $\mathbb{P}_1$  and  $\mathbb{P}_2$ . Then, for any  $X \subseteq S$ ,  $\mathbf{a}|_X$  is independent in both  $\mathbb{P}_1^{(X)}$  and  $\mathbb{P}_2^{(X)}$  (the restrictions of  $\mathbb{P}_1$  and  $\mathbb{P}_2$  to  $X$ , respectively), and from Theorem 3 and (7) it follows that  $\mathbf{a}(X) \leq \varrho X$ . Thus  $\mathbf{a}$  is independent in  $\mathbb{P}$ .

Finally, if  $\varrho', \varrho_i$  ( $i \in I$ ) are integral then also  $\varrho$  is integral.  $\square$

Now we generalize Theorem 1 to  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_1, \mathbb{P}_2)$ -systems of representatives.

**Theorem 4.** *Let  $S, T$  be finite sets, let  $\mathcal{P}_S = (\mathbb{P}_s = (T, \varrho_s): s \in S)$ ,  $\mathcal{P}_T = (\mathbb{P}_t = (S, \varrho_t): t \in T)$  be systems of (integral) polymatroids, let  $\mathbb{P}_1 = (S, \varrho_1)$ ,  $\mathbb{P}_2 = (T, \varrho_2)$  be (integral) polymatroids and let  $k \in \mathbb{R}_+$ . Then there exists an (integral)  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_1, \mathbb{P}_2)$ -system of representatives with modulus at least  $k$  if and only if*

$$\min_{X \subseteq S, J \subseteq T} \left( \varrho_1(S \setminus X) + \varrho_2(T \setminus J) \right. \\ \left. + \min_{Z \subseteq X \times J} \left( \sum_{s \in X} \varrho_s(Z/s) + \sum_{t \in J} \varrho_t((X \times J) \setminus Z)/t \right) \right) \geq k.$$

**Proof.** Take  $\mathbb{P}'_s = (\{s\} \times T, \varrho'_s)$  such that  $\varrho'_s(\{s\} \times J) = \varrho_s(J)$  ( $s \in S, J \subseteq T$ ) and  $\mathbb{P}'_t = (S \times \{t\}, \varrho'_t)$  such that  $\varrho'_t(X \times \{t\}) = \varrho_t(X)$  ( $t \in T, X \subseteq S$ ). Take the polymatroids  $\mathbb{P}_S, \mathbb{P}_T$  on  $S \times T$  such that  $\mathbb{P}_S = \mathbb{P}_1 | \prod_{s \in S} \mathbb{P}'_s$  and  $\mathbb{P}_T = \mathbb{P}_2 | \prod_{t \in T} \mathbb{P}'_t$ . Then  $\mathbf{a} \in \mathbb{R}_+^{S \times T}$  is an  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_1, \mathbb{P}_2)$ -SR iff  $\mathbf{a}$  is independent in both  $\mathbb{P}_S$  and  $\mathbb{P}_T$ , and the theorem follows from Theorem 3.  $\square$

The following theorem is a generalization of Theorem 2 but also results from [7] and [9].

**Theorem 5.** *Let  $S, T$  be finite sets, let  $\mathcal{P}_S = (\mathbb{P}_s = (T, \varrho_s): s \in S)$ ,  $\mathcal{P}_T = (\mathbb{P}_t = (S, \varrho_t): t \in T)$  be systems of polymatroids and let  $\mathbb{P}_2 = (S, \varrho_2)$  be a polymatroid. Then  $\mathbf{u} = (u_s: s \in S) \in \mathbb{R}_+^S$  is an  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal if and*

only if it is an independent vector of the polymatroid  $\mathbb{P} = (S, \varrho)$  such that for any  $X \subseteq S$ ,

$$\varrho(X) = \min_{J \subseteq T} \left( \varrho_2(T \setminus J) + \min_{Z \subseteq X \times J} \left( \sum_{s \in X} \varrho_s(Z/s) + \sum_{t \in J} \varrho_t(((X \times J) \setminus Z)/t) \right) \right).$$

Furthermore, if  $\mathbb{P}_2, \mathbb{P}_s, \mathbb{P}_t$  ( $s \in S, t \in T$ ) are integral then also  $\mathbb{P}$  is integral. If also  $\mathbf{u} \in \mathbb{Z}_+^S$  is an integral  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal, then  $\mathbf{u}$  has an integral  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -origin.

**P r o o f.** It is easy to check that  $\varrho$  is monotone and  $\varrho(\emptyset) = 0$ . We prove submodularity.

Let  $X, Y \subseteq S$ . Choose  $J \subseteq T, K \subseteq T, Z \subseteq X \times J, V \subseteq Y \times K$  such that

$$\begin{aligned} \varrho(X) &= \varrho_2(T \setminus J) + \sum_{s \in X} \varrho_s(Z/s) + \sum_{t \in J} \varrho_t(((X \times J) \setminus Z)/t), \\ \varrho(Y) &= \varrho_2(T \setminus K) + \sum_{s \in Y} \varrho_s(V/s) + \sum_{t \in K} \varrho_t(((Y \times K) \setminus V)/t). \end{aligned}$$

Take the partition of  $(X \cup Y) \times (J \cup K)$  into ten sets  $A_1, A_2, \dots, A_{10}$ :


$$\begin{aligned} A_1 &= Z \cap V \subseteq (X \cap Y) \times (J \cap K), \\ A_2 &= ((X \cap Y) \times (J \cap K)) \setminus A_1, \\ A_3 &= (X \setminus Y) \times (J \cap K), \\ A_4 &= (Y \setminus X) \times (J \cap K), \\ A_5 &= (X \cap Y) \times (J \setminus K), \\ A_6 &= (X \setminus Y) \times (J \setminus K), \\ A_7 &= (Y \setminus X) \times (J \setminus K), \\ A_8 &= (X \cap Y) \times (K \setminus J), \\ A_9 &= (X \setminus Y) \times (K \setminus J), \\ A_{10} &= (Y \setminus X) \times (K \setminus J). \end{aligned}$$

Denote, for any  $i \in \{1, 2, \dots, 10\}$ ,

$$\begin{aligned} Z_i &= Z \cap A_i, \\ V_i &= V \cap A_i. \end{aligned}$$

Clearly  $Z = Z_1 \cup Z_2 \cup Z_3 \cup Z_5 \cup Z_6, V = V_1 \cup V_2 \cup V_4 \cup V_8 \cup V_{10}$ .  $V_1 = Z_1 = A_1$ , but  $V_i \cap Z_i = \emptyset$  for any  $i \in \{2, 3, \dots, 10\}$ . (See a symbolic representation of  $A_1, A_2, \dots, A_{10}$  in Fig. 1. Subsets of  $S$  and  $T$  are expressed as segments of the axis





$A_1 = Z_1 = V_1$      $R = \equiv$      $W = \text{||||}$   
 Fig. 1

and subsets of  $S \times T$  are depicted as parts of the plane in this figure. For instance  $Z$  and  $V$  are depicted as circles,  $A_1$  as the intersection of the circles and  $A_3, \dots, A_{10}$  as squares.)

Choose  $R, W$  and  $B_i, i \in \{2, 3, \dots, 10\}$ , such that

$$\begin{aligned}
 R &= Z_1 \cup Z_2 \cup Z_5 \cup V_2 \cup V_8 \quad (\subseteq (X \cap Y) \times (J \cup K)), \\
 W &= Z_1 \cup Z_3 \cup V_4 \quad (\subseteq (X \cup Y) \times (J \cap K)), \\
 B_i &= A_i \setminus (Z_i \cup V_i).
 \end{aligned}$$

Then we can check (see e.g. Fig. 1) that

$$\begin{aligned}
 ((X \cap Y) \times (J \cup K)) \setminus R &= B_2 \cup B_5 \cup B_8, \\
 ((X \cup Y) \times (J \cap K)) \setminus W &= B_2 \cup B_3 \cup B_4 \cup Z_2 \cup V_2.
 \end{aligned}$$

The sets  $A_1 (= Z_1 = V_1), Z_2, Z_3, Z_5, Z_6, V_2, V_4, V_8, V_{10}, B_2, \dots, B_{10}$  are pairwise disjoint. Using this fact and the submodularity and monotonicity of  $\varrho_2, \varrho_3, \varrho_t$  ( $s \in S$ ,

$t \in T$ ) we get

$$\begin{aligned}
& \varrho(X) + \varrho(Y) \\
&= \varrho_2(T \setminus J) + \sum_{s \in X} \varrho_s(Z/s) + \sum_{t \in J} \varrho_t(((X \times J) \setminus Z)/t) \\
&\quad + \varrho_2(T \setminus K) + \sum_{s \in Y} \varrho_s(V/s) + \sum_{t \in K} \varrho_t(((Y \times K) \setminus V)/t) \\
&= \varrho_2(T \setminus J) + \varrho_2(T \setminus K) + \sum_{s \in X \setminus Y} \varrho_s((Z_3 \cup Z_6)/s) + \sum_{s \in Y \setminus X} \varrho_s((V_4 \cup V_{10})/s) \\
&\quad + \sum_{s \in X \cap Y} \varrho_s((Z_1 \cup Z_2 \cup Z_5)/s) + \sum_{s \in X \cap Y} \varrho_s((V_1 \cup V_2 \cup V_8)/s) \\
&\quad + \sum_{t \in J \setminus K} \varrho_t((B_5 \cup B_8)/t) + \sum_{t \in K \setminus J} \varrho_t((B_8 \cup B_{10})/t) \\
&\quad + \sum_{t \in J \cap K} \varrho_t((B_2 \cup B_3 \cup V_2)/t) + \sum_{t \in J \cap K} \varrho_t((B_2 \cup B_4 \cup Z_2)/t) \\
&\geq \varrho_2(T \setminus (J \cap K)) \varrho_2(T \setminus (J \cup K)) + \sum_{s \in X \setminus Y} \varrho_s((Z_3)/s) + \sum_{s \in Y \setminus X} \varrho_s((V_4)/s) \\
&\quad + \sum_{s \in X \cap Y} \varrho_s((Z_1)/s) + \sum_{s \in X \cap Y} \varrho_s((Z_1 \cup Z_2 \cup Z_5 \cup V_2 \cup V_8)/s) \\
&\quad + \sum_{t \in J \setminus K} \varrho_t((B_5)/t) + \sum_{t \in K \setminus J} \varrho_t((B_8)/t) \\
&\quad + \sum_{t \in J \cap K} \varrho_t((B_2)/t) + \sum_{t \in J \cap K} \varrho_t((B_2 \cup B_3 \cup B_4 \cup Z_2 \cup V_2)/t) \\
&= \varrho_2(T \setminus (J \cup K)) + \sum_{s \in X \cap Y} \varrho_s(R/s) + \sum_{t \in J \cup K} \varrho_t(((X \cap Y) \times (J \cup K) \setminus R)/t) \\
&\quad + \varrho_2(T \setminus (J \cap K)) + \sum_{s \in X \cup Y} \varrho_s(W/s) \\
&\quad + \sum_{t \in J \cap K} \varrho_t(((X \cup Y) \times (J \cap K) \setminus W)/t) \\
&\geq \varrho(X \cup Y) + \varrho(X \cap Y).
\end{aligned}$$

Thus  $\varrho$  is submodular and  $\mathbb{P} = (S, \varrho)$  is a polymatroid.

Replace the polymatroid  $\mathbb{P}_1$  in Theorem 4 by  $\mathbb{U}_{k,S}$  where  $k$  is sufficiently large (e.g., let  $k = \sum_{s \in S} \varrho_s(T) + \sum_{t \in T} \varrho_t(S)$ ). Then it is easy to check that any  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{U}_{k,S}, \mathbb{P}_2)$ -SR is just an  $(X, J, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -SR and that  $\varrho X$  is the maximal modulus of an  $(X, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -SR (polytransversal).

Therefore, if  $\mathbf{u} \in \mathbb{R}_+^S$  is an  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal, then  $\mathbf{u}|_X$  is an  $(X, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal, and  $\mathbf{u}(X) \leq \varrho X$  for any  $X \subseteq S$ . Thus  $\mathbf{u}$  is independent in  $\mathbb{P}$ .

Let  $\mathbf{u} = (u_s : s \in S) \in \mathbb{R}_+^S$  be independent in  $\mathbb{P}$ . Then denote by  $\mathcal{P}_S^{(\mathbf{u})}$  the system of polymatroids  $(\mathbb{P}_s^{(u_s)} = (T, \varrho_s^{(u_s)}): s \in S)$ . Note that  $\varrho_s^{(u_s)}(J) = \min\{u_s, \varrho_s(J)\}$  ( $J \subseteq T, s \in S$ ). Let  $m$  denote the maximal modulus of an  $(S, T, \mathcal{P}_S^{(\mathbf{u})}, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal. Then replacing  $\mathbb{P}_1$  by  $\mathbb{U}_{k,S}$  ( $k$  sufficiently large) and  $\mathcal{P}_S$  by  $\mathcal{P}_S^{(\mathbf{u})}$  in Theorem 4 and applying the above argument we get

$$\begin{aligned} m &= \min_{J \subseteq T} \left( \varrho_2(T \setminus J) + \min_{Z \subseteq S \times J} \left( \sum_{s \in S} \varrho_s^{(u_s)}(Z/s) + \sum_{t \in J} \varrho_t(((S \times J) \setminus Z)/t) \right) \right) \\ &= \min_{J \subseteq T} \left( \varrho_2(T \setminus J) + \min_{Z \subseteq S \times J} \left( \sum_{s \in S} \min\{u_s, \varrho_s(Z/s)\} + \sum_{t \in J} \varrho_t(((S \times J) \setminus Z)/t) \right) \right). \end{aligned}$$

Let  $X = \{s \in S; u_s > \varrho_s(Z/s)\}$ . Then we can easily check that

$$\begin{aligned} m &= \mathbf{u}(S \setminus X) + \min_{J \subseteq T} \left( \varrho_2(T \setminus J) \right. \\ &\quad \left. + \min_{Z \subseteq X \times J} \left( \sum_{s \in X} \varrho_s(Z/s) + \sum_{t \in J} \varrho_t(((X \times J) \setminus Z)/t) \right) \right) \\ &= \mathbf{u}(S \setminus X) + \varrho_1(X) \geq \mathbf{u}(S \setminus X) + \mathbf{u}(X) = |\mathbf{u}|. \end{aligned}$$

Since  $\varrho_s^{(u_s)}(T) \leq u_s$  ( $s \in S$ ) then the inequality  $m \geq |\mathbf{u}|$  is possible iff  $\mathbf{u}$  is a  $(S, T, \mathcal{P}_S^{(\mathbf{u})}, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal with modulus  $m$ . Thus  $\mathbf{u}$  is also an  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal.

Therefore  $\mathbf{u} \in \mathbb{R}_+^S$  is an  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal if and only if  $\mathbf{u}$  is independent in  $\mathbb{P}$ .

Furthermore, if  $\mathbb{P}_2, \mathbb{P}_s, \mathbb{P}_t$  ( $s \in S, t \in T$ ) are integral then  $\mathbb{P}$  is integral. If also  $\mathbf{u}$  is integer valued then all polymatroids we have dealt with in the proof are integral. Thus, by Theorem 4, we can take an integer valued  $(S, T, \mathcal{P}_S^{(\mathbf{u})}, \mathcal{P}_T, \mathbb{P}_2)$ -SR of modulus  $m$ , then it must be an  $(S, T, \mathcal{P}_S^{(\mathbf{u})}, \mathcal{P}_T, \mathbb{P}_2)$ -origin of  $\mathbf{u}$ , and also an  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -origin of  $\mathbf{u}$ , concluding the proof.  $\square$

#### 4. CONCLUDING REMARKS

Note that Theorem 4 is equivalent with the intersection theorem of Edmonds (Theorem 3). But as pointed out by Schrijver [19], Theorem 3 is equivalent to many problems from combinatorial optimization. Thus our results are equivalent to them, too.

As pointed out by Poljak [16], especially interesting is the similarity with the following flow model of Lawler and Martel [11]. By a *polymatroidal flow network*  $\mathcal{F}$  we mean a directed multigraph  $G = (V, E)$  with a *source*  $s$ , a *sink*  $t$  and a collection

of polymatroids  $\mathbb{P}_v^+ = (\Delta_v^+, \varrho_v^+)$ ,  $\mathbb{P}_v^- = (\Delta_v^-, \varrho_v^-)$  where  $v \in V$  and  $\Delta_v^+$  ( $\Delta_v^-$ ) denotes the set of arcs directed into (out of)  $v$ . By a *flow* in  $\mathcal{F}$  we mean any vector  $f \in \mathbb{R}_+^E$ . A flow  $f$  is called *feasible* in  $\mathcal{F}$  if it satisfies the following conditions:

$$\begin{aligned} f|_{\Delta_v^+} &= f|_{\Delta_v^-} \text{ for any } v \in V \setminus \{s, t\}, \\ f|_{\Delta_v^+} &\text{ is independent in } \mathbb{P}_v^+ \text{ for any } v \in V, \\ f|_{\Delta_v^-} &\text{ is independent in } \mathbb{P}_v^- \text{ for any } v \in V. \end{aligned}$$

By a *value* of a feasible flow  $f$  we mean the quantity  $v = f(\Delta_v^-) - f(\Delta_v^+) = f(\Delta_v^+) - f(\Delta_v^-)$ . A polymatroidal network flow is called *integral* if  $\mathbb{P}_v^+$  and  $\mathbb{P}_v^-$  are integral for any  $v \in V$ .

An *arc-partitioned cut*  $(S, T, L, U)$  is defined by a partition of vertices into two sets  $S$  and  $T$  with  $s \in S$ ,  $t \in T$  and by a partition of the arcs directed from  $S$  to  $T$  into two sets  $L$  and  $U$ . The *capacity* of an arc partitioned cut is defined as

$$c(S, T, L, U) = \sum_{v \in S} \varrho_v^-(U \cap \Delta_v^-) + \sum_{v \in T} \varrho_v^+(L \cap \Delta_v^+).$$

In [11] it is shown that this flow model has the max-flow min-cut property.

**Theorem 6.** *Let  $\mathcal{F}$  be an (integral) polymatroidal flow network. Then the maximal value of an (integral) feasible flow is equal to the minimum capacity of an arc-partitioned cut.*

It is easy to check that Theorem 4 follows from Theorem 6. On the other hand Theorem 6 follows from Theorem 3 (see e.g. [19]) and, thus, also from Theorem 4. Therefore Theorems 3, 4, and 6 are pairwise equivalent.

Note that there exists no analogue of Theorem 5 in flow theory. On the other hand it presents a very natural generalization of results from transversal theory, especially those of Edmonds and Fulkerson [4], Mirsky and Perfect [14] and Perfect [15].

Theorem 5 describes in fact an operation on polymatroids. This “transversal” operation creates the polymatroid  $\mathbb{P}$  from a polymatroid  $\mathbb{P}_2$  and finite systems of polymatroids  $\mathcal{P}_S$  and  $\mathcal{P}_T$  (thus we can call  $\mathbb{P}$  the *polymatroid of*  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversals). It is easy to check that the operation from Lemma 1, the operations truncation and restriction on polymatroids, the polymatroid sum (see [18, pages 351–352]) and thus also the product of polymatroids can be characterized as special cases of the “transversal” operation.

It is well known (see e.g. [20], [1] for more details) that not every matroid can be characterized as a transversal matroid. Nevertheless, every polymatroid  $\mathbb{P} = (S, \varrho)$  can be characterized as an  $(S, T, \mathcal{P}_S, \mathcal{P}_T, \mathbb{P}_2)$ -polytransversal. It suffices to set  $T = \{1\}$ ,  $\mathcal{P}_T = (\mathbb{P}_1 = \mathbb{P})$ ,  $\mathcal{P}_S = (\mathbb{P}_s = \mathbb{U}_{k,S} : s \in S)$  and  $\mathbb{P}_2 = \mathbb{U}_{k,T}$ ,

where  $k$  is sufficiently large. Then it is routine to check that the polymatroid of  $(S, T, \mathcal{P}_S, \mathcal{P}_T, P_2)$ -polytransversals is equal to  $\mathcal{P}$ . The situation could change if we required some restrictions for polymatroids from  $\mathcal{P}_S$  and  $\mathcal{P}_T$ . For instance what will happen if  $\mathcal{P}_S$  and  $\mathcal{P}_T$  are systems of polymatroids of uniform matroids? This could generalize the characterization of transversal matroids (see [20] for more details).

#### References

- [1] *M. Aigner*: Combinatorial Theory. Springer, Berlin, 1979.
- [2] *R. A. Brualdi*: Common transversals and strong exchange systems. J. Combin. Theory Ser. B 3 (1970), 307–329.
- [3] *J. Edmonds*: Submodular functions, matroids and certain polyhedra. Combinatorial structures and their applications (R. Guy, H. Hanani, N. Sauer, J. Schönheim, eds.). Gordon and Breach, New York, 1970, pp. 69–87.
- [4] *J. Edmonds, D. R. Fulkerson*: Transversals and matroid partitions. J. Res. Nat. Bur. Standards Sect. B 69 (1965), 147–153.
- [5] *M. Grötschel, L. Lovász, A. Schrijver*: Geometric Algorithms and Combinatorial Optimization. Springer, Berlin, 1988.
- [6] *P. Hall*: On representatives of subsets. J. London Math. Soc. 10 (1935), 26–30.
- [7] *P. Horák*: Transversals and matroids. Topics in Combinatorics and Graph Theory (R. Bodendiek, R. Henn, eds.). Physica-Verlag, Heidelberg, 1990, pp. 381–389.
- [8] *M. Kochol*: The notion and basic properties of  $M$ -transversals. Discrete Math. 104 (1991), 191–196.
- [9] *M. Kochol*: About a generalization of transversals. Math. Bohem. 119 (1994), 143–149.
- [10] *M. Kochol*: Some generalizations of transversal theory. CSc. thesis, Bratislava, 1990. (In Slovak.)
- [11] *E. L. Lawler, C. U. Martel*: Computing maximal “polymatroidal” network flows. Math. Oper. Research 7 (1982), 334–347.
- [12] *C. J. H. McDiarmid*: Rado’s theorem for polymatroids. Proc. Cambridge Phil. Soc. 78 (1975), 263–281.
- [13] *L. Mirsky*: Transversal Theory. Academic Press, London, 1971.
- [14] *L. Mirsky, H. Perfect*: Applications of the notion of independence to combinatorial analysis. J. Combin. Theory 2 (1967), 327–357.
- [15] *H. Perfect*: Independence spaces and combinatorial problems. Proc. London Math. Soc. 19 (1969), 17–30.
- [16] *S. Poljak*: Personal communication.
- [17] *R. Rado*: A theorem on independence relations. Quart. J. Math. (Oxford) 13 (1942), 83–89.
- [18] *A. Recski*: Matroid Theory and its Applications in Electric Network Theory and in Statics. Springer, Berlin, 1989.
- [19] *A. Schrijver*: Total dual integrality from directed graphs, crossing families, and sub- and supermodular functions. Progress in Combinatorial Optimization (W. R. Pulleyblank, ed.). Academic Press, Toronto, 1984, pp. 315–362.
- [20] *D. J. A. Welsh*: Matroid Theory. Academic Press, London, 1976.
- [21] *D. R. Woodall*: Vector transversals. J. Combin. Theory Ser. B 32 (1982), 189–205.

*Author’s address*: Martin Kochol, Institute for Informatics, Slovak Academy of Sciences, P. O. Box 56, Dúbravská cesta 9, 840 00 Bratislava 4, Slovakia, e-mail: kochol@savba.sk.