

Miroslav Dont

A note on the parabolic variation

Mathematica Bohemica, Vol. 125 (2000), No. 3, 257–268

Persistent URL: <http://dml.cz/dmlcz/126129>

Terms of use:

© Institute of Mathematics AS CR, 2000

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON THE PARABOLIC VARIATION

MIROSLAV DONT, Praha

(Received May 14, 1998)

Abstract. A condition for solvability of an integral equation which is connected with the first boundary value problem for the heat equation is investigated. It is shown that if this condition is fulfilled then the boundary considered is $\frac{1}{2}$ -Hölder. Further, some simple concrete examples are examined.

Keywords: heat equation, boundary value problem, integral equations, parabolic variation

MSC 1991: 31A25, 31A20

Let φ be a continuous function with bounded variation on a compact interval (a, b) ,

$$K = \{[\varphi(t), t]; t \in (a, b)\}.$$

For $[x, t] \in \mathbb{R}^2$, $\alpha, r > 0$, $\alpha < +\infty$, let $n_{x,t}(r, \alpha)$ denote the number (finite or $+\infty$) of points of the set

$$K \cap \left\{ [\xi, \tau] \in \mathbb{R}^2; t - \tau = \left(\frac{\xi - x}{2\alpha} \right)^2, 0 < t - \tau < r \right\}.$$

For fixed $[x, t] \in \mathbb{R}^2$, $r > 0$, the function $n_{x,t}(r, \alpha)$ is a measurable function of the variable $\alpha \in (0, +\infty)$ and thus one can define

$$(1) \quad V_K(r; x, t) = \int_0^{+\infty} e^{-\alpha^2} n_{x,t}(r, \alpha) d\alpha.$$

Note that the function $V_K(+\infty; \cdot, \cdot)$ is called the parabolic variation of the set (curve) K . Parabolic variation was defined in [2] in connection with the study of a

Support of the Research Project J04/98/210000010 of Ministry of Education of the Czech Republic is gratefully acknowledged.

heat potential in \mathbb{R}^2 . Note that boundedness of the parabolic variation is a condition for the existence of limits of this potential on K and for the validity of an analogue of the jump formula.

For $[x, t] \in \mathbb{R}^2$, $t > a$, let $\alpha_{x,t}$ be the function defined on the interval $(a, \min\{t, b\})$ by

$$(2) \quad \alpha_{x,t}(\tau) = \frac{x - \varphi(\tau)}{2\sqrt{t - \tau}}.$$

The function $\alpha_{x,t}$ has locally finite variation on $(a, \min\{t, b\})$ (under the assumption that φ is of finite variation on (a, b)) and

$$(3) \quad V_K(r; x, t) = \int_{\max\{a, t-r\}}^{\min\{t, b\}} e^{-\alpha_{x,t}^2(\tau)} d(\text{var } \alpha_{x,t}(\tau))$$

whenever $\max\{a, t-r\} < \min\{t, b\}$ [otherwise $V_K(r; x, t) = 0$].

Let $[x, t] \in K$, $t > a$. It is known that if $V_K(r; x, t) < +\infty$ for some $r > 0$ then there is a limit (finite or infinite)

$$(4) \quad \alpha_{x,t}(t) = \lim_{\tau \rightarrow t-} \alpha_{x,t}(\tau)$$

[by this equality the value $\alpha_{x,t}(t)$ is defined]. Further, let G be the function defined on \mathbb{R}^1 by

$$(5) \quad G(t) = \begin{cases} 0, & t = -\infty, \\ \int_{-\infty}^t e^{-x^2} dx, & t > -\infty. \end{cases}$$

The condition

$$(6) \quad \lim_{\tau \rightarrow 0+} \sup_{t \in (a, b)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] < 1$$

plays a role in connection with the first boundary value problem for the heat equation on the set

$$\{[x, t] \in \mathbb{R}^2; t \in (a, b), x > \varphi(t)\}$$

or on the set

$$\{[x, t] \in \mathbb{R}^2; t \in (a, b), x < \varphi(t)\}.$$

Analogous conditions appear also in connection with the first boundary value problem for the heat equation on sets of the form

$$\{[x, t] \in \mathbb{R}^2; t \in (a, b), \varphi_1(t) < x < \varphi_2(t)\},$$

where φ_1, φ_2 are continuous functions of bounded variation on (a, b) . Under the condition (6) the relevant integral equation has a solution and the Fourier problem can be represented by the heat potential. It was shown in [1] that under the condition (6) the just mentioned integral equation is solvable also in the space of all bounded Baire functions and that the corresponding Neumann series converges.

Let us prove the following two assertions.

Proposition 1. *Let φ be a Lipschitz function on (a, b) . Then*

$$(7) \quad \lim_{r \rightarrow 0^+} \sup_{t \in (a, b)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] = 0.$$

Let φ be a continuous function of bounded variation on (a, b) , $a < a' < b' \leq b$ and suppose that φ is Lipschitz on (a', b') . Then

$$(8) \quad \lim_{r \rightarrow 0^+} \sup_{t \in (a' + \delta, b')} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] = 0$$

for any $\delta > 0$, $\delta < b' - a'$.

Proof. If φ is Lipschitz on (a, b) then clearly $\alpha_{\varphi(t), t}(t) = 0$ and thus

$$\left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| = 0$$

for each $t \in (a, b)$.

Recall a well known fact that if F is a non-negative continuous function on an interval $J \subset \mathbb{R}^1$, f, g continuous functions of locally bounded variation on J then

$$\int_J F \, d \operatorname{var}(fg) \leq \int_J F|f| \, d \operatorname{var} g + \int_J F|g| \, d \operatorname{var} f.$$

Let $k \in \mathbb{R}^1$ be such that for any $t_1, t_2 \in (a, b)$

$$|\varphi(t_1) - \varphi(t_2)| \leq k|t_1 - t_2|;$$

then $|\varphi'(\tau)| \leq k$ for almost all $\tau \in (a, b)$. Using the expression (3) for V_K we get for $t \in (a, b)$, $r > 0$,

$$\begin{aligned} V_K(r; \varphi(t), t) &= \int_{\max\{a, t-r\}}^t e^{-\alpha_{\varphi(t), t}^2(\tau)} d_r \text{var } \alpha_{\varphi(t), t}(\tau) \\ &\leq \int_{\max\{a, t-r\}}^t d_r \text{var } \frac{\varphi(t) - \varphi(\tau)}{2\sqrt{t-\tau}} \\ &\leq \int_{\max\{a, t-r\}}^t \frac{1}{2\sqrt{t-\tau}} d_r \text{var}(\varphi(t) - \varphi(\tau)) \\ &\quad + \int_{\max\{a, t-r\}}^t |\varphi(t) - \varphi(\tau)| d_r \text{var } \frac{1}{2\sqrt{t-\tau}} \\ &\leq \int_{\max\{a, t-r\}}^t \frac{|\varphi'(\tau)|}{2\sqrt{t-\tau}} d\tau + \int_{\max\{a, t-r\}}^t k(t-\tau) \frac{1}{4(t-\tau)^{3/2}} d\tau \\ &\leq \frac{3}{2}k \int_{\max\{a, t-r\}}^t \frac{d\tau}{2\sqrt{t-\tau}} \leq \frac{3}{2}k\sqrt{r}. \end{aligned}$$

Now we see that (7) is valid.

Denote for a while

$$K' = \{[\varphi(t), t]; t \in \langle a', b' \rangle\}.$$

Fix $\delta > 0$, $\delta < b' - a'$. Then for $r > 0$, $r < \delta$ and $t \in \langle a' + \delta, b' \rangle$

$$V_K(r; \varphi(t), t) = V_{K'}(r; \varphi(t), t).$$

The second part of the assertion follows now from the first part applied to the interval $\langle a', b' \rangle$. \square

If the condition (6) is fulfilled then the absolute values of limits

$$\alpha_{\varphi(t), t}(t) = \lim_{\tau \rightarrow t-} \frac{\varphi(t) - \varphi(\tau)}{2\sqrt{t-\tau}}$$

$[t \in \langle a, b \rangle]$ are bounded by a finite constant. This fact alone does not imply that φ is $\frac{1}{2}$ -Hölder. Nevertheless, the following assertion is valid.

Proposition 2. *Let φ be a continuous function of bounded variation on $\langle a, b \rangle$. If the condition (6) is fulfilled then φ is $\frac{1}{2}$ -Hölder on $\langle a, b \rangle$.*

Proof. Let $r > 0$ be such that

$$(9) \quad \sup_{t \in \langle a, b \rangle} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] = \lambda < 1.$$

Then, of course,

$$\left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \leq \lambda$$

for each $t \in (a, b)$ and hence there is a $c_0 \in \mathbb{R}^1$ such that for each $t \in (a, b)$

$$|\alpha_{\varphi(t), t}(t)| = \lim_{\tau \rightarrow t^-} \left| \frac{\varphi(t) - \varphi(\tau)}{2\sqrt{t - \tau}} \right| \leq c_0.$$

Choose $k \in \mathbb{R}^1$, $k > c_0$ such that

$$(10) \quad \frac{2}{\sqrt{\pi}} \int_0^k e^{-\alpha^2} d\alpha > \lambda.$$

Suppose that φ is not $\frac{1}{2}$ -Hölder continuous on (a, b) . Then there are $t_1, t_2 \in (a, b)$, $|t_1 - t_2| < r$, such that

$$(11) \quad |\varphi(t_1) - \varphi(t_2)| > 2k\sqrt{|t_1 - t_2|}.$$

We can assume that $t_1 > t_2$. Put

$$c = |\alpha_{\varphi(t_1), t_1}(t_1)| = \lim_{\tau \rightarrow t_1^-} \left| \frac{\varphi(t_1) - \varphi(\tau)}{2\sqrt{t_1 - \tau}} \right|.$$

We have $c < k$, of course. Let $\delta > 0$ be such that

$$c + \delta < k.$$

Then there is an $\varepsilon > 0$, $\varepsilon < t_1 - t_2$ such that

$$(12) \quad |\varphi(t_1) - \varphi(\tau)| < 2(c + \delta)\sqrt{t_1 - \tau}$$

for any $\tau \in (t_1 - \varepsilon, t_1)$. Let us show that whenever $\alpha \in \mathbb{R}^1$,

$$c + \delta < \alpha < k,$$

then

$$(13) \quad n_{\varphi(t_1), t_1}(r, \alpha) \geq 1;$$

recall that for $[x, t] \in \mathbb{R}^2$, $\alpha, r > 0$ the symbol $n_{x, t}(r, \alpha)$ stands for the number of points of the intersection

$$K \cap \left\{ [\xi, \tau] \in \mathbb{R}^2; t - \tau = \left(\frac{\xi - x}{2\alpha} \right)^2, 0 < t - \tau < r \right\}.$$

Consider for example the case $\varphi(t_2) > \varphi(t_1)$. Given $\alpha > c + \delta$ put

$$p = \left\{ [\xi, \tau] \in \mathbb{R}^2; t_1 - \tau = \left(\frac{\xi - \varphi(t_1)}{2\alpha} \right)^2, 0 < t_1 - \tau < r, \xi > \varphi(t_1) \right\}.$$

If $[\xi, \tau] \in p$, $\tau \in (t_1 - \varepsilon, t_1)$ then (12) yields

$$\xi - \varphi(t_1) = 2\alpha\sqrt{t_1 - \tau} > 2(c + \delta)\sqrt{t_1 - \tau} > \varphi(\tau) - \varphi(t_1),$$

that is

$$(14) \quad \xi > \varphi(\tau).$$

If $[\xi, \tau] \in p$, $\tau = t_2$, then (11) yields (if $\alpha < k$)

$$\xi - \varphi(t_1) = 2\alpha\sqrt{t_1 - t_2} < 2k\sqrt{t_1 - t_2} < \varphi(t_2) - \varphi(t_1),$$

that is

$$\xi < \varphi(t_2).$$

This, together with the fact that for $[\xi, \tau] \in p$, $\tau \in (t_1 - \varepsilon, t_1)$ the inequality (14) holds, implies that

$$K \cap p \neq \emptyset,$$

which means that $n_{\varphi(t_1), t_1}(r, \alpha) \geq 1$ for any $\alpha \in (c + \delta, k)$, indeed. Since $\delta > 0$ was arbitrary we see that (13) is valid for any $\alpha \in (c, k)$. Similarly in the case $\varphi(t_2) < \varphi(t_1)$.

Since

$$\left| 1 - \frac{2}{\sqrt{\pi}} G(q) \right| = \frac{2}{\sqrt{\pi}} \int_0^{|q|} e^{-\alpha^2} d\alpha$$

for $q \in {}^*\mathbb{R}^1$ we now obtain [using (1) and (10)]

$$\begin{aligned} & \frac{2}{\sqrt{\pi}} V_K(r; \varphi(t_1), t_1) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t_1), t_1}(t_1)) \right| \\ & \geq \frac{2}{\sqrt{\pi}} \int_c^k e^{-\alpha^2} d\alpha + \frac{2}{\sqrt{\pi}} \int_0^c e^{-\alpha^2} d\alpha \\ & = \frac{2}{\sqrt{\pi}} \int_0^k e^{-\alpha^2} d\alpha > \lambda, \end{aligned}$$

but this contradicts (9). □

Now let us evaluate the values of the term

$$\lim_{r \rightarrow 0^+} \sup_{t \in (a,b)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t(t)}) \right| \right]$$

in the following three simple particular cases.

Example 1. Let $\langle a, b \rangle = \langle 0, 1 \rangle$ and

$$\varphi(t) = k\sqrt{t}$$

on $\langle a, b \rangle$, where $k > 0$.




Figure 1

Denote

$$K = \{[\xi, \tau] \in \mathbb{R}^2; \tau \in \langle 0, 1 \rangle, \xi = k\sqrt{\tau}\}$$

and consider a point $[k\sqrt{\beta}, \beta] \in K$, where $\beta \in \langle 0, 1 \rangle$. For $\alpha > 0$ denote further

$$p_\alpha = \left\{ [\xi, \tau] \in \mathbb{R}^2; \beta - \tau = \left(\frac{k\sqrt{\beta} - \xi}{2\alpha} \right)^2, \tau < \beta \right\}.$$

It is easy to see that $K \cap p_\alpha \neq \emptyset$ if and only if $\alpha \leq \frac{k}{2}$. The dashed parabola in Figure 1 corresponds to the case $\alpha = \frac{k}{2}$ and the solid parabola p_α to the case $\alpha < \frac{k}{2}$. For $\alpha \leq \frac{k}{2}$ the intersection $K \cap p_\alpha$ contains just one point. Now we see that if $r > 0$, $\beta < r$, then


$$n_{k\sqrt{\beta}, \beta}(r, \alpha) = \begin{cases} 1 & \text{for } \alpha \in \langle 0, \frac{k}{2} \rangle, \\ 0 & \text{for } \alpha > \frac{k}{2}, \end{cases}$$

and thus for such $\epsilon, \delta \in \mathbb{R}^+$ always hold,

... i k.i

$$\frac{\epsilon}{\delta} > 0$$

...2. vs fi -V ...; ; * >n. i ... sly iffcti i>; <(\cdot) < \cdot C>



it is (Visy to see hát for pach *

• 5

... 10 for a^1.

ik ii'."

I_h(r; 0, 0) = 0.

264

"M

...-%:lmfm

ifKsr hancí,

*s i € (- i;

$$P_{\alpha} = \{K/i\}^G K^2; \rightarrow \alpha \cdot \alpha$$

A ; . < - - -

Altogether,

$$\lim_{r \rightarrow 0^+} \sup_{t \in (-1, 0)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] = \frac{2}{\sqrt{\pi}} \int_0^{\frac{k}{2}} e^{-\alpha^2} d\alpha < 1.$$

Example 3. Consider the case $(a, b) = (-1, 1)$,

$$\varphi(\tau) = k\sqrt{|\tau|},$$

where $k > 0$. By example 2 we have

$$\lim_{r \rightarrow 0^+} \sup_{t \in (-1, 0)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] = \frac{2}{\sqrt{\pi}} \int_0^{\frac{k}{2}} e^{-\alpha^2} d\alpha.$$

It follows from Proposition 1 that for any $\varepsilon > 0$, $\varepsilon < 1$, we have

$$\lim_{r \rightarrow 0^+} \sup_{t \in (\varepsilon, 1)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] = 0.$$

Thus it suffices to examine points $[k\sqrt{\beta}, \beta]$ for $\beta > 0$ near zero.




Figure 3

Let $r \in (0, 1)$, $\beta > 0$, $\beta < \frac{r}{2}$. Denote

$$\begin{aligned} p_\alpha &= \left\{ [\xi, \tau] \in \mathbb{R}^2; \beta - \tau = \left(\frac{\xi - k\sqrt{\beta}}{2\alpha} \right)^2, \tau < \beta \right\}, \\ p_\alpha^+ &= \{ [\xi, \tau] \in p_\alpha; \xi > k\sqrt{\beta} \}, \\ p_\alpha^- &= \{ [\xi, \tau] \in p_\alpha; \xi < k\sqrt{\beta} \}, \\ K_\infty &= K \cup \{ [\xi, \tau] \in \mathbb{R}^2; \tau \in (-\infty, 0), \xi = k\sqrt{-\tau} \}. \end{aligned}$$

It is easy to see that $K \cap p_\alpha^- = \emptyset$ for any $\alpha > \frac{k}{2}$, for $\alpha = \frac{k}{2}$ the intersection $K \cap p_\alpha^-$ is a singleton, and $K \cap p_\alpha^-$ consists of two points whenever $\alpha \in (0, \frac{k}{2})$.

Now consider $K_\infty \cap p_\alpha^+$. Elementary calculation yields that $K_\infty \cap p_\alpha^+ \neq \emptyset$ if and only if $\alpha \in (0, \frac{k}{2})$, and for $\alpha \in (0, \frac{k}{2})$ the intersection $K_\infty \cap p_\alpha^+$ is a singleton. If $\alpha \in (0, \frac{k}{2})$, $\{[\xi, \tau]\} = K_\infty \cap p_\alpha^+$ then

$$\xi = k\sqrt{\beta} \frac{k^2 + 2\sqrt{2}\alpha\sqrt{k^2 - 2\alpha^2}}{k^2 - 4\alpha^2}.$$

Thus we see that $n_{k\sqrt{\beta}, \beta}(r, \alpha) = 0$ for $\alpha > \frac{k}{2}$, $n_{k\sqrt{\beta}, \beta}(r, \frac{k}{2}) = 1$ and

$$2 \leq n_{k\sqrt{\beta}, \beta}(r, \alpha) \leq 3$$

for $\alpha \in (0, \frac{k}{2})$. Here $n_{k\sqrt{\beta}, \beta}(r, \alpha) = 3$ if and only if

$$(*) \quad \sqrt{\beta} \frac{k^2 + 2\sqrt{2}\alpha\sqrt{k^2 - 2\alpha^2}}{k^2 - 4\alpha^2} < \sqrt{r - \beta}.$$

Given $\alpha_0 \in (0, \frac{k}{2})$ then there is a $\beta > 0$, $\beta < r/2$ such that for each $\alpha \in (0, \alpha_0)$ the condition $(*)$ is fulfilled. Now one can see that for $\varepsilon \in (0, 1)$

$$\sup_{\beta \in (0, \varepsilon)} V_K(r; k\sqrt{\beta}, \beta) = 3 \int_0^{\frac{k}{2}} e^{-\alpha^2} d\alpha.$$

Since $\alpha_{\varphi(t), t}(t) = 0$ for $t \in (0, 1)$ we obtain

$$\lim_{r \rightarrow 0^+} \sup_{t \in (-1, 1)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] = 3 \frac{2}{\sqrt{\pi}} \int_0^{\frac{k}{2}} e^{-\alpha^2} d\alpha.$$

As

$$\frac{2}{\sqrt{\pi}} \int_0^c e^{-\alpha^2} d\alpha = \frac{1}{3}$$

for

$$c \approx 0.304570194 \dots$$

the condition

$$\lim_{r \rightarrow 0^+} \sup_{t \in (-1, 1)} \left[\frac{2}{\sqrt{\pi}} V_K(r; \varphi(t), t) + \left| 1 - \frac{2}{\sqrt{\pi}} G(\alpha_{\varphi(t), t}(t)) \right| \right] < 1$$

is not fulfilled for all $k > 0$ but only if

$$k < 0.609140388 \dots$$

References

- [1] *M. Dont*: Fourier problem with bounded Baire data. *Math. Bohem.* 22 (1997), 405–441.
- [2] *M. Dont*: On a heat potential. *Czechoslovak Math. J.* 25 (1975), 84–109.
- [3] *M. Dont*: A note on a heat potential and the parabolic variation. *Čas. Pěst. Mat.* 101 (1976), 28–44.
- [4] *J. Král*: *Teorie potenciálu I*. SPN, Praha, 1965.

Author's address: *Miroslav Dont*, Department of Mathematics, Faculty of Electrical Engineering, Czech Technical University, Technická 2, 166 27 Prague, Czech Republic, e-mail: dont@math.feld.cvut.cz.