

Ján Jakubík

Graph automorphisms of a finite modular lattice

Czechoslovak Mathematical Journal, Vol. 49 (1999), No. 2, 443–447

Persistent URL: <http://dml.cz/dmlcz/127500>

Terms of use:

© Institute of Mathematics AS CR, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

GRAPH AUTOMORPHISMS OF A FINITE MODULAR LATTICE

JÁN JAKUBÍK, Košice

(Received June 25, 1997)

G. Birkhoff ([2], Problem 6) proposed the following problem:

To find all finite lattices L such that each automorphism of the unoriented graph corresponding to L turns out to be a lattice automorphism.

Let us denote by \mathcal{C} the class of all lattices which satisfy the condition mentioned.

In the present note we give a partial solution to this problem concerning modular lattices. By applying the methods and the results of [3] and [4] we prove

(*) Let L be a finite modular lattice. Then the following conditions are equivalent:

- (i) L belongs to \mathcal{C} .
- (ii) No direct factor of L having more than one element is self-dual.

Let us remark that the related Problem 5 in [2] (proposed already in [1] as Problem 8 and dealing with unoriented graphs of finite lattices) was solved in [3] for the particular case of modular lattices and remains unsolved for the general case.

1. PRELIMINARIES

In the whole paper L denotes a finite lattice. For $a, b \in L$ we put $a \prec b$ or $b \succ a$ if $a < b$ and the interval $[a, b]$ of L is a two-element set.

Let $G(L)$ be the unoriented graph such that

- (i) L is the set of all vertices of $G(L)$;
- (ii) a pair $(x, y) \in L \times L$ is an edge in $G(L)$ if and only if either $x \prec y$ or $x \succ y$.

For each lattice A we denote by A^\sim the lattice which is dual to A . If there exists an isomorphism of A onto A^\sim , then A is called self-dual.

Let us have a direct product $A \times B$ of finite lattices A and B . Then for $(a_1, b_1), (a_2, b_2) \in A \times B$ the relation

$$(a_1, b_1) \prec (a_2, b_2)$$

Supported by Grant GA SAV 1230/97.

is valid if and only if either $a_1 \prec a_2$ and $b_1 = b_2$, or $a_1 = a_2$ and $b_1 \prec b_2$. From this we conclude

1.1. Lemma. *Let ψ be an isomorphism of L onto the direct product $A \times B$. Further suppose that χ is an isomorphism of B onto B^\sim . For each $x \in L$ we put $\varphi(x) = y$, where*

$$\psi(x) = (a, b), \quad y = \psi^{-1}((a, \chi(b))).$$

Then φ is an automorphism of the graph $G(L)$.

1.2. Lemma. *Let the assumptions of 1.1 be satisfied. Further suppose that B has more than one element. Then φ fails to be a lattice automorphism on L .*

P r o o f. Choose $a \in A$. There exist $b_1, b_2 \in B$ with $b_1 \prec b_2$. Put

$$x = \psi^{-1}((a, b_1)), \quad y = \psi^{-1}((a, b_2)).$$

Then $x \prec y$. We have

$$\varphi(x) = \psi^{-1}((a, \chi(b_1))), \quad \varphi(y) = \psi^{-1}((a, \chi(b_2)))$$

and $\chi(b_1) \succ \chi(b_2)$. Therefore $\varphi(x) \succ \varphi(y)$. □

1.3. Corollary. *If L belongs to \mathcal{C} , then no direct factor of L having more than one element is self-dual.*

2. INTERNAL DIRECT PRODUCT DECOMPOSITIONS

Let A, B be lattices and let

$$\psi: L \rightarrow A \times B$$

be an isomorphism of L onto the direct product $A \times B$. For $x \in L$ with $\psi(x) = (a, b)$ we put $a = x_A, b = x_B$.

Let x^0 be a fixed element of L . We denote

$$A_0 = \{x \in L: x_B = x_B^0\}, \quad B_0 = \{x \in L: x_A = x_A^0\}.$$

Then A_0 and B_0 are convex sublattices of L with $A_0 \cap B_0 = \{x^0\}$. Moreover, A_0 is isomorphic to A and B_0 is isomorphic to B .

Consider the mapping

$$(1) \quad \psi_0: L \rightarrow A_0 \times B_0$$

defined by

$$\psi(x) = (x(A_0), x(B_0)),$$

where $x(A_0)$ is an element of A_0 such that

$$(x(A_0))_A = x_A;$$

similarly, $x(B_0)$ is an element of B_0 such that

$$(x(B_0))_B = x_B.$$

Then the mapping ψ_0 is an isomorphism of L onto the lattice $A_0 \times B_0$. We say that ψ_0 is an internal direct product decomposition of L with the central element x^0 . The lattices A_0 and B_0 are called internal direct factors of L . (Cf. [4].)

2.1. Lemma. (Cf. [4], Lemma 2.4.) *Suppose that (1) is an internal direct product decomposition of L with the central element x^0 and that, moreover,*

$$\psi_1: L \rightarrow A_0 \times C_0$$

is also an internal direct product decomposition of L with the central element x^0 . Then $B_0 = C_0$.

Now suppose that L_1 and L_2 are finite modular lattices and that φ is an isomorphism of $G(L_1)$ onto $G(L_2)$. Such situation was investigated in [3].

We denote by \mathcal{A}_1 the set of all intervals $[x, y]$ of L_1 such that

$$x \prec y \quad \text{and} \quad \varphi(x) \prec \varphi(y).$$

Further let \mathcal{B}_1 be the set of all intervals $[u, v]$ of L_1 such that

$$u \prec v \quad \text{and} \quad \varphi(u) \succ \varphi(v).$$

Analogously we define the sets \mathcal{A}_2 and \mathcal{B}_2 of intervals of L_2 (with φ^{-1} instead of φ).

Let x_1^0 be a fixed element of L_1 . We denote by A_1^0 the set of all elements $x \in L_1$ such that either $x = x_1^0$, or there exist $y_1, y_2, \dots, y_n \in L_1$ which satisfy the following conditions:

- (i) $y_1 = x_1^0, y_n = x$,
- (ii) if $i \in \{1, 2, \dots, n-1\}$, then the elements y_i, y_{i+1} are comparable and the corresponding interval of L_1 belongs to \mathcal{A}_1 .

Similarly we define the set $B_1^0 \subseteq L_1$ (taking \mathcal{B}_1 instead of \mathcal{A}_1).

Further let x_2^0 be an arbitrary element of L_2 . In an analogous way we define the subsets A_2^0 and B_2^0 of L_2 (taking φ^{-1} instead of φ).

Looking at the construction performed in [3] (cf. the lemmas used for proving Theorem 1 in [3]) and applying the notion of the internal direct product decomposition we arrive at the following lemma:

2.2. Lemma. *Under the assumptions as above, there exist internal direct product decompositions*

$$\begin{aligned} \psi_1: L_1 &\rightarrow A_1^0 \times B_1^0 \quad (\text{with the central element } x_1^0), \\ \psi_2: L_2 &\rightarrow A_2^0 \times B_2^0 \quad (\text{with the central element } x_2^0) \end{aligned}$$

such that

- (i) the lattices A_1^0 and A_2^0 are isomorphic,
- (ii) the lattice B_1^0 is isomorphic to $(B_2^0)^\sim$.

3. PROOF OF (*)

Suppose that no direct factor of L having more than one element is self-dual.

Let φ be an automorphism of the graph $G(L)$. We put $L = L_1 = L_2$ and apply Lemma 2.2 above. Choose x^0 in L and put $x^0 = x_1^0 = x_2^0$. Then under the notation as in Section 2 we have

$$\mathcal{A}_1 = \mathcal{A}_2, \quad \mathcal{B}_1 = \mathcal{B}_2.$$

Thus, in the set-theoretical sense, we get $A_1^0 = A_2^0$. Further, since A_1^0 and A_2^0 are sublattices of L , we obtain that A_1^0 and A_2^0 are equal as lattices. Put $A_1^0 = A = A_2^0$. Then in view of 2.2 we obtain internal direct product decompositions

$$\begin{aligned} \psi_1: L &\rightarrow A \times B_1^0, \\ \psi_2: L &\rightarrow A \times B_2^0 \end{aligned}$$

with the same central element x^0 . Thus according to 2.1,

$$B_1^0 = B_2^0.$$

Moreover, in view of 2.2 (ii), B_1^0 is dually isomorphic to B_2^0 , hence B_1^0 is self-dual. Then the assumption yields that B_1^0 is a one element set.

Since the element x^0 of L was arbitrarily chosen, we conclude that the set \mathcal{B}_1 must be empty and thus all prime intervals of L belong to \mathcal{A}_1 .

Let $x, y \in L$. If $x < y$, then there are y_1, y_2, \dots, y_n in L such that $x = y_1 \prec y_2 \prec \dots \prec y_n = y$, whence $\varphi(x) = \varphi(y_1) \prec \varphi(y_2) \prec \dots \prec \varphi(y_n) = \varphi(y)$, thus $\varphi(x) < \varphi(y)$. Conversely, by applying φ^{-1} instead of φ we get that $\varphi(x) < \varphi(y)$ implies $x < y$. Hence φ is a lattice isomorphism.

Therefore we have

3.1. Lemma. *Suppose that L is a modular lattice such that none of its direct factors having more than one element is self-dual. Then each automorphism of $G(L)$ is an automorphism of the lattice L .*

Now, (*) is a consequence of 1.3 and 3.1.

We conclude by remarking that all the above considerations remain valid if the assumption that L is a finite modular lattice is replaced by the assumption that L is a modular lattice such that each bounded chain in L is finite.

References

- [1] *G. Birkhoff*: Lattice Theory. Second Edition, Providence, 1948.
- [2] *G. Birkhoff*: Lattice Theory. Third Edition, Providence, 1967.
- [3] *J. Jakubík*: On graph isomorphism of modular lattices. Czechoslovak Math. J. 4 (1954), 131–141. (In Russian.)
- [4] *J. Jakubík, M. Csontóová*: Convex isomorphisms of directed multilattices. Math. Bohemica 118 (1993), 359–379.

Author's address: Matematický ústav SAV, Grešáková 6, 040 01 Košice, Slovakia, email: musavke@mail.saske.sk.