

Thiruvaiyaru V. Panchapagesan

A Borel extension approach to weakly compact operators on  $C_0(T)$

*Czechoslovak Mathematical Journal*, Vol. 52 (2002), No. 1, 97–115

Persistent URL: <http://dml.cz/dmlcz/127705>

## Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A BOREL EXTENSION APPROACH TO WEAKLY COMPACT  
OPERATORS ON  $C_0(T)$

T. V. PANCHAPAGESAN, Mérida

(Received January 29, 1999)

*Dedicated to Professor K. S. Padmanabhan on the occasion  
of his seventieth birthday.*

*Abstract.* Let  $X$  be a quasicomplete locally convex Hausdorff space. Let  $T$  be a locally compact Hausdorff space and let  $C_0(T) = \{f: T \rightarrow \mathbb{C}, f \text{ is continuous and vanishes at infinity}\}$  be endowed with the supremum norm. Starting with the Borel extension theorem for  $X$ -valued  $\sigma$ -additive Baire measures on  $T$ , an alternative proof is given to obtain all the characterizations given in [13] for a continuous linear map  $u: C_0(T) \rightarrow X$  to be weakly compact.

*MSC 2000:* 47B38, 46G10, 28B05

1. INTRODUCTION

Let  $T$  be a locally compact Hausdorff space and let  $C_0(T)$  be the Banach space of all complex valued continuous functions vanishing at infinity in  $T$ , endowed with the supremum norm. Then its dual  $M(T)$  is the Banach space of all bounded complex Radon measures  $\mu$  on  $T$  with the norm given by  $\|\mu\| = \text{var}(\mu, \mathcal{B}(T))(T)$ . Let  $X$  be a locally convex Hausdorff space (briefly, an lcHs) which is quasicomplete and let  $u: C_0(T) \rightarrow X$  be a continuous linear map. When  $X$  is complete and  $T$  is compact, Grothendieck gave in Theorem 6 of [6] some necessary and sufficient conditions for  $u$  to be weakly compact. As observed in [14], Grothendieck's techniques, contrary to

---

Supported by the project C-845-97-05-B of the C.D.C.H.T. of the Universidad de los Andes, Mérida, Venezuela.

Remark 2 on p. 161 of [6], are not powerful enough to extend his characterizations when  $T$  is a non  $\sigma$ -compact locally compact Hausdorff space.

In [13], using the Baire and  $\sigma$ -Borel characterizations of weakly compact subsets of  $M(T)$  as given in [12], we obtained 35 characterizations for the continuous linear map  $u: C_0(T) \rightarrow X$  to be weakly compact, where  $X$  is a quasicomplete lchS. These include the characterizations mentioned in Remark 2 on p. 161 of Grothendieck [6] and in Theorem 9.4.10 of [5], whose proof as given in [5] is incorrect without the hypothesis of  $\sigma$ -compactness of  $T$  (see [14]). In [13] we also obtained a theorem on regular Borel and  $\sigma$ -Borel extensions of  $X$ -valued  $\sigma$ -additive Baire measures on  $T$  (briefly, the Borel extension theorem) and Theorem 5.3 of Thomas [16] (dispensing with the technique of reduction to the metrizable compact case) as a consequence of these characterizations.

The Riesz representation theorem was used in [9], [10] to obtain the regular Borel and  $\sigma$ -Borel extensions of a complex Baire measure on  $T$ . The paper [13] can be considered to be its analogue for  $X$ -valued Baire measures on  $T$  with the Riesz representation theorem being replaced by the Bartle-Dunford-Schwartz representation of weakly compact operators, since the Borel extension theorem for such Baire measures was deduced there from the characterizations of weakly compact operators on  $C_0(T)$ .

On the other hand, the regular  $\sigma$ -Borel extension of positive Baire measures on  $T$  was used in Halmos [7] to derive the Riesz representation theorem for positive linear forms on  $C_0(T)$ . In this context the following question arises: Is it possible to obtain all the characterizations given in [13] for a continuous linear map  $u: C_0(T) \rightarrow X$  to be weakly compact, starting with the Borel extension theorem for  $X$ -valued Baire measures on  $T$ ? Recently, in our joint work with Dobrakov ([4]), combining the Borel extension theorem with the first part of Theorem 1 of [13] and Lemma 1 and Theorem 2 of [6], we answered the question in the affirmative when  $c_0 \not\subset X$  and  $X$  is a quasicomplete lchS (namely, Theorem 5.3 of [16]). In the present paper, we also answer the question in the affirmative for arbitrary quasicomplete lchS  $X$  and for this, along with the Borel extension theorem, we use the quoted results of [13] and [6], Lemmas 1–7 of Section 2 below and Theorem 1 of [11]. Thus the present paper can be considered to be the vector analogue of the treatment of Halmos [7].

## 2. PRELIMINARIES

In this section we fix the notation and terminology. For the convenience of the reader we also give some definitions and results from literature.

In the sequel  $T$  will denote a locally compact Hausdorff space and  $C_0(T)$  the Banach space of all complex valued continuous functions vanishing at infinity in  $T$ , endowed with the supremum norm  $\|f\|_T = \sup_{t \in T} |f(t)|$ .

Let  $\mathcal{K}$  ( $\mathcal{K}_0$ ) be the family of all compacts (compact  $G_\delta$ s) in  $T$ .  $\mathcal{B}_0(T)$ ,  $\mathcal{B}_c(T)$  and  $\mathcal{B}(T)$  are the  $\sigma$ -rings generated by  $\mathcal{K}_0$ ,  $\mathcal{K}$  and the class of all open sets in  $T$ , respectively. The members of  $\mathcal{B}_0(T)$  ( $\mathcal{B}_c(T)$ ,  $\mathcal{B}(T)$ ) are called *Baire sets* ( $\sigma$ -*Borel sets*, *Borel sets*, respectively) of  $T$ . Since a subset  $E$  of  $T$  belongs to  $\mathcal{B}_c(T)$  if and only if  $E$  is a  $\sigma$ -bounded Borel set, the members of  $\mathcal{B}_c(T)$  are called  $\sigma$ -Borel sets.

$M(T)$  is the Banach space of all bounded complex Radon measures on  $T$  with their domain restricted to  $\mathcal{B}(T)$ . Thus each  $\mu \in M(T)$  is a Borel regular (bounded) complex measure on  $\mathcal{B}(T)$  and has the norm given by  $\|\mu\| = \text{var}(\mu, \mathcal{B}(T))(T)$ . For  $\mu \in M(T)$ ,  $|\mu|(E) = \text{var}(\mu, \mathcal{B}(T))(E)$ ,  $E \in \mathcal{B}(T)$ .

We recall the following result from [12, Lemma 1].

**Proposition 1.** For  $\mu \in M(T)$ ,

$$|\mu|_{\mathcal{B}_0(T)}(\cdot) = \text{var}(\mu|_{\mathcal{B}_0(T)}, \mathcal{B}_0(T))(\cdot)$$

and

$$|\mu|_{\mathcal{B}_c(T)}(\cdot) = \text{var}(\mu|_{\mathcal{B}_c(T)}, \mathcal{B}_c(T))(\cdot).$$

A vector measure is an additive set function defined on a ring of sets with values in an lchS. In the sequel  $X$  will denote an lchS with a topology  $\tau$ . Let  $\Gamma$  be the set of all  $\tau$ -continuous seminorms on  $X$ . The dual of  $X$  is denoted by  $X^*$ .

The strong topology  $\beta(X^*, X)$  of  $X^*$  is the locally convex topology induced by the seminorms  $\{p_B: B \text{ bounded in } X\}$ , where  $p_B(x^*) = \sup_{x \in B} |x^*(x)|$ .  $X^{**}$  denotes the dual of  $(X^*, \beta(X^*, X))$  and is endowed with the locally convex topology  $\tau_e$  of uniform convergence in equicontinuous subsets of  $X^*$ . Note that  $(X^*, \beta(X^*, X))$  and  $(X^{**}, \tau_e)$  are lchS.

It is well known that the canonical injection  $J: X \rightarrow X^{**}$  given by  $\langle Jx, x^* \rangle = \langle x, x^* \rangle$  for all  $x \in X$  and  $x^* \in X^*$ , is linear. Identifying  $X$  with  $JX \subset X^{**}$  one has  $\tau_e|_{JX} = \tau_e|_X = \tau$ .

**Definition 1.** A linear map  $u: C_0(T) \rightarrow X$  is called a *weakly compact operator* on  $C_0(T)$  if  $\{uf: \|f\|_T \leq 1\}$  is relatively weakly compact in  $X$ .

The following result (Corollary 9.3.2 of [5], which is essentially a consequence of Lemma 1 of [6]) plays a key role in Section 4.

**Proposition 2.** *Let  $E$  and  $F$  be lchS with  $F$  quasicomplete. If  $u: E \rightarrow F$  is linear and continuous, then the following conditions are equivalent.*

- (i)  $u$  maps bounded subsets of  $E$  into relatively weakly compact subsets of  $F$ .
- (ii)  $u^*(A)$  is relatively  $\sigma(E^*, E^{**})$ -compact for each equicontinuous subset  $A$  of  $F^*$ .
- (iii)  $u^{**}(E^{**}) \subset F$ .

The following result is due to Theorem 2 of [6], which is the same as Theorem 4.22.1 of [5].

**Proposition 3.** *Let  $A$  be a bounded set in  $M(T)$ . Then the following assertions are equivalent.*

- (i)  $A$  is relatively weakly compact.
- (ii) For each disjoint sequence  $(U_n)_1^\infty$  of open sets in  $T$ ,

$$\lim_n \sup_{\mu \in A} |\mu(U_n)| = 0.$$

- (iii) For  $(U_n)$  as in (ii),  $\lim_n \sup_{\mu \in A} |\mu|(U_n) = 0$ .
- (iv) Let  $\varepsilon > 0$ .

- (a) For each compact  $K$  in  $T$ , there exists an open set  $U$  in  $T$  such that  $K \subset U$  and  $\sup_{\mu \in A} |\mu|(U \setminus K) < \varepsilon$ ; and
- (b) there exists a compact  $C$  such that  $\sup_{\mu \in A} |\mu|(T \setminus C) < \varepsilon$ .

For each  $\tau$ -continuous seminorm  $p$  on  $X$ , let  $p(x) = \|x\|_p$ ,  $x \in X$ , and let  $X_p = (X, \|\cdot\|_p)$  be the associated seminormed space. The completion of the quotient normed space  $X_p/p^{-1}(0)$  is denoted by  $\tilde{X}_p$ . Let  $\Pi_p: X_p \rightarrow X_p/p^{-1}(0) \subset \tilde{X}_p$  be the canonical quotient map.

Let  $\mathcal{S}$  be a  $\sigma$ -ring of subsets of a non empty set  $\Omega$ . Given a vector measure  $m: \mathcal{S} \rightarrow X$ , for each  $\tau$ -continuous seminorm  $p$  on  $X$ , let  $m_p: \mathcal{S} \rightarrow \tilde{X}_p$  be given by  $m_p(E) = (\Pi_p \circ m)(E)$  for  $E \in \mathcal{S}$ . Then  $m_p$  is a Banach space valued vector measure on  $\mathcal{S}$ . We define the  $p$ -semivariation  $\|m\|_p$  of  $m$  by

$$\|m\|_p(E) = \|m_p\|(E) \text{ for } E \in \mathcal{S}$$

and

$$\|m\|_p(\Omega) = \|m_p\|(\Omega) = \sup_{E \in \mathcal{S}} \|m_p\|(E)$$

where  $\|m_p\|$  is the semivariation of the vector measure  $m_p$  and is given by  $\|m_p\|(E) = \sup\{\|x^* \circ m\|(E) : x^* \in \tilde{X}_p^*, \|x^*\| \leq 1\}$  (see p. 2 of [1]).

An  $X$ -valued vector measure  $m$  on a  $\sigma$ -ring  $\mathcal{S}$  of subsets of  $\Omega$  is said to be *bounded* if  $\{m(E) : E \in \mathcal{S}\}$  is bounded in  $X$  and equivalently, if  $\|m\|_p(\Omega) < \infty$  for each  $\tau$ -continuous seminorm  $p$  on  $X$ . When  $m$  is  $\sigma$ -additive, then  $m_p$  is a Banach space valued  $\sigma$ -additive vector measure on the  $\sigma$ -ring  $\mathcal{S}$  and hence by Corollary I.1.19 of [1],  $\|m\|_p(\Omega) = \|m_p\|(\Omega) \leq 4 \sup_{E \in \mathcal{S}} \|m(E)\|_p < \infty$ .

For the theory of integration of bounded  $\mathcal{S}$ -measurable scalar functions with respect to a bounded quasicomplete lchS-valued vector measure on the  $\sigma$ -ring  $\mathcal{S}$ , the reader is referred to [11] or [13]. We need the following results from Lemma 6 of [11] and Proposition 7 of [13].

**Proposition 4.** *Let  $X$  be a quasicomplete lchS and let  $\mathcal{S}$  be a  $\sigma$ -ring of subsets of  $\Omega$ . Then:*

- (i) *If  $f$  is a bounded  $\mathcal{S}$ -measurable scalar function and  $m$  is an  $X$ -valued bounded vector measure on  $\mathcal{S}$ , then  $f$  is  $m$ -integrable and*

$$x^* \left( \int_{\Omega} f \, dm \right) = \int_{\Omega} f d(x^* \circ m)$$

*for each  $x^* \in X^*$ .*

- (ii) *(Lebesgue bounded convergence theorem) If  $m$  is an  $X$ -valued  $\sigma$ -additive vector measure on  $\mathcal{S}$  and  $(f_n)$  is a bounded sequence of  $\mathcal{S}$ -measurable scalar functions with  $\lim_n f_n(w) = f(w)$  for each  $w \in \Omega$ , then  $f$  is  $m$ -integrable and*

$$\int_E f \, dm = \lim_n \int_E f_n \, dm$$

*for each  $E \in \mathcal{S}$ .*

The following result follows from the first part of Theorem 1 of [13], and is analogous to Theorem VI.2.1 of [1] for lchS-valued continuous linear maps on  $C_0(T)$ . It plays a key role in Sections 3 and 4.

**Proposition 5.** *Let  $X$  be an lchS. Let  $u : C_0(T) \rightarrow X$  be a continuous linear map. Then there exists a unique  $X^{**}$ -valued vector measure  $m$  on  $\mathcal{B}(T)$  possessing the following properties:*

- (i)  *$x^* \circ m \in M(T)$  for each  $x^* \in X^*$  and consequently,  $m : \mathcal{B}(T) \rightarrow X^{**}$  is  $\sigma$ -additive in the  $\sigma(X^{**}, X^*)$ -topology.*
- (ii) *The mapping  $x^* \rightarrow x^* \circ m$  of  $X^*$  into  $M(T)$  is weak\*-weak\* continuous. Moreover,  $u^* x^* = x^* \circ m$ ,  $x^* \in X^*$ .*

- (iii)  $x^*uf = \int_T f d(x^* \circ m)$  for each  $f \in C_0(T)$  and  $x^* \in X^*$ .
- (iv)  $\{m(E) : E \in \mathcal{B}(T)\}$  is  $\tau_e$ -bounded in  $X^{**}$ .
- (v)  $m(E) = u^{**}(\chi_E)$  for  $E \in \mathcal{B}(T)$ .

**Definition 2.** Let  $u : C_0(T) \rightarrow X$  be a continuous linear map. Then the vector measure  $m$  as given in Proposition 5 is called the *representing measure* of  $u$ .

**Definition 3.** A  $\sigma$ -additive vector measure  $m : \mathcal{B}_0(T) \rightarrow X$  ( $\mathcal{B}(T) \rightarrow X$ ,  $\mathcal{B}_c(T) \rightarrow X$ ) is called an  $X$ -valued *Baire (Borel,  $\sigma$ -Borel) measure* on  $T$ .

**Definition 4.** Let  $\mathcal{S}$  be a  $\sigma$ -ring of sets in  $T$  with  $\mathcal{S} \supset \mathcal{K}$  or  $\mathcal{K}_0$ . Let  $m : \mathcal{S} \rightarrow X$  be a vector measure. Then  $m$  is said to be  $\mathcal{S}$ -*regular* ( $\mathcal{S}$ -*outer regular*,  $\mathcal{S}$ -*inner regular*) in  $E \in \mathcal{S}$  if, given a  $p$  in  $\Gamma$  and  $\varepsilon > 0$ , there exist a compact  $K \in \mathcal{S}$  and an open set  $U \in \mathcal{S}$  with  $K \subset E \subset U$  (an open set  $U \in \mathcal{S}$  with  $E \subset U$ , a compact  $K \in \mathcal{S}$  with  $K \subset E$ ) such that  $\|m\|_p(U \setminus K) < \varepsilon$  ( $\|m\|_p(U \setminus E) < \varepsilon$ ,  $\|m\|_p(E \setminus K) < \varepsilon$ , respectively). Even though  $T$  does not belong to  $\mathcal{S}$  one can define  $\mathcal{S}$ -*inner regularity* of  $m$  in  $T$  as follows. Given  $p \in \Gamma$  and  $\varepsilon > 0$ , there exists a compact  $K \in \mathcal{S}$  such that  $\|m\|_p(B) < \varepsilon$  for all  $B \in \mathcal{S}$  with  $B \subset T \setminus K$ . The vector measure  $m$  is said to be  $\mathcal{S}$ -*regular* ( $\mathcal{S}$ -*outer regular*,  $\mathcal{S}$ -*inner regular*) if it is so in each  $E \in \mathcal{S}$ . When  $\mathcal{S} = \mathcal{B}(T)$  ( $\mathcal{B}_0(T)$ ,  $\mathcal{B}_c(T)$ ), we use the term *Borel (Baire,  $\sigma$ -Borel) regularity* or *outer regularity* or *inner regularity*.

**Remark 1.** In the above definition one can replace  $\Gamma$  by any other family of continuous seminorms on  $X$  which induces the topology  $\tau$ .

The following proposition on regular Borel and  $\sigma$ -Borel extensions of an  $X$ -valued Baire measure on  $T$  is well known and plays a key role in Section 4. It was first proved in [3], [8] for Banach space valued Baire measures on  $T$  and extended to group valued measures in [15]. For a simple and direct proof of the proposition see [4]. Note that a highly technical operator theoretic proof is given in [13] as mentioned in the introduction.

**Proposition 6.** Let  $m$  be an  $X$ -valued Baire measure on  $T$  and let  $X$  be a quasicomplete *lchS*. Then  $m$  is Baire regular in  $T$ . Moreover, there exists a unique  $X$ -valued Borel ( $\sigma$ -Borel) regular  $\sigma$ -additive extension  $\hat{m}$  ( $\hat{m}_c$ ) of  $m$  on  $\mathcal{B}(T)$  ( $\mathcal{B}_c(T)$ , respectively). Moreover,  $\hat{m}|_{\mathcal{B}_c(T)} = \hat{m}_c$ .

### 3. SOME LEMMAS

Throughout this section  $X$  denotes a quasicomplete lcHs with the topology  $\tau$ . Let  $u: C_0(T) \rightarrow X$  be a continuous linear map with the representing measure  $m$ . Let  $m_0 = m|_{\mathcal{B}_0(T)}$  and  $m_c = m|_{\mathcal{B}_c(T)}$ .

Let  $\mathcal{E} = \{A \subset X^*: A \text{ equicontinuous}\}$ , and let  $p_A(x) = \sup_{x^* \in A} |x^*(x)|$  and  $p_A(x^{**}) = \sup_{x^* \in A} |x^{**}(x^*)|$  for  $A \in \mathcal{E}$ ,  $x \in X$  and  $x^{**} \in X^{**}$ . Then the family of seminorms  $\Gamma_{\mathcal{E}} = \{p_A: A \in \mathcal{E}\}$  induces the topology  $\tau$  of  $X$  and  $\tau_e$  of  $X^{**}$ .

Let  $X_A = X_{p_A}/p_A^{-1}(0)$  and let  $Y_A = \widetilde{X_A}$ , the completion of the normed space  $X_A$ . For  $E \in \mathcal{B}(T)$ ,

$$\|m_{p_A}\|(E) = \sup\{|y^* \circ m|(E): y^* \in Y_A^*, \|y^*\| \leq 1\}.$$

**Lemma 1.** *Let  $A \in \mathcal{E}$ . Then:*

(i) *For  $E \in \mathcal{B}(T)$*

$$\|m_{p_A}\|(E) = \|m\|_{p_A}(E) = \sup\{|x^* \circ m|(E): x^* \in A\}.$$

(ii) *For  $E \in \mathcal{B}_c(T)$*

$$\begin{aligned} \|(m_c)_{p_A}\|(E) &= \|m_c\|_{p_A}(E) = \sup\{|x^* \circ m_c|(E): x^* \in A\} \\ &= \sup\{|x^* \circ m|(E): x^* \in A\} \end{aligned}$$

where  $|x^* \circ m_c|(E) = \text{var}(x^* \circ m_c, \mathcal{B}_c(T))(E)$ .

(iii) *For  $E \in \mathcal{B}_0(T)$*

$$\begin{aligned} \|(m_0)_{p_A}\|(E) &= \|m_0\|_{p_A}(E) = \sup\{|x^* \circ m_0|(E): x^* \in A\} \\ &= \sup\{|x^* \circ m|(E): x^* \in A\} \end{aligned}$$

where  $|x^* \circ m_0|(E) = \text{var}(x^* \circ m_0, \mathcal{B}_0(T))(E)$ .

**Proof.** Each element  $\tilde{x} \in X_A$  is of the form  $\tilde{x} = x + p_A^{-1}(0)$  for some  $x \in X$  and it is easy to show that the quotient norm  $\|\tilde{x}\|_{p_A} = p_A(x)$ . For  $x^* \in A$ , let  $\Psi_{x^*}(x + p_A^{-1}(0)) = x^*(x)$ . Then  $\Psi_{x^*}: X_A \rightarrow \mathbb{C}$  is well defined and linear. Moreover, for  $x^* \in A$ ,

$$|\Psi_{x^*}(x + p_A^{-1}(0))| = |x^*(x)| \leq p_A(x) = \|x + p_A^{-1}(0)\|_{p_A}$$

and hence  $\|\Psi_{x^*}\| \leq 1$ . Then by continuity  $\Psi_{x^*}$  has a unique continuous linear extension to the whole of  $Y_A$  with the norm less than or equal to one and we denote



this extension again by  $\Psi_{x^*}$ . Clearly, the mapping  $x^* \rightarrow \Psi_{x^*}$  of  $A$  into  $Y_A^*$  is injective. For  $\tilde{x} = x + p_A^{-1}(0) \in X_A$  with  $x \in X$  we have

$$\begin{aligned} \|\tilde{x}\|_{p_A} &= \|x + p_A^{-1}(0)\|_{p_A} = p_A(x) = \sup_{x^* \in A} |x^*(x)| \\ &= \sup_{x^* \in A} |\Psi_{x^*}(\tilde{x})| \leq \sup_{y^* \in Y_A^*, \|y^*\| \leq 1} |y^*(\tilde{x})| = \|\tilde{x}\|_{p_A} \end{aligned}$$

and hence

$$(1) \quad \|\tilde{x}\|_{p_A} = \sup_{x^* \in A} |\Psi_{x^*}(\tilde{x})|.$$

Let us write  $\Psi_{x^*}(y) = x^*(y)$  for  $x^* \in A$  and  $y \in Y_A$ . Let  $y \in Y_A$  and let  $\varepsilon > 0$ . Since  $X_A$  is dense in  $Y_A$ , there exists  $\tilde{x} \in X_A$  such that  $\|y - \tilde{x}\|_{p_A} < \varepsilon$ . Then by (1) we have

$$\begin{aligned} \|y\|_{p_A} &< \varepsilon + \|\tilde{x}\|_{p_A} = \varepsilon + \sup_{x^* \in A} |\Psi_{x^*}(\tilde{x})| \\ &\leq \varepsilon + \sup_{x^* \in A} |\Psi_{x^*}(\tilde{x} - y)| + \sup_{x^* \in A} |\Psi_{x^*}(y)| \\ &\leq \varepsilon + \|\tilde{x} - y\|_{p_A} + \sup_{x^* \in A} |x^*(y)| < 2\varepsilon + \sup_{x^* \in A} |x^*(y)| \end{aligned}$$

and hence

$$\|y\|_{p_A} = \sup_{x^* \in A} |\Psi_{x^*}(y)| = \sup_{x^* \in A} |x^*(y)|$$

for  $y \in Y_A$ . Thus  $\{\Psi_{x^*} : x^* \in A\}$  is a norm determining subset of  $\{y^* \in Y_A^* : \|y^*\| \leq 1\}$ . Using this result and writing  $\Psi_{x^*}(y) = x^*(y)$  for all  $x^* \in A$  and  $y \in Y_A$  in the proof of the first part of Proposition 11 of [1], one can show that

$$(2) \quad \begin{aligned} \|m\|_{p_A}(E) &= \|m_{p_A}\|(E) = \sup\{|\Psi_{x^*} \circ m|(E) : x^* \in A\} \\ &= \sup\{|x^* \circ m|(E) : x^* \in A\} \end{aligned}$$

for  $E \in \mathcal{B}(T)$ . Thus (i) holds.

Replacing  $m$  by  $m_c$  (by  $m_0$ ) and  $\mathcal{B}(T)$  by  $\mathcal{B}_c(T)$  (by  $\mathcal{B}_0(T)$ ) in the above argument, similarly we have

$$\|(m_c)_{p_A}\|(E) = \|m_c\|_{p_A}(E) = \sup\{|x^* \circ m_c|(E) : x^* \in A\}$$

for  $E \in \mathcal{B}_c(T)$  and

$$\|(m_0)_{p_A}\|(E) = \|m_0\|_{p_A}(E) = \sup\{|x^* \circ m_0|(E) : x^* \in A\}$$

for  $E \in \mathcal{B}_0(T)$ . Now a reference to Proposition 1 completes the proofs of (ii) and (iii) of the lemma.  $\square$

The following result is the same as Lemma 2 of [13].

**Lemma 2.**  $u^*A$  is bounded in  $M(T)$  for each  $A \in \mathcal{E}$ .

**Notation 1.**  $\mathcal{U}_0$  denotes the family of all open Baire sets in  $T$ .

**Lemma 3.** Suppose  $m_0(\mathcal{U}_0) \subset X$ . Then:

- (i)  $m_0$  is  $\sigma$ -additive in  $\mathcal{U}_0$  in  $\tau$ . That is, given a disjoint sequence  $(U_n)_1^\infty$  in  $\mathcal{U}_0$ , then  $m_0(\bigcup_1^\infty U_n) = \sum_1^\infty m_0(U_n)$  (in the topology  $\tau$ ).
- (ii) If  $(U_n)_1^\infty$  is a disjoint sequence in  $\mathcal{U}_0$ , then, for each  $A \in \mathcal{E}$ ,  $\lim_n \|m_0\|_{p_A}(U_n) = 0$ .

**P r o o f.** (i) By Proposition 5 (i),  $x^* \circ m \in M(T)$  for  $x^* \in X^*$  and hence

$$(x^* \circ m_0)\left(\bigcup_1^\infty U_n\right) = \sum_1^\infty (x^* \circ m_0)(U_n)$$

for each  $x^* \in X^*$ . By hypothesis,  $m_0(\mathcal{U}_0) \subset X$  and hence by the Orlicz-Pettis theorem we conclude that  $m_0(\bigcup_1^\infty U_n) = \sum_1^\infty m_0(U_n)$  in the topology  $\tau$ . Thus (i) holds.

(ii) If possible, let  $\inf_n \|m_0\|_{p_A}(U_n) > 4\delta > 0$  for some  $A \in \mathcal{E}$ . Then by Lemma 1 we have  $\sup_{x^* \in A} |x^* \circ m_0|(U_n) > 4\delta$  for all  $n$ . Then there exists an  $x_n^* \in A$  such that  $|x_n^* \circ m_0|(U_n) > 4\delta$ . Consequently,  $\sup_{B \in \mathcal{B}_0(T), B \subset U_n} |(x_n^* \circ m_0)(B)| > \delta$  and hence there exists  $B_n \subset U_n$  in  $\mathcal{B}_0(T)$  such that  $|(x_n^* \circ m_0)(B_n)| > \delta$ . Since  $x_n^* \circ m_0$  is a ( $\sigma$ -additive) scalar Baire measure, it is Baire regular and hence there exists an open Baire set  $G_n$  with  $B_n \subset G_n \subset U_n$  such that  $|(x_n^* \circ m_0)(G_n)| > \delta$ . Consequently,  $\inf_n |(x_n^* \circ m_0)(G_n)| > \delta$ . This is absurd, since  $|(x_n^* \circ m_0)(G_n)| \leq \|m_0(G_n)\|_{p_A} \rightarrow 0$  by (i) as  $(G_n)$  is a disjoint sequence in  $\mathcal{U}_0$ .  $\square$

**Lemma 4.**  $m_0$  is Baire inner regular in  $E \in \mathcal{B}_0(T)$  if and only if, for each  $A \in \mathcal{E}$  and  $\varepsilon > 0$ , there exists a compact  $K \in \mathcal{K}_0$  with  $K \subset E$  such that  $\sup_{\mu \in u^*A} |\mu|(E \setminus K) < \varepsilon$ ; i.e. if and only if, for each  $A \in \mathcal{E}$ ,  $u^*A$  is uniformly Baire inner regular in  $E$  in the sense of Definition 1 of [12].

**P r o o f.** Let  $m_0$  be Baire inner regular in  $E \in \mathcal{B}_0(T)$ . Given  $A \in \mathcal{E}$  and  $\varepsilon > 0$ , by Definition 4 there exists  $K \in \mathcal{K}_0$  with  $K \subset E$  such that  $\|m_0\|_{p_A}(E \setminus K) < \varepsilon$ . Then by Lemma 1 and Proposition 5 (ii) we have

$$\|m_0\|_{p_A}(E \setminus K) = \sup_{x^* \in A} |x^* \circ m|(E \setminus K) = \sup_{\mu \in u^*A} |\mu|(E \setminus K) < \varepsilon.$$

The converse is immediate from Definition 4 and Lemma 1 as  $u^*A = \{x^* \circ m : x^* \in A\}$  by Proposition 5 (ii) and  $\Gamma_{\mathcal{E}} = \{p_A : A \in \mathcal{E}\}$  induces the topology  $\tau_e$  of  $X^{**}$ .  $\square$

In the proofs of Lemmas 5 and 6 below we use, respectively, the implications (iii)  $\Rightarrow$  (iv) and (iv)  $\Rightarrow$  (v) of Theorem 1 of [12].

**Lemma 5.** *Let  $m_0(\mathcal{U}_0) \subset X$ . Then:*

- (i)  $m_0$  is Baire inner regular (in  $\tau_e$ ) in each  $U \in \mathcal{U}_0$ .
- (ii) For each  $\varepsilon > 0$  and for each  $A \in \mathcal{E}$ , there exists a  $K \in \mathcal{X}_0$  such that  $\|m\|_{p_A}(T \setminus K) = \sup_{x^* \in A} |x^* \circ m|(T \setminus K) < \varepsilon$ .

**Proof.** Let  $A \in \mathcal{E}$ . Then by Proposition 5 (ii),  $u^*A = \{x^* \circ m : x^* \in A\}$  and by Lemma 2,  $u^*A$  is bounded in  $M(T)$ . By Proposition 1, Lemma 1 and Lemma 3 (ii), for each disjoint sequence  $(U_n)$  of open Baire sets we have  $\lim_n \sup_{x^* \in A} |x^* \circ m|(U_n) = \lim_n \sup_{\mu \in u^*A} |\mu|(U_n) = 0$ . Thus by the implication (iii)  $\Rightarrow$  (iv) of Theorem 1 of [12] the result holds.  $\square$

**Lemma 6.** *Suppose  $m_0$  is Baire inner regular in each  $U \in \mathcal{U}_0$  with respect to the topology  $\tau_e$  of  $X^{**}$  and, for each  $\varepsilon > 0$  and for each  $A \in \mathcal{E}$ , suppose there exists  $K \in \mathcal{X}_0$  such that  $\|m_0\|_{p_A}(T \setminus K) = \sup_{x^* \in A} \{|x^* \circ m|(B) : B \subset T \setminus K, B \in \mathcal{B}_0(T)\} < \varepsilon$  (note that the range of  $m_0$  is contained in  $X^{**}$ ). Then  $m_0$  is Baire inner regular in  $\mathcal{B}_0(T)$  with respect to  $\tau_e$ .*

**Proof.** Let  $A \in \mathcal{E}$ . Then by Lemma 2,  $u^*A$  is bounded in  $M(T)$ . Since  $m_0$  is Baire inner regular in each open Baire set, Lemma 4 implies that  $u^*A$  is uniformly Baire inner regular (in the sense of Definition 1 of [12]) in each open Baire set.

**Claim 1.**

$$(3) \quad \|m\|_{p_A}(T \setminus K) = \sup_{x^* \in A} |x^* \circ m|(T \setminus K) = \sup_{\mu \in u^*A} |\mu|(T \setminus K) < \varepsilon.$$

In fact, by the second hypothesis, by the Borel regularity of  $|x^* \circ m|$ , by Theorem 50.D of [7] and by Lemma 1 (i), Proposition 1 and Proposition 5 (ii), we have

$$\begin{aligned} \|m\|_{p_A}(T \setminus K) &= \sup_{x^* \in A} |x^* \circ m|(T \setminus K) \\ &= \sup_{\mu \in u^*A} \sup_{C \in \mathcal{X}, C \subset T \setminus K} |\mu|(C) \\ &= \sup_{\mu \in u^*A} \sup_{C \in \mathcal{X}_0, C \subset T \setminus K} |\mu|(C) \\ &= \sup_{x^* \in A} \sup_{C \in \mathcal{X}_0, C \subset T \setminus K} |x^* \circ m_0|(C) \\ &< \varepsilon. \end{aligned}$$

Hence the claim holds.

Thus, in virtue of (3), the hypotheses of the lemma show that  $u^*A$  satisfies the hypothesis of the statement (iv) of Theorem 1 of [12]. Consequently, by (iv)  $\Rightarrow$  (v) of Theorem 1 of [12],  $u^*A$  is uniformly Baire inner regular in each  $E \in \mathcal{B}_0(T)$ . Since this holds for all  $A \in \mathcal{E}$  and since  $\Gamma_{\mathcal{E}}$  induces the topology  $\tau_e$ , Lemma 4 yields that  $m_0$  is Baire inner regular in  $\mathcal{B}_0(T)$ .  $\square$

**Lemma 7.** *Suppose  $m$  ( $m_c, m_0$ ) is Borel ( $\sigma$ -Borel, Baire) inner regular (in  $\tau_e$ ) in  $\mathcal{B}(T)$  ( $\mathcal{B}_c(T), \mathcal{B}_0(T)$ ). Then  $m$  ( $m_c, m_0$ , respectively) is  $\sigma$ -additive in  $\tau_e$ .*

**Proof.** Let  $A \in \mathcal{E}$  and let  $\varepsilon > 0$ . Let  $\mathcal{S} = \mathcal{B}(T)$  and  $\gamma = m$  ( $\mathcal{S} = \mathcal{B}_c(T)$  and  $\gamma = m_c$ ;  $\mathcal{S} = \mathcal{B}_0(T)$  and  $\gamma = m_0$ , respectively). Since  $\|\gamma(E)\|_{p_A} \leq \|\gamma\|_{p_A}(E)$  for  $E \in \mathcal{S}$ , it suffices to show that  $\lim_n \|\gamma\|_{p_A}(E_n) = 0$  whenever  $(E_n)$  is a decreasing sequence in  $\mathcal{S}$  with  $\bigcap_1^\infty E_n = \emptyset$ . By hypothesis, for each  $n$  there exists a compact  $K_n \in \mathcal{S}$  with  $K_n \subset E_n$  such that  $\|\gamma\|_{p_A}(E_n \setminus K_n) < \varepsilon/2^n$ . Then adapting suitably the proof at the end of p. 158 and at the top of p. 159 of [1], we can show that there exists  $n_0$  such that  $\|\gamma\|_{p_A}(E_n) < \varepsilon$  for  $n \geq n_0$ . Hence the lemma holds.  $\square$

#### 4. CHARACTERIZATIONS OF WEAKLY COMPACT OPERATORS ON $C_0(T)$

Let  $X$  be a quasicomplete lcHs. Using Propositions 1–6 and Lemmas 1–7 of the preceding sections and Theorem 1 of [11] we will obtain below all the 35 characterizations given in [13] for a continuous linear map  $u: C_0(T) \rightarrow X$  to be weakly compact. As mentioned at the outset, the Borel extension theorem (Proposition 6) for  $\sigma$ -additive  $X$ -valued Baire measures on  $T$  plays a key role in the present proof in contrast to the proofs of the characterization theorems of [13].

**Theorem 1.** *Let  $u: C_0(T) \rightarrow X$  be a continuous linear map, where  $X$  is a quasicomplete lcHs. Let  $m$  be the representing measure of  $u$  and let  $m_c = m|_{\mathcal{B}_c(T)}$  and  $m_0 = m|_{\mathcal{B}_0(T)}$ . Then the following assertions are equivalent.*

- (i)  $u$  is weakly compact.
- (ii) The range of  $m$  is contained in  $X$ .
- (iii) The range of  $m_c$  is contained in  $X$ .
- (iv) The range of  $m_0$  is contained in  $X$ .
- (v)  $m(U) \in X$  for all open sets  $U$  in  $T$ .
- (vi)  $m(F) \in X$  for all closed sets  $F$  in  $T$ .
- (vii)  $m(U) \in X$  for all  $\sigma$ -Borel open sets  $U$  in  $T$ .
- (viii)  $m(U) \in X$  for all open Baire sets  $U$  in  $T$ .
- (ix)  $m(U) \in X$  for all open sets  $U$  in  $T$  which are  $\sigma$ -compact.
- (x)  $m(F) \in X$  for all closed sets  $F$  in  $T$  which are  $G_\delta$ .

- (xi)  $m(U) \in X$  for all open sets  $U$  in  $T$  which are  $F_\sigma$ .
- (xii) For each increasing sequence  $(f_n)_1^\infty \subset C_0(T)$  with  $0 \leq f_n \leq 1$ ,  $(uf_n)$  converges weakly in  $X$ .
- (xiii)  $m$  is  $\sigma$ -additive in the topology  $\tau_e$  of  $X^{**}$ .
- (xiv)  $m_c$  is  $\sigma$ -additive in the topology  $\tau_e$  of  $X^{**}$ .
- (xv)  $m_0$  is  $\sigma$ -additive in the topology  $\tau_e$  of  $X^{**}$ .
- (xvi)  $m$  is strongly additive in the topology  $\tau_e$  of  $X^{**}$ .
- (xvii)  $m_c$  is strongly additive in the topology  $\tau_e$  of  $X^{**}$ .
- (xviii)  $m_0$  is strongly additive in the topology  $\tau_e$  of  $X^{**}$ .
- (xix)  $m$  is Borel regular in  $\tau_e$  of  $X^{**}$ .
- (xx)  $m$  is Borel inner regular in  $\tau_e$  of  $X^{**}$ .
- (xxi)  $m$  is Borel inner regular (in  $\tau_e$ ) in each open set  $U$  in  $T$ .
- (xxii)  $m$  is Borel outer regular (in  $\tau_e$ ) in each compact set  $K$  in  $T$  and Borel inner regular (in  $\tau_e$ ) in the set  $T$ .
- (xxiii)  $m_c$  is  $\sigma$ -Borel regular in  $\tau_e$  of  $X^{**}$ .
- (xxiv)  $m_c$  is  $\sigma$ -Borel inner regular in  $\tau_e$  of  $X^{**}$ .
- (xxv)  $m_c$  is  $\sigma$ -Borel inner regular (in  $\tau_e$ ) in each  $\sigma$ -Borel open set  $U$  in  $T$  and in the set  $T$ .
- (xxvi)  $m_c$  is  $\sigma$ -Borel outer regular (in  $\tau_e$ ) in each compact set  $K$  in  $T$  and  $\sigma$ -Borel inner regular (in  $\tau_e$ ) in the set  $T$ .
- (xxvii)  $m_0$  is Baire regular in  $\tau_e$  of  $X^{**}$ .
- (xxviii)  $m_0$  is Baire inner regular in  $\tau_e$  of  $X^{**}$ .
- (xxix)  $m_0$  is Baire inner regular (in  $\tau_e$ ) in each open Baire set  $U$  in  $T$  and in the set  $T$ .
- (xxx)  $m_0$  is Baire outer regular (in  $\tau_e$ ) in each compact  $G_\delta$  in  $T$  and Baire inner regular (in  $\tau_e$ ) in the set  $T$ .
- (xxxii) All bounded Borel measurable scalar functions  $f$  on  $T$  are  $m$ -integrable and  $\int_T f \, dm \in X$ .
- (xxxiii) All bounded  $\mathcal{B}_c(T)$ -measurable scalar functions  $f$  on  $T$  are  $m_c$ -integrable and  $\int_T f \, dm_c \in X$ .
- (xxxiiii) All bounded Baire measurable scalar functions  $f$  on  $T$  are  $m_0$ -integrable and  $\int_T f \, dm_0 \in X$ .
- (xxxv) All bounded scalar functions  $f$  belonging to the first Baire class in  $T$  are  $m_0$ -integrable and  $\int_T f \, dm_0 \in X$ .
- (xxxvi)  $u^{**}f \in X$  for all bounded scalar functions  $f$  belonging to the first Baire class in  $T$ .

Proof. In the sequel we will prove only those implications which are not obvious.

(i)  $\Rightarrow$  (ii): By (i) and Proposition 2,  $u^{**}C_0^{**}(T) \subset X$  and by Proposition 5 (v),  $m(E) = u^{**}(\chi_E)$  for  $E \in \mathcal{B}(T)$ . As  $\mathcal{B}(T) \subset C_0^{**}(T)$ , (ii) holds.

(viii)  $\Rightarrow$  (iv): In fact, by hypothesis (viii) and by Lemmas 5 and 6,  $m_0$  is Baire inner regular in  $\tau_e$  of  $X^{**}$ . Given  $K \in \mathcal{K}_0$ , by Theorem 50.D of Halmos [7] there exists  $U \in \mathcal{U}_0$  such that  $K \subset U$  and hence  $m_0(K) = m_0(U) - m_0(U \setminus K) \in X$ . Thus  $m_0(\mathcal{K}_0) \subset X$ . Let  $E \in \mathcal{B}_0(T)$ . Let  $D(E) = \{K \in \mathcal{K}_0: K \subset E\}$  and let  $K_1 \supseteq K_2$  for  $K_1, K_2 \in D(E)$  if  $K_1 \supset K_2$ . Then by the Baire inner regularity of  $m_0$  in  $E$ ,  $\lim_{D(E)} m_0(K) = m_0(E)$  so that the net  $\{m_0(K): K \in D(E)\}$  is  $\tau_e$ -Cauchy with the limit  $m_0(E)$ . Since by Proposition 5 (iv),  $m$  has  $\tau_e$ -bounded range in  $X^{**}$ ,  $m_0(\mathcal{K}_0)$  is  $\tau$ -bounded in  $X$ . Thus there exists a  $\tau$ -bounded closed set  $H$  in  $X$  such that  $m_0(\mathcal{K}_0(T)) \subset H$ . Since  $X$  is quasicomplete, we conclude that  $m_0(E) \in H \subset X$ . Thus  $m_0$  has the range in  $X$ .

(iv)  $\Rightarrow$  (i): In fact, by hypothesis, Proposition 5 (i) and the Orlicz-Pettis theorem,  $m_0$  is  $\sigma$ -additive in  $\tau$ . Then by Proposition 6 there exists a unique  $X$ -valued Borel regular  $\sigma$ -additive extension  $\hat{m}$  of  $m_0$  on  $\mathcal{B}(T)$ . As each  $f \in C_0(T)$  is a bounded Baire measurable function by Theorem 51.B of Halmos [7], by Proposition 5 (iii) we have

$$x^*uf = \int_T f d(x^* \circ m) = \int_T f d(x^* \circ m_0) = \int_T f d(x^* \circ \hat{m}), \quad f \in C_0(T).$$

Since  $x^* \circ m \in M(T)$  by Proposition 5 (i) and since  $x^* \circ \hat{m} \in M(T)$  as  $\hat{m}$  is Borel regular and  $\sigma$ -additive, it follows by the uniqueness part of the Riesz representation theorem that  $x^* \circ m = x^* \circ \hat{m}$  for each  $x^* \in X^*$ . Since  $m$  has the range in  $X^{**}$  and  $\hat{m}$  has the range in  $X$  we conclude that  $m = \hat{m}$  and hence  $m$  not only has the range in  $X$  but also is  $\sigma$ -additive in  $\mathcal{B}(T)$  in  $\tau$ . Thus, given a disjoint sequence  $(U_n)$  of open sets in  $T$ ,  $m(\bigcup_1^\infty U_n) = \sum_1^\infty m(U_n)$  and in particular,  $\lim_n m(U_n) = 0$ . Thus, for each equicontinuous subset  $A$  of  $X^*$ , Proposition 5 (ii) yields  $\lim_n \|m(U_n)\|_{p_A} = \lim_n \sup_{x^* \in A} |(x^* \circ m)(U_n)| = \lim_n \sup_{\mu \in u^*A} |\mu(U_n)| = 0$ . Moreover, by Lemma 2,  $u^*A$  is bounded in  $M(T)$ . Therefore, by Proposition 3,  $u^*A$  is relatively weakly compact in  $M(T)$ . Consequently, by Proposition 2,  $u$  is weakly compact. Thus (i) holds.

(x)  $\Rightarrow$  (xi): Let  $U$  be an open set in  $T$  such that it is a countable union of closed sets. Then  $T \setminus U$  is a closed set which is  $G_\delta$  and hence by hypothesis (x) we have  $m(U) = m(T) - m(T \setminus U) \in X$ . Hence (xi) holds.

(ix)  $\Rightarrow$  (viii): by § 14, Chapter III of Dinculeanu [2].

(ii)  $\Rightarrow$  (xii): Let  $(f_n)$  be as in (xii). Then  $\lim_n f_n(t) = f(t)$  exists in  $[0,1]$  for each  $t \in T$  and  $f$  is Borel measurable. Then the hypothesis (ii) combined with Proposition 5 (i) and the Orlicz-Pettis theorem implies that  $m$  is  $\sigma$ -additive in  $\mathcal{B}(T)$ .

Consequently, by Proposition 4 we obtain

$$\lim_n \int_T f_n \, dm = \int_T f \, dm \in X.$$

Then by Propositions 4 (i) and 5 (iii) we have

$$\lim_n x^* u f_n = \lim_n \int_T f_n \, d(x^* \circ m) = x^* \left( \lim_n \int_T f_n \, dm \right) = x^* \left( \int_T f \, dm \right)$$

for all  $x^* \in X^*$ . Thus (xii) holds.

(xii)  $\Rightarrow$  (viii): Let  $U \in \mathcal{U}_0$ . Then by § 14, Chapter III of Dinculeanu [2], there exists a sequence  $(K_n) \subset \mathcal{K}_0$  such that  $K_n \nearrow U$ . By Urysohn's lemma we can choose an increasing sequence  $g_n$  of non negative continuous functions with compact supports such that  $g_n \nearrow \chi_U$ . Then by hypothesis there exists a vector  $x_0 \in X$  such that  $\lim_n x^* u g_n = x^* x_0$  for all  $x^* \in X^*$ . Therefore, by the Lebesgue bounded convergence theorem and by Proposition 5 we have  $x^* x_0 = \lim_n \int_T g_n \, d(x^* \circ m) = x^* m(U)$  for all  $x^* \in X^*$ . Since  $m(U) \in X^{**}$ , it follows that  $m(U) = x_0 \in X$ . Hence (viii) holds.

(ii)  $\Rightarrow$  (xiii): By (ii)  $m$  has the range in  $X$  and hence by Proposition 5 (i) and the Orlicz-Pettis theorem  $m$  is  $\sigma$ -additive in  $\tau$ . Since  $\tau_e|_X = \tau$ , (xiii) holds.

(xv)  $\Rightarrow$  (i): Let  $Y$  be the completion of  $(X^{**}, \tau_e)$ . Then by hypothesis  $m_0: \mathcal{B}_0(T) \rightarrow Y$  is  $\sigma$ -additive in  $\tau_e$  and hence by Proposition 6 there exists a unique  $Y$ -valued Borel regular  $\sigma$ -additive (in  $\tau_e$ ) extension  $\tilde{m}$  of  $m_0$  on  $\mathcal{B}(T)$ . Each  $f \in C_0(T)$  is a bounded Baire measurable function by Theorem 51.B of Halmos [7] and consequently, by Proposition 5 (iii) we have

$$x^* u f = \int_T f \, d(x^* \circ m) = \int_T f \, d(x^* \circ m_0) = \int_T f \, d(x^* \circ \tilde{m})$$

for each  $f \in C_0(T)$ . By Proposition 5 (i),  $x^* \circ m \in M(T)$ . Since each  $x^* \in X^*$  is  $\tau_e$ -continuous in  $X^{**}$ , it follows that  $x^* \circ \tilde{m}$  is a  $\sigma$ -additive regular Borel complex measure on  $T$  and hence  $x^* \circ \tilde{m} \in M(T)$ . Thus the continuous linear functional  $x^* u$  on  $C_0(T)$  is represented by both  $x^* \circ m$  and  $x^* \circ \tilde{m}$  belonging to  $M(T)$  and hence  $x^* \circ m = x^* \circ \tilde{m}$  for all  $x^* \in X^*$ . Since  $m$  takes values in  $X^{**}$  and  $\tilde{m}$  takes values in  $Y$ , it follows that  $m = \tilde{m}$  so that  $\tilde{m}$  has values in  $X^{**}$ . Moreover,  $m (= \tilde{m})$  is  $\sigma$ -additive in  $\tau_e$ . Consequently, given a disjoint sequence  $(U_n)$  of open sets in  $T$ , by Proposition 5 (ii) we have  $\lim_n \|m(U_n)\|_{p_A} = \lim_n \sup_{x^* \in A} |(x^* \circ m)(U_n)| = \lim_n \sup_{\mu \in u^* A} |\mu(U_n)| = 0$  for each  $A \in \mathcal{E}$ . Moreover, for such  $A$ , by Lemma 2,  $u^* A$  is bounded in  $M(T)$ . Then by an argument similar to that in the end of the proof of (iv)  $\Rightarrow$  (i) we conclude that  $u$  is weakly compact. Hence (i) holds.

(xviii)  $\Rightarrow$  (i): Let  $\Sigma(\mathcal{B}_0(T))$  be the Banach space of all bounded complex functions which are uniform limits of sequences of  $\mathcal{B}_0(T)$ -simple functions, with pointwise addition and scalar multiplication and with the supremum norm  $\|\cdot\|_T$ . Let

$$Vf = \int_T f \, dm_0, \quad f \in \Sigma(\mathcal{B}_0(T)).$$

By Proposition 5 (iv),  $m_0$  is a  $\tau_e$ -bounded vector measure and hence, by Lemma 6 of [11],  $V$  is a well defined  $X^{**}$ -valued continuous linear map. Then as the representing measure  $m_0$  of  $V$  (see Definition 2 of [11]) is strongly additive by hypothesis (xviii), by Theorem 1 of [11]  $V$  is a weakly compact operator. By Theorem 51.B of Halmos [7] each  $f \in C_0(T)$  is Baire measurable and bounded and hence is the uniform limit of a sequence of Baire simple functions. Hence  $C_0(T) \subset \Sigma(\mathcal{B}_0(T))$ . In particular,  $V|_{C_0(T)}$  is weakly compact. Moreover, by Propositions 4 (i) and 5 (iii), we have

$$x^*Vf = \int_T f \, d(x^* \circ m_0) = \int_T f \, d(x^* \circ m) = x^*uf, \quad f \in C_0(T)$$

for each  $x^* \in X^*$ . Since  $Vf \in X^{**}$  and  $uf \in X$ , we conclude that  $Vf = uf$  for each  $f \in C_0(T)$ . Consequently,  $u = V|_{C_0(T)}$  and hence  $\{uf: \|f\|_T \leq 1\}$  is relatively  $\sigma(X^{**}, X^{***})$ -compact. Since  $u(C_0(T)) \subset X$ , it follows that  $\{uf: \|f\|_T \leq 1\}$  is relatively weakly compact in  $X$ . Thus  $u$  is weakly compact. Hence (i) holds.

(ii)  $\Rightarrow$  (xix): By (ii), Proposition 5 (i) and the Orlicz-Pettis theorem,  $m$  is  $\sigma$ -additive in  $\mathcal{B}(T)$  in the topology  $\tau$  of  $X$ . Then  $m_0$  is  $\sigma$ -additive in  $\mathcal{B}_0(T)$  and has the range in  $X$ . Therefore, by Proposition 6 there exists a unique Borel regular  $X$ -valued  $\sigma$ -additive (in  $\tau$ ) extension  $\hat{m}$  of  $m_0$  on  $\mathcal{B}(T)$ . Then by Proposition 5 (iii) and by the fact that each  $f \in C_0(T)$  is bounded and Baire measurable (by Theorem 51.B of [7]), we have

$$x^*uf = \int_T f \, d(x^* \circ m) = \int_T f \, d(x^* \circ m_0) = \int_T f \, d(x^* \circ \hat{m})$$

for each  $x^* \in X^*$  and  $f \in C_0(T)$ . Since  $x^* \circ m \in M(T)$  by Proposition 5 (i) and since  $x^* \circ \hat{m} \in M(T)$  as  $\hat{m}$  is Borel regular and  $\sigma$ -additive in  $\tau$  with values in  $X$ , we conclude that  $x^* \circ m = x^* \circ \hat{m}$  for each  $x^* \in X^*$ . Since by hypothesis  $m$  has the range in  $X$  and  $\hat{m}$  in  $X$ , it follows that  $m = \hat{m}$ . Thus  $m$  is Borel regular in  $\tau$  and hence  $m$  is Borel regular in  $\tau_e$  as  $\tau_e|_X = \tau$ . Thus (xix) holds.

(xxi) (or (xxv), (xxix))  $\Rightarrow$  (xxviii): Let  $U \in \mathcal{U}_0$  or let  $U = T$ . Let  $A \in \mathcal{E}$  and  $\varepsilon > 0$ . Then by hypothesis and by Theorem 50.D of Halmos [7] there exists a compact  $G_\delta$   $K$  such that  $K \subset U$  and  $\|m\|_{p_A}(U \setminus K) < \varepsilon$  ( $\|m_c\|_{p_A}(U \setminus K) < \varepsilon$ ,



$\|m_0\|_{p_A}(U \setminus K) < \varepsilon$ , respectively). Thus, in particular,  $\|m_0\|_{p_A}(E) < \varepsilon$  for all  $E \in \mathcal{B}_0(T)$  with  $E \subset U \setminus K$ . Since this holds for all  $U \in \mathcal{U}_0$  and for  $U = T$ , the conditions of Lemma 6 are satisfied by  $m_0$ . Therefore,  $m_0$  is Baire inner regular in  $\mathcal{B}_0(T)$ . Hence (xxviii) holds.

(xxviii)  $\Rightarrow$  (xv): by Lemma 7.

(xxii)  $\Rightarrow$  (i): Let  $K \in \mathcal{K}$  and let  $A \in \mathcal{E}$ . Given  $\varepsilon > 0$ , by hypothesis there exists an open set  $U$  in  $T$  such that  $\|m\|_{p_A}(U \setminus K) < \varepsilon$ . Then by Lemma 1 (i) and Proposition 5 (ii) we have  $\sup\{|x^* \circ m|(U \setminus K) : x^* \in A\} = \sup_{\mu \in u^*A} |\mu|(U \setminus K) < \varepsilon$  and by Lemma 2,  $u^*A$  is bounded in  $M(T)$ . Thus condition (iv) (a) of Proposition 3 is satisfied by  $u^*A$ . Since  $m$  is inner regular in  $T$ , there exists a compact set  $C$  such that  $\|m\|_{p_A}(T \setminus C) < \varepsilon$  so that by an argument similar to that above we have  $\sup_{\mu \in u^*A} |\mu|(T \setminus C) < \varepsilon$ . Therefore, condition (iv) (b) of Proposition 3 is also satisfied by  $u^*A$ . Hence by Proposition 3,  $u^*A$  is relatively weakly compact in  $M(T)$  and consequently, by Proposition 2,  $u$  is weakly compact. Thus (i) holds.

(ii)  $\Rightarrow$  (xxiii): Proceeding as in the proof of (ii)  $\Rightarrow$  (xix), we have  $m = \hat{m}$  on  $\mathcal{B}(T)$ . Since  $\hat{m}|_{\mathcal{B}_c(T)}$  is  $\sigma$ -Borel regular by Proposition 6, we conclude that  $m_c$  is  $\sigma$ -Borel regular in  $\tau$  and hence in  $\tau_e$ . Thus (xxiii) holds.

(xxiv)  $\Rightarrow$  (xiv): by Lemma 7.

(xxiii) implies the first part of (xxv) and (xix) implies the second part of (xxv). As (xxv)  $\Rightarrow$  (xxviii), it follows that (i)  $\Leftrightarrow$  (xxv).

(xix)  $\Rightarrow$  (xxvi): Given  $K \in \mathcal{K}$ ,  $A \in \mathcal{E}$  and  $\varepsilon > 0$ , then by hypothesis there exists an open set  $U$  with  $U \supset K$  such that  $\|m\|_{p_A}(U \setminus K) < \varepsilon$ . By Theorem 50.D of Halmos [7] we can choose a  $V \in \mathcal{U}_0$  such that  $K \subset V \subset U$  so that  $\|m_c\|_{p_A}(V \setminus K) < \varepsilon$ . Thus  $m_c$  is  $\sigma$ -Borel outer regular in  $K$ . Clearly,  $m_c$  is  $\sigma$ -Borel inner regular in  $T$  as  $m$  is, by hypothesis, Borel inner regular in  $T$ . Hence (xxvi) holds.

(xxvi)  $\Rightarrow$  (i): Let  $K \in \mathcal{K}$ . Proceeding as in the proof of (xxii)  $\Rightarrow$  (i), we have  $\|m_c\|_{p_A}(U \setminus K) < \varepsilon$ , where  $U$  is a  $\sigma$ -Borel open set containing  $K$ . Thus by Lemma 1 (ii) and Proposition 1 we have  $\|m_c\|_{p_A}(U \setminus K) = \sup_{x^* \in A} |x^* \circ m_c|(U \setminus K) = \sup_{x^* \in A} |x^* \circ m|(U \setminus K) < \varepsilon$ . Hence, by Proposition 5 (ii),  $\sup_{\mu \in u^*A} |\mu|(U \setminus K) < \varepsilon$ . Since  $u^*A$  is bounded in  $M(T)$  by Lemma 2, condition (iv) (a) of Proposition 3 is satisfied by  $u^*A$ . Again by hypothesis, there exists a compact  $C$  such that  $\|m_c\|_{p_A}(T \setminus C) < \varepsilon$ . Thus for each compact  $K \subset T \setminus C$ , by Lemma 1 (ii) we have  $\sup_{x^* \in A} |x^* \circ m|(K) < \varepsilon$ . As  $|x^* \circ m|$  is Borel regular by Proposition 5 (i) for each  $x^* \in A$ , and  $x^* \circ m = u^*x^*$  by Proposition 5 (ii), it follows that  $\sup_{x^* \in A} |x^* \circ m|(T \setminus C) = \sup_{\mu \in u^*A} |\mu|(T \setminus C) \leq \varepsilon$ . Thus condition (iv) (b) of Proposition 3 is also satisfied by  $u^*A$ . Therefore,  $u^*A$  is relatively weakly compact in  $M(T)$  for each  $A \in \mathcal{E}$ . Now by Proposition 2 we conclude that  $u$  is weakly compact. Hence (i) holds.

(xv)  $\Rightarrow$  (xxvii): Since  $m_0$  is  $\sigma$ -additive in  $\tau_e$ , by the first part of Proposition 6,  $m_0$  is Baire regular in  $\tau_e$ . Thus (xxvii) holds.

(xxix)  $\Rightarrow$  (xxviii): by Lemma 6.

(xix)  $\Rightarrow$  (xxix): Let  $U \in \mathcal{U}_0$ ,  $A \in \mathcal{E}$  and  $\varepsilon > 0$ . By hypothesis, there exists a compact  $K \subset U$  such that  $\|m\|_{p_A}(U \setminus K) < \varepsilon$ . By Theorem 50.D of Halmos [7] there exists a compact  $C \in \mathcal{X}_0$  such that  $K \subset C \subset U$ . Then  $\|m_0\|_{p_A}(U \setminus C) < \varepsilon$ . Hence  $m_0$  is Baire inner regular in  $U$ . As  $m$  is Borel inner regular in  $T$ , there exists  $K \in \mathcal{K}$  such that  $\|m\|_{p_A}(T \setminus K) < \varepsilon$ . By Theorem 50.D of Halmos [7] there exists  $C \in \mathcal{X}_0$  such that  $K \subset C$  and hence  $\|m_0\|_{p_A}(B) < \varepsilon$  for all  $B \in \mathcal{B}_0(T)$  with  $B \subset T \setminus C$ . Thus  $m_0$  is Baire inner regular in  $T$ . Hence (xxix) holds.

(xix)  $\Rightarrow$  (xxx): Let  $K \in \mathcal{X}_0$ ,  $A \in \mathcal{E}$  and  $\varepsilon > 0$ . By hypothesis and by Theorem 50.D of Halmos [7] there exists  $U \in \mathcal{U}_0$  with  $K \subset U$  such that  $\|m\|_{p_A}(U \setminus K) < \varepsilon$  so that by (i) and (iii) of Lemma 1 we have  $\|m_0\|_p(U \setminus K) < \varepsilon$ . Similarly, we can show that  $m_0$  is Baire inner regular in  $T$ . Hence (xxx) holds.

(xxx)  $\Rightarrow$  (xxix): Clearly, it suffices to show that  $m_0$  is Baire inner regular in each open Baire set. Given  $A \in \mathcal{E}$  and  $\varepsilon > 0$ , by the hypothesis of Baire inner regularity of  $m_0$  in  $T$  and by Theorem 50.D of Halmos [7] there exists a compact  $\Omega \in \mathcal{X}_0$  such that  $\|m_0\|_{p_A}(T \setminus \Omega) < \varepsilon/2$ . Let  $U \in \mathcal{U}_0$  such that  $U$  is relatively compact.

**Claim 1.**  $m_0$  is Baire inner regular in  $U$ .

In fact, by Theorem 50.D of Halmos [7] we can choose a compact  $C \in \mathcal{X}_0$  such that  $\bar{U} \subset C$ . Then  $U = C \setminus (C \setminus U)$  and  $C \setminus U \in \mathcal{X}_0$  by Theorem 51.D of Halmos [7]. Therefore, by hypothesis there exists  $W \in \mathcal{U}_0$  with  $W \supset C \setminus U$  such that  $\|m_0\|_{p_A}(W \setminus (C \setminus U)) < \varepsilon$ . Now  $U = C \setminus (C \setminus U) \supset C \setminus W$  and  $C \setminus W \in \mathcal{X}_0$  again by Theorem 51.D of Halmos [7]. Moreover,  $U \setminus (C \setminus W) = U \cap ((T \setminus C) \cup W) = U \cap W$ . On the other hand,  $W \setminus (C \setminus U) \supset W \cap U$ . Therefore,  $\|m_0\|_{p_A}(U \setminus (C \setminus W)) < \varepsilon$ . Thus the claim holds.

Now let  $U \in \mathcal{U}_0$ . Choose by Theorem 50.D of Halmos [7] a relatively compact open Baire set  $V$  such that  $\Omega \subset V$ . Then  $U \cap V$  is relatively compact and belongs to  $\mathcal{U}_0$ . Therefore, by Claim 1,  $m_0$  is Baire inner regular in  $U \cap V$  and hence there exists a compact  $K \in \mathcal{X}_0$  with  $K \subset U \cap V$  such that  $\|m_0\|_{p_A}((U \cap V) \setminus K) < \varepsilon/2$ . Then  $K \subset U$  and  $\|m_0\|_{p_A}(U \setminus K) \leq \|m_0\|_{p_A}((U \cap V) \setminus K) + \|m_0\|_{p_A}(U \setminus \Omega) < \varepsilon$ . Therefore,  $m_0$  is Baire inner regular in each open Baire set and hence (xxix) holds.

(ii)  $\Rightarrow$  (xxx), (xxxii) and (xxxiii): By (ii), Proposition 5(i) and the Orlicz-Pettis theorem  $m$  is  $X$ -valued and  $\sigma$ -additive in  $\tau$ . Since every bounded Borel ( $\sigma$ -Borel, Baire) measurable scalar function is the uniform limit of a sequence of Borel ( $\sigma$ -Borel, Baire) simple functions and  $m$  is a  $\tau$ -bounded  $X$ -valued vector measure,  $f$  is  $m$ -integrable (see Definition 1 of [11]) and  $\int_T f dm \in X$  ( $f$  is  $m_c$ -integrable and  $\int_T f dm_c \in X$ ,  $f$  is  $m_0$ -integrable and  $\int_T f dm_0 \in X$ , respectively).

(xxxii) (or xxxiii), (xxxiii)  $\Rightarrow$  (ii) ((iii), (iv)): Let  $E \in \mathcal{B}(T)$  ( $E \in \mathcal{B}_c(T)$ ,  $E \in \mathcal{B}_0(T)$ ). Then by hypothesis,  $m(E)$  ( $m_c(E)$ ,  $m_0(E)$ ) belongs to  $X$ . Thus (ii) ((iii), (iv), respectively) holds.

(xxxiv)  $\Rightarrow$  (viii): Let  $U$  be an open Baire set. Then by § 14, Chapter III of Dinculeanu [2], there exists an increasing sequence  $K_n$  of compact  $G_\delta$  sets such that  $U = \bigcup_1^\infty K_n$ . Then by Urysohn's lemma we can choose non negative continuous functions  $g_n$  with compact supports such that  $g_n \nearrow \chi_U$ . Thus  $\chi_U$  belongs to the first Baire class and is bounded. Then by hypothesis,  $m_0(U) \in X$ . Thus (viii) holds.

(i)  $\Rightarrow$  (xxxv): If  $u$  is weakly compact, then by Proposition 2,  $u^{**}$  has the range in  $X$ . Since the bounded scalar functions of the first Baire class belong to  $C_0^{**}(T)$ , (xxxv) holds.

(xxxv)  $\Rightarrow$  (viii): By Proposition 5 (v),  $u^{**}(\chi_U) = m(U)$  for  $U \in \mathcal{U}_0$ . As observed in the proof of (xxxiv)  $\Rightarrow$  (viii),  $\chi_U$  is bounded and belongs to the first Baire class. Hence, by hypothesis,  $m(U) \in X$ . Thus (viii) holds.

This completes the proof of the theorem. □

**Remark 2.** As in [13], the strict Dunford-Pettis property of  $C_0(T)$  is an immediate consequence of the above theorem and the proof of the latter is not based on this property unlike the proof of Theorem 6 of Grothendieck [6]. Theorem 5.3 of Thomas [16] is also deducible from the above theorem by the same argument as that used in the proof of Theorem 13 in [13].

**Remark 3.** All these 35 characterizations are given in [13] in Theorems 2–9. Some of the proofs given here are the same as those in [13] (for example, (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii) of Theorem 2 of [13], (i)  $\Leftrightarrow$  (xi) of Theorem 3 of [13] and Theorem 9 of [13]) but, for the sake of completeness, we have given the proofs of all non obvious equivalences of these 35 characterizations. In the present proof the use of Theorems 1 and 2 of [12] has been dispensed with unlike the proof in [13] and instead, the Borel extension theorem has been used along with the first part of Theorem 1 of [13], Lemma 1 and Theorem 2 of [6], Theorem 1 of [11] and Lemmas 1–7.

### References

- [1] *J. Diestel and J. J. Uhl: Vector Measures. Survey No.15. Amer. Math. Soc., Providence, RI., 1977.*
- [2] *N. Dinculeanu: Vector Measures. Pergamon Press, New York, 1967.*
- [3] *N. Dinculeanu and I. Kluvánek: On vector measures. Proc. London Math. Soc. 17(1967), 505–512.*
- [4] *I. Dobrakov and T. V. Panchapagesan: A simple proof of the Borel extension theorem and weak compactness of operators. To appear in Czechoslovak Math. J.*
- [5] *R. E. Edwards: Functional Analysis, Theory and Applications. Holt, Rinehart and Winston, New York, 1965.*

- [6] *A. Grothendieck*: Sur les applications lineares faiblement compactes d'espaces du type  $C(K)$ . *Canad. J. Math.* 5 (1953), 129–173.
- [7] *P. R. Halmos*: *Measure Theory*. Van Nostrand, New York, 1950.
- [8] *I. Kluvánek*: Characterizations of Fourier-Stieltjes transform of vector and operator valued measures. *Czechoslovak Math. J.* 17 (1967), 261–277.
- [9] *T. V. Panchapagesan*: On complex Radon measures I. *Czechoslovak Math. J.* 42 (1992), 599–612.
- [10] *T. V. Panchapagesan*: On complex Radon measures II. *Czechoslovak Math. J.* 43 (1993), 65–82.
- [11] *T. V. Panchapagesan*: Applications of a theorem of Grothendieck to vector measures. *J. Math. Anal. Appl.* 214 (1997), 89–101.
- [12] *T. V. Panchapagesan*: Baire and  $\sigma$ -Borel characterizations of weakly compact sets in  $M(T)$ . *Trans. Amer. Math. Soc.* (1998), 4539–4547.
- [13] *T. V. Panchapagesan*: Characterizations of weakly compact operators on  $C_0(T)$ . *Trans. Amer. Math. Soc.* (1998), 4549–4567.
- [14] *T. V. Panchapagesan*: On the limitations of the Grothendieck techniques. To appear in *Rev. Real Acad. Cienc. Exact. Fis. Natur. Madrid*.
- [15] *M. Sion*: Outer measures with values in a topological group. *Proc. London Math. Soc.* 19 (1969), 89–106.
- [16] *E. Thomas*: L'integration par rapport a une mesure de Radon vectorielle. *Ann. Inst. Fourier (Grenoble)* 20 (1970), 55–191.

*Author's address*: Departamento de Matematicas, Facultad de Ciencias, Universidad de los Andes, Merida, Venezuela, e-mail: [panchapa@ciens.ula.ve](mailto:panchapa@ciens.ula.ve).