

A. D. Gadjiev; R. O. Efendiyev; E. Ibikli

On Korovkin type theorem in the space of locally integrable functions

Czechoslovak Mathematical Journal, Vol. 53 (2003), No. 1, 45–53

Persistent URL: <http://dml.cz/dmlcz/127779>

Terms of use:

© Institute of Mathematics AS CR, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON KOROVKIN TYPE THEOREM IN THE SPACE
OF LOCALLY INTEGRABLE FUNCTIONS

A. D. GADJIEV, R. O. EFENDIYEV, Baku, and E. İBIKLİ, Ankara

(Received November 24, 1999)

Abstract. It is shown that a Korovkin type theorem for a sequence of linear positive operators acting in weighted space $L_{p,w}(\text{loc})$ does not hold in all this space and is satisfied only on some subspace.

Keywords: linear positive operators, Korovkin type theorem, weighted $L_p(\text{loc})$ spaces

MSC 2000: 41A36, 41A25

1.

A Korovkin type theorem for linear positive operators acting from $L_p(a, b)$ to $L_p(a, b)$ was studied in [4] and then some new results in this direction were established. We refer to the papers [1], [2], [3], [8], [10], [11], [12] [13]. Note that all the results just mentioned are devoted to the case of a finite interval $[a, b]$.

We consider the space of locally integrable functions on the entire real axis, and the sequences of linear positive operators defined in this space.

For $w(x) = 1 + x^2$, $-\infty < x < \infty$ and any fixed $h > 0$, we will denote by $L_{p,w}(\text{loc})$ the space of measurable functions f satisfying the inequality

$$(1) \quad \left(\frac{1}{2h} \int_{x-h}^{x+h} |f(t)|^p dt \right)^{\frac{1}{p}} \leq M_f w(x), \quad -\infty < x < \infty,$$

where $p \geq 1$ and M_f is a positive constant which depends on the function f . Obviously, $L_{p,w}(\text{loc})$ is a linear normed space with norm

$$\|f\|_{p,w} = \sup_{-\infty < x < \infty} \frac{\left(\frac{1}{2h} \int_{x-h}^{x+h} |f(t)|^p dt \right)^{\frac{1}{p}}}{w(x)}$$

($\|f\|_{p,w}$ may depend also on h). To simplify notation, we need the following. For any finite real numbers a and b put

$$(2) \quad \|f; L_p(a, b)\| = \left(\frac{1}{b-a} \int_a^b |f(t)|^p dt \right)^{\frac{1}{p}},$$

$$(3) \quad \|f; L_{p,w}(a, b)\| = \sup_{a \leq x \leq b} \frac{\|f; L_p(x-h, x+h)\|}{w(x)},$$

$$(4) \quad \|f; L_{p,w}(|x| \geq a)\| = \sup_{|x| \geq a} \frac{\|f; L_p(x-h, x+h)\|}{w(x)}.$$

It follows that the norm in $L_{p,w}(\text{loc})$ may be written in the form

$$\|f\|_{p,w} = \sup_{-\infty < x < \infty} \frac{\|f; L_p(x-h, x+h)\|}{w(x)}.$$

Let $L_{p,w}^k(\text{loc})$ be the subspace of all functions $f \in L_{p,w}(\text{loc})$ for which there exists a constant k_f such that

$$(5) \quad \lim_{|x| \rightarrow \infty} \frac{\|f - k_f w; L_p(x-h, x+h)\|}{w(x)} = 0.$$

In the case of $k_f = 0$ we will write $L_{p,w}^0(\text{loc})$. Korovkin type theorems for sequence of linear positive operators acting in weighted space of continuous functions, defined on the entire real line, were studied in [5]¹, [6]. We will study these theorems in the space $L_{p,w}(\text{loc})$. The set of all linear positive operators acting from $L_{p,w}(\text{loc})$ to $L_{p,w}(\text{loc})$ will be denoted by $(L_{p,w}(\text{loc}) \rightarrow L_{p,w}(\text{loc}))^+$.

2.

We shall deal with the following problem.

Let the sequence of operators $L_n \in (L_{p,w}(\text{loc}) \rightarrow L_{p,w}(\text{loc}))^+$ satisfy the conditions:

i) The norms of these operators are uniformly bounded, that is,

$$(6) \quad \|L_n\| \leq C < \infty,$$

where C is a constant independent of n ;

¹ A. D. Gadjiev = A. D. Gadźiev (also in other translation papers A. D. Gadzhiev, A. D. Gadźiev).

ii) For $m = 0, 1, 2$

$$(7) \quad \lim_{n \rightarrow \infty} \|L_n(t^m; x) - x^m\|_{p,w} = 0,$$

where $L_n(t^m; x) := L_n(t^m)(x)$.

Is it possible to assert then that for each function $f \in L_{p,w}(\text{loc})$

$$\lim_{n \rightarrow \infty} \|L_n(f; x) - f(x)\|_{p,w} = 0?$$

We show that the answer to this question is negative.

Our main result is the following.

Theorem 1. *There exists a sequence of operators $L_n \in (L_{p,w}(\text{loc}) \rightarrow L_{p,w}(\text{loc}))^+$ satisfying conditions (6), (7) and there exists a function f^* in $L_{p,w}(\text{loc})$ for which*

$$\overline{\lim}_{n \rightarrow \infty} \|L_n f^* - f^*\|_{p,w} \geq 2^{1-\frac{1}{p}}.$$

Proof. We define a sequence of operators L_n by the formulas

$$L_n(f; x) = \begin{cases} \frac{x^2}{(x+h)^2} f(x+h) & \text{for } (2n-2)h \leq x \leq (2n+1)h, \\ f(x) & \text{otherwise.} \end{cases}$$

It is easy to see that

$$\|L_n f\|_{p,w} \leq 4\|f\|_{p,w},$$

that is, L_n are bounded operators belonging to $(L_{p,w}(\text{loc}) \rightarrow L_{p,w}(\text{loc}))^+$ and (6) holds.

Now, for $m = 0, 1$,

$$\|L_n(t^m; x) - x^m\|_{p,w} \leq \sup_{(2n-1)h \leq x \leq 2nh} \frac{(x+h)^m}{1+x^2} \leq \frac{(2n+1)^m h^m}{1+4h^2(2n-1)^2}$$

and therefore

$$\lim_{n \rightarrow \infty} \|L_n(t^m; x) - x^m\|_{p,w} = 0, \quad m = 0, 1.$$

Also, since $L_n(t^2; x) = x^2$, the conditions (7) hold.

Consider the function

$$f^*(x) = \begin{cases} x^2 & \text{if } x \in \bigcup_{k=1}^{\infty} [(2k-1)h, 2kh), \\ -x^2 & \text{if } x \in \bigcup_{k=0}^{\infty} (2kh, (2k+1)h], \\ 0 & \text{if } x < 0. \end{cases}$$

Then $f^* \in L_{p,w}(\text{loc})$ and we get

$$\begin{aligned} \|L_n f^* - f^*\|_{p,w} &= \sup_{(2n-1)h < x < (2n+1)h} \frac{\|L_n f^* - f^*; L_p((2n-1)h, (2n+1)h)\|}{w(x)} \\ &\geq \frac{1}{w(2nh)} \left(\frac{1}{2h} \int_{(2n-1)h}^{2nh} \left| \frac{y^2}{(y+h)^2} f^*(y+h) - f^*(y) \right|^p dy \right)^{\frac{1}{p}} \\ &\geq 2^{1-\frac{1}{p}} \frac{(2n-1)^2 h^2}{1+4n^2 h^2}. \end{aligned}$$

The theorem is proved. □

3.

Now we show that the above mentioned problem has a positive solution in the subset $L_{p,w}^k(\text{loc})$. First we give the following simple proposition.

Lemma. *Let the sequence of operators $A_n \in (L_{p,w}(\text{loc}) \rightarrow L_{p,w}(\text{loc}))^+$ satisfy the three conditions:*

$$\lim_{n \rightarrow \infty} \|A_n(t^m; x) - x^m\|_{p,w} = 0 \quad (m = 0, 1, 2).$$

Then, for any continuous and bounded function f on the real axis,

$$\lim_{n \rightarrow \infty} \|A_n f - f; L_{p,w}(a, b)\| = 0$$

holds, where a and b are any real numbers.

The proof of this Lemma is conducted in the same way as in the case of the space $C(a, b)$.

Since f is uniformly continuous function on any closed interval, given $\varepsilon > 0$ there exists a positive $\delta = \delta(\varepsilon)$ such that

$$|f(t) - f(x)| < \varepsilon \quad \text{if } |t - x| < \delta, \quad x \in [a, b], \quad t \in \mathbb{R}.$$

Also, setting $M = \sup_{x \in \mathbb{R}} |f(x)|$, we can write

$$|f(t) - f(x)| < 2M \quad \text{if } |t - x| \geq \delta \quad x \in [a, b], \quad t \in \mathbb{R}.$$

Therefore, from the basic inequality

$$|f(t) - f(x)| < \varepsilon + \frac{2M}{\delta^2} (t - x)^2,$$

where $-\infty < t < \infty$, $x \in [a, b]$, it follows that

$$\begin{aligned} \|A_n f - f; L_{p,w}(a, b)\| &\leq \varepsilon + M \|A_n(1; x) - 1; L_{p,w}(a, b)\| \\ &\quad + \frac{2M}{\delta^2} \|A_n((t-x)^2; x); L_{p,w}(a, b)\| \end{aligned}$$

and the last two terms tend to zero as $n \rightarrow \infty$ by the conditions of the Lemma.

Theorem 2. *If $A_n \in (L_{p,w}(\text{loc}) \rightarrow L_{p,w}(\text{loc}))^+$ is a sequence of operators satisfying the conditions (6) and (7), then*

$$\lim_{n \rightarrow \infty} \|A_n f - f\|_{p,w} = 0$$

for each function $f \in L_{p,w}^k(\text{loc})$.

Proof. Since $f \in L_{p,w}^k(\text{loc})$ implies that $f - k_f, w \in L_{p,w}^0(\text{loc})$, it is sufficient to prove the theorem for the function $f \in L_{p,w}^0(\text{loc})$. For $\varepsilon > 0$, there exists a point x_0 such that the inequality

$$(8) \quad \left(\frac{1}{2h} \int_{x-h}^{x+h} |f(t)|^p dt \right)^{\frac{1}{p}} < \varepsilon w(x)$$

holds for all x , $|x| \geq x_0$.

By the well known Luzin Theorem (see, for example [7]), there exists a continuous function φ on the finite interval $[-x_0 - h, x_0 + h]$ such that the inequality

$$(9) \quad \|f - \varphi; L_p(-x_0, x_0)\| < \varepsilon$$

is fulfilled. Setting

$$(10) \quad \delta < \min \left\{ \frac{2h\varepsilon^p}{M^p(x_0)}, h \right\},$$

where $M(x_0) = \max \left\{ \max_{|x| \leq x_0+h} |\varphi(x)|, 1 \right\}$, we can construct a continuous function g by the formulas

$$g(x) = \begin{cases} \varphi(x) & \text{if } |x| \leq x_0 + h, \\ 0 & \text{if } |x| \geq x_0 + h + \delta, \\ \text{linear} & \text{otherwise.} \end{cases}$$

Then, by (8), (9), (10) and the Minkowski inequality, we obtain

$$\begin{aligned}
\|f - g\|_{p,w} &\leq \|f - g; L_{p,w}(-x_0, x_0)\| + \|f - g; L_{p,w}(|x| \geq x_0 + h + \delta)\| \\
&\quad + \|f - g; L_{p,w}(x_0, x_0 + h + \delta)\| \\
&\quad + \|f - g; L_{p,w}(-x_0 - h - \delta, -x_0)\| \\
&< 2\varepsilon + \|f - g; L_p(x_0 - h, x_0 + h)\| + \frac{1}{w(x_0)} \|f; L_p(x_0 + h, x_0 + 2h + \delta)\| \\
&\quad + \|g; L_p(x_0 + h, x_0 + 2h + \delta)\| + \frac{1}{w(x_0)} \|f; L_p(-x_0 - 2h - \delta, -x_0 - h)\| \\
&\quad + \|g; L_p(-x_0 - h - \delta, -x_0 - h)\| \\
&\quad + \|f; L_p(-x_0 - h, -x_0 + h)\| \\
&< 4\varepsilon + 2M(x_0) \left(\frac{\delta}{2h}\right)^{\frac{1}{p}} + \frac{1}{w(x_0)} \|f; L_p(x_0 + h, x_0 + 3h)\| \\
&\quad + \frac{1}{w(x_0)} \|f; L_p(-x_0 - 3h, -x_0 - h)\|
\end{aligned}$$

and, on using the inequality (1),

$$\|f - g\|_{p,w} \leq 6\varepsilon + 2\varepsilon \frac{w(x_0 + 2h)}{w(x_0)} < C_1\varepsilon,$$

where $C_1 = 6 + 2(1 + 2h)^2$, since $w(x) = 1 + x^2$.

Consequently, for each $f \in L_{p,w}^0(\text{loc})$, there exists a continuous and bounded function g such that

$$(11) \quad \|f - g\|_{p,w} < C_1\varepsilon$$

for any $\varepsilon > 0$.

Now we can find a point $x_1 > x_0$ such that

$$(12) \quad w(x_1) > \frac{M(x_0)}{\varepsilon} \quad \text{and} \quad g(x) = 0 \quad \text{for} \quad |x| > x_1,$$

where $M(x_0)$ is defined above. Then, by (6), (11) and the Lemma (cf., e.g., [9], pp. 28, 29),

$$\begin{aligned}
\|L_n f - f\|_{p,w} &\leq \|L_n(f - g)\|_{p,w} + \|L_n g - g\|_{p,w} + \|f - g\|_{p,w} \\
&\leq (C + 1)\|f - g\|_{p,w} + \|L_n g - g\|_{p,w} \\
&\leq (C + 1)\varepsilon + \|L_n g - g; L_{p,w}(-x_1, x_1)\| + \|L_n g; L_{p,w}(|x| \geq x_1)\| \\
&\leq (C + 2)\varepsilon + \|L_n g; L_{p,w}(|x| \geq x_1)\|.
\end{aligned}$$

Since $|g(x)| \leq M(x_0)$ for all $x \in \mathbb{R}$, we can write

$$\begin{aligned} \|L_n g; L_{p,w}(|x| \geq x_1)\| &\leq M(x_0) \|L_n 1; L_{p,w}(|x| \geq x_1)\| \\ &\leq M(x_0) \|L_n 1 - 1\|_{p,w} + M(x_0) \|1; L_{p,w}(|x| \geq x_1)\|. \end{aligned}$$

Therefore

$$\|L_n f - f\|_{p,w} \leq (C + 2)\varepsilon + M(x_0) \|L_n 1 - 1\|_{p,w} + \frac{M(x_0)}{w(x_1)}.$$

In view of (7) and (12), the proof is completed. \square

4.

Theorem 2 gives a way of approximating all functions in $L_p(\text{loc})$. Namely, we have the following result.

Theorem 3. *Let a sequence of linear operators $A_n \in (L_{p,w}(\text{loc}) \rightarrow L_{p,w}(\text{loc}))^+$ satisfy the conditions i) and ii). Then, for all functions $f \in L_{p,w}(\text{loc})$,*

$$\lim_{n \rightarrow \infty} \sup_{-\infty < x < \infty} \frac{\|A_n f - f; L_p(x-h, x+h)\|}{1 + |x|^{2-\varepsilon}} = 0,$$

where $\varepsilon > 0$ is any positive number (ε cannot be zero).

P r o o f. By the condition (6) for any fixed x_0

$$a_n = \sup_{|x| > x_0} \frac{\|A_n f - f; L_p(x-h, x+h)\|}{1 + x^2}$$

is bounded, if $f \in L_{p,w}(\text{loc})$.

Also by the Lemma for any fixed x_0 ,

$$b_n = \sup_{|x| \leq x_0} \frac{\|A_n f - f; L_p(x-h, x+h)\|}{1 + x^2}$$

tends to zero as $n \rightarrow \infty$. Indeed, by the Luzin Theorem there exists a continuous function φ_1 on the interval $[-x_0 - h, x_0 + h]$ such that the inequality

$$(13) \quad \|f - \varphi_1; L_p(-x_0 - h, x_0 + h)\| < \varepsilon$$

is fulfilled. Moreover, setting

$$\begin{aligned}\varphi_1(x) &= \varphi_1(-x_0 - h) & \text{if } x \leq x_0 - h, \\ \varphi_1(x) &= \varphi_1(x_0 + h) & \text{if } x \geq x_0 + h,\end{aligned}$$

we see that φ_1 is continuous and bounded function on the whole real axis, for which the Lemma holds. Now,

$$\begin{aligned}b_n &\leq \|A_n f - f; L_p(-x_0 - h, x_0 + h)\| \\ &\leq (1 + \|A_n\|) \|f - \varphi_1; L_p(-x_0 - h, x_0 + h)\| + \|A_n \varphi_1 - \varphi_1; L_p(-x_0 - h, x_0 + h)\|\end{aligned}$$

and $\lim_{n \rightarrow \infty} b_n = 0$ by (6), (13) and the Lemma.

All that remains to see is that

$$\sup_{-\infty < x < \infty} \frac{\|A_n f - f; L_p(x - h, x + h)\|}{1 + |x|^{2+\varepsilon}} \leq (1 + x_0^2) b_n + a_n \sup_{|x| \geq x_0} \frac{1 + x^2}{1 + |x|^{2+\varepsilon}}$$

and the proof is completed since the last statement concerning ε follows from Theorem 1.

Acknowledgment. We are very thankful to the referee for the attentive reading of the manuscript and for valuable comments and suggestions.

References

- [1] *H. Berens and A. DeVore*: Quantitative Korovkin theorems for positive linear operators in L_p space. *Trans. Amer. Math. Soc.* 245 (1978), 349–361.
- [2] *S. J. Bernau*: Theorems of Korovkin type for L^p spaces. *Pacific J. Math.* 53 (1974), 11–19.
- [3] *K. Donner*: Korovkin Theorems in L^p -spaces. *J. Funct. Anal.* 42 (1981), 12–28.
- [4] *V. K. Dzyadik*: On the approximation of functions by linear positive operators and singular integrals. *Mat. Sbornik* 70 (1966), 508–517. (In Russian.)
- [5] *A. D. Gadziev*: The convergence problem for a sequence of positive linear operators on unbounded sets, and Theorems analogous to that of P. P. Korovkin. *Dokl. Akad. Nauk SSSR* 218, no. 5; *English Translated Soviet Math. Dokl.* 15 (1974), no. 5.
- [6] *A. D. Gadziev*: On P. P. Korovkin type theorems. *Math. Zametki*, Vol. 20 (1976). (In Russian.)
- [7] *N. B. Haaser and J. A. Sullivan*: *Real Analysis*. Dover Publications, INC, New York, 1991.
- [8] *W. Kitto and D. E. Walbert*: Korovkin approximations in L^p -spaces. *Pacific J. Math.* 63 (1976), 153–167.
- [9] *A. Kufner, O. John and S. Fučík*: *Function Spaces*. Academia, Prague, 1977.
- [10] *M. W. Muller*: L_p -approximation by the method of integral Meyer-König and Zeller operators. *Studia Math.* 63 (1978), 81–88.

- [11] *J. J. Swetits and B. Wood*: Quantitative estimates for L_p -approximation with positive linear operators. *J. Approx. Theory* 38 (1983), 81–89.
- [12] *J. J. Swetits and B. Wood*: On degree of L_p -approximation with positive linear operators. *J. Approx. Theory* 87 (1996), 239–241.
- [13] *B. Wood*: Degree of L_p -approximation with certain positive convolution operators. *J. Approx. Theory* 23 (1978), 354–363.

Authors' addresses: A. D. Gadjiev, Institut of Mathematics and Mechanics, Academy of Sciences of The Azerbaijan Republic, Azerbaijan Academy of National Aviation, Baku, Azerbaijan; currently: Ankara University Faculty of Sciences, Department of Mathematics, 06100 Tandogan, Ankara, Turkey, e-mail: haciyev@science.ankara.edu.tr; R. O. Efendiyev, Institut of Mathematics and Mechanics, Academy of Sciences of the Azerbaijan Republic, Baku, Azerbaijan, e-mail: frteb@aaas.ab.az; E. İbikli, Ankara University, Faculty of Sciences, Department of Mathematics, 06100 Tandogan, Ankara, Turkey, e-mail: ibikli@science.ankara.edu.tr.