

Yin-Zhu Gao

LJ-spaces

Czechoslovak Mathematical Journal, Vol. 57 (2007), No. 4, 1223–1237

Persistent URL: <http://dml.cz/dmlcz/128235>

Terms of use:

© Institute of Mathematics AS CR, 2007

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

LJ-SPACES

YIN-ZHU GAO, Nanjing

(Received October 29, 2005)

Abstract. In this paper *LJ*-spaces are introduced and studied. They are a common generalization of Lindelöf spaces and *J*-spaces researched by E. Michael. A space X is called an *LJ*-space if, whenever $\{A, B\}$ is a closed cover of X with $A \cap B$ compact, then A or B is Lindelöf. Semi-strong *LJ*-spaces and strong *LJ*-spaces are also defined and investigated. It is demonstrated that the three spaces are different and have interesting properties and behaviors.

Keywords: *LJ*-spaces, Lindelöf, *J*-spaces, *L*-map, (countably) compact, perfect map, order topology, connected, topological linear spaces

MSC 2000: 54D20, 54D30, 54F05, 54F65

1. INTRODUCTION

The Jordan curve theorem is one of the classical theorems of mathematics; it says that if C is a simple closed curve in the plane \mathbb{R}^2 , then $\mathbb{R}^2 \setminus C$ has precisely two components W_1 and W_2 , of which C is the common boundary [M].

Generalizing these properties, E. Michael [3] introduced and studied the following *J*-spaces.

A space X is a *J*-space if, whenever $\{A, B\}$ is a closed cover of X with $A \cap B$ compact, then A or B is compact.

A compact space is a *J*-space, but a *J*-space need not be compact.

We wonder whether in the definition of the *J*-space, “ A or B is compact” is equivalent to “ A or B is Lindelöf”. If not, what properties would the following space have?

Definition 1. A space X is an *LJ*-space if, whenever $\{A, B\}$ is a closed cover of X with $A \cap B$ compact, then A or B is Lindelöf.

The project is supported by NSFC (No. 10571081).

Obviously, both the Lindelöf spaces and J -spaces are LJ -spaces. In this note, we show that the LJ -space is different from the J -space or the Lindelöf space. Related spaces—strong LJ -spaces and semi-strong LJ -spaces are also introduced and studied. That the three classes of spaces are different is demonstrated by examples; their characterizations and relationships are investigated. They have interesting properties and behavior.

Throughout the note, spaces are topological spaces which are Hausdorff. A space X is Lindelöf if every open cover of X has a countable subcover. All maps are continuous. A map $f: X \rightarrow Y$ is boundary-perfect ([3]) if f is closed and $\partial(f^{-1}(y))$ is compact for any $y \in Y$. For a subset A of the space X , we reserve ∂A and A° for the boundary and interior of A respectively, and the symbols \mathbb{R} and \mathbb{Z}^+ for the sets of all real numbers and all non-negative integers respectively. Further, $\mathbb{R}^+ = \{x \in \mathbb{R}: x \geq 0\}$ and $\mathbb{R}^- = \{x \in \mathbb{R}: x \leq 0\}$. The cardinality of a set A is denoted by $|A|$. As a space, every ordinal has the usual order topology unless specifically stated otherwise. Other terms and symbols will be found in [1].

2. DEFINITIONS AND IMPLICATIONS

The following two spaces are related to J -spaces. A space X is a *strong J -space* [3] if every compact $K \subset X$ is contained in a compact subset M of X such that $X \setminus M$ is connected. A space X is a *semi-strong J -space* [3] if every compact $K \subset X$ is contained in a compact subset M of X such that $M \cup C = X$ for some connected $C \subset X \setminus K$. In [3], it is shown that the following implications hold while the inverses are not true:

$$\text{compactness} \Rightarrow \text{strong } J \Rightarrow \text{semi-strong } J \Rightarrow J.$$

We are naturally interested in the properties introduced below.

Definition 2. A space X is a *strong LJ -space* if every compact $K \subset X$ is contained in a closed Lindelöf $L \subset X$ such that $X \setminus L$ is connected.

Definition 3. A space X is a *semi-strong LJ -space* if every compact $K \subset X$ is contained in a closed Lindelöf $L \subset X$ such that $L \cup C = X$ for some connected $C \subset X \setminus K$.

Clearly, Lindelöf spaces are strong LJ -spaces and LJ -spaces. So \mathbb{R}^+ , \mathbb{R}^- , \mathbb{R}^n ($n > 1$), the real line \mathbb{R} and the Sorgenfrey line S are strong LJ -spaces. In [3], it is shown that \mathbb{R}^+ , \mathbb{R}^- and \mathbb{R}^n ($n > 1$) are also strong J -spaces while \mathbb{R} is not a J -space.

Proposition 1. *The Sorgenfrey line S is not a J -space.*

Proof. The closed cover $\{(-\infty, 0], [0, \infty)\}$ of S satisfies that $(-\infty, 0] \cap [0, \infty) = \{0\}$ is compact, but neither $(-\infty, 0]$ nor $[0, \infty)$ is compact. \square

It was shown that every topological linear space $X \neq \mathbb{R}$ is a strong J -space (Proposition 2.6 of [3]). Since strong J -spaces are strong LJ -spaces and the real line \mathbb{R} is a strong LJ -space, we have

Proposition 2. *All topological linear spaces are strong LJ -spaces.*

The long line Z (see [8] and [3]) (that is, $Z = [0, \omega_1) \times [0, 1)$ with the order topology generated by the lexicographical order) is connected, non-compact, countably compact and locally compact.

Proposition 3.

- (1) *The long line Z is a strong J -space (so a strong LJ -space), but not a Lindelöf space.*
- (2) *The product $\{0, 1\} \times Z$ is not an LJ -space.*

Proof. Let $K \subset Z$ be compact. Then K is bounded and so there exists an $\alpha_0 \in [0, \omega_1)$ such that $K \subset [0, \alpha_0) \times [0, 1)$. Then $L = [0, \alpha_0) \times [0, 1)$ is compact and $Z \setminus L$ is connected. Thus Z is a strong J -space. Clearly, Z is not Lindelöf.

(2) Put $A = \{0\} \times Z$, $B = \{1\} \times Z$. Then the closed cover $\{A, B\}$ of $\{0, 1\} \times Z$ is the desired one. \square

Proposition 4.

- (1) *$[0, \omega_1)$ is a J -space (so an LJ -space), but not a semi-strong LJ -space. Moreover, any closed subspace of $[0, \omega_1)$ is a J -space.*
- (2) *The product $[0, \omega_1) \times [0, \omega_1)$ is not an LJ -space (so not a J -space).*

Proof. (1) Let $\{A, B\}$ be a closed cover of $[0, \omega_1)$ and let $A \cap B$ be compact. Then A or B is bounded in $[0, \omega_1)$. In fact, assume that both A and B are unbounded in $[0, \omega_1)$; then $A \cap B$ is unbounded, which contradicts the compactness of $A \cap B$. Without loss of generality, we assume that A is bounded in $[0, \omega_1)$, then there exists a $\beta \in [0, \omega_1)$ such that $A \subset [0, \beta]$. Thus A is compact since $[0, \beta]$ is compact. So $[0, \omega_1)$ is a J -space.

Let us note that if $A \subset [0, \omega_1)$ with $|A| \geq 2$, then A is not connected. For the compact $K = \{0\} \subset [0, \omega_1)$, if $L \supset K$ is closed, Lindelöf, and $C \subset ([0, \omega_1) \setminus K)$ is connected, then $L \cup C \neq [0, \omega_1)$, so the LJ -space $[0, \omega_1)$ is not a semi-strong LJ -space.

Let F be a closed subspace of $[0, \omega_1)$. If F is compact, then it is a J -space. If F is not compact, then F is also a J -space since F and $[0, \omega_1)$ are homeomorphic.

(2) Put $A = \{0\} \times [0, \omega_1)$, $B = [1, \omega_1) \times [0, \omega_1)$, then $\{A, B\}$ is a closed cover of $[0, \omega_1) \times [0, \omega_1)$ with $A \cap B$ compact, however, neither A nor B is Lindelöf. \square

In Example 5 we present an ω_1 -broom space $Y(\omega_1)$ and show that it is a semi-strong LJ -space and has interesting properties, but it is not a strong LJ -space.

Theorem 1. *Let X be a space and let us consider the following assertions:*

- (A) X is a strong LJ -space; (a) X is a strong J -space;
 (B) X is a semi-strong LJ -space; (b) X is a semi-strong J -space;
 (C) X is an LJ -space; (c) X is a J -space.

Then $(A) \Rightarrow (B) \Rightarrow (C)$, $(a) \Rightarrow (A)$, $(b) \Rightarrow (B)$, $(c) \Rightarrow (C)$ and the implications are not reversible.

Proof. $(A) \Rightarrow (B)$ is clear. To show $(B) \Rightarrow (C)$, let $\{A, B\}$ be a closed cover of X with $A \cap B$ compact. Then there is a closed Lindelöf $L \subset X$ and a connected $C \subset X \setminus (A \cap B)$ such that $A \cap B \subset L$ and $L \cup C = X$. Since $\{A \cap C, B \cap C\}$ is a disjoint closed cover of the connected set C , so $A \cap C = \emptyset$ or $B \cap C = \emptyset$. Thus $A \subset X \setminus C \subset L$ or $B \subset X \setminus C \subset L$ is Lindelöf. So X is an LJ -space.

The other implications are obvious.

The Sorgenfrey line S satisfies the conditions (A), (B) and (C), but by Proposition 1, it does not satisfy (c), (b) or (a). $(C) \not\Rightarrow (B)$ follows by Proposition 4 (1); $(B) \not\Rightarrow (A)$ follows by Example 5; $(A) \not\Rightarrow$ Lindelöf follows by Proposition 3 (1). \square

3. INTERNAL CHARACTERIZATIONS

Proposition 5. *The following conditions are equivalent for a space X .*

- (1) X is a strong LJ -space (or a strong J -space).
- (2) If \mathscr{W} is a disjoint open cover of $X \setminus K$ with K compact, then there is a $W \in \mathscr{W}$ and a connected open $C \subset W$ such that $X \setminus C$ is Lindelöf (compact, respectively).
- (3) Same as (2), but with $|\mathscr{W}| = 2$.

Proof. (1) \Rightarrow (2). By (1), X has a connected open $C \subset X \setminus K$ with $X \setminus C$ Lindelöf (compact). So $C \subset \cup \mathscr{W}$. Since C is connected and \mathscr{W} is disjoint and open, we have a $W \in \mathscr{W}$ such that $C \subset W$. (2) \Rightarrow (3) and (3) \Rightarrow (1) are obvious. \square

Proposition 6. *The following conditions are equivalent for a space X .*

- (1) X is a semi-strong LJ -space (a semi-strong J -space).
- (2) If \mathscr{W} is a disjoint open cover of $X \setminus K$ with K compact, then there is a $W \in \mathscr{W}$ and a connected $C \subset W$ such that $\overline{X \setminus C}$ is Lindelöf (compact).
- (3) Same as (2), but with $|\mathscr{W}| = 2$.

Proof. (1) \Rightarrow (2). By (1), X has a connected $C \subset X \setminus K$ and a closed Lindelöf (a compact) $L \supset K$ with $C \cup L = X$. So $C \subset W$ for a $W \in \mathscr{W}$ since $C \subset \cup \mathscr{W}$ and $\overline{X \setminus C} \subset L$ is Lindelöf (compact). (2) \Rightarrow (3) and (3) \Rightarrow (1) are obvious. \square

Lemma 1. *If B is a closed non-Lindelöf subset of X and $C \subset B$ is Lindelöf, then there is a closed non-Lindelöf $D \subset B$ with $D \cap C = \emptyset$.*

Proof. Let \mathscr{U} be an open cover of B with no countable subcover. Pick a countable $\mathscr{F} \subset \mathscr{U}$ covering C . Then $D = B \setminus \bigcup \mathscr{F}$ has the required properties. \square

Theorem 2. *The following conditions are equivalent for a space X .*

- (1) X is an LJ -space.
- (2) For any $A \subset X$ with compact ∂A , \overline{A} or $\overline{X - A}$ is Lindelöf.
- (3) If A and B are disjoint closed subsets of X with ∂A or ∂B compact, then A or B is Lindelöf.
- (4) If \mathscr{W} is a disjoint open cover of $X \setminus K$ with K compact, then $X \setminus W$ is Lindelöf for some $W \in \mathscr{W}$.
- (5) Same as (4), but with $|\mathscr{W}| = 2$.

Proof. (1) \Rightarrow (2) is clear since $\partial A = \overline{A} \cap \overline{X - A}$ and $\{\overline{A}, \overline{X - A}\}$ covers X .

(2) \Rightarrow (3). Let A, B be disjoint closed subsets of X and let ∂A be compact, then A or $\overline{X \setminus A}$ is Lindelöf by (2). Since $B \subset \overline{X \setminus A}$, A or B is Lindelöf.

(3) \Rightarrow (1). Let $\{A, B\}$ be a closed cover of X with $A \cap B$ compact. Suppose B is not Lindelöf. By Lemma 1 there is a closed non-Lindelöf $D \subset B$ with $D \cap (A \cap B) = \emptyset$. Thus A and D are disjoint closed subsets of X and $\partial A \subset A \cap B$ is compact. So A or D is Lindelöf. Since D is not Lindelöf, A must be Lindelöf.

(1) \Leftrightarrow (5) and (4) \Rightarrow (5) are obvious.

(5) \Rightarrow (4). Assume (5). If for some $W_0 \in \mathscr{W}$, $W_0 \cup K$ is not Lindelöf, that is, $\{W_0, W^*\}$, where $W^* = \cup \{W \in \mathscr{W} : W \neq W_0\}$, has W^* such that $X \setminus W^* = W_0 \cup K$ is not Lindelöf, so by (5), $X \setminus W_0$ is Lindelöf. If for any $W \in \mathscr{W}$, $W \cup K$ is Lindelöf, then $\overline{W} \subset W \cup K$ is Lindelöf and X is Lindelöf. To show this, for any open cover \mathscr{U} of X take a finite $\mathscr{F} \subset \mathscr{U}$ covering K . Put $U = \bigcup \mathscr{F}$. It is enough to show that $\mathscr{W}' = \{W \in \mathscr{W} : W \not\subset U\}$ is countable. Suppose not. Then $\mathscr{W}' = \mathscr{W}_1 \cup \mathscr{W}_2$ with $\mathscr{W}_1 \cap \mathscr{W}_2 = \emptyset$, $\mathscr{W}_1 \cap \mathscr{W}'$ and $\mathscr{W}_2 \cap \mathscr{W}'$ both uncountable. Let $V_i = \bigcup \mathscr{W}_i$ ($i = 1, 2$). Then $\{V_1, V_2\}$ is a disjoint open cover of $X \setminus K$. By (5), $X \setminus V_1$ or $X \setminus V_2$ is Lindelöf.

Let $X \setminus V_2$ be Lindelöf. Then $\overline{V_1} \subset V_1 \cup K = X \setminus V_2$ is Lindelöf and so $C = \overline{V_1} \setminus U$ is also Lindelöf. Put $\mathscr{W}'_1 = \mathscr{W}_1 \cap \mathscr{W}'$. Then \mathscr{W}'_1 covers C and each $W \in \mathscr{W}'_1$ intersects C . This is a contradiction since C is Lindelöf and \mathscr{W}'_1 is uncountable and disjoint. So for any $W \in \mathscr{W}$, $X \setminus W$ is Lindelöf. \square

Theorem 3. *Let $\{X_1, X_2\}$ be a closed cover of X with $X_1 \cap X_2$ compact, then the following conditions are equivalent.*

- (1) X is an (resp. a semi-strong, a strong) LJ -space.
- (2) One of X_1 and X_2 is Lindelöf and the other is an (resp. a semi-strong, a strong, respectively) LJ -space.

Proof. (a) *For the LJ -space.* (1) \Rightarrow (2). By (1), X_1 or X_2 is Lindelöf. Let X_2 be Lindelöf. Let $\{A, B\}$ be a closed cover of X_1 with $A \cap B$ compact. Then X has a closed cover $\{A, B \cup X_2\}$ with $A \cap (B \cup X_2)$ compact. Hence A or $B \cup X_2$ is Lindelöf and so A or B is Lindelöf. (2) \Rightarrow (1). Let X_2 be Lindelöf, X_1 an LJ -space and $\{A, B\}$ a closed cover of X with $A \cap B$ compact. Put $A_i = A \cap X_i$ and $B_i = B \cap X_i$ ($i = 1, 2$). Then $\{A_1, B_1\}$ is a closed cover of X_1 with $A_1 \cap B_1$ compact. So A_1 or B_1 is Lindelöf. Let B_1 be Lindelöf. Then $B = B_1 \cup B_2$ is also Lindelöf.

(b) *For the semi-strong LJ -space.* (1) \Rightarrow (2). By (1) and Theorem 1 ((B) \Rightarrow (C)), let X_2 be Lindelöf and $K_1 \subset X_1$ compact. Then $K = K_1 \cup (X_1 \cap X_2)$ is compact. So $K \subset L$ for a closed Lindelöf $L \subset X$ and $L \cup C = X$ for a connected $C \subset X \setminus K$. Let $L_1 = L \cap X_1$. Put $M_i = C \cap X_i$, $i = 1, 2$, then $C = M_1 \cup M_2$. So $M_1 = \emptyset$ or $M_2 = \emptyset$ since C is connected. If $M_2 = \emptyset$, then $C \cup L_1 = X_1$ with $C \subset X_1$ and the Lindelöf $L_1 \supset K_1$. If $M_1 = \emptyset$, then the Lindelöf $X_1 = L_1$ is a semi-strong LJ -space. (2) \Rightarrow (1). Let X_2 be Lindelöf, $K \subset X$ be compact and $K_1 = K \cap X_1$. Then X_1 has a closed Lindelöf $L_1 \supset K_1$ and a connected $C \subset X_1 \setminus K_1$ such that $L_1 \cup C = X_1$. Put $L = L_1 \cup X_2$, then is closed Lindelöf, $L \supset K$, $L \cup C = X$ and $C \subset X \setminus K$.

(c) *For the strong LJ -space.* (1) \Rightarrow (2). By (1), let X_2 be Lindelöf. Let $K_1 \subset X_1$ be compact. Then $K = K_1 \cup (X_1 \cap X_2)$ is compact, so $K \subset L$ for a closed Lindelöf $L \subset X$ with $X \setminus L$ connected. Put $L_1 = L \cap X_1$, $M_i = (X \setminus L) \cap X_i$, $i = 1, 2$, then $X \setminus L = M_1 \cup M_2$. So $M_1 = \emptyset$ or $M_2 = \emptyset$. If $M_2 = \emptyset$, then $X_1 \setminus L_1 = X \setminus L$ is connected with $L_1 \subset X_1$ Lindelöf and $L_1 \supset K_1$. If $M_1 = \emptyset$, then the Lindelöf $X_1 = L_1$ is a strong LJ -space. (2) \Rightarrow (1). Let X_2 be Lindelöf and X_1 a strong LJ -space. Let $K \subset X$ be compact. Then $K_1 = (K \cup X_2) \cap X_1$ is compact, so $K_1 \subset L_1$ for a closed Lindelöf $L_1 \subset X_1$ with $X_1 \setminus L_1$ connected. Put $L = L_1 \cup X_2$, then $L \supset K$ is Lindelöf and $X \setminus L = X_1 \setminus L_1$ is connected. \square

Corollary 1. *Let $A \subset X$ be closed with ∂A compact. Then if X is an (a semi-strong, a strong) LJ -space, so is A .*

Proof. Put $X_1 = A$, $X_2 = \overline{X \setminus A}$. Then the conclusion follows from Theorem 3. \square

Corollary 2. Let $\{X_1, X_2\}$ be a closed cover of X with X_2 Lindelöf. Then

- (1) if X_1 is an (a semi-strong) LJ -space, so is X .
- (2) if X_1 is a strong LJ -space with $\partial(X_1)$ compact, so is X .

Proof. (1) See Theorem 3 (case (a), (2) \Rightarrow (1) and case (b), (2) \Rightarrow (1)).

(2) Let $K \subset X$ be compact and $K_1 = (K \cap X_1) \cup \partial(X_1)$. Then X_1 has a closed Lindelöf $L_1 \supset K_1$ with $X_1 \setminus L_1$ connected. Put $B = X_2 \setminus X_1^\circ$ and $L = L_1 \cup B$, then the closed Lindelöf $L \supset K$ and $X \setminus L = X_1 \setminus L_1$ is connected. \square

Corollary 3. Let $X = E \cup U$ with U open in X and \overline{U} compact. Then if E is an (a semi-strong, a strong) LJ -space, so is X .

Proof. The closed $A = X \setminus U \subset E$ has a compact boundary in X and thus in E , so A is an LJ -space by Corollary 1 since E is an LJ -space. X has a closed cover $\{A, \overline{U}\}$ with \overline{U} compact, so by Theorem 3, X is an LJ -space. The proofs of the other cases are similar. \square

Remark 1. (1) (a) If $\{X_1, X_2\}$ is a closed cover of X with $X_1 \cap X_2$ compact, then X is a semi-strong J -space iff one of X_1 and X_2 is compact and the other is a semi-strong J -space (since semi-strong $J \Rightarrow J$, the proof is similar to Theorem 3 (b)). (b) Corollaries 1 and 3 are also true for a semi-strong J -space (this follows from (a)).

(2) In Theorem 3 and Corollary 2, the “Lindelöf” cannot be removed. In fact, the long line Z is a strong J -space, but not a Lindelöf one (see Proposition 3), but the topological sum $Z \oplus Z$ is not an LJ -space.

(3) In Corollary 1, the “ ∂A compact” cannot be omitted (see Theorem 6(2)).

Proposition 7. Let E be a component of X . If X is a (semi-)strong LJ -space, so is E . Moreover, if a closed subset A is a union of components of X , so is A .

Proof. Let $K \subset E$ be compact, then X has a closed Lindelöf $L \supset K$ with $X \setminus L$ connected since X is a strong LJ -space. If $L \supset E$, then the Lindelöf E is a strong LJ -space. If $L \not\supset E$, then the connected set $X \setminus L$ intersects E and hence $X \setminus L \subset E$. So E has a closed Lindelöf $L' = L \cap E \supset K$ and $E \setminus L' = X \setminus L$ is connected. The proof for a semi-strong LJ -space is similar. \square

Theorem 4. Let $\{X_1, X_2\}$ be a closed cover of X with $X_1 \cap X_2$ non-Lindelöf. If X_1 and X_2 are (semi-strong) LJ -spaces, so is X .

Proof. To show that X is an LJ -space, let $\{A, B\}$ be a closed cover of X with $A \cap B$ compact. For $i = 1, 2$, let $A_i = A \cap X_i$ and $B_i = B \cap X_i$. Then $\{A_i, B_i\}$ is a closed cover of the LJ -space X_i with $A_i \cap B_i$ compact, so either A_i or B_i is Lindelöf. Note that $X_1 \cap X_2 = (A_1 \cup B_1) \cap (A_2 \cup B_2) \subset (A \cap B) \cup B_1 \cup A_2$. If B_1 is Lindelöf, A_2 cannot be Lindelöf since $A \cap B$ is compact while $X_1 \cap X_2$ is not Lindelöf. Hence B_2 is Lindelöf, so $B = B_1 \cup B_2$ is also Lindelöf. The case for A_1 being Lindelöf is similar.

To show that X is a semi-strong LJ -space, let $K \subset X$ be compact and $K_i = K \cap X_i$ for $i = 1, 2$. Then K_i is compact, and so there is a closed Lindelöf $L_i \supset K_i$ in X_i and connected $C_i \subset X_i \setminus K_i$ with $C_i \cup L_i = X_i$ for $i = 1, 2$. Let $L = L_1 \cup L_2$ and $C = C_1 \cup C_2$. Clearly $L \supset K$ is closed Lindelöf and $C \cup L = X$. Since $X_1 \cap X_2$ is non-Lindelöf, $(X_1 \cap X_2) \setminus L \neq \emptyset$. Also $X_i \setminus L \subset X_i \setminus L_i \subset C_i$ for $i = 1, 2$, so $(X_1 \cap X_2) \setminus L \subset (C_1 \cap C_2)$. Hence $C_1 \cap C_2 \neq \emptyset$ and thus C is connected. Clearly $C \subset X \setminus K$. \square

Remark 2. Theorem 4 is not true for strong LJ -spaces (see Example 5(2)) and is not reversible (in fact, the semi-strong LJ -space Y in Example 5 has a closed cover $\{Y, F\}$ with $Y \cap F = F$ non-Lindelöf. Y is a semi-strong LJ -space, but F is not an LJ -space since it is discrete and uncountable). In Theorem 4, the assumption that $X_1 \cap X_2$ is non-Lindelöf is also needed (see Remark 1 (1)).

4. EXTERNAL CHARACTERIZATIONS

To characterize the LJ -space, we introduce the notion of an L -map.

Definition 4. A map $f: X \rightarrow Y$ is an L -map if f is closed and $f^{-1}(y)$ is Lindelöf for any $y \in Y$.

Clearly, a perfect map is an L -map and is boundary-perfect (for the definition, see Introduction). A boundary-perfect map need not be an L -map (see the map g in Remark 6). Example 1 shows that an L -map need not be perfect or boundary-perfect.

Theorem 5. The following conditions are equivalent for a space X .

- (1) X is an LJ -space.
- (2) If a closed $f: X \rightarrow Y$ has $\partial(f^{-1}(y_0))$ compact and $f^{-1}(y_0)$ non-Lindelöf for a $y_0 \in Y$, then $f^{-1}(y)$ is Lindelöf for any $y \in Y \setminus \{y_0\}$.
- (3) Every boundary-perfect map $f: X \rightarrow Y$ onto a non-Lindelöf space Y is an L -map.

Proof. (1) \Rightarrow (2). For any $y \in Y \setminus \{y_0\}$, $A_0 = f^{-1}(y_0)$ and $A = f^{-1}(y)$ are disjoint closed subsets of X with $\partial(A_0)$ compact. Since $A_0 = f^{-1}(y_0)$ is not Lindelöf, by Theorem 2, $A = f^{-1}(y)$ is Lindelöf.

(2) \Rightarrow (1). Let A_1, A_2 be disjoint closed subsets of X with $\partial(A_1)$ or $\partial(A_2)$ compact. Suppose that $\partial(A_1)$ is compact. Let Y be the quotient space obtained from X by identifying A_i with a point y_i for $i = 1, 2$, and let $f: X \rightarrow Y$ be the quotient map. Clearly f is closed and $\partial(A_1) = \partial(f^{-1}(y_1))$ is compact. If $A_1 = f^{-1}(y_1)$ is not Lindelöf, then since $y_2 \in Y \setminus \{y_1\}$, by (2), $A_2 = f^{-1}(y_2)$ is Lindelöf. So by Theorem 2, X is an LJ -space.

(1) \Rightarrow (3). Let $f: X \rightarrow Y$ be as in the assumption and $y \in Y$. Since $\partial(f^{-1}(y))$ is compact, by Theorem 2, $f^{-1}(y)$ or $\overline{X - f^{-1}(y)}$ is Lindelöf. But $\overline{X - f^{-1}(y)}$ is not Lindelöf because Y is not Lindelöf, so $f^{-1}(y)$ is Lindelöf. Hence f is an L -map.

(3) \Rightarrow (1). Let $\{A, B\}$ be a closed cover of X with $A \cap B$ compact and let $Y = X/B$, let $f: X \rightarrow Y$ be the quotient map and $y_0 = f(B)$. Then f is closed, and if $y \in Y$, then $\partial(f^{-1}(y))$ is compact. So f is boundary-perfect. If Y is non-Lindelöf, then f is an L -map by the given condition, so $B = f^{-1}(y_0)$ is Lindelöf. If Y is Lindelöf, then the closed $f(A)$ is also Lindelöf. Then $f|_A: A \rightarrow f(A)$ is perfect. Hence $A = f|_A^{-1}(f(A))$ is Lindelöf. \square

Remark 3. Theorem 5 is false if the assumption that Y is non-Lindelöf is omitted. Indeed, $f: X \rightarrow Y$, where X is the non-Lindelöf LJ -space Z in Proposition 3 and Y is a singleton, is such an example.

Corollary 4. Every closed map $f: X \rightarrow Y$ from a paracompact LJ -space X onto a non-Lindelöf q -space Y is an L -map.

Proof. This follows from Theorem 5 and the result that every closed map $f: X \rightarrow Y$ from a paracompact space X onto a q -space Y is boundary-perfect (see [4]). \square

Remark 4. (1) Example 2 shows that the paracompactness of X in Corollary 4 cannot be omitted.

(2) In Corollary 4 the assumption that X is an LJ -space cannot be deleted. In fact, let \mathbb{R} be discrete, $X = \mathbb{R} \times \mathbb{R}$ and $Y = \mathbb{R}$. Let $f: X \rightarrow Y$ be the projection, then f is a closed map, but not an L -map.

Proposition 8. Let $f: X \rightarrow Y$ be a perfect map onto Y . Then

- (1) if X is an (a semi-strong) LJ -space, so is Y .
- (2) when f is open, if X is a strong LJ -space, so is Y .

Proof. (1) is obvious since the inverse image of a compact set is compact for a perfect map.

(2) Let $K \subset Y$ be compact. Then X has a closed Lindelöf $L' \supset f^{-1}(K)$ with $X \setminus L'$ connected. Put $L = Y \setminus f(X \setminus L')$, then $L \supset K$ and $Y \setminus L = f(X \setminus L')$ is connected. Since $f^{-1}(L) \subset L'$, $f^{-1}(L)$ is Lindelöf and thus L is also Lindelöf. \square

Remark 5. In Proposition 8 (2) the “open” cannot be omitted (see Example 5 (4)).

Recall that a continuous map $f: X \rightarrow Y$ is monotone if all fibers $f^{-1}(y)$ are connected.

Proposition 9. *Let $f: X \rightarrow Y$ be a monotone L -map onto Y . Then, if Y is an (a semi-strong, a strong) LJ -space, so is X .*

Proof. (a) Let $\{A, B\}$ be a closed cover of X with $A \cap B$ compact. Then $\{f(A), f(B)\}$ is a closed cover of Y . By Lemma 5.5 of [3], $f(A) \cap f(B) = f(A \cap B)$ is compact. So $f(A)$ or $f(B)$ is Lindelöf. Thus $f^{-1}(f(A))$ or $f^{-1}(f(B))$ is Lindelöf since f is an L -map. So A or B is Lindelöf and X is an LJ -space.

(b) Let $K \subset X$ be compact. Then Y has a closed Lindelöf $L' \supset f(K)$ and a connected $C' \subset Y \setminus f(K)$ with $C' \cup L' = Y$. Then $L = f^{-1}(L')$ is closed Lindelöf. Since f is closed and monotone, $C = f^{-1}(C')$ is connected by Theorem 6.1.29 of [1]. Clearly $L \supset K$, $C \subset X \setminus K$ and $L \cup C = X$. Thus X is a semi-strong LJ -space.

(c) Let $K \subset X$ be compact. Then Y has a closed Lindelöf $L \supset f(K)$ with $Y \setminus L$ connected. So $f^{-1}(L) \supset K$ is Lindelöf and $X \setminus f^{-1}(L) = f^{-1}(Y \setminus L)$ is connected since f is closed and monotone. So X is a strong LJ -space. \square

Remark 6. (1) Let $f: X \rightarrow Y$ be a monotone perfect map onto Y . Then, if Y is a semi-strong J -space, so is X (the proof is similar to the case (b) of Proposition 9).

(2) In Proposition 9 the “monotone” cannot be deleted: let Y be the long line Z which is a connected, non-Lindelöf, strong J -space and let $X = Y \oplus Y$. Then the obvious map $f: X \rightarrow Y$ is perfect, but clearly X is not an LJ -space. Also, the assumption in Proposition 9 that f is an L -map cannot be omitted or replaced by f being boundary-perfect. Indeed, if X is as above and E is a two-point space, then the obvious map $g: X \rightarrow E$ is boundary-perfect and monotone with each $f^{-1}(e)(e \in E)$ being a strong J -space, but X is not an LJ -space.

Proposition 10. *The following conditions are equivalent for a space Y .*

- (1) Y is an (a semi-strong, a strong) LJ -space.
- (2) $Y \times Z$ is an (a semi-strong, a strong) LJ -space for every connected and compact space Z .
- (3) $Y \times Z$ is an (a semi-strong, a strong) LJ -space for some connected and compact space Z .

Proof. (1) \Rightarrow (2) is by Proposition 9 with $X = Y \times Z$ and $f: X \rightarrow Y$ the projection. (2) \Rightarrow (3) is obvious. (3) \Rightarrow (1) is by Proposition 8 with $X = Y \times Z$ and $f: X \rightarrow Y$ the projection. \square

Proposition 11. *Each of the following conditions implies that $Y \times Z$ is an (a semi-strong, a strong) LJ-space.*

- (1) Y and Z are connected (semi-strong, strong) LJ-spaces.
- (2) Y is a connected, non-compact (semi-strong, strong) LJ-space and Z is connected.

Proof. (1) If Y or Z is compact, this follows from Proposition 10. If neither Y nor Z is compact, by Proposition 2.5 of [3], $Y \times Z$ is a strong J -space and it follows from Theorem 1.

(2) If Z is compact, this follows from Proposition 10. If Z is not compact, it follows from Propositions 2.5 of [3] and Theorem 1. \square

Remark 7. (1) Propositions 10 and 11 are true for semi-strong J -spaces (by Proposition 8.5 of [3] and Remark 6 (1)).

(2) In (1) and in Proposition 10 (2), (3) (Proposition 5.7(b), (c) of [3]), the connectedness cannot be omitted: by Proposition 3 (2), the long line Z is a strong J -space, but $Z \times \{0, 1\}$ is not an LJ-space.

5. RELATIONSHIPS

Recall a space X is called *hereditarily disconnected* if X does not contain any connected subsets of cardinality larger than one.

Theorem 6. *Let (A), (B), (C), (a), (b) and (c) be the same as in Theorem 1. Then*

- (1) *for a locally connected space X ; $(A) \Leftrightarrow (B) \Leftrightarrow (C)$.*
- (2) *none of the six properties is productive (additive, preserved by the quotient mapping, hereditary with respect to closed subspaces);*
- (3) *for a countably compact space X , $(A) \Leftrightarrow (a)$, $(B) \Leftrightarrow (b)$, $(C) \Leftrightarrow (c)$, $(D) \Leftrightarrow (d)$ and $(E) \Leftrightarrow (e)$;*
- (4) *for a hereditarily disconnected space X , “ X is Lindelöf” $\Leftrightarrow (A) \Leftrightarrow (B)$ and “ X is compact” $\Leftrightarrow (a) \Leftrightarrow (b)$.*

Proof. (1) $(A) \Rightarrow (B) \Rightarrow (C)$ follows by Theorem 1. $(C) \Rightarrow (A)$. Let $K \subset X$ be compact. Since X is locally connected, there is a disjoint open cover \mathscr{W} of $X \setminus K$

with each $W \in \mathscr{W}$ connected. By Theorem 2, there exists a $W_0 \in \mathscr{W}$ such that $L = X \setminus W_0$ is Lindelöf. Clearly $L \supset K$ and $X \setminus L$ is connected.

(2) Not productive: by Proposition 3 (2). Not additive: by Remark 1 (1). Not preserved by the quotient mapping: by Example 4.

Not hereditary with respect to closed subspaces.

For (a), (b) and (c): the strong J -space \mathbb{R}^+ has a closed discrete subspace \mathbb{Z}^+ which is not a J -space.

For (A): the long line Z is a strong LJ -space having a closed subspace $[0, \omega_1) \times \{0\}$ homeomorphic to $[0, \omega_1)$ which is not a strong LJ -space by Proposition 4.

For (B) and (C): in Example 5, the semi-strong LJ -space Y has a discrete closed subspace F which is uncountable, so F is not an LJ -space.

(3) is obvious since in a countably compact space Lindelöfness \Leftrightarrow compactness.

(4) Clearly, “ X is Lindelöf” \Rightarrow (A) \Rightarrow (B) and “ X is compact” \Rightarrow (a) \Rightarrow (b). To show that (B) \Rightarrow “ X is Lindelöf” ((b) \Rightarrow “ X is compact”), let $K \subset X$ be compact. By (B) ((b)) X has a closed Lindelöf (a compact) $L \supset K$ and a connected $C \subset X \setminus K$ such that $L \cup C = X$. Since X is hereditarily disconnected, $C = \emptyset$ or C is a one-point set. So X is Lindelöf (compact). \square

6. EXAMPLES

Example 1. An L -map which is not boundary-perfect (so not perfect).

Let $I_i = [o_i, 1_i]$ ($i \in \omega$) be the copy of the unit closed interval $I = [0, 1]$ and let $X = \bigoplus \{I_i : i \in \omega\}$ be the topological sum. Define an equivalence relation \mathscr{R} on X as follows: for each $x_i \in I_i$, if $x_i \neq o_i$, then $x_i \mathscr{R} x_i$; if $x_i = o_i$, then $o_i \mathscr{R} o_j$, $j \in \omega$. Then the natural map $f: X \rightarrow Y = X/\mathscr{R}$ is an L -map, but not a boundary-perfect map.

Example 2. A closed and open map $f: X \rightarrow Y$ from a locally compact strong J -space (so a strong LJ -space) X onto a non-Lindelöf q -space Y which is not an L -map.

P r o o f. Let Z be the long line. Then Z is non-Lindelöf and first countable (so a q -space). Let $X = Z \times Z$, $Y = Z$ and let $f: X \rightarrow Y$ be the projection onto the first coordinate. Then f is open. Let us show that f is also closed. Note that X is countably compact since Z is countably compact and first countable (see Theorem 3.10.36 of [1]). Let $F \subset X$ be closed, then F is countably compact and therefore $f(F)$ is countably compact in Z and thus closed in Z . Since Z is connected non-compact, $X = Z \times Z$ is a strong J -space by Proposition 2.5 of [3]. Clearly f is not an L -map. \square

Example 3. The Niemytzki plane X is not an LJ -space.

Proof. Let $A = [0, 1] \times [0, 1]$, $B = X \setminus (0, 1) \times [0, 1]$. Then $\{A, B\}$ is a closed cover with $A \cap B$ compact, but neither A nor B is Lindelöf. \square

Example 4. A strong J -space X whose quotient space Q is not an LJ -space.

Proof. Let Q be the Niemytzki plane. Put $X = Q \times \mathbb{R}$, where \mathbb{R} is the real line. By Proposition 2.5 of [3], the product space X is a strong J -space. Clearly Q is a quotient space and the projection $p: Q \times \mathbb{R} \rightarrow Q$ is the quotient map. \square

The following ω_1 -broom space $Y(\omega_1)$ is an interesting space. From Theorem 1, Theorem 6 (2), Remarks 2 and 5, we have seen that it plays an important role in this note.

Let Z be the long line and $X = Z \times \mathbb{R}^+$ with the product topology, where \mathbb{R}^+ is with the usual topology. For $\alpha \in [0, \omega_1)$ and integer $i \geq 1$, let $E_{\alpha,i}$ be the closed segment joining $\langle\langle \alpha, 0 \rangle, 0 \rangle$ to $\langle\langle \alpha + 1, 0 \rangle, \frac{1}{i} \rangle$, where $\langle \alpha, 0 \rangle$ and $\langle \alpha + 1, 0 \rangle$ are points of Z . Put

$$E_\alpha = \left(\bigcup_{i=1}^{\infty} E_{\alpha,i} \right) \cup (\langle\langle \alpha, 0 \rangle, \langle \alpha + 1, 0 \rangle \rangle \times \{0\}),$$

where $\langle\langle \alpha, 0 \rangle, \langle \alpha + 1, 0 \rangle \rangle$ is a closed interval of Z .

We define $Y(\omega_1) = \bigcup \{E_\alpha : \alpha \in [0, \omega_1)\}$ to be a subspace of X and call $Y(\omega_1)$ the ω_1 -broom space; we also write Y instead of $Y(\omega_1)$.

Example 5. The ω_1 -broom space Y is a semi-strong LJ -space such that

- (1) Y is not a strong LJ -space;
- (2) Y has a closed cover $\{A, B\}$ with $A \cap B$ non-Lindelöf and both A and B are strong LJ -spaces;
- (3) Y has a closed discrete subspace F which is uncountable;
- (4) there is a perfect map $f: M \rightarrow Y$ from a strong LJ -space M onto Y .

Proof. For any $\alpha \in [0, \omega_1)$, let $L_\alpha = \{\langle y_1, y_2 \rangle \in Y : y_1 \leq \langle \alpha, 0 \rangle\}$ and $C_\alpha = \overline{Y \setminus L_\alpha}$. Then L_α is Lindelöf, C_α is connected and $L_\alpha \cup C_\alpha = Y$. Now for any compact $K \subset Y$, pick α such that $K \subset L_\alpha$. Then $K \subset L_{\alpha+1}$, $C_{\alpha+1} \subset Y \setminus K$ and $L_{\alpha+1} \cup C_{\alpha+1} = Y$. So Y is a semi-strong LJ -space.

(1) Y is not a strong LJ -space. In fact, for the “beginning point” $\langle\langle 0, 0 \rangle, 0 \rangle$ of Y , let the compact subset H be the one-point set $\{\langle\langle 0, 0 \rangle, 0 \rangle\}$. If $L \subset Y$ is closed, Lindelöf and $H \subset L$, then we can see that $Y \setminus L$ is not connected.

(2) Put $A = (Z \times \{0\}) \cup \left(\bigcup \{E_\alpha : \alpha \in [0, \omega_1), \alpha \text{ is a successor ordinal}\} \right)$, $B = (Z \times \{0\}) \cup \left(\bigcup \{E_\alpha : \alpha \in [0, \omega_1), \alpha \text{ is a limit ordinal}\} \right)$.

Then $\{A, B\}$ is a closed cover of Y with $A \cap B = Z \times \{0\}$ non-Lindelöf.

Let us show that A is a strong LJ -space. For a limit ordinal α , put $L_\alpha^A = \{\langle z, y \rangle \in A : z \leq \langle \alpha, 0 \rangle\}$, then L_α^A is closed Lindelöf, $A \setminus L_\alpha^A$ is connected and each compact $K \subset A$ is a subset of some L_α^A .

Similarly, B is also a strong LJ -space.

(3) Put $F = \{\langle \alpha + 1, 0 \rangle, 1 \rangle : \alpha \in [0, \omega_1)\}$, then the uncountable F is a closed discrete subspace of Y .

(4) Let $M = B$ be a subspace of Y and $D = \{\alpha \in [0, \omega_1) : \alpha \text{ is a limit ordinal}\}$. Then D with the order topology is homeomorphic to $[0, \omega_1)$. So there exists an order preserving homeomorphic map $\varphi : D \rightarrow [0, \omega_1)$. For any $\alpha \in D$, let $f_\alpha : E_\alpha \rightarrow E_{\varphi(\alpha)}$ be a homeomorphic map. Now we define $f : M \rightarrow Y$ as follows.

For any $\langle z, y \rangle \in M$,

$$f(\langle z, y \rangle) = \begin{cases} f(\langle z, y \rangle) = f_\alpha(\langle z, y \rangle), & \langle z, y \rangle \in E_\alpha, \alpha \in D, \\ f(\langle z, 0 \rangle) = \langle \langle \alpha + 1, 0 \rangle, 0 \rangle, & \langle z, 0 \rangle \in [\langle \alpha + 1, 0 \rangle, \langle \alpha^+, 0 \rangle] \times \{0\}, \end{cases}$$

where α^+ is the smallest of the limit ordinals greater than α . Then f is a perfect map. □


Corollary 5.

- (1) *The ω_1 -broom space $Y(\omega_1)$ cannot be the image under an open perfect map of the long line Z .*
- (2) *Under CH, the Niemytzki plane cannot be the the image under an perfect map of the long line Z or the ω_1 -broom space $Y(\omega_1)$.*

Proof. (1) The long line Z is a strong LJ -space, thus by Proposition 8 (2), so is its open perfect image. But by Example 5, the ω_1 -broom space $Y(\omega_1)$ is not a strong LJ -space.

(2) The long line Z and the ω_1 -broom space $Y(\omega_1)$ are LJ -spaces by Proposition 8 (1), so their perfect images, but the Niemytzki plane is not an LJ -spaces. □

Now we illustrate the harmonious relationships with a diagram.



Acknowledgment. The author would like to thank the referee for the suggestion of using the present title of the paper instead of the former title “L-spaces” which was used before for denoting regular, hereditarily Lindelöf and nonseparable spaces.

References

- [1] *R. R. Engelking*: General Topology. Revised and completed edition, Heldermann Verlag, Berlin, 1989. zbl
- [2] *Y. Kodama and K. Nagami*: Theory of General Topology. Iwanami, Tokyo, 1974. (In Japanese.)
- [3] *E. Michael*: J -spaces. *Top. Appl.* 102 (2000), 315–339. zbl
- [4] *E. Michael*: A note on closed maps and compact sets. *Israel Math. J.* 2 (1964), 173–176. zbl
- [5] *E. Michael*: A survey of J -spaces. *Proceeding of the Ninth Prague Topological Symposium Contributed papers from the Symposium held in Prague Czech Republic, August 19–25, 2001*, pp. 191–193. zbl
- [6] *J. R. Munkres*: Topology. Prentice-Hall, Englewood Cliffs, NJ, 1975. zbl
- [7] *K. Nowinski*: Closed mappings and the Freudenthal compactification. *Fund. Math.* 76 (1972), 71–83. zbl
- [8] *L. A. Steen and J. A. Seebach, Jr*: Counterexamples in Topology. Springer-Verlag, New York, 1978. zbl

Author's address: Yin-Zhu Gao, Department of Mathematics, Nanjing University, Nanjing 210093, P.R. China, e-mail: yzgao@jssmail.com.cn.