

Sanjib Kumar Datta; Arindam Jha
Some results on the growth properties of Wronskians

Archivum Mathematicum, Vol. 45 (2009), No. 1, 59--69

Persistent URL: <http://dml.cz/dmlcz/128287>

Terms of use:

© Masaryk University, 2009

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

SOME RESULTS ON THE GROWTH PROPERTIES OF WRONSKIANs

SANJIB KUMAR DATTA AND ARINDAM JHA

ABSTRACT. The aim of this paper is to study the comparative growth properties of the composition of entire and meromorphic functions and wronskians generated by them improving some earlier results.

1. INTRODUCTION, DEFINITIONS AND NOTATIONS

We denote by \mathbb{C} the set of all finite complex numbers. Let f be a meromorphic function and g be an entire function defined on \mathbb{C} . We use the standard notations and definitions in the theory of meromorphic functions which are available in [3]. In the sequel we use the following notation:

$$\log^{[k]} x = \log \left(\log^{[k-1]} x \right) \text{ for } i = 1, 2, 3, \dots \quad \text{and} \quad \log^{[0]} x = x.$$

We recall the following definitions:

Definition 1. The order $\rho(f)$ and lower order $\mu(f)$ of a meromorphic function f are defined as

$$\rho(f) = \limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r} \quad \text{and} \quad \mu(f) = \liminf_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r}.$$

If f is entire, one can easily verify that,

$$\rho(f) = \limsup_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f)}{\log r} \quad \text{and} \quad \mu(f) = \liminf_{r \rightarrow \infty} \frac{\log^{[2]} M(r, f)}{\log r}.$$

Definition 2. The hyper order $\rho_2(f)$ and hyper lower order $\mu_2(f)$ of a meromorphic function f are defined as follows:

$$\rho_2(f) = \limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f)}{\log r} \quad \text{and} \quad \mu_2(f) = \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f)}{\log r}.$$

If f is entire, then

$$\rho_2(f) = \limsup_{r \rightarrow \infty} \frac{\log^{[3]} M(r, f)}{\log r} \quad \text{and} \quad \mu_2(f) = \liminf_{r \rightarrow \infty} \frac{\log^{[3]} M(r, f)}{\log r}.$$

2000 *Mathematics Subject Classification*: primary 30D35; secondary 30D30.

Key words and phrases: transcendental entire and meromorphic function, Wronskian, proximate lower order, composition, growth.

Received April 2, 2008, revised December 2008. Editor O. Došlý.

Definition 3. The type $\tau(f)$ of a meromorphic function f is defined as :

$$\tau(f) = \limsup_{r \rightarrow \infty} \frac{T(r, f)}{r^{\rho(f)}}, \quad 0 < \rho(f) < \infty.$$

When f is entire, then

$$\tau(f) = \limsup_{r \rightarrow \infty} \frac{\log M(r, f)}{r^{\rho(f)}}, \quad 0 < \rho(f) < \infty.$$

Definition 4. A function $\mu_f(r)$ is called a lower proximate order of a meromorphic function f of finite lower order $\mu(f)$ if

- (i) $\mu_f(r)$ is non negative and continuous for $r \geq r_0$, say,
- (ii) $\mu_f(r)$ is differentiable for $r \geq r_0$ except possibly at isolated points at which $\mu'_f(r+0)$ and $\mu'_f(r-0)$ exists,
- (iii) $\lim_{r \rightarrow \infty} \mu_f(r) = \mu(f)$,
- (iv) $\lim_{r \rightarrow \infty} r\mu'_f(r) \log r = 0$ and
- (v) $\liminf_{r \rightarrow \infty} \frac{T(r, f)}{r^{\mu_f(r)}} = 1$.

Definition 5. Let a be a complex number, finite or infinite. The Nevanlinna deficiency and Valiron deficiency of ' a ' with respect to a meromorphic function f is defined as

$$\delta(a; f) = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, a; f)}{T(r, f)} = \liminf_{r \rightarrow \infty} \frac{m(r, a; f)}{T(r, f)}$$

and

$$\Delta(a; f) = 1 - \liminf_{r \rightarrow \infty} \frac{N(r, a; f)}{T(r, f)} = \limsup_{r \rightarrow \infty} \frac{m(r, a; f)}{T(r, f)}.$$

From the second fundamental theorem it follows that the set of values of $a \in C \cup \{\infty\}$ for which $\delta(a; f) > 0$ is countable and $\sum_{a \neq \infty} \delta(a; f) + \delta(\infty; f) \leq 2$ (cf. [3, p. 43]). If in particular $\sum_{a \neq \infty} \delta(a; f) + \delta(\infty; f) = 2$, we say that f has the maximal deficiency sum.

Definition 6. A meromorphic function $a = a(z)$ is called small with respect to f if $T(r, a) = S(r, f)$.

Definition 7. Let a_1, a_2, \dots, a_k be linearly independent meromorphic functions and small with respect to f . We denote by $L(f) = W(a_1, a_2, \dots, a_k, f)$ the Wronskian determinant of a_1, a_2, \dots, a_k, f i.e.,

$$L(f) = \begin{vmatrix} a_1 & a_2 & \dots & a_k & f \\ a'_1 & a'_2 & \dots & a'_k & f' \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_1^{(k)} & a_2^{(k)} & \dots & a_k^{(k)} & f^{(k)} \end{vmatrix}.$$

Since the natural extension of a derivative is a differential polynomial, in this paper we prove our results for a special type of linear differential polynomials *viz.*, the Wronskians. In the paper we prove some new results depending on the comparative growth properties of composite entire or meromorphic functions and Wronskians generated by one of the factors which improve some earlier theorems.

2. LEMMAS

In this section we present some lemmas which will be needed in the sequel.

Lemma 1 ([1]). *If f is meromorphic and g is entire then for all sufficiently large values of r*

$$T(f \circ g) \leq \{1 + o(1)\} \frac{T(r, g)}{\log M(r, g)} T(M(r, g), f) .$$

Lemma 2 ([2]). *Let f be meromorphic and g be entire and suppose that $0 < \mu \leq \rho_g \leq \infty$. Then for a sequence of values of r tending to infinity,*

$$T(r, f \circ g) \geq T(\exp(r^\mu), f) .$$

Lemma 3 ([5]). *Let f be a transcendental meromorphic function having maximal deficiency sum. Then*

$$\lim_{r \rightarrow \infty} \frac{T(r, L(f))}{T(r, f)} = 1 + k - k\delta(\infty; f) .$$

Lemma 4. *If f be a transcendental meromorphic function with the maximal deficiency sum, then*

$$\rho(L(f)) = \rho(f)$$

and

$$\mu(L(f)) = \mu(f) .$$

Also

$$\tau(L(f)) = \{1 + k - k\delta(\infty; f)\} \tau(f) .$$

Proof. By Lemma 3, $\lim_{r \rightarrow \infty} \frac{\log T(r, L(f))}{\log T(r, f)}$ exists and equal to 1. So

$$\begin{aligned} \rho(L(f)) &= \limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r} \cdot \lim_{r \rightarrow \infty} \frac{\log T(r, L(f))}{\log T(r, f)} \\ &= \rho(f) \cdot 1 = \rho(f) . \end{aligned}$$

In a similar manner, $\mu(L(f)) = \mu(f)$.

Again

$$\begin{aligned} \tau(L(f)) &= \limsup_{r \rightarrow \infty} \frac{T(r, L(f))}{r^{\rho(L(f))}} \\ &= \limsup_{r \rightarrow \infty} \frac{T(r, f)}{r^{\rho(f)}} \cdot \lim_{r \rightarrow \infty} \frac{T(r, L(f))}{T(r, f)} \\ &= \{1 + k - k\delta(\infty; f)\} \tau(f) . \end{aligned}$$

This proves the lemma. \square

Lemma 5. *Let f be a transcendental meromorphic function with the maximal deficiency sum. Then the hyper order and (hyper lower order) of $L(f)$ and f are equal.*

The proof of Lemma 5 is omitted because it can be carried out in the line of Lemma 4.

Lemma 6. *For a meromorphic function f of finite lower order, lower proximate order exists.*

The lemma can be proved in the line of Theorem 1 [4] and so the proof is omitted. For further reference see also [8] and [9].

Lemma 7. *Let f be a meromorphic function of finite lower order $\mu(f)$. Then for $\delta (> 0)$ the function $r^{\mu(f)+\delta-\mu_f(r)}$ is ultimately an increasing function of r .*

Proof. Since

$$\frac{d}{dr} r^{\mu(f)+\delta-\mu_f(r)} = \{\mu(f) + \delta - \mu_f(r) - r\mu'_f(r) \log r\} \frac{r^{\mu(f)+\delta-\mu_f(r)}}{r} > 0$$

for all sufficiently large values of r , the lemma follows. \square

Lemma 8 ([6]). *Let f be an entire function of finite lower order. If there exists entire functions a_i ($i = 1, 2, \dots, n; n \leq \infty$) satisfying $T(r, a_i) = o\{T(r, f)\}$ and $\sum_{i=1}^n \delta(a_i; f) = 1$, then $\lim_{r \rightarrow \infty} \frac{T(r, f)}{\log M(r, f)} = \frac{1}{\pi}$.*

3. THEOREMS

In this section we present the main results of the paper.

Theorem 1. *Let f be a meromorphic function and g be a transcendental entire function with maximal deficiency sum. If $\mu(f)$ and $\mu(g)$ are both finite, then*

$$\liminf_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(g))} \leq \frac{3 \cdot \rho(f) \cdot 2^{\mu(g)}}{1 + k - k\delta(\infty; g)}.$$

Proof. If $\rho(f) = \infty$, the theorem is obvious. So we suppose that $\rho(f) < \infty$. Since $T(r, g) \leq \log^+ M(r, g)$, in view of Lemma 1, we get for all sufficiently large values of r ,

$$\begin{aligned} T(f \circ g) &\leq \{1 + o(1)\} T(M(r, g), f), \\ \text{i.e., } \log T(r, f \circ g) &\leq \log \{1 + o(1)\} + \log T(M(r, g), f) \\ &\leq o(1) + (\rho(f) + \varepsilon) \log M(r, g), \\ \text{i.e., } \liminf_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, g)} &\leq (\rho(f) + \varepsilon) \liminf_{r \rightarrow \infty} \frac{\log M(r, g)}{T(r, g)}. \end{aligned}$$

Since $\varepsilon (> 0)$ is arbitrary, it follows that

$$(1) \quad \liminf_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, g)} \leq \rho(f) \cdot \liminf_{r \rightarrow \infty} \frac{\log M(r, g)}{T(r, g)}.$$

As $\liminf_{r \rightarrow \infty} \frac{T(r, f)}{r^{\mu_f(r)}} = 1$, so for given ε ($0 < \varepsilon < 1$) we get for a sequence of values of r tending to infinity,

$$(2) \quad T(r, g) \leq (1 + \varepsilon) r^{\mu_g(r)}$$

and for all sufficiently large values of r ,

$$(3) \quad T(r, g) \geq (1 - \varepsilon) r^{\mu_g(r)} .$$

Since $\log M(r, g) \leq 3T(2r, g)$, we have by (2), for a sequence of values of r tending to infinity,

$$(4) \quad \log M(r, g) \leq 3T(2r, g) \leq 3(1 + \varepsilon) (2r)^{\mu_g(2r)} .$$

Combining (3) and (4) we obtain for a sequence of values of r tending to infinity,

$$\frac{\log M(r, g)}{T(r, g)} \leq \frac{3(1 + \varepsilon)}{(1 - \varepsilon)} \cdot \frac{(2r)^{\mu_g(2r)}}{r^{\mu_g(r)}} .$$

Now for any δ (> 0), for a sequence of values of r tending to infinity,

$$(5) \quad \begin{aligned} \frac{\log M(r, g)}{T(r, g)} &\leq \frac{3(1 + \varepsilon)}{(1 - \varepsilon)} \cdot \frac{(2r)^{\mu(g) + \delta}}{(2r)^{\mu(g) + \delta - \mu_g(2r)}} \cdot \frac{1}{r^{\mu_g(r)}} \\ \text{i.e., } \frac{\log M(r, g)}{T(r, g)} &\leq \frac{3(1 + \varepsilon)}{(1 - \varepsilon)} \cdot 2^{\mu(g) + \delta} \end{aligned}$$

because $r^{\mu(g) + \delta - \mu_g(2r)}$ is ultimately an increasing function of r by Lemma 7. Since ε (> 0) and δ (> 0) are arbitrary, it follows from (5) that

$$(6) \quad \liminf_{r \rightarrow \infty} \frac{\log M(r, g)}{T(r, g)} \leq 3 \cdot 2^{\mu(g)} .$$

Thus from (1) and (6) we obtain that

$$(7) \quad \liminf_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, g)} \leq 3 \cdot \rho(f) \cdot 2^{\mu(g)} .$$

Now in view of Lemma 3 and (7) we get

$$\begin{aligned} \liminf_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(g))} &= \liminf_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, g)} \cdot \lim_{r \rightarrow \infty} \frac{T(r, g)}{T(r, L(g))} \\ &\leq \frac{3 \cdot \rho(f) \cdot 2^{\mu(g)}}{1 + k - k\delta(\infty; g)} . \end{aligned}$$

This proves the theorem. □

Theorem 2. *Let f be a meromorphic function of finite order and g be transcendental entire of finite lower order with maximal deficiency sum. Then*

$$\liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f \circ g)}{\log T(r, L(g))} \leq 1 .$$

Proof. Since $T(r, g) \leq \log^+ M(r, g)$, in view of Lemma 1, we get for all sufficiently large values of r ,

$$\begin{aligned} \log T(r, f \circ g) &\leq \log \{1 + o(1)\} + \log T(M(r, g), f) \\ &\leq o(1) + (\rho(f) + \varepsilon) \log M(r, g) \end{aligned}$$

(8) i.e., $\log^{[2]} T(r, f \circ g) \leq \log^{[2]} M(r, g) + O(1)$.

It is well known that for any entire function g , $\log M(r, g) \leq 3T(2r, g)$ (cf. [3]). Then for $0 < \varepsilon < 1$ and $\delta (> 0)$, for a sequence of values of r tending to infinity it follows from (5) that

$$(9) \quad \log^{[2]} M(r, g) \leq \log T(r, g) + O(1).$$

Now combining (8) and (9) we obtain for a sequence of values of r tending to infinity,

$$\begin{aligned} \log^{[2]} T(r, f \circ g) &\leq \log T(r, g) + O(1) \\ \text{i.e., } \frac{\log^{[2]} T(r, f \circ g)}{\log T(r, g)} &\leq 1. \end{aligned}$$

(10)

As by Lemma 3, $\lim_{r \rightarrow \infty} \frac{\log T(r, g)}{\log T(r, L(g))}$ exists and is equal to 1, from (10) we obtain that

$$\begin{aligned} \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f \circ g)}{\log T(r, L(g))} &= \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f \circ g)}{\log T(r, g)} \cdot \lim_{r \rightarrow \infty} \frac{\log T(r, g)}{\log T(r, L(g))} \\ &\leq 1 \cdot 1 = 1. \end{aligned}$$

Thus the theorem is established. \square

Theorem 3. Let f and g be two transcendental entire functions each having maximal deficiency sum such that $\mu(f) > 0$, $\rho(g) < \mu(f) \leq \rho(f) < \infty$. Also let there exist entire functions a_i ($i = 1, 2, \dots, n; n \leq \infty$) with

- (i) $T(r, a_i) = o\{T(r, g)\}$ as $r \rightarrow \infty$ for $i = 1, 2, \dots, n$ and
(ii) $\sum_{i=1}^n \delta(a_i; g) = 1$. Then

$$\lim_{r \rightarrow \infty} \frac{\{\log T(r, f \circ g)\}^2}{T(r, L(f))T(r, L(g))} = 0.$$

Proof. In view of the inequality $T(r, g) \leq \log^+ M(r, g)$ and Lemma 1 we obtain for all sufficiently large values of r ,

$$\begin{aligned} T(f \circ g) &\leq \{1 + o(1)\} T(M(r, g), f) \\ \text{i.e., } \log T(r, f \circ g) &\leq \log \{1 + o(1)\} + \log T(M(r, g), f) \\ &\leq o(1) + (\rho(f) + \varepsilon) \log M(r, g) \\ &\leq o(1) + (\rho(f) + \varepsilon) r^{\rho(g) + \varepsilon}. \end{aligned}$$

(11)

Again in view of Lemma 4, we get for all sufficiently large values of r ,

$$\begin{aligned} \log T(r, L(f)) &> (\mu(L(f)) - \varepsilon) \log r \\ \text{i.e., } \log T(r, L(f)) &> (\mu(f) - \varepsilon) \log r \\ (12) \quad \text{i.e., } T(r, L(f)) &> r^{\mu(f) - \varepsilon}. \end{aligned}$$

Now combining (11) and (12) it follows for all sufficiently large values of r ,

$$(13) \quad \frac{\log T(r, f \circ g)}{T(r, L(f))} \leq \frac{o(1) + (\rho(f) + \varepsilon) r^{\rho(g) + \varepsilon}}{r^{\mu(f) - \varepsilon}}.$$

Since $\rho(g) < \mu(f)$, we can choose $\varepsilon (> 0)$ in such a way that

$$(14) \quad \rho(g) + \varepsilon < \mu(f) - \varepsilon.$$

So in view of (13) and (14) it follows that

$$(15) \quad \lim_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(f))} = 0.$$

Again from Lemma 3 and Lemma 8 we get for all sufficiently large values of r ,

$$\begin{aligned} \frac{\log T(r, f \circ g)}{T(r, L(g))} &\leq \frac{o(1) + (\rho(f) + \varepsilon) \log M(r, g)}{T(r, L(g))} \\ \text{i.e., } \limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(g))} &\leq (\rho(f) + \varepsilon) \limsup_{r \rightarrow \infty} \frac{\log M(r, g)}{T(r, L(g))} \\ \text{i.e., } \limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(g))} &\leq (\rho(f) + \varepsilon) \limsup_{r \rightarrow \infty} \frac{\log M(r, g)}{T(r, g)} \cdot \lim_{r \rightarrow \infty} \frac{T(r, g)}{T(r, L(g))} \\ \text{i.e., } \limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(g))} &\leq (\rho(f) + \varepsilon) \cdot \pi \cdot \frac{1}{1 + k - k\delta(\infty; g)}. \end{aligned}$$

Since $\varepsilon (> 0)$ is arbitrary it follows from above that

$$(16) \quad \limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(g))} \leq \rho(f) \cdot \pi \cdot \frac{1}{1 + k - k\delta(\infty; g)}.$$

In view of (15) i.e., $\lim_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(f))} = 0$ and using (16) we obtain that

$$\begin{aligned} \limsup_{r \rightarrow \infty} \frac{\{\log T(r, f \circ g)\}^2}{T(r, L(f)) T(r, L(g))} &= \lim_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(f))} \cdot \limsup_{r \rightarrow \infty} \frac{\log T(r, f \circ g)}{T(r, L(g))} \\ &\leq 0 \cdot \frac{\pi \rho(f)}{1 + k - k\delta(\infty; g)} = 0, \\ \text{i.e., } \lim_{r \rightarrow \infty} \frac{\{\log T(r, f \circ g)\}^2}{T(r, L(f)) T(r, L(g))} &= 0. \end{aligned}$$

This proves the theorem. \square

Theorem 4. *If f and g be two entire functions with f to be transcendental having maximal deficiency sum satisfying the following conditions:*

(i) $\mu(f) > 0$, (ii) $\rho_2(f) < \infty$ and (iii) $0 < \mu(g) \leq \rho(g)$. Then

$$\limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f \circ g)}{\log^{[2]} T(r, L(f))} \geq \max \left\{ \frac{\mu(g)}{\mu_2(f)}, \frac{\rho(g)}{\rho_2(f)} \right\}.$$

Proof. We know that for $r > 0$ (cf. [7]) and for all sufficiently large values of r ,

$$(17) \quad T(r, f \circ g) \geq \frac{1}{3} \log M \left\{ \frac{1}{8} M \left(\frac{r}{4}, g \right) + o(1), f \right\}.$$

Since $\mu(f)$ and $\mu(g)$ are the lower orders of f and g respectively then for given ε (> 0) and for large values of r we obtain that

$$\log M(r, f) > r^{\mu(f)-\varepsilon} \quad \text{and} \quad \log M(r, g) > r^{\mu(g)-\varepsilon}$$

where $0 < \varepsilon < \min \{ \mu(f), \mu(g) \}$. So from (17) we have for all sufficiently large values of r ,

$$\begin{aligned} T(r, f \circ g) &\geq \frac{1}{3} \left\{ \frac{1}{8} M \left(\frac{r}{4}, g \right) + o(1) \right\}^{\mu(f)-\varepsilon} \\ \text{i.e., } T(r, f \circ g) &\geq \frac{1}{3} \left\{ \frac{1}{9} M \left(\frac{r}{4}, g \right) \right\}^{\mu(f)-\varepsilon} \\ \text{i.e., } \log T(r, f \circ g) &\geq O(1) + (\mu(f) - \varepsilon) \log M \left(\frac{r}{4}, g \right) \\ \text{i.e., } \log T(r, f \circ g) &\geq O(1) + (\mu(f) - \varepsilon) \left(\frac{r}{4} \right)^{\mu(g)-\varepsilon} \\ (18) \quad \text{i.e., } \log^{[2]} T(r, f \circ g) &\geq O(1) + (\mu(g) - \varepsilon) \log r. \end{aligned}$$

Again in view of Lemma 5, we get for a sequence of values of r tending to infinity,

$$\begin{aligned} \log^{[2]} T(r, L(f)) &\leq (\mu_2(L(f)) + \varepsilon) \log r \\ (19) \quad \text{i.e., } \log^{[2]} T(r, L(f)) &\leq (\mu_2(f) + \varepsilon) \log r. \end{aligned}$$

Combining (18) and (19) it follows for a sequence of values of r tending to infinity

$$\frac{\log^{[2]} T(r, f \circ g)}{\log^{[2]} T(r, L(f))} \geq \frac{O(1) + (\mu(g) - \varepsilon) \log r}{(\mu_2(f) + \varepsilon) \log r}.$$

Since ε (> 0) is arbitrary, we obtain that

$$(20) \quad \limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f \circ g)}{\log^{[2]} T(r, L(f))} \geq \frac{\mu(g)}{\mu_2(f)}.$$

Again from (17) we get for a sequence of values of r tending to infinity,

$$\begin{aligned} \log T(r, f \circ g) &\geq O(1) + (\mu(f) - \varepsilon) \left(\frac{r}{4} \right)^{\rho(g)-\varepsilon} \\ (21) \quad \text{i.e., } \log^{[2]} T(r, f \circ g) &\geq O(1) + (\rho(g) - \varepsilon) \log r. \end{aligned}$$

Also in view of Lemma 5, for all sufficiently large values of r , we have

$$(22) \quad \log^{[2]} T(r, L(f)) \leq (\rho_2(L(f)) + \varepsilon) \log r = (\rho_2(f) + \varepsilon) \log r.$$

Now from (21) and (22) it follows for a sequence of values of r tending to infinity that

$$\frac{\log^{[2]} T(r, f \circ g)}{\log^{[2]} T(r, L(f))} \geq \frac{O(1) + (\rho(g) - \varepsilon) \log r}{(\rho_2(f) + \varepsilon) \log r}.$$

As ε ($0 < \varepsilon < \rho(g)$) is arbitrary, we obtain from above that

$$(23) \quad \limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f \circ g)}{\log^{[2]} T(r, L(f))} \geq \frac{\rho(g)}{\rho_2(f)}.$$

Therefore from (20) and (23) we get that

$$\limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, f \circ g)}{\log^{[2]} T(r, L(f))} \geq \max \left\{ \frac{\mu(g)}{\mu_2(f)}, \frac{\rho(g)}{\rho_2(f)} \right\}.$$

Thus the theorem is established. \square

Theorem 5. *Let f be transcendental meromorphic with maximal deficiency sum and g be entire such that (i) $0 < \mu_2(f) < \rho_2(f)$, (ii) $\rho(g) < \infty$. If h be transcendental meromorphic function with finite order then*

$$\liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))} \leq \min \left\{ \frac{\mu(g)}{\mu_2(f)}, \frac{\rho(g)}{\rho_2(f)} \right\}.$$

Proof. In view of Lemma 1 and the inequality $T(r, g) \leq \log^+ M(r, g)$, we obtain for all sufficiently large values of r

$$(24) \quad \log T(r, h \circ g) \leq o(1) + (\rho(h) + \varepsilon) \log M(r, g).$$

Also for a sequence of values of r tending to infinity,

$$(25) \quad \log M(r, g) \leq r^{\mu(g)+\varepsilon}.$$

Combining (24) and (25) it follows for a sequence of values of r tending to infinity,

$$(26) \quad \begin{aligned} \log T(r, h \circ g) &\leq o(1) + (\rho(h) + \varepsilon) r^{\mu(g)+\varepsilon} \\ \text{i.e., } \log T(r, h \circ g) &\leq \{(\rho(h) + \varepsilon) + o(1)\} r^{\mu(g)+\varepsilon} \\ \text{i.e., } \log^{[2]} T(r, h \circ g) &\leq O(1) + (\mu(g) + \varepsilon) \log r. \end{aligned}$$

Again in view of Lemma 5, we have for all sufficiently large values of r ,

$$(27) \quad \log^{[2]} T(r, L(f)) > (\mu_2(L(f)) - \varepsilon) \log r = (\mu_2(f) - \varepsilon) \log r.$$

Now from (26) and (27) we get for a sequence of values of r tending to infinity,

$$\frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))} \leq \frac{O(1) + (\mu(g) + \varepsilon) \log r}{(\mu_2(f) - \varepsilon) \log r}.$$

As ε (> 0) is arbitrary, it follows that

$$(28) \quad \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))} \leq \frac{\mu(g)}{\mu_2(f)}.$$

In view of Lemma 1 we get for all sufficiently large values of r ,

$$(29) \quad \log^{[2]} T(r, h \circ g) \leq O(1) + (\rho(g) + \varepsilon) \log r.$$

Also for a sequence of values of r tending to infinity,

$$(30) \quad \log^{[2]} T(r, L(f)) > (\rho_2(L(f)) - \varepsilon) \log r = (\rho_2(f) - \varepsilon) \log r.$$

Combining (29) and (30) we have for a sequence of values of r tending to infinity,

$$\frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))} \leq \frac{O(1) + (\rho(g) + \varepsilon) \log r}{(\rho_2(f) - \varepsilon) \log r}.$$

Since $\varepsilon (> 0)$ is arbitrary, it follows from above that

$$(31) \quad \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))} \leq \frac{\rho(g)}{\rho_2(f)}.$$

Now from (28) and (29) we get that

$$\liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))} \leq \min \left\{ \frac{\mu(g)}{\mu_2(f)}, \frac{\rho(g)}{\rho_2(f)} \right\}.$$

This proves the theorem. \square

The following theorem is a natural consequence of Theorem 4 and Theorem 5.

Theorem 6. *Let f and h be transcendental entire functions and g be an entire function such that (i) $0 < \mu_2(f) \leq \rho_2(f) < \infty$, (ii) $0 < \mu(h) \leq \rho(h) < \infty$, (iii) $0 < \mu(g) \leq \rho(g) < \infty$. Also let f has a maximal deficiency sum. Then*

$$\begin{aligned} \liminf_{r \rightarrow \infty} \frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))} &\leq \min \left\{ \frac{\mu(g)}{\mu_2(f)}, \frac{\rho(g)}{\rho_2(f)} \right\} \leq \max \left\{ \frac{\mu(g)}{\mu_2(f)}, \frac{\rho(g)}{\rho_2(f)} \right\} \\ &\leq \limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(r, h \circ g)}{\log^{[2]} T(r, L(f))}. \end{aligned}$$

Taking $f = h$ in Theorem 4 the above theorem can be proved in view of Theorem 5.

Theorem 7. *Let f be transcendental meromorphic with maximal deficiency sum and g be entire such that $0 < \mu(f) \leq \rho(f) < \infty$. Then*

$$\limsup_{r \rightarrow \infty} \frac{\log^{[2]} T(\exp(r^{\rho(g)}), f \circ g)}{\log^{[2]} T(\exp(r^\mu), L(f))} = \infty.$$

Proof. Let $0 < \mu' < \rho(g)$. Then in view of Lemma 2 we get for a sequence of values of r tending to infinity,

$$\begin{aligned} \log T(r, f \circ g) &\geq \log T(\exp(r^{\mu'}), f) \\ &\geq (\mu(f) - \varepsilon) \log(\exp(r^{\mu'})) \\ &\geq (\mu(f) - \varepsilon) r^{\mu'} \\ \text{i.e., } \log^{[2]} T(r, f \circ g) &\geq O(1) + \mu' \log r. \end{aligned}$$

So for a sequence of values of r tending to infinity,

$$(32) \quad \begin{aligned} & \log^{[2]} T(\exp(r^{\rho(g)}), f \circ g) \geq O(1) + \mu' \log(\exp(r^{\rho(g)})) \\ \text{i.e., } & \log^{[2]} T(\exp(r^{\rho(g)}), f \circ g) \geq O(1) + \mu' r^{\rho(g)}. \end{aligned}$$

Again in view of Lemma 4, we have for all sufficiently large values of r ,

$$(33) \quad \begin{aligned} & \log T(\exp(r^\mu), L(f)) \leq (\rho(L(f)) + \varepsilon) \log(\exp(r^\mu)) \\ \text{i.e., } & \log T(\exp(r^\mu), L(f)) \leq (\rho(f) + \varepsilon) r^\mu \\ \text{i.e., } & \log^{[2]} T(\exp(r^\mu), L(f)) \leq O(1) + \mu \log r. \end{aligned}$$

Now combining (32) and (33) we obtain for a sequence of values of r tending to infinity,

$$\frac{\log^{[2]} T(\exp(r^{\rho(g)}), f \circ g)}{\log^{[2]} T(\exp(r^\mu), L(f))} \geq \frac{O(1) + \mu' r^{\rho(g)}}{O(1) + \mu \log r},$$

from which the theorem follows. \square

Acknowledgement. The authors are thankful to the referee for his / her valuable suggestions towards the improvement of the paper.

REFERENCES

- [1] Bergweiler, W., *On the Nevanlinna characteristic of a composite function*, Complex Variables **10** (1988), 225–236.
- [2] Bergweiler, W., *On the growth rate of composite meromorphic functions*, Complex Variables **14** (1990), 187–196.
- [3] Hayman, W. K., *Meromorphic Functions*, The Clarendon Press, Oxford, 1964.
- [4] Lahiri, I., *Generalised proximate order of meromorphic functions*, Mat. Vesnik **41** (1989), 9–16.
- [5] Lahiri, I., Banerjee, A., *Value distribution of a Wronskian*, Port. Math. (N.S.) **61** (2) (2004), 161–175.
- [6] Lin, Q., Dai, C., *On a conjecture of Shah concerning small functions*, Kexue Tongbao (Chinese) **31** (4) (1986), 220–224, (English ed.).
- [7] Niino, K., Yang, C. C., *Some growth relationships on factors of two composite entire functions*, Factorization theory of meromorphic functions and related topics, Marcel Dekker, Inc. (New York and Basel), 1982, pp. 95–99.
- [8] Shah, S. M., *In proximate orders of integral functions*, Bull. Amer. Math. Soc. **52** (1946), 326–328.
- [9] Shah, S. M., *A note on lower proximate orders*, J. Indian Math. Soc. (N.S.) **12** (1948), 31–32.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH BENGAL
PIN: 734013, WEST BENGAL, INDIA
E-mail: sanjib_kr_datta@yahoo.co.in

TARANGAPUR N. K. HIGH SCHOOL
POST: TARANGAPUR, UTTAR DINAJPUR
PIN: 733129, WEST BENGAL, INDIA
E-mail: d_ajha@yahoo.co.in