

Martin Čadek; Jiří Vanžura

On the classification of oriented vector bundles over 5-complexes

*Czechoslovak Mathematical Journal*, Vol. 43 (1993), No. 4, 753–764

Persistent URL: <http://dml.cz/dmlcz/128427>

## Terms of use:

© Institute of Mathematics AS CR, 1993

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE CLASSIFICATION OF ORIENTED VECTOR BUNDLES  
OVER 5-COMPLEXES

MARTIN ČADEK, JIŘÍ VANŽURA, Brno

(Received July 7, 1992)

## 1. INTRODUCTION

The effort to classify vector bundles over a fixed CW-complex has a long history. The first result in this direction is the assertion that every two-dimensional oriented vector bundle is uniquely determined by its Euler class. Complete characterization of oriented vector bundles over a 4-dimensional CW-complex was given in [2] using the difference cocycles. In [8] E. Thomas found conditions for a mapping  $f \in [X, Y]$  to be uniquely determined by its cohomology homomorphism  $f^* \in \text{Hom}(H^*(Y), H^*(X))$  under the assumptions that  $X$  is a suspension or  $Y$  is an H-space. He also applied the result to  $Y = BO$ , the classifying space for the group  $O$ , and so he obtained conditions on  $H^*(X)$  under which stable vector bundles over  $X$  are determined by their Stiefel-Whitney and Pontrjagin classes. A further progress was made in [3] where the question how many  $n$ -dimensional vector bundles over a CW-complex of the same dimension are determined by a stable vector bundle  $\xi$ . The results are given in terms of  $\xi$  and they allow successful application for  $n = 3$  and  $7$ . Earlier results concerning characterization of oriented vector bundles over low dimensional complexes were summarized and completed in [13]. Using elementary homotopy theoretic methods and relations among characteristic classes L. M. Woodward has given the classification of stable oriented vector bundles over CW-complexes of dimension  $\leq 8$  and the classification of  $n$ -dimensional oriented vector bundles over CW-complexes of dimension  $n$  for  $n = 3, 4, 6, 7, 8$ , both in terms of characteristic classes. A typical condition on a CW-complex  $X$  to admit such a classification is:  $H^4(X, \mathbf{Z})$  has no element of order 4.

In dimension 5 the situation is much more complicated as can be seen on the example of the sphere  $S^5$ . Both the trivial and the tangent bundle over  $S^5$  have all characteristic classes equal to zero. Moreover, all conditions of Woodward's type are satisfied. The aim of our paper is to derive necessary and sufficient conditions

on a 5-dimensional CW-complex  $X$  which make the classification of 5-dimensional oriented vector bundles over  $X$  in terms of characteristic classes possible. This is carried out in Section 3 using a combination of the method of Postnikov tower and the Woodward method (see [9] and [13]).

The maximal number of linearly independent sections in a vector bundle  $\xi$  is defined to be the span of  $\xi$ . As a consequence of the classification described above we compute the span of 5-dimensional oriented vector bundles over CW-complexes of the same dimension. These results complete computations of Thomas for tangent bundles over 5-dimensional manifolds given in [12] and also our results for the dimensions 6 and 7 obtained in [1]. Together with results on the existence of a 2-distribution and a 4-distribution with a complex structure they form the contents of Section 4.

## 2. PRELIMINARIES

All vector bundles will be considered over a connected CW-complex  $X$  and will be oriented. The letter  $\varepsilon$  will stand for the trivial one-dimensional vector bundle. The mapping  $\beta_k : H^*(X, \mathbf{Z}_k) \rightarrow H^*(X, \mathbf{Z})$  is the Bockstein homomorphism associated with the exact sequence  $0 \rightarrow \mathbf{Z} \rightarrow \mathbf{Z} \rightarrow \mathbf{Z}_k \rightarrow 0$ . The mappings  $i_* : H^*(X, \mathbf{Z}_2) \rightarrow H^*(X, \mathbf{Z}_4)$  and  $\varrho_k : H^*(X, \mathbf{Z}) \rightarrow H^*(X, \mathbf{Z}_k)$  are induced from the inclusion  $\mathbf{Z}_2 \rightarrow \mathbf{Z}_4$  and reduction mod  $k$ , respectively.

An important role in our considerations is played by the Pontrjagin square  $\mathfrak{P}$ , a cohomology operation from  $H^{2k}(X, \mathbf{Z}_2)$  into  $H^{4k}(X, \mathbf{Z}_4)$  satisfying the relations

$$\begin{aligned} (1) \quad & \mathfrak{P}\varrho_2 x = \varrho_4 x^2, \\ (2) \quad & \mathfrak{P}(u + v) = \mathfrak{P}u + \mathfrak{P}v + i_*(u \cdot v), \end{aligned}$$

for  $x \in H^{2k}(X, \mathbf{Z})$  and  $u, v \in H^{2k}(X, \mathbf{Z}_2)$ . See [5], chapter 2.

We will use  $w_j(\xi)$  for the  $j$ -th Stiefel-Whitney class of the vector bundle  $\xi$ ,  $p_1(\xi)$  for the first Pontrjagin class, and  $e(\xi)$  for the Euler class. For a complex vector bundle  $\zeta$  the symbol  $c_j(\zeta)$  denotes the  $j$ -th Chern class. The letters  $w_j, p_1, e$  stand for characteristic classes of the universal oriented  $n$ -dimensional vector bundle over the classifying space  $BSO(n)$ . Our results given below are based on the following relations among the characteristic classes:

$$\begin{aligned} (3) \quad & \varrho_4 p_1(\xi) = \mathfrak{P}w_2(\xi) + i_* w_4(\xi), \\ (4) \quad & w_6(\xi) = Sq^2 w_4(\xi) + w_2(\xi)w_4(\xi), \end{aligned}$$

the former being proved in [4] and [7] and the latter being a special case of the Wu formula.

The Eilenberg-MacLane space with the  $n$ -th homotopy group  $G$  will be denoted by  $K(G, n)$  and  $\iota_n$  will stand for the fundamental class in  $H^n(K(G, n), G)$ . Writing the fundamental class it will be always clear which group  $G$  we have in mind.

In the proof of Theorem 1 we will need suspension. Being defined for every fibration  $F \xrightarrow{j} E \xrightarrow{p} B$ , it is a natural mapping from a subgroup of  $H^{k+1}(B)$  into  $H^k(F)/\text{im } j^*$  which commutes with the Steenrod squares and  $i_*$  (see [5]).

We say that  $x \in H^*(X, \mathbf{Z})$  is an element of order  $k$  ( $k = 2, 3, 4, \dots$ ) if and only if  $x \neq 0$  and  $k$  is the least positive integer such that  $kx = 0$  (if it exists). Some results will involve the following hypotheses:

**Condition (A).**  $H^4(X, \mathbf{Z})$  has no element of order 4.

**Condition (B).**  $Sq^2 H^3(X, \mathbf{Z}_2) = H^5(X, \mathbf{Z}_2)$ .

**Remark.** An important example of a CW-complex which satisfies Condition (B) is a 5-dimensional oriented smooth manifold  $M$  with  $w_2(M) \neq 0$ . The Poincaré duality and the fact that the second Wu class is equal to  $w_2(M)$  yields

$$Sq^2 H^3(M, \mathbf{Z}_2) = w_2(M) H^3(M, \mathbf{Z}_2) = H^5(M, \mathbf{Z}_2).$$

### 3. CLASSIFICATION THEOREM

Let  $X$  be a connected CW-complex of dimension  $\leq 5$ . Our problem consists in finding conditions on  $X$  such that for every  $a \in H^2(X, \mathbf{Z}_2)$ ,  $b \in H^4(X, \mathbf{Z}_2)$ ,  $c \in H^4(X, \mathbf{Z})$  there is at most one oriented 5-dimensional vector bundle  $\xi$  with  $w_2(\xi) = a$ ,  $w_4(\xi) = b$ ,  $p_1(\xi) = c$ . A necessary and sufficient condition on  $a$ ,  $b$ ,  $c$  for the existence of such a vector bundle derived in [W] is given by the relation  $\varrho_4 c = \mathfrak{P}a + i_* b$  (see (3)). Up to homotopy there is just one mapping  $f: X \rightarrow K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4) \times K(\mathbf{Z}, 4)$  such that  $f^*(\iota_2 \otimes 1 \otimes 1) = a$ ,  $f^*(1 \otimes \iota_4 \otimes 1) = b$ ,  $f^*(1 \otimes 1 \otimes \iota_4) = c$ . Similarly,  $w_2, w_4, p_1$ , the cohomology classes of  $BSO(5)$ , determine a mapping  $\alpha: BSO(5) \rightarrow K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4) \times K(\mathbf{Z}, 4)$  which can be considered to be a fibration. Now the problem described above can be formulated as a problem of lifting: *when every mapping  $f: X \rightarrow K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4) \times K(\mathbf{Z}, 4)$  has at most one lifting  $\xi: X \rightarrow BSO(5)$  in the fibration  $\alpha$ .*

$$\begin{array}{ccc}
 & & BSO(5) \\
 & \nearrow \xi & \downarrow \alpha \\
 X & \xrightarrow{f} & K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4) \times K(\mathbf{Z}, 4)
 \end{array}$$

To solve this problem we will construct a Postnikov tower for the fibration  $\alpha: BSO(5) \rightarrow K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4) \times K(\mathbf{Z}, 4)$ . Put  $K = K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4) \times K(\mathbf{Z}, 4)$  and denote the fibre of  $\alpha$  by  $V$ . Let us recall that  $\pi_k(BSO(5)) \cong 0$  for  $k = 1, 3$ ,  $\pi_k(BSO(5)) \cong \mathbf{Z}_2$  for  $k = 2, 5$  and  $\pi_4(BSO(5)) \cong \mathbf{Z}$ . Considering the characteristic classes as mappings from  $BSO(5)$  into appropriate Eilenberg-MacLane spaces, we get  $w_{2*} = id: \pi_2(BSO(5)) \rightarrow \mathbf{Z}_2$ ,  $w_{4*} = \varrho_2: \pi_4(BSO(5)) \rightarrow \mathbf{Z}_2$ , and  $p_{1*}: \pi_4(BSO(5)) \rightarrow \mathbf{Z}$  is a multiplication by 2. See [13]. From the long exact homotopy sequence we compute:  $\pi_1(V) \cong \pi_2(V) \cong 0$ ,  $\pi_3(V) \cong \mathbf{Z}_4$ ,  $\pi_4(V) \cong 0$ , and  $\pi_5(V) = \mathbf{Z}_2$ . The first invariant in the Postnikov tower is the transgression of a fundamental class in  $H^3(V, \mathbf{Z}_4)$ . It is a generator of  $\ker \alpha^* \subset H^4(K, \mathbf{Z}_4)$ . Hence it is equal to

$$\varrho_4(1 \otimes 1 \otimes \iota_4) - \mathfrak{P}\iota_2 \otimes 1 \otimes 1 - 1 \otimes i_* \iota_4 \otimes 1.$$

Let  $E_1$  be the first stage of the Postnikov tower and let the new mappings be denoted according to the diagram.

$$\begin{array}{ccccc} \bar{F}_1 & \longrightarrow & V & \xrightarrow{\bar{\beta}_1} & K(\mathbf{Z}_4, 3) \\ & & \downarrow & & \downarrow i_1 \\ F_1 & \longrightarrow & BSO(5) & \xrightarrow{\beta_1} & E_1 \\ & & \downarrow \alpha & & \downarrow \pi_1 \\ & & K & \xlongequal{\quad} & K \xrightarrow{\varrho_4(1 \otimes 1 \otimes \iota_4) - \mathfrak{P}\iota_2 \otimes 1 \otimes 1 - 1 \otimes i_* \iota_4 \otimes 1} K(\mathbf{Z}_4, 4) \end{array}$$

Consider  $\beta_1: BSO(5) \rightarrow E_1$  as a fibration with a fibre  $F_1$ . This fibre is homotopy equivalent to the homotopy fibre  $\bar{F}_1$  of the mapping  $\bar{\beta}_1$  (see [9]). Hence computing the homotopy groups of  $\bar{F}_1$  we get that  $F_1$  is 4-connected and  $\pi_5(F_1) \cong \mathbf{Z}_2$ . Consequently,  $\beta_1$  is a 5-equivalence.

The next invariant  $\varphi \in H^6(E_1, \mathbf{Z}_2)$  is the transgression of the generator of  $H^5(F_1, \mathbf{Z}_2)$  in the Serre exact sequence for the fibration  $\beta_1$ .  $E_1$  is also the first stage in the Postnikov tower for the fibration  $\hat{\alpha}: BSO(6) \rightarrow K$  determined by  $w_2$ ,  $w_4$  and  $p_1$ . The mapping  $\hat{\beta}_1: BSO(6) \rightarrow E_1$  in this Postnikov tower is a 6-equivalence (since  $\pi_5(BSO(6)) \cong 0$ ). Using the Serre exact sequence for the fibration  $\hat{\beta}_1$ , we get that  $\hat{\beta}_1^*$  is an isomorphism between  $H^6(E_1, \mathbf{Z}_2)$  and  $H^6(BSO(6), \mathbf{Z}_2)$ . The latter group has generators  $w_2^3$ ,  $w_3^2$ ,  $w_2 w_4$  and  $Sq^2 w_4 (= w_6 + w_2 w_4)$ . Hence the generators of  $H^6(E_1, \mathbf{Z}_2)$  are  $\pi_1^*(\iota_2^3 \otimes 1 \otimes 1)$ ,  $\pi_1^*((Sq^1 \iota_2)^2 \otimes 1 \otimes 1)$ ,  $\pi_1^*(\iota_2 \otimes \iota_4 \otimes 1)$ ,  $\pi_1^*(1 \otimes Sq^2 \iota_4 \otimes 1)$ . The mapping  $\beta_1^*: H^6(E_1, \mathbf{Z}_2) \rightarrow H^6(BSO(5), \mathbf{Z}_2)$  maps them into  $w_2^3$ ,  $w_3^2$ ,  $w_2 w_4$  and  $Sq^2 w_4 = w_2 w_4$ , respectively. Consequently, using the Serre exact sequence for the fibration  $\beta_1$  we get  $\varphi = \pi_1^*(\iota_2 \otimes \iota_4 \otimes 1 + 1 \otimes Sq^2 \iota_4 \otimes 1)$ . So we can build the second

stage  $E_2$  of our Postnikov tower.

$$\begin{array}{ccccc}
 \bar{F}_2 & \longrightarrow & F_1 & \xrightarrow{\bar{\beta}_2} & K(\mathbf{Z}_2, 5) \\
 & & \downarrow & & \downarrow i_2 \\
 F_2 & \longrightarrow & BSO(5) & \xrightarrow{\beta_2} & E_2 \\
 & & \downarrow \beta_1 & & \downarrow \pi_2 \\
 & & E_1 & \xlongequal{\quad} & E_1 \xrightarrow{\pi_1^*(\iota_2 \otimes \iota_4 \otimes 1 + 1 \otimes Sq^2 \iota_4 \otimes 1)} K(\mathbf{Z}_2, 6)
 \end{array}$$

Let the notation of new mappings accord with the diagram. We can consider  $\beta_2$  to be a fibration with a fibre  $F_2$ . Similarly as for the first stage, we can compute the homotopy groups of  $F_2$ . So we get that  $F_2$  is 5-connected and  $\beta_2$  is a 6-equivalence.

Let  $C = K(\mathbf{Z}_4, 4) \times K(\mathbf{Z}_2, 6)$ . Up to homotopy there is just one mapping  $k = (k_1, k_2): K \rightarrow C$  given by

$$\begin{aligned}
 k_1^*(\iota_4) &= 1 \otimes 1 \otimes \varrho_4 \iota_4 - \mathfrak{P} \iota_2 \otimes 1 \otimes 1 + 1 \otimes i_* \iota_4 \otimes 1 \\
 k_2^*(\iota_6) &= \iota_2 \otimes \iota_4 \otimes 1 + 1 \otimes Sq^2 \iota_4 \otimes 1.
 \end{aligned}$$

Due to Lemma 8.1 in [10], there is a homeomorphism  $g: E_2 \rightarrow E$  where  $\pi: E \rightarrow K$  is a principal fibration with the classifying map  $k: K \rightarrow C$ . Moreover,  $\pi_1 \circ \pi_2 = \pi \circ g$  and the fibration  $\beta = g \circ \beta_2: BSO(5) \rightarrow E$  is a 6-equivalence. Hence, we can consider the situation

$$\begin{array}{ccc}
 BSO(5) & \xrightarrow{\beta = 6\text{-equiv}} & E \\
 \downarrow \alpha & & \downarrow \pi \\
 K & \xlongequal{\quad} & K \xrightarrow{k = (k_1, k_2)} C
 \end{array}$$

which allows us to prove our main result.

**Theorem 1.** *Let  $X$  be a connected CW-complex of dimension  $\leq 5$  and suppose*

$$\gamma: [X, BSO(5)] \rightarrow H^2(X, \mathbf{Z}_2) \oplus H^4(X, \mathbf{Z}_2) \oplus H^4(X, \mathbf{Z})$$

is defined by  $\gamma(\xi) = (w_2(\xi), w_4(\xi), p_1(\xi))$ . Then

- (i)  $\text{im } \gamma = \{(a, b, c) \mid \varrho_4 c = \mathfrak{P}a + i_* b\}$ ,
- (ii)  $\gamma$  is injective if and only if Conditions (A) and (B) are satisfied.

**Proof.** (i) follows immediately from the fact that a mapping  $f: X \rightarrow K$  can be lifted in the fibration  $\alpha$  into  $BSO(5)$  if and only if  $f^*(1 \otimes 1 \otimes \varrho_4 \iota_4 - \mathfrak{P} \iota_2 \otimes 1 \otimes 1 - 1 \otimes i_* \iota_4 \otimes 1) = 0$ . (See similar proofs in [1].)

(ii) Since the space  $E$  is a homotopy fibre of the mapping  $k: K \rightarrow C$ , the Puppe sequence

$$\Omega K \xrightarrow{\Omega k} \Omega C \xrightarrow{q} E \xrightarrow{\pi} K \xrightarrow{k} C$$

yields the exact sequence

$$\rightarrow [X, \Omega K] \xrightarrow{(\Omega k)_*} [X, \Omega C] \xrightarrow{q_*} [X, E] \xrightarrow{\pi_*} [X, K] \xrightarrow{k_*} [X, C].$$

Moreover,  $\beta$  being a 6-equivalence,  $\beta_*: [X, BSO(5)] \rightarrow [X, E]$  is a bijection for every CW-complex of dimension  $\leq 5$ . The following statements are equivalent:

- (1)  $\gamma = \alpha_* = \pi_* \circ \beta_*: [X, BSO(5)] \rightarrow [X, K]$  is injective.
- (2)  $\pi_*: [X, E] \rightarrow [X, K]$  is injective.
- (3)  $q_* = 0$
- (4)  $(\Omega k)_*: [X, \Omega K] \rightarrow [X, \Omega C]$  is surjective.

Hence we need to compute  $(\Omega k_1)^*: H^3(K(\mathbf{Z}_4, 3), \mathbf{Z}_4) \rightarrow H^3(\Omega K, \mathbf{Z}_4)$  and  $(\Omega k_2)^*: H^5(K(\mathbf{Z}_2, 5), \mathbf{Z}_2) \rightarrow H^5(\Omega K, \mathbf{Z}_2)$ .

First, let us consider  $k_1$ .

$$\begin{array}{ccccc} \Omega K & \xlongequal{\quad} & K(\mathbf{Z}_2, 1) \times K(\mathbf{Z}_2, 3) \times K(\mathbf{Z}, 3) & \xrightarrow{\Omega k_1} & K(\mathbf{Z}_4, 3) \\ \downarrow & & \downarrow & & \downarrow \\ PK & \xlongequal{\quad} & PK(\mathbf{Z}_2, 2) \times PK(\mathbf{Z}_2, 4) \times PK(\mathbf{Z}, 4) & \xrightarrow{Pk_1} & PK(\mathbf{Z}_4, 4) \\ \downarrow & & \downarrow & & \downarrow \\ K & \xlongequal{\quad} & K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4) \times K(\mathbf{Z}, 4) & \xrightarrow{k_1} & K(\mathbf{Z}_4, 4) \end{array}$$

Every element in  $H^*(K, \mathbf{Z}_4)$  is suspensive. If we denote all suspensions by  $\sigma$ , we get

$$\begin{aligned} (\Omega k_1)^*(\iota_3) &= (\Omega k_1)^*(\sigma \iota_4) = \sigma(k_1^* \iota_4) = \sigma(1 \otimes 1 \otimes \varrho_4 \iota_4) - \sigma(\mathfrak{P} \iota_2 \otimes 1 \otimes 1) \\ &\quad - \sigma(1 \otimes i_* \iota_4 \otimes 1) = 1 \otimes 1 \otimes \sigma(\varrho_4 \iota_4) - \sigma(\mathfrak{P} \iota_2) \otimes 1 \otimes 1 - 1 \otimes \sigma(i_* \iota_4) \otimes 1 \end{aligned}$$

the last equality being a consequence of the definition of suspension and coboundary operator. In the fibration  $K(\mathbf{Z}, 3) \rightarrow PK(\mathbf{Z}, 4) \rightarrow K(\mathbf{Z}, 4)$  we get

$$\sigma(\varrho_4 \iota_4) = \varrho_4(\sigma \iota_4) = \varrho_4 \iota_3.$$

In the fibration  $K(\mathbf{Z}_2, 3) \rightarrow PK(\mathbf{Z}_2, 4) \rightarrow K(\mathbf{Z}_2, 4)$  we have

$$\sigma(i_*\iota_4) = i_*(\sigma\iota_4) = i_*\iota_3$$

and finally, in the fibration  $K(\mathbf{Z}_2, 1) \rightarrow PK(\mathbf{Z}_2, 2) \rightarrow K(\mathbf{Z}_2, 2)$  we obtain

$$(5) \quad \sigma(\mathfrak{P}\iota_2) = i_*\iota_1^3.$$

Since this fact is not generally known, we will prove it at the end of this section. As a result of these computations we get

$$(\Omega k_1)_* : [X, \Omega K] \rightarrow [X, K(\mathbf{Z}_4, 3)] : (a, b, c) \mapsto \varrho_4 c - i_* a^3 - i_* b.$$

Hence  $(\Omega k_1)_*$  is surjective if and only if

$$(6) \quad H^3(X, \mathbf{Z}_4) = \varrho_4 H^3(X, \mathbf{Z}) + i_* H^3(X, \mathbf{Z}_2).$$

We show that (6) is equivalent to the condition (A).

(A)  $\Rightarrow$  (6). Let  $x \in H^3(X, \mathbf{Z}_4)$ , then  $4\beta_4 x = 0$ . (A) implies that  $2\beta_4 x = 0$ . Consequently, there is a  $y \in H^3(X, \mathbf{Z}_2)$  such that  $\beta_4 x = \beta_2 y = \beta_4 i_* y$ . That is why  $\beta_4(x - i_* y) = 0$ , which implies  $x = i_* y + \varrho_4 z$  for some  $z \in H^3(X, \mathbf{Z})$ .

(6)  $\Rightarrow$  (A). Let  $v \in H^4(X, \mathbf{Z})$  satisfy  $4v = 0$ . Then  $v = \beta_4 x$  where  $x = \varrho_4 z + i_* y \in H^3(X, \mathbf{Z}_4)$  so that  $v = \beta_4 \varrho_4 z + \beta_4 i_* y = \beta_4 i_* y = \beta_2 y$ . Hence  $2v = 0$  and  $v$  is not an element of order 4.

Now consider the mapping  $k_2$ . The computation of  $(\Omega k_2)^* : H^5(K(\mathbf{Z}_2, 5), \mathbf{Z}_2) \rightarrow H^5(\Omega K, \mathbf{Z}_2)$  gives

$$\begin{aligned} (\Omega k_2)^*(\iota_5) &= (\Omega k_2)^*(\sigma\iota_6) = \sigma k_2^*(\iota_6) = 1 \otimes \sigma(Sq^2\iota_4) \otimes 1 + \sigma(\iota_2 \otimes \iota_4) \otimes 1 \\ &= 1 \otimes Sq^2\iota_3 \otimes 1 + \sigma(\iota_2 \otimes \iota_4) \otimes 1. \end{aligned}$$

We are going to prove that  $\sigma(\iota_2 \otimes \iota_4) = 0$ . Consider the fibration

$$\Omega B \rightarrow PB \xrightarrow{p} B$$

where  $B = K(\mathbf{Z}_2, 2) \times K(\mathbf{Z}_2, 4)$ . Let  $\hat{p}^* : H^6(B, \mathbf{Z}_2) \rightarrow H^6(PB, \Omega B; \mathbf{Z}_2)$  be determined by the mapping  $p$ . It is sufficient to show  $\hat{p}^*(\iota_2 \otimes \iota_4) = 0$ . Using the Serre spectral sequence with coefficients  $\mathbf{Z}_2$  for the above fibration, we have

$$\hat{p}^* : H^6(B, \mathbf{Z}_2) \cong E_2^{6,0} \rightarrow E_6^{6,0} \hookrightarrow H^6(PB, \Omega B; \mathbf{Z}_2).$$



We compute  $d_2: E_2^{4,1} \rightarrow E_2^{6,0}$ . Since  $E_2^{4,1} \cong E_2^{4,0} \otimes E_2^{0,1}$ , for the generators of  $E_2^{4,1}$  we obtain

$$\begin{aligned} d_2(\iota_2^2 \otimes \iota_1) &= d_2(\iota_2^2) \cdot \iota_1 + \iota_2^2 \cdot d_2(\iota_1) = \iota_2^2 \cdot \iota_2 = \iota_2^3, \\ d_2(\iota_4 \otimes \iota_1) &= d_2(\iota_4) \cdot \iota_1 + \iota_4 \cdot d_2(\iota_1) = \iota_4 \cdot \iota_2. \end{aligned}$$

Hence  $\iota_4 \cdot \iota_2$  vanishes in  $E_3^{6,0}$  and  $\hat{p}^*(\iota_2 \otimes \iota_4) = 0$ .

So we conclude that

$$(\Omega k_2)_*[X, \Omega K] \rightarrow [X, K(\mathbf{Z}_2, 5)]: (a, b, c) \mapsto Sq^2 b$$

and its surjectivity is given directly by Condition (B).

It remains to prove the relation (5). Consider the Serre spectral sequence for the fibration  $K(\mathbf{Z}_2, 1) \rightarrow PK(\mathbf{Z}_2, 2) \rightarrow K(\mathbf{Z}_2, 2)$  with coefficients  $\mathbf{Z}_4$ . For brevity we will again denote this fibration by  $\Omega B \rightarrow PB \xrightarrow{p} B$ . It is not difficult to show that  $H^4(B, \mathbf{Z}_4) \cong \mathbf{Z}_4$  with the generator  $\mathfrak{P}\iota_2$  and  $H^3(\Omega B, \mathbf{Z}_4) \cong \mathbf{Z}_2$  with the generator  $i_*\iota_1^3$ . The coboundary operator in the long exact sequence for the couple  $(PB, \Omega B)$  is an isomorphism, hence it is sufficient to prove that  $\hat{p}^*(\mathfrak{P}\iota_2) \neq 0$ ,  $\hat{p}^*: H^4(B, \mathbf{Z}_4) \rightarrow H^4(PB, \Omega B; \mathbf{Z}_4)$  being induced by  $p$ . Since

$$E_4^{4,0} \cong H^4(B, \mathbf{Z}_4) / \ker \hat{p}^*,$$

it is sufficient to show that  $E_4^{4,0} \neq 0$ . We have

$$\begin{aligned} E_2^{2,1} &\cong H^2(B, H^1(\Omega B, \mathbf{Z}_4)) \cong \mathbf{Z}_2 \cong E_2^{2,0} \otimes E_2^{0,1}, \\ E_2^{4,0} &\cong H^4(B, H^0(\Omega B, \mathbf{Z}_4)) \cong \mathbf{Z}_4. \end{aligned}$$

Moreover,  $d_2: E_2^{2,1} \rightarrow E_2^{4,0}$  is injective because

$$\begin{aligned} d_2(i_*\iota_2 \otimes i_*\iota_1) &= d_2(i_*\iota_2) \cdot i_*\iota_1 + i_*\iota_2 \cdot d_2(i_*\iota_1) = \\ &= i_*\iota_2 \cdot \tau(i_*\iota_1) = i_*\iota_2^2 \end{aligned}$$

where  $\tau$  is a transgression. Hence  $E_3^{4,0} \cong \mathbf{Z}_2$ . Further,  $E_3^{1,2} \cong 0$ ,  $E_3^{7,-2} \cong 0$  and consequently,  $E_4^{4,0} \cong \mathbf{Z}_2$ .  $\square$

#### 4. SPAN AND THE EXISTENCE OF DISTRIBUTIONS

In this section we compute the span of oriented 5-dimensional vector bundles over a 5-dimensional CW-complex satisfying Conditions (A) and (B) of Theorem 1. Under the same conditions we find all oriented 5-dimensional vector bundles which admit a 2-distribution, i.e. an oriented 2-dimensional subbundle, and all oriented 5-dimensional vector bundles which admit a 4-distribution endowed with a complex structure, i.e. a complex 2-dimensional subbundle. For these purposes we need

**Theorem 2.** *Let  $X$  be a connected CW-complex of dimension  $\leq 5$  and let  $W \in H^2(X, \mathbf{Z}_2)$ ,  $P \in H^4(X, \mathbf{Z})$ . Then there exists an oriented 3-dimensional vector bundle  $\xi$  over  $X$  with*

$$w_2(\xi) = W, \quad p_1(\xi) = P$$

if and only if

$$\varrho_4 P = \mathfrak{P}W.$$

*Proof* is very similar to the proof of the first part of Theorem 1. See also [13].

□

**Corollary 1.** *Let  $X$  be a connected CW-complex of dimension  $\leq 5$  satisfying Conditions (A) and (B). Then an oriented 5-dimensional vector bundle  $\xi$  has a 2-distribution with Euler class  $U$  if and only if*

$$(7) \quad \varrho_2 U^2 + w_2(\xi)\varrho_2 U + w_4(\xi) = 0.$$

*Proof.* ( $\Rightarrow$ ) Let  $\xi = \zeta \oplus \tau$  where  $\tau$  is an oriented 2-dimensional vector bundle over  $X$  with the Euler class  $U$  and  $\zeta$  is an oriented 3-dimensional vector bundle over  $X$ . Then

$$\begin{aligned} w_2(\xi) &= w_2(\zeta) + w_2(\tau) = w_2(\zeta) + \varrho_2 U, \\ w_4(\xi) &= w_2(\zeta) \cdot w_2(\tau) = w_2(\zeta) \cdot \varrho_2 U. \end{aligned}$$

Substituting from here into the expression  $\varrho_2 U^2 + w_2(\xi) \cdot \varrho_2 U + w_4(\xi)$ , we get (7).

( $\Leftarrow$ ) Let  $U \in H^2(X, \mathbf{Z})$  satisfy (7). There is an oriented 2-dimensional vector bundle  $\tau$  over  $X$  with the Euler class  $U$ . Put

$$W = w_2(\xi) + \varrho_2 U, \quad P = p_1(\xi) - U^2.$$

Then

$$\begin{aligned} \varrho_4 P - \mathfrak{P}W &= \varrho_4 p_1(\xi) - \varrho_4 U^2 - \mathfrak{P}(w_2(\xi) + \varrho_2 U) = \\ &= \varrho_4 p_1(\xi) - \varrho_4 U^2 - \mathfrak{P}w_2(\xi) - \mathfrak{P}\varrho_2 U - i_*(w_2(\xi)\varrho_2 U) = \\ &= i_*(\varrho_2 U^2 + w_2(\xi)\varrho_2 U + w_4(\xi)) = 0. \end{aligned}$$

According to Theorem 2, there is an oriented 3-dimensional vector bundle  $\zeta$  over  $X$  with  $w_2(\zeta) = W$  and  $p_1(\zeta) = P$ . We compute the characteristic classes of the vector bundle  $\zeta \oplus \tau$ .

$$\begin{aligned} w_2(\zeta \oplus \tau) &= w_2(\zeta) + w_2(\tau) = W + \varrho_2 U = w_2(\xi), \\ w_4(\zeta \oplus \tau) &= w_2(\zeta) \cdot w_2(\tau) = W \cdot \varrho_2 U = w_2(\xi)\varrho_2 U + \varrho_2 U^2 = \\ &= w_4(\xi), \\ p_1(\zeta \oplus \tau) &= p_1(\zeta) + p_1(\tau) = P + U^2 = p_1(\xi). \end{aligned}$$

(See [13] for the additivity of  $p_1$  in this case.) Theorem 1 now implies that  $\xi = \zeta \oplus \tau$ , which completes the proof.  $\square$

**Remark.** As far as it is known to the authors there are only two general results concerning 2-distributions in 5 or  $4k + 1$ -dimensional vector bundles. See [11], Theorems 1.3 and 4.1. The former deals with spin manifolds (i.e.  $w_1(X) = w_2(X) = 0$ ) and tangent bundles while the latter requires  $\text{span} \geq 2$ . Both examine the existence of 2-distributions with the Euler class  $2U \in H^2(X, \mathbf{Z})$ .

**Corollary 2.** *Let  $X$  be a connected CW-complex of dimension  $\leq 5$  and let  $\xi$  be an oriented 5-dimensional vector bundle over  $X$ . Then*

(1)  $\text{span } \xi \geq 1$  if and only if  $e(\xi) = 0$ .

If Conditions (A) and (B) are satisfied then

(2)  $\text{span } \xi \geq 2$  if and only if  $w_4(\xi) = 0$ .

(3)  $\text{span } \xi \geq 3$  if and only if  $w_4(\xi) = 0$  and there is a  $U \in H^2(X, \mathbf{Z})$  such that  $w_2(\xi) = \varrho_2 U$ ,  $p_1(\xi) = U^2$ .

(4)  $\text{span } \xi = 5$  if and only if  $w_2(\xi) = 0$ ,  $w_4(\xi) = 0$ ,  $p_1(\xi) = 0$ .

**Proof.** (1) is well known and is included only for completeness.

(2) is the immediate consequence of Corollary 1 for  $U = 0$ .

(3)( $\Rightarrow$ ) Let  $\xi = \zeta \oplus 3\varepsilon$  where  $\zeta$  is an oriented 2-dimensional vector bundle over  $X$ . Then  $w_4(\xi) = w_4(\zeta) = 0$  and for  $U = e(\zeta)$  we get  $w_2(\xi) = w_2(\zeta) = \varrho_2 U$ ,  $p_1(\xi) = p_1(\zeta) = U^2$ .

( $\Leftarrow$ ) For  $U \in H^2(X, \mathbf{Z})$  there is an oriented 2-dimensional vector bundle  $\zeta$  over  $X$  such that  $e(\zeta) = U$ . Then  $w_2(\zeta \oplus 3\varepsilon) = w_2(\zeta) = \varrho_2 U = w_2(\xi)$ ,  $w_4(\zeta \oplus 3\varepsilon) = w_4(\zeta) =$

$0 = w_4(\xi)$  and  $p_1(\zeta \oplus 3\varepsilon) = p_1(\zeta) = U^2 = p_1(\xi)$ . Theorem 1 implies that  $\zeta \oplus 3\varepsilon = \xi$  since the characteristic classes of both vector bundles are the same.

(4) follows immediately from Theorem 1. □

**Remark.** Statements (3) and (4) of Corollary 2 under a little bit different conditions were already known to E. Thomas [12]. Statement (2) under Conditions (A) and (B) is new. It deals with the cases which are not covered in [12]. The condition  $w_4(\xi) = 0$  coincides with the condition for the stable span of  $4k + 1$ -dimensional vector bundles over a CW-complex of the same dimension to be  $\geq 2$ . See [6], Theorem 2.1.1.

Now we will investigate the existence of distributions with complex structure. The case of 2-distributions is treated in Corollary 1. Here we will deal with 4-distributions. For this purpose we need the following

**Theorem 3.** *Let  $X$  be a connected CW-complex of dimension  $\leq 5$  and let  $C_1 \in H^2(X, \mathbf{Z})$ ,  $C_2 \in H^4(X, \mathbf{Z})$ . Then there exists a 2-dimensional complex vector bundle  $\zeta$  over  $X$  with the Chern classes*

$$c_1(\zeta) = C_1, \quad c_2(\zeta) = C_2.$$

*Proof* of this theorem follows the same lines as in [13]. □

**Corollary 3.** *Let  $X$  be a connected CW-complex of dimension  $\leq 5$  satisfying the conditions (A) and (B). Then an oriented 5-dimensional vector bundle  $\xi$  over  $X$  has a 4-distribution with a complex structure if and only if*

- (i)  $e(\xi) = 0$ ,
- (ii)  $\beta_2 w_2(\xi) = 0$ .

*Proof.* ( $\Rightarrow$ ) Let  $\eta$  be a 4-distribution in  $\xi$  with complex structure. Then obviously  $e(\xi) = 0$  and  $\beta_2 w_2(\xi) = \beta_2 w_2(\eta \oplus \varepsilon) = \beta_2 w_2(\eta) = \beta_2 \varrho_2 c_1(\eta) = 0$ .

( $\Leftarrow$ ) We have  $\beta_2 w_2(\xi) = 0$  and  $\beta_2 w_4(\xi) = e(\xi) = 0$ . Consequently, we can find  $a_1 \in H^2(X, \mathbf{Z})$  and  $a_2 \in H^4(X, \mathbf{Z})$  such that  $\varrho_2 a_1 = w_2(\xi)$  and  $\varrho_2 a_2 = w_4(\xi)$ . Then

$$\varrho_4(a_1^2 - 2a_2) = \mathfrak{P}\varrho_2 a_1 + i_* \varrho_2 a_2 = \mathfrak{P}w_2(\xi) + i_* w_4(\xi) = \varrho_4 p_1(\xi).$$

Hence there is a  $b \in H^4(X, \mathbf{Z})$  such that  $a_1^2 - 2a_2 - 4b = p_1(\xi)$ . Put  $C_1 = a_1$  and  $C_2 = a_2 + 2b$ . According to Theorem 3 there exists a complex vector bundle  $\eta$  over  $X$  of complex dimension 2 with

$$c_1(\eta) = C_1 \quad \text{and} \quad c_2(\eta) = C_2.$$

Let us now consider the 5-dimensional real vector bundle  $\eta \oplus \varepsilon$ . We get

$$\begin{aligned}w_2(\eta \oplus \varepsilon) &= w_2(\eta) = \varrho_2 c_1(\eta) = \varrho_2 C_1 = w_2(\xi), \\w_4(\eta \oplus \varepsilon) &= w_4(\eta) = \varrho_2 c_2(\eta) = \varrho_2 C_2 = w_4(\xi), \\p_1(\eta \oplus \varepsilon) &= p_1(\eta) = c_1(\eta)^2 - 2c_2(\eta) = C_1^2 - 2C_2 = p_1(\xi).\end{aligned}$$

Theorem 1 implies that  $\xi = \eta \oplus \varepsilon$ . This completes the proof.  $\square$

**Remark.** Let us recall that an  $f$ -structure on a vector bundle  $\xi$  is an endomorphism  $f: \xi \rightarrow \xi$  satisfying the polynomial equation  $f^3 + f = 0$  with  $\dim \ker f$  constant. It can be easily seen that if  $f$  is an  $f$ -structure then  $\xi = \zeta \oplus \eta$  where  $\zeta = \ker f$  and  $\eta = \ker(f^2 + \text{id})$ . This means that on a vector bundle  $\xi$  there exists an  $f$ -structure if and only if there exists a distribution  $\eta \subset \xi$  endowed with a complex structure. If  $\xi$  is an oriented 5-dimensional vector bundle over a connected CW-complex  $X$  of dimension 5, we can distinguish two cases. In the first case of  $\dim \eta = 2$  the existence problem for an  $f$ -structure is covered by Corollary 1. The second case of  $\dim \eta = 4$  is treated in Corollary 3.

#### References

- [1] *M. Čadek, J. Vanžura*: On oriented vector bundles over CW-complex of dimension 6 and 7, preprint (1991).
- [2] *A. Dold, H. Whitney*: Classification of oriented sphere bundles over a 4-complex, *Ann. of Math.* **69** (1959), 667–677.
- [3] *I. James, E. Thomas*: An approach to the enumeration problem for non-stable vector bundles, *J. Math. Mech.* **14** (1965), 485–506.
- [4] *J. Milnor*: Some consequences of a theorem of Bott, *Ann. of Math.* **68** (1958), 444–449.
- [5] *R. E. Mosher, M. C. Tangora*: Cohomology operations and applications in homotopy theory, Harper & Row, Publishers, New York, Evanston and London, 1968.
- [6] *Tze-Beng Ng*: On the geometric dimension of vector bundles, span of a manifold and immersion of manifolds in manifolds, *Exposition. Math.* **8** (1990), 193–226.
- [7] *E. Thomas*: On the cohomology of the real Grassmann complexes, *Trans. Amer. Math. Soc.* **96** (1960), 67–89.
- [8] *E. Thomas*: Homotopy classification of maps by cohomology homomorphisms, *Trans. Amer. Math. Soc.* **111** (1964), 138–151.
- [9] *E. Thomas*: Seminar on fibre bundles, *Lecture Notes in Math.*, no. 13, Springer, Berlin-Heidelberg-New York, 1966.
- [10] *E. Thomas*: Postnikov invariants and higher order cohomology operation, *Ann. of Math.* **85** (1967), 184–217.
- [11] *E. Thomas*: Fields of tangent  $k$ -planes on manifolds, *Invent. Math.* **3** (1967), 334–347.
- [12] *E. Thomas*: Vector fields on low dimensional manifolds, *Math. Z.* **103** (1968), 85–93.
- [13] *L. M. Woodward*: The classification of orientable vector bundles over CW-complexes of small dimension, *Proc. Roy. Soc. Edinburgh* **92A** (1982), 175–179.

*Author's address*: Academy of Sciences of the Czech Republic, Institute of Mathematics, Žitkova 22, 616 62 Brno, Czech Republic.