

Hermann Render

Spaces with a Borel-complete Stone-Čech compactification

Czechoslovak Mathematical Journal, Vol. 45 (1995), No. 2, 325–326

Persistent URL: <http://dml.cz/dmlcz/128522>

Terms of use:

© Institute of Mathematics AS CR, 1995

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SPACES WITH A BOREL-COMPLETE
STONE-ČECH COMPACTIFICATION

HERMANN RENDER, Duisburg

(Received April 15, 1993)

A topological space is called *Borel-complete* if every two-valued Borel measure is a Dirac measure. In this short note we want to investigate the Tychonoff spaces X whose Stone-Čech compactification βX is Borel-complete. In [3] it was proved that pseudocompactness is a necessary condition; moreover a conjecture was stated that X is even compact. In other words, the conjecture claims that the following class

$$(1) \quad \mathcal{B} := \{\beta X \setminus X; \beta X \text{ is Borel-complete}\}$$

consists only of the empty set. Our main result shows that the conjecture is not true; e.g. it shows that every separable metric space is in \mathcal{B} .

Theorem. *A Tychonoff space is in the class \mathcal{B} (up to homeomorphy) iff it is a subset of a Borel-complete compact space.*

Let Ψ be the space described in [6, p. 272]. As pointed out in [3] Ψ is a Borel-complete pseudocompact space. Moreover one can assume that $\Psi^* := \beta\Psi \setminus \Psi$ consists exactly of one element, cf. [4]. Consequently $\beta\Psi$ is the union of two Borel-complete closed subsets; by Theorem 2.4 in [3] $\beta\Psi$ is Borel-complete. This example already shows that the above conjecture is not valid.

Proof of Theorem. The necessity is clear since Borel-completeness is hereditary. Let now Y be a subset of a Borel-complete compact Hausdorff space K . Define $S := \beta\Psi \times K \setminus (\Psi^* \times Y)$. At first we show that $\beta S = \beta\Psi \times K$. Note that $\Psi \times K$ is a pseudocompact space. By Glicksberg's Theorem (see [6, p. 199]) $\Psi \times K$ is C^* -embedded in $\beta\Psi \times K$. Since S contains $\Psi \times K$ the space S is C^* embedded in $\beta\Psi \times K$ and therefore $\beta S = \beta\Psi \times K$. As a product of two Borel-complete spaces βS is Borel-complete. On the other side the space $\beta S \setminus S = \Psi^* \times Y$ is homeomorphic to Y . The proof is complete. \square

As pointed out in [3] it is unclear which topological spaces X admit a Borel-complete compact extension. By Theorem 3.1 in [3] every space with a perfectly normal compactification and by Corollary 2.5 every discrete space of non-measurable cardinality have this property.

References

- [1] *W. Adamski*: Complete spaces and zero-one measures. *Manuscripta Math.* 18 (1976), 343–352.
- [2] *R.L. Blair*: Closed-completeness in spaces with weak covering properties. In *Set-theoretic topology* (G.M. Reed, ed.). Academic Press, New York, 1977, pp. 17–45.
- [3] *A.W. Hager, G.D. Reynolds, M.D. Rice*: Borel-complete topological spaces. *Fund. Math.* 75 (1972), 135–143.
- [4] *S. Mrówka*: Some set-theoretic constructions in topology. *Fund. Math.* 94 (1977), 83–92.
- [5] *M.D. Rice, G.R. Reynolds*: Weakly Borel-complete topological spaces. *Fund. Math.* 105 (1980), 179–185.
- [6] *R.C. Walker*: *The Stone-Čech compactification*. Springer-Verlag, New York, 1974.

Author's address: Universität Duisburg, Fachbereich Mathematik, Lotharstr. 65, D-47057 Duisburg, Federal Republic of Germany.