

Applications of Mathematics

Sever Silvestru Dragomir

A sequence of mappings associated with the Hermite-Hadamard inequalities and applications

Applications of Mathematics, Vol. 49 (2004), No. 2, 123–140

Persistent URL: <http://dml.cz/dmlcz/134563>

Terms of use:

© Institute of Mathematics AS CR, 2004

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A SEQUENCE OF MAPPINGS ASSOCIATED WITH THE HERMITE-HADAMARD INEQUALITIES AND APPLICATIONS

SEVER S. DRAGOMIR, Melbourne City

(Received January 1, 2002, in revised version May 16, 2003)

Abstract. New properties for some sequences of functions defined by multiple integrals associated with the Hermite-Hadamard integral inequality for convex functions and some applications are given.

Keywords: Hermite-Hadamard inequality

MSC 2000: 26D15, 26D10, 26D99

1. INTRODUCTION

The integral inequality

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a)+f(b)}{2},$$

which holds for any convex function $f: [a, b] \rightarrow \mathbb{R}$, is well known in literature as the Hermite-Hadamard inequality.

There is an extensive amount of literature devoted to this simple and nice result which has many applications in the theory of special means and in information theory for divergence measures, from which we would like to refer the reader to [1]–[54].

The main aim of this paper is to consider some natural sequences of functions defined by multiple integrals and study their properties in relation to the Hermite-Hadamard inequality.

2. PROPERTIES OF THE SEQUENCE OF MAPPINGS H_n

Let $I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be an interval of real numbers and $a, b \in I$ with $a < b$, and let $f: I \rightarrow \mathbb{R}$ be a mapping which is integrable on $[a, b]$. Then we can define a sequence of mappings $H_n: [0, 1] \rightarrow \mathbb{R}$ by

$$H_n(t) := \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n$$

for $n \geq 1$ and $t \in [0, 1]$.

Some properties of this sequence of mappings are embodied in the following theorem.

Theorem 1. *Let $f: I \subseteq \mathbb{R}$ be a convex mapping on I and let $a, b \in I$ with $a < b$. Then*

- (i) H_n are convex on $[0, 1]$ for all $n \geq 1$;
- (ii) the following inequalities hold:

$$(2.1) \quad f\left(\frac{a+b}{2}\right) \leq H_n(t) \leq \frac{1}{(b-a)^{n+1}} \times \int_a^b \dots \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t)x_{n+1}\right) dx_1 \dots dx_{n+1}$$

and

$$(2.2) \quad H_n(t) \leq t \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n + (1-t)f\left(\frac{a+b}{2}\right) \leq \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n$$

for all $t \in [0, 1]$;

- (iii) the mapping H_n is monotonic nondecreasing on $[0, 1]$ for all $n \geq 1$ and one has the bounds

$$(2.3) \quad \min_{t \in [0,1]} H_n(t) = f\left(\frac{a+b}{2}\right) = H_n(0) \quad \text{for all } n \geq 1$$

and

$$(2.4) \quad \max_{t \in [0,1]} H_n(t) = \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n = H_n(1) \quad \text{for } n \geq 1.$$

Proof. (i) Follows by the convexity of f .

(ii) Applying Jensen's integral inequality, we obtain

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t)x_{n+1}\right) dx_{n+1} \\ & \geq f\left[\frac{1}{b-a} \int_a^b \left(t \frac{x_1 + \dots + x_n}{n} + (1-t)x_{n+1}\right) dx_{n+1}\right] \\ & = f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \end{aligned}$$

for all $x_1, \dots, x_n \in [a, b]$ and $t \in [0, 1]$.

Taking an integral mean over $[a, b]^n$ we deduce the second inequality in (2.1).

By Jensen's integral inequality for multiple integrals we have

$$\begin{aligned} & \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n \\ & \geq f\left[\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n\right] \\ & = f\left(\frac{a+b}{2}\right), \end{aligned}$$

and the inequality (2.1) is completely proved.

By the convexity of f on $[a, b]$, we can write

$$f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \leq t f\left(\frac{x_1 + \dots + x_n}{n}\right) + (1-t) f\left(\frac{a+b}{2}\right)$$

for all $x_1, \dots, x_n \in [a, b]$ and $t \in [0, 1]$. Taking an integral mean over $[a, b]^n$, we deduce

$$H_n(t) \leq t \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n + (1-t) f\left(\frac{a+b}{2}\right),$$

and the first inequality in (2.2) is proved.

As we know (see for example [26]) that

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n,$$

we obtain the last part of (2.2).

(iii) Let $0 < t_1 < t_2 \leq 1$. By the convexity of H_n , which follows by (i) now proved, we have that

$$\frac{H_n(t_2) - H_n(t_1)}{t_2 - t_1} \geq \frac{H_n(t_1) - H_n(0)}{t_1},$$

but $H_n(t_1) \geq H_n(0)$ (see the first inequality in (2.1)) and hence we get that $H_n(t_2) - H_n(t_1) \geq 0$ for all $0 \leq t_1 < t_2 \leq 1$, which shows that the mapping $H_n(\cdot)$ is monotonic nondecreasing on $[0, 1]$. The bounds (2.3) and (2.4) follow by the inequalities (2.1) and (2.2). We omit the details. \square

We now give another result on monotonicity which, in a sense, completes the above theorem.

Theorem 2. *Let $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping on I and let $a, b \in I$ with $a < b$. Then*

$$(2.5) \quad f\left(\frac{a+b}{2}\right) \leq H_{n+1}(t) \leq H_n(t) \leq \dots \leq H_1(t) = H(t),$$

where

$$H(t) := \frac{1}{b-a} \int_a^b f\left(tx + (1-t)\frac{a+b}{2}\right) dx$$

for all $n \geq 1$ and $t \in [0, 1]$. That is, the sequence of mappings $(H_n)_{n \geq 1}$ is monotonically nonincreasing.

Proof. Let us define real numbers belonging to $[a, b]$:

$$\begin{aligned} y_1 &:= t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}; \\ y_2 &:= t \frac{x_2 + x_3 + \dots + x_{n+1}}{n} + (1-t) \frac{a+b}{2}; \\ &\vdots \\ y_{n+1} &:= t \frac{x_{n+1} + x_1 + \dots + x_{n-1}}{n} + (1-t) \frac{a+b}{2}, \end{aligned}$$

where $x_1, \dots, x_{n+1} \in [a, b]$.

Using Jensen's discrete inequality, we may state that

$$\begin{aligned} &\frac{1}{n+1} \left[f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \right. \\ &\quad + f\left(t \frac{x_2 + \dots + x_{n+1}}{n} + (1-t) \frac{a+b}{2}\right) + \dots \\ &\quad \left. + f\left(t \frac{x_{n+1} + x_1 + \dots + x_{n-1}}{n} + (1-t) \frac{a+b}{2}\right) \right] \\ &\geq f\left(t \frac{x_1 + \dots + x_{n+1}}{n+1} + (1-t) \frac{a+b}{2}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_{n+1} \in [a, b]$.

Taking an integral mean over $[a, b]^{n+1}$, we deduce

$$\begin{aligned} & \frac{1}{n+1} \left[\frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1} \right. \\ & \quad \left. + \cdots + \frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_{n+1} + x_1 + \cdots + x_{n-1}}{n} \right. \right. \\ & \quad \quad \left. \left. + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1} \right] \\ & \geq \frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_{n+1}}{n+1} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1}. \end{aligned}$$

However, it is easy to see that

$$\begin{aligned} & \frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1} \\ & = \cdots = \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_n \end{aligned}$$

and thus, by the above inequality, we conclude

$$H_n(t) \geq H_{n+1}(t) \quad \text{for all } t \in [0, 1] \text{ and } n \geq 1.$$

The proof is thus completed. \square

It is natural to ask what happens with the difference $H_n(t) - f\left(\frac{1}{2}(a+b)\right)$ which is clearly non-negative for all $t \in [0, 1]$.

The following theorem contains an upper bound for this difference.

Theorem 3. *Let $f: I \subseteq \mathbb{R} \rightarrow R$ be a convex mapping and f'_+ its right derivative which exists on $\overset{\circ}{I}$ and is monotonic nondecreasing on $\overset{\circ}{I}$. If $a, b \in \overset{\circ}{I}$ with $a < b$, then the inequalities*

$$\begin{aligned} (2.6) \quad & 0 \leq H_n(t) - f\left(\frac{a+b}{2}\right) \\ & \leq \frac{t}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \\ & \quad \times \left(x_1 - \frac{a+b}{2} \right) dx_1 \cdots dx_n \\ & \leq \frac{t}{\sqrt{n}} \frac{b-a}{2\sqrt{3}} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} \right. \right. \right. \\ & \quad \left. \left. \left. + (1-t) \frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}} \end{aligned}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

Proof. As f is convex on I , we can write

$$f(x) - f(y) \geq f'_+(y)(x - y) \quad \text{for all } x, y \in I.$$

Choosing in this inequality

$$x = \frac{a+b}{2} \quad \text{and} \quad y = t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2},$$

we deduce the inequality

$$\begin{aligned} & f\left(\frac{a+b}{2}\right) - f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \\ & \geq t f'_+\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \left(\frac{a+b}{2} - \frac{x_1 + \dots + x_n}{n}\right). \end{aligned}$$

Taking an integral mean over $[a, b]^n$, we derive that

$$\begin{aligned} (2.7) \quad & f\left(\frac{a+b}{2}\right) - H_n(t) \\ & \geq t \left[\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \frac{a+b}{2} f'_+\left(t \frac{x_1 + \dots + x_n}{n} \right. \right. \\ & \qquad \qquad \qquad \left. \left. + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n \right. \\ & \quad \left. - \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \right. \\ & \qquad \qquad \qquad \left. \times \left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n \right], \end{aligned}$$

from where we deduce the second part of (2.6).

Now, let us observe that the right hand side in the inequality (2.7) is the integral

$$\begin{aligned} I = & - \frac{t}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \\ & \times \left(\frac{x_1 + \dots + x_n}{n} - \frac{a+b}{2}\right) dx_1 \dots dx_n. \end{aligned}$$

By the well-known Cauchy-Buniakowsky-Schwartz integral inequality for multiple integrals, we deduce the last part of the inequality (2.6).

The proof of the theorem is thus completed. □

Corollary 1. Let $f: I \subseteq \mathbb{R} \rightarrow R$ be a convex mapping and $a, b \in I$ with $a < b$. Put $M := \sup_{x \in [a, b]} |f'(x)| < \infty$. Then we have the inequality

$$(2.8) \quad 0 \leq H_n(t) - f\left(\frac{a+b}{2}\right) \leq \frac{t(b-a)M}{2\sqrt{3}\sqrt{n}}$$

for all $t \in [0, 1]$ and $n \geq 1$.

In particular, we have

$$\lim_{n \rightarrow \infty} H_n(t) = f\left(\frac{a+b}{2}\right) \quad \text{uniformly on } [0, 1].$$

The following result also holds:

Theorem 4. Under the assumptions as in Theorem 3, we have

$$(2.9) \quad \begin{aligned} & 0 \leq tH_n(1) + (1-t)H_n(0) - H_n(t) \\ & \leq \frac{t(1-t)}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \left(x_1 - \frac{a+b}{2}\right) dx_1 \cdots dx_n \\ & \leq \frac{t(1-t)(b-a)}{2\sqrt{3}\sqrt{n}} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}} \end{aligned}$$

for all $n \geq 1$ and $t \in [0, 1]$.

Proof. By the convexity of f we can write

$$(2.10) \quad \begin{aligned} & f\left(t\frac{x_1 + \cdots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f\left(\frac{x_1 + \cdots + x_n}{n}\right) \\ & \geq (1-t)f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \left(\frac{a+b}{2} - \frac{x_1 + \cdots + x_n}{n}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_n \in [a, b]$.

Similarly, we have

$$(2.11) \quad \begin{aligned} & f\left(t\frac{x_1 + \cdots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f\left(\frac{a+b}{2}\right) \\ & \geq -tf'_+\left(\frac{a+b}{2}\right) \left(\frac{a+b}{2} - \frac{x_1 + \cdots + x_n}{n}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_n \in [a, b]$.

If we multiply the inequality (2.10) by t and (2.11) by $(1-t)$ and add the obtained inequalities, we deduce

$$\begin{aligned} & f\left(t\frac{x_1 + \dots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - tf\left(\frac{x_1 + \dots + x_n}{n}\right) - (1-t)f\left(\frac{a+b}{2}\right) \\ & \geq t(1-t)\left[f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) - f'_+\left(\frac{a+b}{2}\right)\right]\left(\frac{a+b}{2} - \frac{x_1 + \dots + x_n}{n}\right). \end{aligned}$$

That is,

$$\begin{aligned} & tf\left(\frac{x_1 + \dots + x_n}{n}\right) + (1-t)f\left(\frac{a+b}{2}\right) - f\left(t\frac{x_1 + \dots + x_n}{n} + (1-t)\frac{a+b}{2}\right) \\ & \leq t(1-t)\left[f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) - f'_+\left(\frac{a+b}{2}\right)\right]\left(\frac{x_1 + \dots + x_n}{n} - \frac{a+b}{2}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_n \in [a, b]$.

Taking an integral mean over $[a, b]^n$, we have

$$\begin{aligned} 0 & \leq tH_n(1) + (1-t)H_n(0) - H_n(t) \\ & \leq t(1-t)\left[\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) x_1 \, dx_1 \dots dx_n \right. \\ & \quad \left. - \frac{a+b}{2} \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) \, dx_1 \dots dx_n\right], \end{aligned}$$

because a simple calculation shows us that

$$\begin{aligned} & \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) \left(\frac{x_1 + \dots + x_n}{n}\right) \, dx_1 \dots dx_n \\ & = \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) x_1 \, dx_1 \dots dx_n \end{aligned}$$

and

$$\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \frac{x_1 + \dots + x_n}{n} \, dx_1 \dots dx_n = \frac{a+b}{2}.$$

Thus, the second inequality in (2.9) is proved.

Now, applying the Cauchy-Buniakowsky-Schwartz integral inequality, we deduce the last part of the theorem. We omit the details. \square

Corollary 2. Under the assumptions as in Theorem 3 and provided $M := \sup_{x \in [a, b]} |f'(x)| < \infty$, we have the inequality

$$(2.12) \quad 0 \leq tH_n(1) + (1-t)H_n(0) - H_n(t) \leq \frac{t(1-t)}{2\sqrt{3}\sqrt{n}}M$$

for all $n \geq 1$ and $t \in [0, 1]$.

In particular,

$$\lim_{n \rightarrow \infty} [tH_n(1) + (1-t)H_n(0) - H_n(t)] = 0$$

uniformly on $[0, 1]$.

The following corollary is interesting as well.

Corollary 3. Under the assumptions as in Theorem 3 and provided there exists a constant $K > 0$ such that

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$(2.13) \quad 0 \leq tH_n(1) + (1-t)H_n(0) - H_n(t) \leq \frac{Kt(1-t)}{12n}(b-a)^2$$

for all $t \in [0, 1]$ and $n \geq 1$.

In addition, it is natural to ask for an upper bound for the difference $H_n(1) - H_n(t)$, $n \geq 1$ for all $t \in [0, 1]$.

Theorem 5. Let $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping on the interval I and let $a, b \in \overset{\circ}{I}$ with $a < b$. Then we have the inequalities

$$(2.14) \quad \begin{aligned} & \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f\left(\frac{x_1 + \cdots + x_n}{n}\right) dx_1 \cdots dx_n - H_n(t) \\ & \leq (1-t) \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \left(x_1 - \frac{a+b}{2}\right) dx_1 \cdots dx_n \\ & \leq \frac{(1-t)(b-a)}{2\sqrt{3}\sqrt{n}} \left(\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b [f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right)]^2 dx_1 \cdots dx_n \right)^{\frac{1}{2}} \end{aligned}$$

for all $t \in [0, 1]$ and $n \geq 1$.

Proof. By the convexity of f , we have that

$$\begin{aligned} & f\left(t\frac{x_1 + \dots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f\left(\frac{x_1 + \dots + x_n}{n}\right) \\ & \geq (1-t)f'_+\left(\frac{x_1 + \dots + x_n}{n}\right)\left[\frac{a+b}{2} - \frac{x_1 + \dots + x_n}{n}\right] \end{aligned}$$

for all $x_1, \dots, x_n \in [a, b]$ and $t \in [0, 1]$.

Now, the argument proceeds as above and we omit the details. \square

Corollary 4. *Under the assumptions as in Theorem 3 and provided $M := \sup_{x \in [a, b]} |f'(x)| < \infty$, we have the inequality*

$$(2.15) \quad 0 \leq H_n(1) - H_n(t) \leq \frac{(1-t)(b-a)M}{2\sqrt{3}\sqrt{n}}.$$

In particular,

$$\lim_{n \rightarrow \infty} [H_n(1) - H_n(t)] = 0$$

uniformly on $[0, 1]$.

Corollary 5. *Under the assumptions as in Theorem 3 and provided there exists a constant $K > 0$ such that*

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$(2.16) \quad 0 \leq H_n(1) - H_n(t) \leq \frac{K(1-t)(b-a)^2}{12n}$$

for all $n \geq 1$ and $t \in [0, 1]$.

Now we establish an upper bound for the difference $H_n(t) - H_{n+1}(t)$, $n \geq 1$ which is non-negative for all $t \in [0, 1]$ (cf. Theorem 2):

Theorem 6. Let $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function on the interval I and let $a, b \in \overset{\circ}{I}$ with $a < b$. Then the inequality

$$\begin{aligned}
 (2.17) \quad 0 &\leq H_n(t) - H_{n+1}(t) \\
 &\leq \frac{t}{n+1} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \right. \\
 &\quad \left. \times \left(x_1 - \frac{a+b}{2} \right) dx_1 \cdots dx_n \right] \\
 &\leq \frac{t(b-a)}{2\sqrt{3}\sqrt{n}(n+1)} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} \right. \right. \right. \\
 &\quad \left. \left. \left. + (1-t) \frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}}
 \end{aligned}$$

holds for all $t \in [0, 1]$ and $n \geq 1$.

Proof. By the convexity of f , we have that

$$\begin{aligned}
 &f \left(t \frac{x_1 + \cdots + x_{n+1}}{n+1} + (1-t) \frac{a+b}{2} \right) - f \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \\
 &\geq \frac{t}{n(n+1)} f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) [nx_{n+1} - (x_1 + \cdots + x_n)]
 \end{aligned}$$

for all $x_1, \dots, x_{n+1} \in [a, b]$ and $t \in [0, 1]$.

Taking an integral mean on $[a, b]^{n+1}$, we derive

$$\begin{aligned}
 0 &\leq H_n(t) - H_{n+1}(t) \\
 &\leq \frac{t}{n+1} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) x_1 dx_1 \cdots dx_n \right. \\
 &\quad \left. - \frac{a+b}{2} \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_n \right],
 \end{aligned}$$

and the second inequality in (2.17) is proved.

Now, let us observe that

$$\begin{aligned}
& \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \\
& \qquad \qquad \qquad \times \left(x_1 - \frac{a+b}{2} \right) dx_1 \cdots dx_n \\
& = \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \\
& \qquad \qquad \qquad \times \left(\frac{x_1 + \cdots + x_n}{n} - \frac{a+b}{2} \right) dx_1 \cdots dx_n \\
& \leq \left(\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right)^{\frac{1}{2}} \\
& \qquad \times \left(\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left(\frac{x_1 + \cdots + x_n}{n} - \frac{a+b}{2} \right)^2 dx_1 \cdots dx_n \right)^{\frac{1}{2}},
\end{aligned}$$

and thus the last inequality is also proved. \square

Corollary 6. Under the assumptions as in Theorem 3, given that $M := \sup_{x \in [a,b]} |f'(x)| < \infty$, we have

$$(2.18) \quad 0 \leq H_n(t) - H_{n+1}(t) \leq \frac{Mt(b-a)}{2\sqrt{3}\sqrt{n}(n+1)}$$

for all $t \in [0, 1]$ and $n \geq 1$.

In particular,

$$\lim_{n \rightarrow \infty} [H_n(t) - H_{n+1}(t)] = 0$$

uniformly on $[0, 1]$.

The following theorem also holds.

Theorem 7. Under the assumptions as in Theorem 3, we also have the bound

$$\begin{aligned}
(2.19) \quad 0 \leq H_n(t) - H_{n+1}(t) \\
\leq \frac{t(b-a)}{2\sqrt{3}\sqrt{n}(n+1)} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \right. \right. \\
\left. \left. - f'_+ \left(\frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}}
\end{aligned}$$

for all $t \in [0, 1]$ and $n \geq 1$.

Proof. The proof is similar to the one of Theorem 6 and we omit the details. \square

Corollary 7. Under the assumptions as in Theorem 3, given that there exists a $K > 0$ such that

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$(2.20) \quad 0 \leq H_n(t) - H_{n+1}(t) \leq \frac{t^2(b-a)^2 K^2}{12n(n+1)}$$

for all $t \in [0, 1]$ and $n \geq 1$.

Finally, note that, by a similar argument to that in the proof of Theorem 6, we can give the following result which completes, in a sense, the estimate in Theorem 3.

Theorem 8. Under the above assumptions the inequality

$$(2.21) \quad 0 \leq H_n(t) - f\left(\frac{a+b}{2}\right) \leq \frac{t(b-a)}{2\sqrt{3}\sqrt{n}} \left[\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \left[f'_+\left(t\frac{x_1 + \dots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f'_+\left(\frac{a+b}{2}\right) \right]^2 dx_1 \dots dx_n \right]^{\frac{1}{2}}$$

holds for all $t \in [0, 1]$ and $n \geq 1$.

Corollary 8. Under the above assumptions, given that there exists a $K > 0$ such that

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$0 \leq H_n(t) - f\left(\frac{a+b}{2}\right) \leq \frac{t^2(b-a)^2 K^2}{12n}$$

for all $t \in [0, 1]$ and $n \geq 1$.

3. APPLICATIONS TO SPECIAL MEANS

Let $0 \leq a < b$ and $p \in (-\infty, 0) \cup [1, \infty) \setminus \{-1\}$. Let us define sequence of mappings

$$h_{p,n}(t) := \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2} \right)^p dx_1 \dots dx_n,$$

where $n \geq 1$, $n \in \mathbb{N}$, and $t \in [0, 1]$.

By virtue of the above results, we can establish the following properties:

- (i) $h_{p,n}(t)$ are convex and monotonic nondecreasing on $[0, 1]$;
- (ii) $h_{p,n}(t) \geq h_{p,n+1}(t)$ for all $n \geq 1$ and $t \in [0, 1]$;
- (iii) the inequalities

$$\begin{aligned} [A(a, b)]^p &\leq h_{p,n}(t) \\ &\leq \frac{1}{(b-a)^{n+1}} \int_a^b \dots \int_a^b \left(t \frac{x_1 + \dots + x_n}{n} + (1-t)x_{n+1} \right)^p dx_1 \dots dx_{n+1} \end{aligned}$$

and

$$\begin{aligned} h_{p,n}(t) &\leq t \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \left(\frac{x_1 + \dots + x_n}{n} \right)^p dx_1 \dots dx_n \\ &\quad + (1-t)[A(a, b)]^p \\ &\leq \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \left(\frac{x_1 + \dots + x_n}{n} \right)^p dx_1 \dots dx_n \end{aligned}$$

hold for all $n \geq 1$ and $t \in [0, 1]$, where $A(a, b) := \frac{a+b}{2}$.

(iv) If $p \geq 1$, then the inequalities

$$0 \leq h_{p,n}(t) - [A(a, b)]^p \leq \frac{t(b-a)pb^{p-1}}{2\sqrt{3}\sqrt{n}}$$

and

$$\begin{aligned} 0 &\leq \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \left(\frac{x_1 + \dots + x_n}{n} \right)^p dx_1 \dots dx_n - h_{p,n}(t) \\ &\leq \frac{(1-t)(b-a)pb^{p-1}}{2\sqrt{3}\sqrt{n}} \end{aligned}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

(v) If $p \geq 1$, the inequalities

$$0 \leq th_{p,n}(1) + (1-t)h_{p,n}(0) - h_{p,n}(t) \leq \frac{t(1-t)pb^{p-1}}{2\sqrt{3}\sqrt{n}}$$

and

$$0 \leq h_{p,n}(t) - h_{p,n+1}(t) \leq \frac{t(b-a)pb^{p-1}}{2\sqrt{3}\sqrt{n}(n+1)}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

(vi) If $p \geq 2$, then the inequalities

$$0 \leq h_{p,n}(t) - [A(a, b)]^p \leq \frac{t^2(b-a)^2p(p-1)b^{p-2}}{12n}$$

and

$$0 \leq h_{p,n}(t) - h_{p,n+1}(t) \leq \frac{t^2(b-a)^2p(p-1)b^{p-2}}{12n(n+1)}$$

and

$$\begin{aligned} 0 &\leq th_{p,n}(1) + (1-t)h_{p,n}(0) - h_{p,n}(t) \\ &\leq \frac{t(1-t)(b-a)^2p(p-1)b^{p-2}}{12n} \end{aligned}$$

and

$$0 \leq h_{p,n}(1) - h_{p,n}(t) \leq \frac{(1-t)(b-a)^2p(p-1)b^{p-2}}{12n}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

Acknowledgement. The author would like to thank the anonymous referee for many valuable comments and remarks that have been incorporated in the final version of the paper.

References

- [1] G. Allasia, C. Giordano, J. Pečarić: Hadamard-type inequalities for $(2r)$ -convex functions with applications. *Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur.* 133 (1999), 187–200.
- [2] H. Alzer: A note on Hadamard's inequalities. *C. R. Math. Rep. Acad. Sci. Canada* 11 (1989), 255–258.
- [3] H. Alzer: On an integral inequality. *Anal. Numér. Théor. Approx.* 18 (1989), 101–103.

- [4] *A. G. Azpeitia*: Convex functions and the Hadamard inequality. *Rev. Colombiana Mat.* **28** (1994), 7–12.
- [5] *D. Barbu, S. S. Dragomir and C. Buşu*: A probabilistic argument for the convergence of some sequences associated to Hadamard's inequality. *Studia Univ. Babeş-Bolyai Math.* **38** (1993), 29–33.
- [6] *C. Buşu, S. S. Dragomir and D. Barbu*: The convergence of some sequences connected to Hadamard's inequality. *Demonstratio Math.* **29** (1996), 53–59.
- [7] *S. S. Dragomir*: A mapping in connection to Hadamard's inequalities. *Anz. Österreich. Akad. Wiss. Math.-Natur. Kl.* **128** (1991), 17–20.
- [8] *S. S. Dragomir*: A refinement of Hadamard's inequality for isotonic linear functionals. *Tamkang J. Math.* **24** (1993), 101–106.
- [9] *S. S. Dragomir*: On Hadamard's inequalities for convex functions. *Mat. Balkanica (N. S.)* **6** (1992), 215–222.
- [10] *S. S. Dragomir*: On Hadamard's inequality for the convex mappings defined on a ball in the space and applications. *Math. Inequal. Appl.* **3** (2000), 177–187.
- [11] *S. S. Dragomir*: On Hadamard's inequality on a disk. *JIPAM. J. Inequal. Pure Appl. Math.* **1** (2000), <http://jipam.vu.edu.au/>. Electronic.
- [12] *S. S. Dragomir*: Some integral inequalities for differentiable convex functions. *Makedon. Akad. Nauk Umet. Oddel. Mat.-Tehn. Nauk. Prilozi* **13** (1992), 13–17.
- [13] *S. S. Dragomir*: Some remarks on Hadamard's inequalities for convex functions. *Extracta Math.* **9** (1994), 88–94.
- [14] *S. S. Dragomir*: Two mappings in connection to Hadamard's inequalities. *J. Math. Anal. Appl.* **167** (1992), 49–56.
- [15] *S. S. Dragomir, R. P. Agarwal*: Two new mappings associated with Hadamard's inequalities for convex functions. *Appl. Math. Lett.* **11** (1998), 33–38.
- [16] *S. S. Dragomir, C. Buşu*: Refinements of Hadamard's inequality for multiple integrals. *Utilitas Math.* **47** (1995), 193–198.
- [17] *S. S. Dragomir, Y. J. Cho and S. S. Kim*: Inequalities of Hadamard's type for Lipschitzian mappings and their applications. *J. Math. Anal. Appl.* **245** (2000), 489–501.
- [18] *S. S. Dragomir, S. Fitzpatrick*: Hadamard inequality for s -convex functions in the first sense and applications. *Demonstratio Math.* **31** (1998), 633–642.
- [19] *S. S. Dragomir, S. Fitzpatrick*: The Hadamard's inequality for s -convex functions in the second sense. *Demonstratio Math.* **32** (1999), 687–696.
- [20] *S. S. Dragomir, N. M. Ionescu*: On some inequalities for convex-dominated functions. *Anal. Numér. Théor. Approx.* **19** (1990), 21–27.
- [21] *S. S. Dragomir, D. S. Milošević and J. Sándor*: On some refinements of Hadamard's inequalities and applications. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.* **4** (1993), 3–10.
- [22] *S. S. Dragomir, B. Mond*: On Hadamard's inequality for a class of functions of Godunova and Levin. *Indian J. Math.* **39** (1997), 1–9.
- [23] *S. S. Dragomir, C. E. M. Pearce*: Quasi-convex functions and Hadamard's inequality. *Bull. Austral. Math. Soc.* **57** (1998), 377–385.
- [24] *S. S. Dragomir, C. E. M. Pearce, and J. E. Pečarić*: On Jensen's related inequalities for isotonic sublinear functionals. *Acta Sci. Math.* **61** (1995), 373–382.
- [25] *S. S. Dragomir, J. E. Pečarić, and L. E. Persson*: Some inequalities of Hadamard type. *Soochow J. Math.* **21** (1995), 335–341.
- [26] *S. S. Dragomir, J. E. Pečarić, and J. Sándor*: A note on the Jensen-Hadamard inequality. *Anal. Numér. Théor. Approx.* **19** (1990), 29–34.
- [27] *S. S. Dragomir, G. H. Toader*: Some inequalities for m -convex functions. *Studia Univ. Babeş-Bolyai Math.* **38** (1993), 21–28.

- [28] *A. M. Fink*: A best possible Hadamard inequality. *Math. Inequal. Appl.* *1* (1998), 223–230.
- [29] *A. M. Fink*: Toward a theory of best possible inequalities. *Nieuw Arch. Wisk.* *12* (1994), 19–29.
- [30] *A. M. Fink*: Two inequalities. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.* *6* (1995), 48–49.
- [31] *B. Gavrea*: On Hadamard's inequality for the convex mappings defined on a convex domain in the space. *JIPAM. J. Inequal. Pure Appl. Math.* *1* (2000), Article 9, <http://jipam.vu.edu.au/>. Electronic.
- [32] *P. M. Gill, C. E. M. Pearce and J. Pečarić*: Hadamard's inequality for r -convex functions. *J. Math. Anal. Appl.* *215* (1997), 461–470.
- [33] *G. H. Hardy, J. E. Littlewood, and G. Pólya*: *Inequalities*. 2nd ed. Cambridge University Press, 1952.
- [34] *K.-C. Lee, K.-L. Tseng*: On weighted generalization of Hadamard's inequality for g functions. *Tamsui Oxf. J. Math. Sci.* *16* (2000), 91–104.
- [35] *A. Lupaş*: The Jensen-Hadamard inequality for convex functions of higher order. *Octagon Math. Mag.* *5* (1997), 8–9.
- [36] *A. Lupaş*: A generalization of Hadamard's inequality for convex functions. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* *544–576* (1976), 115–121.
- [37] *D. M. Maksimović*: A short proof of generalized Hadamard's inequalities. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* *634–677* (1979), 126–128.
- [38] *D. S. Mitrinović, I. Lacković*: Hermite and convexity. *Aequationes Math.* *28* (1985), 229–232.
- [39] *D. S. Mitrinović, J. E. Pečarić, and A. M. Fink*: *Classical and New Inequalities in Analysis*. Kluwer Academic Publishers, Dordrecht, 1993.
- [40] *E. Neuman*: Inequalities involving generalized symmetric means. *J. Math. Anal. Appl.* *120* (1986), 315–320.
- [41] *E. Neuman, J. E. Pečarić*: Inequalities involving multivariate convex functions. *J. Math. Anal. Appl.* *137* (1989), 514–549.
- [42] *E. Neuman*: Inequalities involving multivariate convex functions. II. *Proc. Amer. Math. Soc.* *109* (1990), 965–974.
- [43] *C. P. Niculescu*: A note on the dual Hermite-Hadamard inequality. *The Math. Gazette* (July 2000).
- [44] *C. P. Niculescu*: Convexity according to the geometric mean. *Math. Inequal. Appl.* *3* (2000), 155–167.
- [45] *C. E. M. Pearce, J. Pečarić, and V. Šimić*: Stolarsky means and Hadamard's inequality. *J. Math. Anal. Appl.* *220* (1998), 99–109.
- [46] *C. E. M. Pearce, A. M. Rubinov*: P -functions, quasi-convex functions and Hadamard-type inequalities. *J. Math. Anal. Appl.* *240* (1999), 92–104.
- [47] *J. E. Pečarić*: Remarks on two interpolations of Hadamard's inequalities. *Makedon. Akad. Nauk. Umet. Oddel. Mat.-Tehn. Nauk. Prilozi* *13* (1992), 9–12.
- [48] *J. Pečarić, S. S. Dragomir*: A generalization of Hadamard's inequality for isotonic linear functionals. *Rad. Mat.* *7* (1991), 103–107.
- [49] *J. Pečarić, F. Proschan, and Y. L. Tong*: *Convex Functions, Partial Orderings and Statistical Applications*. Academic Press, Boston, 1992.
- [50] *J. Sándor*: An application of the Jensen-Hadamard inequality. *Nieuw Arch. Wisk.* *8* (1990), 63–66.
- [51] *J. Sándor*: On the Jensen-Hadamard inequality. *Studia Univ. Babeş-Bolyai, Math.* *36* (1991), 9–15.

- [52] *P. M. Vasić, I. B. Lacković, and D. M. Maksimović*: Note on convex functions. IV. On Hadamard's inequality for weighted arithmetic means. Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. 678–715 (1980), 199–205.
- [53] *G. S. Yang, M. C. Hong*: A note on Hadamard's inequality. Tamkang J. Math. 28 (1997), 33–37.
- [54] *G. S. Yang, K. L. Tseng*: On certain integral inequalities related to Hermite-Hadamard inequalities. J. Math. Anal. Appl. 239 (1999), 180–187.

Author's address: S. S. Dragomir, School of Computer Science & Mathematics, Victoria University, P.O. Box 14428, Melbourne City, MC 8001, Australia, e-mail: **sever**
matilda.vu.edu.au.

A SEQUENCE OF MAPPINGS ASSOCIATED WITH THE HERMITE-HADAMARD INEQUALITIES AND APPLICATIONS

SEVER S. DRAGOMIR, Melbourne City

(Received January 1, 2002, in revised version May 16, 2003)

Abstract. New properties for some sequences of functions defined by multiple integrals associated with the Hermite-Hadamard integral inequality for convex functions and some applications are given.

Keywords: Hermite-Hadamard inequality

MSC 2000: 26D15, 26D10, 26D99

1. INTRODUCTION

The integral inequality

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a)+f(b)}{2},$$

which holds for any convex function $f: [a, b] \rightarrow \mathbb{R}$, is well known in literature as the Hermite-Hadamard inequality.

There is an extensive amount of literature devoted to this simple and nice result which has many applications in the theory of special means and in information theory for divergence measures, from which we would like to refer the reader to [1]–[54].

The main aim of this paper is to consider some natural sequences of functions defined by multiple integrals and study their properties in relation to the Hermite-Hadamard inequality.

2. PROPERTIES OF THE SEQUENCE OF MAPPINGS H_n

Let $I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be an interval of real numbers and $a, b \in I$ with $a < b$, and let $f: I \rightarrow \mathbb{R}$ be a mapping which is integrable on $[a, b]$. Then we can define a sequence of mappings $H_n: [0, 1] \rightarrow \mathbb{R}$ by

$$H_n(t) := \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n$$

for $n \geq 1$ and $t \in [0, 1]$.

Some properties of this sequence of mappings are embodied in the following theorem.

Theorem 1. *Let $f: I \subseteq \mathbb{R}$ be a convex mapping on I and let $a, b \in I$ with $a < b$. Then*

- (i) H_n are convex on $[0, 1]$ for all $n \geq 1$;
- (ii) the following inequalities hold:

$$(2.1) \quad f\left(\frac{a+b}{2}\right) \leq H_n(t) \leq \frac{1}{(b-a)^{n+1}} \times \int_a^b \dots \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t)x_{n+1}\right) dx_1 \dots dx_{n+1}$$

and

$$(2.2) \quad H_n(t) \leq t \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n + (1-t)f\left(\frac{a+b}{2}\right) \leq \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n$$

for all $t \in [0, 1]$;

- (iii) the mapping H_n is monotonic nondecreasing on $[0, 1]$ for all $n \geq 1$ and one has the bounds

$$(2.3) \quad \min_{t \in [0,1]} H_n(t) = f\left(\frac{a+b}{2}\right) = H_n(0) \quad \text{for all } n \geq 1$$

and

$$(2.4) \quad \max_{t \in [0,1]} H_n(t) = \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n = H_n(1) \quad \text{for } n \geq 1.$$

Proof. (i) Follows by the convexity of f .

(ii) Applying Jensen's integral inequality, we obtain

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t)x_{n+1}\right) dx_{n+1} \\ & \geq f\left[\frac{1}{b-a} \int_a^b \left(t \frac{x_1 + \dots + x_n}{n} + (1-t)x_{n+1}\right) dx_{n+1}\right] \\ & = f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \end{aligned}$$

for all $x_1, \dots, x_n \in [a, b]$ and $t \in [0, 1]$.

Taking an integral mean over $[a, b]^n$ we deduce the second inequality in (2.1).

By Jensen's integral inequality for multiple integrals we have

$$\begin{aligned} & \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n \\ & \geq f\left[\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n\right] \\ & = f\left(\frac{a+b}{2}\right), \end{aligned}$$

and the inequality (2.1) is completely proved.

By the convexity of f on $[a, b]$, we can write

$$f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \leq t f\left(\frac{x_1 + \dots + x_n}{n}\right) + (1-t) f\left(\frac{a+b}{2}\right)$$

for all $x_1, \dots, x_n \in [a, b]$ and $t \in [0, 1]$. Taking an integral mean over $[a, b]^n$, we deduce

$$H_n(t) \leq t \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n + (1-t) f\left(\frac{a+b}{2}\right),$$

and the first inequality in (2.2) is proved.

As we know (see for example [26]) that

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n,$$

we obtain the last part of (2.2).

(iii) Let $0 < t_1 < t_2 \leq 1$. By the convexity of H_n , which follows by (i) now proved, we have that

$$\frac{H_n(t_2) - H_n(t_1)}{t_2 - t_1} \geq \frac{H_n(t_1) - H_n(0)}{t_1},$$

but $H_n(t_1) \geq H_n(0)$ (see the first inequality in (2.1)) and hence we get that $H_n(t_2) - H_n(t_1) \geq 0$ for all $0 \leq t_1 < t_2 \leq 1$, which shows that the mapping $H_n(\cdot)$ is monotonic nondecreasing on $[0, 1]$. The bounds (2.3) and (2.4) follow by the inequalities (2.1) and (2.2). We omit the details. \square

We now give another result on monotonicity which, in a sense, completes the above theorem.

Theorem 2. *Let $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping on I and let $a, b \in I$ with $a < b$. Then*

$$(2.5) \quad f\left(\frac{a+b}{2}\right) \leq H_{n+1}(t) \leq H_n(t) \leq \dots \leq H_1(t) = H(t),$$

where

$$H(t) := \frac{1}{b-a} \int_a^b f\left(tx + (1-t)\frac{a+b}{2}\right) dx$$

for all $n \geq 1$ and $t \in [0, 1]$. That is, the sequence of mappings $(H_n)_{n \geq 1}$ is monotonically nonincreasing.

Proof. Let us define real numbers belonging to $[a, b]$:

$$\begin{aligned} y_1 &:= t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}; \\ y_2 &:= t \frac{x_2 + x_3 + \dots + x_{n+1}}{n} + (1-t) \frac{a+b}{2}; \\ &\vdots \\ y_{n+1} &:= t \frac{x_{n+1} + x_1 + \dots + x_{n-1}}{n} + (1-t) \frac{a+b}{2}, \end{aligned}$$

where $x_1, \dots, x_{n+1} \in [a, b]$.

Using Jensen's discrete inequality, we may state that

$$\begin{aligned} &\frac{1}{n+1} \left[f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \right. \\ &\quad + f\left(t \frac{x_2 + \dots + x_{n+1}}{n} + (1-t) \frac{a+b}{2}\right) + \dots \\ &\quad \left. + f\left(t \frac{x_{n+1} + x_1 + \dots + x_{n-1}}{n} + (1-t) \frac{a+b}{2}\right) \right] \\ &\geq f\left(t \frac{x_1 + \dots + x_{n+1}}{n+1} + (1-t) \frac{a+b}{2}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_{n+1} \in [a, b]$.

Taking an integral mean over $[a, b]^{n+1}$, we deduce

$$\begin{aligned} & \frac{1}{n+1} \left[\frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1} \right. \\ & \quad + \cdots + \frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_{n+1} + x_1 + \cdots + x_{n-1}}{n} \right. \\ & \quad \quad \quad \left. \left. + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1} \right] \\ & \geq \frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_{n+1}}{n+1} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1}. \end{aligned}$$

However, it is easy to see that

$$\begin{aligned} & \frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_{n+1} \\ & = \cdots = \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_n \end{aligned}$$

and thus, by the above inequality, we conclude

$$H_n(t) \geq H_{n+1}(t) \quad \text{for all } t \in [0, 1] \text{ and } n \geq 1.$$

The proof is thus completed. \square

It is natural to ask what happens with the difference $H_n(t) - f(\frac{1}{2}(a+b))$ which is clearly non-negative for all $t \in [0, 1]$.

The following theorem contains an upper bound for this difference.

Theorem 3. *Let $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping and f'_+ its right derivative which exists on $\overset{\circ}{I}$ and is monotonic nondecreasing on $\overset{\circ}{I}$. If $a, b \in \overset{\circ}{I}$ with $a < b$, then the inequalities*

$$\begin{aligned} (2.6) \quad & 0 \leq H_n(t) - f \left(\frac{a+b}{2} \right) \\ & \leq \frac{t}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \\ & \quad \quad \quad \times \left(x_1 - \frac{a+b}{2} \right) dx_1 \cdots dx_n \\ & \leq \frac{t}{\sqrt{n}} \frac{b-a}{2\sqrt{3}} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} \right. \right. \right. \\ & \quad \quad \quad \left. \left. \left. + (1-t) \frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}} \end{aligned}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

Proof. As f is convex on I , we can write

$$f(x) - f(y) \geq f'_+(y)(x - y) \quad \text{for all } x, y \in \overset{\circ}{I}.$$

Choosing in this inequality

$$x = \frac{a+b}{2} \quad \text{and} \quad y = t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2},$$

we deduce the inequality

$$\begin{aligned} & f\left(\frac{a+b}{2}\right) - f\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \\ & \geq t f'_+\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \left(\frac{a+b}{2} - \frac{x_1 + \dots + x_n}{n}\right). \end{aligned}$$

Taking an integral mean over $[a, b]^n$, we derive that

$$\begin{aligned} (2.7) \quad & f\left(\frac{a+b}{2}\right) - H_n(t) \\ & \geq t \left[\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \frac{a+b}{2} f'_+\left(t \frac{x_1 + \dots + x_n}{n} \right. \right. \\ & \qquad \qquad \qquad \left. \left. + (1-t) \frac{a+b}{2}\right) dx_1 \dots dx_n \right. \\ & \quad \left. - \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \right. \\ & \qquad \qquad \qquad \left. \times \left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n \right], \end{aligned}$$

from where we deduce the second part of (2.6).

Now, let us observe that the right hand side in the inequality (2.7) is the integral

$$\begin{aligned} I = & - \frac{t}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(t \frac{x_1 + \dots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \\ & \times \left(\frac{x_1 + \dots + x_n}{n} - \frac{a+b}{2}\right) dx_1 \dots dx_n. \end{aligned}$$

By the well-known Cauchy-Buniakowsky-Schwartz integral inequality for multiple integrals, we deduce the last part of the inequality (2.6).

The proof of the theorem is thus completed. □

Corollary 1. Let $f: I \subseteq \mathbb{R} \rightarrow R$ be a convex mapping and $a, b \in \overset{\circ}{I}$ with $a < b$. Put $M := \sup_{x \in [a, b]} |f'(x)| < \infty$. Then we have the inequality

$$(2.8) \quad 0 \leq H_n(t) - f\left(\frac{a+b}{2}\right) \leq \frac{t(b-a)M}{2\sqrt{3}\sqrt{n}}$$

for all $t \in [0, 1]$ and $n \geq 1$.

In particular, we have

$$\lim_{n \rightarrow \infty} H_n(t) = f\left(\frac{a+b}{2}\right) \quad \text{uniformly on } [0, 1].$$

The following result also holds:

Theorem 4. Under the assumptions as in Theorem 3, we have

$$(2.9) \quad \begin{aligned} 0 &\leq tH_n(1) + (1-t)H_n(0) - H_n(t) \\ &\leq \frac{t(1-t)}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \left(x_1 - \frac{a+b}{2}\right) dx_1 \cdots dx_n \\ &\leq \frac{t(1-t)(b-a)}{2\sqrt{3}\sqrt{n}} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}} \end{aligned}$$

for all $n \geq 1$ and $t \in [0, 1]$.

Proof. By the convexity of f we can write

$$(2.10) \quad \begin{aligned} &f\left(t\frac{x_1 + \cdots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f\left(\frac{x_1 + \cdots + x_n}{n}\right) \\ &\geq (1-t)f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right)\left(\frac{a+b}{2} - \frac{x_1 + \cdots + x_n}{n}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_n \in [a, b]$.

Similarly, we have

$$(2.11) \quad \begin{aligned} &f\left(t\frac{x_1 + \cdots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f\left(\frac{a+b}{2}\right) \\ &\geq -tf'_+\left(\frac{a+b}{2}\right)\left(\frac{a+b}{2} - \frac{x_1 + \cdots + x_n}{n}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_n \in [a, b]$.

If we multiply the inequality (2.10) by t and (2.11) by $(1-t)$ and add the obtained inequalities, we deduce

$$\begin{aligned} & f\left(t\frac{x_1 + \dots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - tf\left(\frac{x_1 + \dots + x_n}{n}\right) - (1-t)f\left(\frac{a+b}{2}\right) \\ & \geq t(1-t)\left[f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) - f'_+\left(\frac{a+b}{2}\right)\right]\left(\frac{a+b}{2} - \frac{x_1 + \dots + x_n}{n}\right). \end{aligned}$$

That is,

$$\begin{aligned} & tf\left(\frac{x_1 + \dots + x_n}{n}\right) + (1-t)f\left(\frac{a+b}{2}\right) - f\left(t\frac{x_1 + \dots + x_n}{n} + (1-t)\frac{a+b}{2}\right) \\ & \leq t(1-t)\left[f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) - f'_+\left(\frac{a+b}{2}\right)\right]\left(\frac{x_1 + \dots + x_n}{n} - \frac{a+b}{2}\right) \end{aligned}$$

for all $t \in [0, 1]$ and $x_1, \dots, x_n \in [a, b]$.

Taking an integral mean over $[a, b]^n$, we have

$$\begin{aligned} 0 & \leq tH_n(1) + (1-t)H_n(0) - H_n(t) \\ & \leq t(1-t)\left[\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) x_1 dx_1 \dots dx_n \right. \\ & \quad \left. - \frac{a+b}{2} \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n \right], \end{aligned}$$

because a simple calculation shows us that

$$\begin{aligned} & \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) \left(\frac{x_1 + \dots + x_n}{n}\right) dx_1 \dots dx_n \\ & = \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f'_+\left(\frac{x_1 + \dots + x_n}{n}\right) x_1 dx_1 \dots dx_n \end{aligned}$$

and

$$\frac{1}{(b-a)^n} \int_a^b \dots \int_a^b \frac{x_1 + \dots + x_n}{n} dx_1 \dots dx_n = \frac{a+b}{2}.$$

Thus, the second inequality in (2.9) is proved.

Now, applying the Cauchy-Buniakowsky-Schwartz integral inequality, we deduce the last part of the theorem. We omit the details. \square

Corollary 2. Under the assumptions as in Theorem 3 and provided $M := \sup_{x \in [a, b]} |f'(x)| < \infty$, we have the inequality

$$(2.12) \quad 0 \leq tH_n(1) + (1-t)H_n(0) - H_n(t) \leq \frac{t(1-t)}{2\sqrt{3}\sqrt{n}}M$$

for all $n \geq 1$ and $t \in [0, 1]$.

In particular,

$$\lim_{n \rightarrow \infty} [tH_n(1) + (1-t)H_n(0) - H_n(t)] = 0$$

uniformly on $[0, 1]$.

The following corollary is interesting as well.

Corollary 3. Under the assumptions as in Theorem 3 and provided there exists a constant $K > 0$ such that

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$(2.13) \quad 0 \leq tH_n(1) + (1-t)H_n(0) - H_n(t) \leq \frac{Kt(1-t)}{12n}(b-a)^2$$

for all $t \in [0, 1]$ and $n \geq 1$.

In addition, it is natural to ask for an upper bound for the difference $H_n(1) - H_n(t)$, $n \geq 1$ for all $t \in [0, 1]$.

Theorem 5. Let $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping on the interval I and let $a, b \in \overset{\circ}{I}$ with $a < b$. Then we have the inequalities

$$(2.14) \quad \begin{aligned} & \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f\left(\frac{x_1 + \cdots + x_n}{n}\right) dx_1 \cdots dx_n - H_n(t) \\ & \leq (1-t) \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \left(x_1 - \frac{a+b}{2}\right) dx_1 \cdots dx_n \\ & \leq \frac{(1-t)(b-a)}{2\sqrt{3}\sqrt{n}} \left(\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+\left(\frac{x_1 + \cdots + x_n}{n}\right) \right]^2 dx_1 \cdots dx_n \right)^{\frac{1}{2}} \end{aligned}$$

for all $t \in [0, 1]$ and $n \geq 1$.

P r o o f. By the convexity of f , we have that

$$\begin{aligned} & f\left(t\frac{x_1 + \dots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f\left(\frac{x_1 + \dots + x_n}{n}\right) \\ & \geq (1-t)f'_+\left(\frac{x_1 + \dots + x_n}{n}\right)\left[\frac{a+b}{2} - \frac{x_1 + \dots + x_n}{n}\right] \end{aligned}$$

for all $x_1, \dots, x_n \in [a, b]$ and $t \in [0, 1]$.

Now, the argument proceeds as above and we omit the details. \square

Corollary 4. *Under the assumptions as in Theorem 3 and provided $M := \sup_{x \in [a, b]} |f'(x)| < \infty$, we have the inequality*

$$(2.15) \quad 0 \leq H_n(1) - H_n(t) \leq \frac{(1-t)(b-a)M}{2\sqrt{3}\sqrt{n}}.$$

In particular,

$$\lim_{n \rightarrow \infty} [H_n(1) - H_n(t)] = 0$$

uniformly on $[0, 1]$.

Corollary 5. *Under the assumptions as in Theorem 3 and provided there exists a constant $K > 0$ such that*

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$(2.16) \quad 0 \leq H_n(1) - H_n(t) \leq \frac{K(1-t)(b-a)^2}{12n}$$

for all $n \geq 1$ and $t \in [0, 1]$.

Now we establish an upper bound for the difference $H_n(t) - H_{n+1}(t)$, $n \geq 1$ which is non-negative for all $t \in [0, 1]$ (cf. Theorem 2):

Theorem 6. Let $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function on the interval I and let $a, b \in \overset{\circ}{I}$ with $a < b$. Then the inequality

$$\begin{aligned}
 (2.17) \quad 0 &\leq H_n(t) - H_{n+1}(t) \\
 &\leq \frac{t}{n+1} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \right. \\
 &\quad \left. \times \left(x_1 - \frac{a+b}{2} \right) dx_1 \cdots dx_n \right] \\
 &\leq \frac{t(b-a)}{2\sqrt{3}\sqrt{n}(n+1)} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} \right. \right. \right. \\
 &\quad \left. \left. \left. + (1-t) \frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}}
 \end{aligned}$$

holds for all $t \in [0, 1]$ and $n \geq 1$.

Proof. By the convexity of f , we have that

$$\begin{aligned}
 &f\left(t \frac{x_1 + \cdots + x_{n+1}}{n+1} + (1-t) \frac{a+b}{2}\right) - f\left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2}\right) \\
 &\geq \frac{t}{n(n+1)} f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) [nx_{n+1} - (x_1 + \cdots + x_n)]
 \end{aligned}$$

for all $x_1, \dots, x_{n+1} \in [a, b]$ and $t \in [0, 1]$.

Taking an integral mean on $[a, b]^{n+1}$, we derive

$$\begin{aligned}
 0 &\leq H_n(t) - H_{n+1}(t) \\
 &\leq \frac{t}{n+1} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) x_1 dx_1 \cdots dx_n \right. \\
 &\quad \left. - \frac{a+b}{2} \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) dx_1 \cdots dx_n \right],
 \end{aligned}$$

and the second inequality in (2.17) is proved.

Now, let us observe that

$$\begin{aligned}
& \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \\
& \qquad \qquad \qquad \times \left(x_1 - \frac{a+b}{2} \right) dx_1 \cdots dx_n \\
& = \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \\
& \qquad \qquad \qquad \times \left(\frac{x_1 + \cdots + x_n}{n} - \frac{a+b}{2} \right) dx_1 \cdots dx_n \\
& \leq \left(\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right)^{\frac{1}{2}} \\
& \qquad \times \left(\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left(\frac{x_1 + \cdots + x_n}{n} - \frac{a+b}{2} \right)^2 dx_1 \cdots dx_n \right)^{\frac{1}{2}},
\end{aligned}$$

and thus the last inequality is also proved. □

Corollary 6. *Under the assumptions as in Theorem 3, given that $M := \sup_{x \in [a,b]} |f'(x)| < \infty$, we have*

$$(2.18) \qquad 0 \leq H_n(t) - H_{n+1}(t) \leq \frac{Mt(b-a)}{2\sqrt{3}\sqrt{n}(n+1)}$$

for all $t \in [0, 1]$ and $n \geq 1$.

In particular,

$$\lim_{n \rightarrow \infty} [H_n(t) - H_{n+1}(t)] = 0$$

uniformly on $[0, 1]$.

The following theorem also holds.

Theorem 7. *Under the assumptions as in Theorem 3, we also have the bound*

$$\begin{aligned}
(2.19) \quad 0 \leq H_n(t) - H_{n+1}(t) \\
\leq \frac{t(b-a)}{2\sqrt{3}\sqrt{n}(n+1)} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+ \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right) \right. \right. \\
\qquad \qquad \qquad \left. \left. - f'_+ \left(\frac{a+b}{2} \right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}}
\end{aligned}$$

for all $t \in [0, 1]$ and $n \geq 1$.

Proof. The proof is similar to the one of Theorem 6 and we omit the details. □

Corollary 7. Under the assumptions as in Theorem 3, given that there exists a $K > 0$ such that

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$(2.20) \quad 0 \leq H_n(t) - H_{n+1}(t) \leq \frac{t^2(b-a)^2 K^2}{12n(n+1)}$$

for all $t \in [0, 1]$ and $n \geq 1$.

Finally, note that, by a similar argument to that in the proof of Theorem 6, we can give the following result which completes, in a sense, the estimate in Theorem 3.

Theorem 8. Under the above assumptions the inequality

$$(2.21) \quad 0 \leq H_n(t) - f\left(\frac{a+b}{2}\right) \leq \frac{t(b-a)}{2\sqrt{3}\sqrt{n}} \left[\frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left[f'_+\left(t\frac{x_1 + \cdots + x_n}{n} + (1-t)\frac{a+b}{2}\right) - f'_+\left(\frac{a+b}{2}\right) \right]^2 dx_1 \cdots dx_n \right]^{\frac{1}{2}}$$

holds for all $t \in [0, 1]$ and $n \geq 1$.

Corollary 8. Under the above assumptions, given that there exists a $K > 0$ such that

$$|f'_+(x) - f'_+(y)| \leq K|x - y| \quad \text{for all } x, y \in [a, b],$$

we have the inequality

$$0 \leq H_n(t) - f\left(\frac{a+b}{2}\right) \leq \frac{t^2(b-a)^2 K^2}{12n}$$

for all $t \in [0, 1]$ and $n \geq 1$.

3. APPLICATIONS TO SPECIAL MEANS

Let $0 \leq a < b$ and $p \in (-\infty, 0) \cup [1, \infty) \setminus \{-1\}$. Let us define sequence of mappings

$$h_{p,n}(t) := \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t) \frac{a+b}{2} \right)^p dx_1 \cdots dx_n,$$

where $n \geq 1$, $n \in \mathbb{N}$, and $t \in [0, 1]$.

By virtue of the above results, we can establish the following properties:

- (i) $h_{p,n}(t)$ are convex and monotonic nondecreasing on $[0, 1]$;
- (ii) $h_{p,n}(t) \geq h_{p,n+1}(t)$ for all $n \geq 1$ and $t \in [0, 1]$;
- (iii) the inequalities

$$\begin{aligned} [A(a, b)]^p &\leq h_{p,n}(t) \\ &\leq \frac{1}{(b-a)^{n+1}} \int_a^b \cdots \int_a^b \left(t \frac{x_1 + \cdots + x_n}{n} + (1-t)x_{n+1} \right)^p dx_1 \cdots dx_{n+1} \end{aligned}$$

and

$$\begin{aligned} h_{p,n}(t) &\leq t \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left(\frac{x_1 + \cdots + x_n}{n} \right)^p dx_1 \cdots dx_n \\ &\quad + (1-t)[A(a, b)]^p \\ &\leq \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left(\frac{x_1 + \cdots + x_n}{n} \right)^p dx_1 \cdots dx_n \end{aligned}$$

hold for all $n \geq 1$ and $t \in [0, 1]$, where $A(a, b) := \frac{a+b}{2}$.

- (iv) If $p \geq 1$, then the inequalities

$$0 \leq h_{p,n}(t) - [A(a, b)]^p \leq \frac{t(b-a)pb^{p-1}}{2\sqrt{3}\sqrt{n}}$$

and

$$\begin{aligned} 0 &\leq \frac{1}{(b-a)^n} \int_a^b \cdots \int_a^b \left(\frac{x_1 + \cdots + x_n}{n} \right)^p dx_1 \cdots dx_n - h_{p,n}(t) \\ &\leq \frac{(1-t)(b-a)pb^{p-1}}{2\sqrt{3}\sqrt{n}} \end{aligned}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

(v) If $p \geq 1$, the inequalities

$$0 \leq th_{p,n}(1) + (1-t)h_{p,n}(0) - h_{p,n}(t) \leq \frac{t(1-t)pb^{p-1}}{2\sqrt{3}\sqrt{n}}$$

and

$$0 \leq h_{p,n}(t) - h_{p,n+1}(t) \leq \frac{t(b-a)pb^{p-1}}{2\sqrt{3}\sqrt{n}(n+1)}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

(vi) If $p \geq 2$, then the inequalities

$$0 \leq h_{p,n}(t) - [A(a, b)]^p \leq \frac{t^2(b-a)^2p(p-1)b^{p-2}}{12n}$$

and

$$0 \leq h_{p,n}(t) - h_{p,n+1}(t) \leq \frac{t^2(b-a)^2p(p-1)b^{p-2}}{12n(n+1)}$$

and

$$\begin{aligned} 0 &\leq th_{p,n}(1) + (1-t)h_{p,n}(0) - h_{p,n}(t) \\ &\leq \frac{t(1-t)(b-a)^2p(p-1)b^{p-2}}{12n} \end{aligned}$$

and

$$0 \leq h_{p,n}(1) - h_{p,n}(t) \leq \frac{(1-t)(b-a)^2p(p-1)b^{p-2}}{12n}$$

hold for all $n \geq 1$ and $t \in [0, 1]$.

Acknowledgement. The author would like to thank the anonymous referee for many valuable comments and remarks that have been incorporated in the final version of the paper.

References

- [1] G. Allasia, C. Giordano, J. Pečarić: Hadamard-type inequalities for $(2r)$ -convex functions with applications. *Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur.* 133 (1999), 187–200.
- [2] H. Alzer: A note on Hadamard's inequalities. *C. R. Math. Rep. Acad. Sci. Canada* 11 (1989), 255–258.
- [3] H. Alzer: On an integral inequality. *Anal. Numér. Théor. Approx.* 18 (1989), 101–103.

- [4] *A. G. Azpeitia*: Convex functions and the Hadamard inequality. *Rev. Colombiana Mat.* 28 (1994), 7–12.
- [5] *D. Barbu, S. S. Dragomir and C. Buşe*: A probabilistic argument for the convergence of some sequences associated to Hadamard's inequality. *Studia Univ. Babeş-Bolyai Math.* 38 (1993), 29–33.
- [6] *C. Buşe, S. S. Dragomir and D. Barbu*: The convergence of some sequences connected to Hadamard's inequality. *Demonstratio Math.* 29 (1996), 53–59.
- [7] *S. S. Dragomir*: A mapping in connection to Hadamard's inequalities. *Anz. Österreich. Akad. Wiss. Math.-Natur. Kl.* 128 (1991), 17–20.
- [8] *S. S. Dragomir*: A refinement of Hadamard's inequality for isotonic linear functionals. *Tamkang J. Math.* 24 (1993), 101–106.
- [9] *S. S. Dragomir*: On Hadamard's inequalities for convex functions. *Mat. Balkanica (N. S.)* 6 (1992), 215–222.
- [10] *S. S. Dragomir*: On Hadamard's inequality for the convex mappings defined on a ball in the space and applications. *Math. Inequal. Appl.* 3 (2000), 177–187.
- [11] *S. S. Dragomir*: On Hadamard's inequality on a disk. *JIPAM. J. Inequal. Pure Appl. Math.* 1 (2000), <http://jipam.vu.edu.au/>. Electronic.
- [12] *S. S. Dragomir*: Some integral inequalities for differentiable convex functions. *Makedon. Akad. Nauk Umet. Oddel. Mat.-Tehn. Nauk. Prilozi* 13 (1992), 13–17.
- [13] *S. S. Dragomir*: Some remarks on Hadamard's inequalities for convex functions. *Extracta Math.* 9 (1994), 88–94.
- [14] *S. S. Dragomir*: Two mappings in connection to Hadamard's inequalities. *J. Math. Anal. Appl.* 167 (1992), 49–56.
- [15] *S. S. Dragomir, R. P. Agarwal*: Two new mappings associated with Hadamard's inequalities for convex functions. *Appl. Math. Lett.* 11 (1998), 33–38.
- [16] *S. S. Dragomir, C. Buşe*: Refinements of Hadamard's inequality for multiple integrals. *Utilitas Math.* 47 (1995), 193–198.
- [17] *S. S. Dragomir, Y. J. Cho and S. S. Kim*: Inequalities of Hadamard's type for Lipschitzian mappings and their applications. *J. Math. Anal. Appl.* 245 (2000), 489–501.
- [18] *S. S. Dragomir, S. Fitzpatrick*: Hadamard inequality for s -convex functions in the first sense and applications. *Demonstratio Math.* 31 (1998), 633–642.
- [19] *S. S. Dragomir, S. Fitzpatrick*: The Hadamard's inequality for s -convex functions in the second sense. *Demonstratio Math.* 32 (1999), 687–696.
- [20] *S. S. Dragomir, N. M. Ionescu*: On some inequalities for convex-dominated functions. *Anal. Numér. Théor. Approx.* 19 (1990), 21–27.
- [21] *S. S. Dragomir, D. S. Milošević and J. Sándor*: On some refinements of Hadamard's inequalities and applications. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.* 4 (1993), 3–10.
- [22] *S. S. Dragomir, B. Mond*: On Hadamard's inequality for a class of functions of Godunova and Levin. *Indian J. Math.* 39 (1997), 1–9.
- [23] *S. S. Dragomir, C. E. M. Pearce*: Quasi-convex functions and Hadamard's inequality. *Bull. Austral. Math. Soc.* 57 (1998), 377–385.
- [24] *S. S. Dragomir, C. E. M. Pearce, and J. E. Pečarić*: On Jensen's related inequalities for isotonic sublinear functionals. *Acta Sci. Math.* 61 (1995), 373–382.
- [25] *S. S. Dragomir, J. E. Pečarić, and L. E. Persson*: Some inequalities of Hadamard type. *Soochow J. Math.* 21 (1995), 335–341.
- [26] *S. S. Dragomir, J. E. Pečarić, and J. Sándor*: A note on the Jensen-Hadamard inequality. *Anal. Numér. Théor. Approx.* 19 (1990), 29–34.
- [27] *S. S. Dragomir, G. H. Toader*: Some inequalities for m -convex functions. *Studia Univ. Babeş-Bolyai Math.* 38 (1993), 21–28.

- [28] *A. M. Fink*: A best possible Hadamard inequality. *Math. Inequal. Appl.* *1* (1998), 223–230.
- [29] *A. M. Fink*: Toward a theory of best possible inequalities. *Nieuw Arch. Wisk.* *12* (1994), 19–29.
- [30] *A. M. Fink*: Two inequalities. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.* *6* (1995), 48–49.
- [31] *B. Gavrea*: On Hadamard’s inequality for the convex mappings defined on a convex domain in the space. *JIPAM. J. Inequal. Pure Appl. Math.* *1* (2000), Article 9, <http://jipam.vu.edu.au/>. Electronic.
- [32] *P. M. Gill, C. E. M. Pearce and J. Pečarić*: Hadamard’s inequality for r -convex functions. *J. Math. Anal. Appl.* *215* (1997), 461–470.
- [33] *G. H. Hardy, J. E. Littlewood, and G. Pólya*: *Inequalities*. 2nd ed. Cambridge University Press, 1952.
- [34] *K.-C. Lee, K.-L. Tseng*: On weighted generalization of Hadamard’s inequality for g functions. *Tamsui Oxf. J. Math. Sci.* *16* (2000), 91–104.
- [35] *A. Lupuş*: The Jensen-Hadamard inequality for convex functions of higher order. *Octogon Math. Mag.* *5* (1997), 8–9.
- [36] *A. Lupuş*: A generalization of Hadamard’s inequality for convex functions. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* *544–576* (1976), 115–121.
- [37] *D. M. Maksimović*: A short proof of generalized Hadamard’s inequalities. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* *634–677* (1979), 126–128.
- [38] *D. S. Mitrinović, I. Lacković*: Hermite and convexity. *Aequationes Math.* *28* (1985), 229–232.
- [39] *D. S. Mitrinović, J. E. Pečarić, and A. M. Fink*: *Classical and New Inequalities in Analysis*. Kluwer Academic Publishers, Dordrecht, 1993.
- [40] *E. Neuman*: Inequalities involving generalized symmetric means. *J. Math. Anal. Appl.* *120* (1986), 315–320.
- [41] *E. Neuman, J. E. Pečarić*: Inequalities involving multivariate convex functions. *J. Math. Anal. Appl.* *137* (1989), 514–549.
- [42] *E. Neuman*: Inequalities involving multivariate convex functions. II. *Proc. Amer. Math. Soc.* *109* (1990), 965–974.
- [43] *C. P. Niculescu*: A note on the dual Hermite-Hadamard inequality. *The Math. Gazette* (July 2000).
- [44] *C. P. Niculescu*: Convexity according to the geometric mean. *Math. Inequal. Appl.* *3* (2000), 155–167.
- [45] *C. E. M. Pearce, J. Pečarić, and V. Šimić*: Stolarsky means and Hadamard’s inequality. *J. Math. Anal. Appl.* *220* (1998), 99–109.
- [46] *C. E. M. Pearce, A. M. Rubinov*: P -functions, quasi-convex functions and Hadamard-type inequalities. *J. Math. Anal. Appl.* *240* (1999), 92–104.
- [47] *J. E. Pečarić*: Remarks on two interpolations of Hadamard’s inequalities. *Makedon. Akad. Nauk. Umet. Oddel. Mat.-Tehn. Nauk. Prilozi* *13* (1992), 9–12.
- [48] *J. Pečarić, S. S. Dragomir*: A generalization of Hadamard’s inequality for isotonic linear functionals. *Rad. Mat.* *7* (1991), 103–107.
- [49] *J. Pečarić, F. Proschan, and Y. L. Tong*: *Convex Functions, Partial Orderings and Statistical Applications*. Academic Press, Boston, 1992.
- [50] *J. Sándor*: An application of the Jensen-Hadamard inequality. *Nieuw Arch. Wisk.* *8* (1990), 63–66.
- [51] *J. Sándor*: On the Jensen-Hadamard inequality. *Studia Univ. Babeş-Bolyai, Math.* *36* (1991), 9–15.

- [52] *P. M. Vasić, I. B. Lacković, and D. M. Maksimović*: Note on convex functions. IV. On Hadamard's inequality for weighted arithmetic means. Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. 678–715 (1980), 199–205.
- [53] *G. S. Yang, M. C. Hong*: A note on Hadamard's inequality. Tamkang J. Math. 28 (1997), 33–37.
- [54] *G. S. Yang, K. L. Tseng*: On certain integral inequalities related to Hermite-Hadamard inequalities. J. Math. Anal. Appl. 239 (1999), 180–187.

Author's address: S. S. Dragomir, School of Computer Science & Mathematics, Victoria University, P.O. Box 14428, Melbourne City, MC 8001, Australia, e-mail: sever@matilda.vu.edu.au.